PROCEEDINGS OF SCIENCE



The model dependence of $m_{ ho}/f_{\pi}$

Daniel Nogradi* and Lorinc Szikszai

Eotvos University, Department of Theoretical Physics, Budapest 1117, Hungary E-mail: nogradi@bodri.elte.hu, szikszail@caesar.elte.hu

Should a strongly coupled composite Higgs boson scenario be realized in Nature the most easily accessible experimental signal would be new particles made up of the same ingredients as the Higgs but with different quantum numbers. The lightest of these hypothetical new particles would probably be the vector mesons. In this contribution we report results on m_{ρ}/f_{π} in the chiral-continuum limit with SU(3) gauge group and $N_f = 2,3,4,5,6$ flavors of fundamental fermions. In addition we compare m_{ρ}/f_{π} results from various models with different gauge groups and fermion content. The main conclusion seems to be that the experimental measurement of this vector meson mass will be able to distinguish between gauge groups but less so between the fermion content.

37th International Symposium on Lattice Field Theory - Lattice2019 16-22 June 2019 Wuhan, China

^{*}Speaker.

[©] Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Introduction

It may very well be the case that the experimentally discovered Higgs boson is the elementary spin-0 particle of the Standard Model and the theoretical description is valid up to very high energies beyond the scope of the LHC or any new accelerator in the near future. If on the other hand the Higgs boson turns out to be a composite particle the bare minimum experimental signal should be the discovery of new particles made up of the same ingredients as the composite Higgs itself but with different quantum numbers. The lightest candidates in a large class of strongly interacting electro-weak models would be the vector mesons in the non-abelian gauge theory responsible for the composite Higgs. The physical scale in these technicolor inspired class of models is given by the decay constant of the Goldstone bosons $f_{\pi} = 246 \text{ GeV}$. Lattice calculations are ideal to obtain the massless ratio m_{ρ}/f_{π} in the chiral-continuum limit from first principles. Conceptually it is a simple task: pick a gauge group G, representation R and flavor number N_f (or potentially several representations and several flavor numbers if not all fermions transform in the same representation), make sure that finite volume effects are small, make sure ρ -decay is handled properly, obtain am_{ρ} and af_{π} in lattice units for various lattice spacings and fermion masses and finally perform a chiral-continuum extrapolation to obtain the ratio. Thus traditional meson spectroscopy will lead to m_{ρ}/f_{π} for every choice (G, R, N_f) . If experiments do detect a new particle in the TeV range the next task would be to find which (G, R, N_f) combination(s) may give rise to that particular mass and quantum number hence it is important to have scanned as many models as possible¹.

Recent lattice studies of the meson spectrum with SU(3) indicate that the fermion content dependence might be very mild for the ratio m_{ρ}/f_{π} . Even though many of the results are at finite lattice spacing, occasionally finite fermion mass and finite volume effects are not always fully controlled the indications are $m_{\rho}/f_{\pi} \sim 8.0$ from a broad range of studies [1, 2, 3, 4, 5, 6, 7, 8, 9, 10]. As long as the gauge group is fixed, it seems then, that the experimental measurement of m_{ρ} will not be able to distinguish the various models as long as the gauge group is SU(3).

In this contribution we report fully controlled simulation results in the infinite volume, chiral and continuum limits with SU(3) and $N_f = 2, 3, 4, 5, 6$ reinforcing the above indications [11]. We obtain $m_{\rho}/f_{\pi} = 7.95(15)$ and no statistically significant N_f -dependence.

The ratio m_{ρ}/f_{π} however is expected to change rapidly with the gauge group based on large-*N* arguments. More precisely if the fermionic degrees of freedom scale as O(N) which is the case with fundamental fermions, we have $m_{\rho} \sim O(1)$ and $f_{\pi} \sim O(\sqrt{N})$ hence the natural combination to consider is $\sqrt{N} m_{\rho}/f_{\pi}$. In the large-*N* limit this combination is finite. Motivated by this we will be comparing various models via the combination $\sqrt{dim(R)} m_{\rho}/f_{\pi}$ where dim(R) is the dimension of the representation; see [12, 13, 14] and references therein.

In the next section the details of the SU(3) calculations with $N_f = 2,3,4,5,6$ fundamental fermions will be given. In section 3 we will compare results from the literature for various (G, R, N_f) by collecting $\sqrt{\dim(R)} m_{\rho}/f_{\pi}$ if possible in the chiral-continuum limit, or at finite lattice spacing and/or finite fermion mass if chiral-continuum limit results are not available. In section 4 we end with notes on possible future studies.

¹Throughout this contribution the normalization of the decay constant is such that $f_{\pi} = 93 \text{ MeV}$ in QCD.

2. Results with SU(3) and fundamental fermions

The lattice discretization is via the tree level improved Symanzik gauge action and stout improved staggered fermions with 4 steps of stout smearing and $\rho = 0.12$ smearing parameter [15]. Ensembles are generated at four lattice spacings with four fermion masses at each, for every N_f . The parameters were chosen such that the ρ is stable and the finite volume corrections from Luscher's formalism [16, 17] predict small finite volume effects. Furthermore, exponential finite volume effects are ensured to be small as well by having $m_{\pi}L > 3.10 + 0.35N_f$ which was shown to lead to less than 1% corrections in low energy observables for $2 \le N_f \le 6$.



Figure 1: Chiral-continuum extrapolation of the topological susceptibility. The ratio $2N_f \chi/(m_\pi^2 f_\pi^2)$ is shown which, at leading order of chiral perturbation theory, is expected to be constant 1. The shaded black region is the result of the chiral-continuum extrapolation. The various colors correspond to various lattice spacings labeled by the bare coupling β in the legend.

Topology change is frequent enough in all runs so that the topological susceptibility can be measured accurately enough. In fact the topological susceptibility was used to test for taste breaking effects via the tree level relation [18],

$$\frac{\langle Q^2 \rangle}{V} = \chi = \frac{1}{2N_f} f_\pi^2 m_\pi^2 + \dots$$
 (2.1)

More precisely, $f_{\pi}^2 m_{\pi}^2/(2\chi)$ can be considered an effective flavor number and if in the chiralcontinuum limit the actual flavor number is obtained within errors, one can be sure that the taste broken Goldstones of the staggered formulation are scaled to zero as $O(a^2)$ as expected [19]. This test is far easier to perform than the full spectroscopy of all taste broken Goldstones directly although relatively long runs are required as the autocorrelation time of the topological charge is typically longer than other quantities. At the same time the topological susceptibility needs to be monitored anyway in order to avoid frozen topology simulations. More precisely, the chiralcontinuum extrapolation is via,

$$\chi w_0^4 = C_0 m_\pi^2 f_\pi^2 w_0^4 + C_1 \frac{a^2}{w_0^2} + C_2 \frac{a^2}{w_0^2} (m_\pi^2 f_\pi^2 w_0^4) , \qquad (2.2)$$

where w_0 is the gradient flow scale [20]. In figure 1 the quantity $2N_f \chi/(m_{\pi}^2 f_{\pi}^2)$ is shown for each N_f for all lattice spacings and masses, together with the chiral-continuum limit, which ought to be consistent with 1. The deviation from 1 is at most 1.5 σ .

The 16 simulation points at each N_f allow for controlled chiral-continuum fits of the quantities f_{π} and m_{ρ} as well. The 4 parameter fits we use are,

$$Xw_0 = C_0 + C_1 m_\pi^2 w_0^2 + C_2 \frac{a^2}{w_0^2} + C_3 \frac{a^2}{w_0^2} m_\pi^2 w_0^2$$
(2.3)

where X is either m_{ρ} or f_{π} . Each fit has thus 12 degrees of freedom. The statistical uncertainty of f_{π} is negligible compared to m_{ρ} and the latter will dominate the final result for the ratio m_{ρ}/f_{π} . With our action both cut-off effects and finite fermion mass effects are rather small for the ratio [11].

The final results are shown in figure 2 as the left most set of points. Interestingly, a constant fit leads to $m_{\rho}/f_{\pi} = 7.95(15)$ with $\chi^2/dof = 0.26$ i.e. the observed N_f -dependence is statistically insignificant.

3. Dependence on the gauge group, representation and flavor number

Quite a large number of models were studied recently, partially or completely motivated by a strongly coupled composite Higgs scenario. Another set of motivations useful for our purposes is the study of the systematics of the large-N expansion because in this context the gauge group dependence can be displayed. The large-N limit is instructive because the observed pattern at SU(3) namely no statistically significant dependence on N_f is in line with large-N expectations. As long as the fermionic degrees of freedom scale with O(N), which is the case in the fundamental, one expects exactly no dependence on N_f in the large-N limit. With the fundamental representation f_{π} scales as \sqrt{N} . If the representation is other than the fundamental it is natural to consider



Figure 2: Top: comparison of $\sqrt{\dim(R)} m_{\rho}/f_{\pi}$ for various models. Bottom: similar comparison of m_{ρ}/f_{π} directly, the labeling of the models is the same as at the top. The SU(3) fundamental $N_f = 2, 3, 4, 5, 6$ are in the chiral-continuum limit [11], the $N_f = 8$ result is at finite lattice spacing and finite fermion mass [10], the SU(3) sextet $N_f = 2$ is at finite lattice spacing and finite fermion mass [7], the SU(N) fundamental quenched results are at finite lattice spacing in the chiral limit [12], the SU(N) fundamental $N_f = 4$ are at finite lattice spacing and finite fermion mass [21, 22], the SU(4) fundamental $N_f = 2$, sextet $N_f = 2$ (both are in the sea simultaneously) results are in the chiral-continuum limit [23], the SU(2) fundamental $N_f = 2$ is in the chiral-continuum limit [24, 25], the SU(2) fundamental $N_f = 4$ is at finite lattice spacing and finite fermion mass [26] and finally with Sp(4) both the quenched [27] and the fundamental $N_f = 2$ are in the chiral-continuum limit [28, 29]. Results at finite lattice spacing and/or finite fermion mass should be interpreted with caution as they naturally contain further systematic errors.

 $\sqrt{\dim(R)} m_{\rho}/f_{\pi}$ where $\dim(R)$ is the dimension of the representation. If the fermionic degrees of freedom scale as $O(N^2)$ then of course the usual large-*N* arguments do not apply.

Starting from SU(2), chiral-continuum results are available with $N_f = 2$ fermions in the fundamental representation [24, 25], $N_f = 4$ at finite lattice spacing and fermion mass [26], and of course the pure gauge case. The aforementioned large-N studies led to results with SU(N), still in the pure gauge case, with N = 2, 3, 4, 5, 6, 7, 17 and the $N = \infty$ in the chiral limit, at finite lattice spacing [12, 13]. Note that with SU(2) all irreducible representations are real. Increasing the gauge group to SU(3) we have of course QCD results (or Nature) and the $N_f = 2, 3, 4, 5, 6$ results of our work in the chiral-continuum limit [11] as well as $N_f = 8$ at finite lattice spacing and finite fermion mass [10]. Still with SU(3) results are available with $N_f = 2$ sextet fermions at finite lattice spacing and finite fermion mass. Further, SU(4) was studied with two species of fermions simultaneously in the sea, $N_f = 2$ fundamental together with $N_f = 2$ sextet (which is real) [23]; see also [30] for results with $N_f = 2$ sextet fermions only. In this case the two species of fermions lead to two different decay constants and two different vector meson masses. Results are available in the chiral-continuum limit [23]. The large-N limit was investigated with $N_f = 4$ fundamental fermions and N = 3, 4, 5, 6, the results are currently at finite lattice spacing and finite fermion mass [21, 22]. Going beyond the unitary groups, the symplectic group Sp(4) (which is sometimes denoted by Sp(2), in any case it is the double covering group of SO(5)) was investigated in the quenched case with both fundamental and 2-index-antisymmetric representation fermions [27] and $N_f = 2$ fundamental fermions [28, 29]; all the Sp(4) results are in the chiral-continuum limit.

A summary of all of these results are shown in figure 2 via the scaled ratio $\sqrt{\dim(R)} m_{\rho}/f_{\pi}$ in the top panel. In order to better display the dependence on the gauge group the unscaled m_{ρ}/f_{π} is also shown in the bottom panel of figure 2. Clearly, the dependence on the fermion content is much less pronounced than the dependence on the gauge group. However when the leading gauge group dependence is factored out, i.e. $\sqrt{\dim(R)} m_{\rho}/f_{\pi}$ is considered, even the gauge group dependence is rather small. Only SU(2) shows considerable deviation from large-N scaling and large N_f -dependence (although the $N_f = 4$ results are at finite lattice spacing and finite fermion mass). Interpretation of results at finite fermion mass and/or finite lattice spacing should be done with care of course.

If one envisions an experimental signal for m_{ρ} the gauge group can perhaps be narrowed down using figure 2 however the information on the fermion content must come from other experimentally accessible quantities unless the gauge group is SU(2).

4. Conclusion and outlook

A strongly interacting composite Higgs boson is clearly an exciting possibility. The best case scenario would be the detection of new particles in the TeV range by the LHC. In order to make full use of this result it would be desirable to obtain fully controlled (volume, lattice spacing, fermion mass) lattice results on the masses of the new particles for as many promising models as possible. Currently the set of fully controlled results is rather limited and even where they are available often the errors are too large for an experimental result to differentiate between models. It does not seem out of reach to obtain fully controlled results with $\sim 3 - 5\%$ errors with SU(2) and $N_f = 2, 3, 4$ and possibly $N_f = 5$ fundamental fermions if the latter is chirally broken. With SU(3) and fundamental fermions $N_f = 7, 8, 9, 10$ seems possible and it is conceivable that the Sp(4) with $N_f = 2$ results [28, 29] can be extended to $N_f = 3, 4$. The closer a model is to the conformal window the more difficult the calculation becomes because of large systematic uncertainties. This is the reason why SU(2) with $N_f = 5, 6$ is difficult and so is SU(3) with $N_f = 2$ sextet [3, 7]. Fixing N_f and increasing N on the other hand is moving the model away from the conformal window so should be relatively straightforward. In this direction m_ρ/f_π is decreasing and can be arbitrarily small for large enough N. Presumably the largest value will occur for SU(2) although this is currently an open question.

Acknowledgements

We would like to thank William Jay, Julius Kuti, Jong-Wan Lee, Fernando Romero Lopez,

Biagio Lucini, Ben Svetitsky and Chik Him Wong for sharing data on m_{ρ}/f_{π} in the various models presented in figure 2. This work was supported by the NKFIH under the grant KKP-126769. We are grateful to Kalman Szabo for code sharing.

References

- [1] Z. Fodor et al., Phys. Lett. B 681, 353 (2009) [arXiv:0907.4562 [hep-lat]].
- [2] X. Y. Jin and R. D. Mawhinney, PoS LAT 2009, 049 (2009) [arXiv:0910.3216 [hep-lat]].
- [3] Z. Fodor et al., Phys. Lett. B 718, 657 (2012) [arXiv:1209.0391 [hep-lat]].
- [4] Y. Aoki et al., Phys. Rev. D 87, no. 9, 094511 (2013) [arXiv:1302.6859 [hep-lat]].
- [5] X. Y. Jin and R. D. Mawhinney, arXiv:1304.0312 [hep-lat].
- [6] G. T. Fleming et al. [LSD Collaboration], arXiv:1312.5298 [hep-lat].
- [7] Z. Fodor et al., PoS LATTICE 2015, 219 (2016) [arXiv:1605.08750 [hep-lat]].
- [8] Z. Fodor et al., Phys. Rev. D 94, no. 1, 014503 (2016) [arXiv:1601.03302 [hep-lat]].
- [9] T. Appelquist et al., Phys. Rev. D 93, no. 11, 114514 (2016) [arXiv:1601.04027 [hep-lat]].
- [10] T. Appelquist et al., Phys. Rev. D 99, no. 1, 014509 (2019) [arXiv:1807.08411 [hep-lat]].
- [11] D. Nogradi and L. Szikszai, JHEP 1905, 197 (2019) [arXiv:1905.01909 [hep-lat]].
- [12] G. S. Bali et al., JHEP 1306, 071 (2013) [arXiv:1304.4437 [hep-lat]].
- [13] L. Castagnini, PhD Thesis, University of Regensburg, https://epub.uni-regensburg.de/32024/1/phd.pdf
- [14] T. DeGrand and Y. Liu, Phys. Rev. D 94, no. 3, 034506 (2016) Erratum: [Phys. Rev. D 95, no. 1, 019902 (2017)] [arXiv:1606.01277 [hep-lat]].
- [15] C. Morningstar and M. J. Peardon, Phys. Rev. D 69, 054501 (2004) [hep-lat/0311018].
- [16] M. Luscher, Nucl. Phys. B 354, 531 (1991).
- [17] M. Luscher, Nucl. Phys. B 364, 237 (1991).
- [18] H. Leutwyler and A. V. Smilga, Phys. Rev. D 46, 5607 (1992).
- [19] B. Billeter, C. E. Detar and J. Osborn, Phys. Rev. D 70, 077502 (2004) [hep-lat/0406032].
- [20] S. Borsanyi et al., JHEP 1209, 010 (2012) [arXiv:1203.4469 [hep-lat]].
- [21] F. Romero-Lopez, A. Donini, P. Hernandez and C. Pena, arXiv:1910.10418 [hep-lat].
- [22] P. Hernandez et al., Eur. Phys. J. C 79, no. 10, 865 (2019) [arXiv:1907.11511 [hep-lat]].
- [23] V. Ayyar et al., Phys. Rev. D 97, no. 7, 074505 (2018) [arXiv:1710.00806 [hep-lat]].
- [24] R. Arthur et al., Phys. Rev. D 94, no. 9, 094507 (2016) [arXiv:1602.06559 [hep-lat]].
- [25] V. Drach, T. Janowski and C. Pica, EPJ Web Conf. 175, 08020 (2018) [arXiv:1710.07218 [hep-lat]].
- [26] A. Amato, V. Leino, K. Rummukainen, K. Tuominen and S. Tahtinen, arXiv:1806.07154 [hep-lat].
- [27] E. J. Bennett et al., arXiv:1912.06505 [hep-lat].
- [28] E. Bennett et al., JHEP 1803, 185 (2018) [arXiv:1712.04220 [hep-lat]].
- [29] E. Bennett et al., arXiv:1909.12662 [hep-lat].
- [30] T. DeGrand et al., Phys. Rev. D 91, 114502 (2015) [arXiv:1501.05665 [hep-lat]].