Feller property of the multiplicative coalescent with linear deletion

Balázs Ráth¹

August 7, 2018

Abstract

We modify the definition of Aldous' multiplicative coalescent process [3] and introduce the multiplicative coalescent with linear deletion (MCLD). A state of this process is a square-summable decreasing sequence of cluster sizes. Pairs of clusters merge with a rate equal to the product of their sizes and clusters are deleted with a rate linearly proportional to their size. We prove that the MCLD is a Feller process. This result is a key ingredient in the description of scaling limits of the evolution of component sizes of the mean field frozen percolation model [22] and the so-called rigid representation of such scaling limits [19].

Keywords: multiplicative coalescent, Feller process

AMS MSC 2010: 60J99, 60B12, 05C80

1 Introduction

Let us define

$$\ell_{\infty}^{\downarrow} = \{ \underline{m} = (m_1, m_2, \dots) : m_1 \geqslant m_2 \geqslant \dots \geqslant 0 \},$$

$$\ell_{2}^{\downarrow} = \{ \underline{m} \in \ell_{\infty}^{\downarrow} : \sum_{i=1}^{\infty} m_i^2 < \infty \},$$

$$\ell_{0}^{\downarrow} = \{ \underline{m} \in \ell_{\infty}^{\downarrow} : \exists i_0 \in \mathbb{N} : m_i = 0 \text{ for any } i \geqslant i_0 \}.$$

¹MTA-BME Stochastics Research Group, Hungary. E-mail: rathb@math.bme.hu

For $\underline{m}, \underline{m'} \in \ell_2^{\downarrow}$ one defines the distance

$$d(\underline{m}, \underline{m}') = \|\underline{m} - \underline{m}'\|_2 = \left(\sum_{i \ge 1} (m_i - m_i')^2\right)^{1/2}.$$
 (1.1)

The metric space $(\ell_2^{\downarrow}, d(\cdot, \cdot))$ is complete and separable.

The multiplicative coalescent process (or briefly MC process), defined in [3, Section 1.5], is a continuous-time Markov process $\mathbf{m}_t, t \geq 0$ with state space ℓ_2^{\downarrow} . The state \mathbf{m}_t represents the ordered sequence of sizes of components, where two components of size m_i and m_j merge with rate $m_i \cdot m_j$. By [3, Proposition 5], the multiplicative coalescent process has the Feller property with respect to the metric $d(\cdot,\cdot)$ on ℓ_2^{\downarrow} . On the other hand, if $\mathbf{m}_0 \in \ell_{\infty}^{\downarrow} \setminus \ell_2^{\downarrow}$, then all of the components instantaneously coagulate and form one component with infinite mass, see [18, Section 2.1]. In Section 2, we collect the basic results about MC relevant for our study.

Let $\lambda \in \mathbb{R}_+$. For any $\underline{m} \in \ell_2^{\downarrow}$ we want to define a continuous time Markov process \mathbf{m}_t with state space ℓ_2^{\downarrow} where $\mathbf{m}_0 = \underline{m}$ and \mathbf{m}_t represents the ordered sequence of sizes of components of a coagulation-deletion process at time t. We want the dynamics of the process $\mathbf{m}_t, t \geq 0$ to satisfy

(i) two components of size
$$m_i$$
 and m_j merge with rate $m_i \cdot m_j$,
(ii) a component of size m_i is deleted with rate $\lambda \cdot m_i$.

We are going to call such a process a multiplicative coalescent with linear deletion with deletion rate λ , and briefly denote it by $MCLD(\lambda)$.

If $\underline{m} \in \ell_0^{\downarrow}$ then the MCLD(λ) process obviously exists and $\mathbf{m}_t \in \ell_0^{\downarrow}$ for any $t \geq 0$. In fact, if $\underline{m} \in \ell_{\infty}^{\downarrow}$ with $\sum_{i=1}^{\infty} m_i < \infty$ then the definition of MCLD(λ) is still quite simple because the time between consecutive coalescences/deletions is always positive. On the other hand, for initial conditions with infinite total mass, the set of times when a coalescence or deletion occurs will be dense in \mathbb{R}_+ , and it is not a priori clear that a well-defined stochastic process satisfying (1.2) exists (see Remark 1.4 below for related non-existence results).

In Section 3 we will give a graphical construction of the process \mathbf{m}_t with initial state $\underline{m} \in \ell_2^{\downarrow}$ and deletion rate λ . This construction of $\mathrm{MCLD}(\lambda)$ is similar to, but not as simple as the graphical construction of the MC given in [3, Section 1.5] because $\mathrm{MCLD}(\lambda)$ lacks the monotonicity properties of MC, see Remark 1.3 below. In Section 3 we also prove the following proposition.

Proposition 1.1. For any $\underline{m} \in \ell_2^{\downarrow}$ our graphical construction of $\mathrm{MCLD}(\lambda)$ (see Section 3) almost surely gives a function $t \mapsto \mathbf{m}_t$ with $\mathbf{m}_0 = \underline{m}$ which is càdlàg with respect to the $d(\cdot, \cdot)$ -metric.

The main result of this paper is that our construction indeed gives rise to a well-behaved continuous-time Markov process on ℓ_2^{\downarrow} :

Theorem 1.2 (Feller property). Let $\underline{m}^{(n)}$, $n \in \mathbb{N}$ be a convergent sequence of elements of ℓ_{+}^{\downarrow} with limit $\underline{m}^{(\infty)}$, i.e., $\lim_{n\to\infty} d(\underline{m}^{(n)},\underline{m}^{(\infty)}) = 0$. For any $t \in \mathbb{R}_{+}$ and $n \in \mathbb{N}_{+} \cup \{\infty\}$, denote by $\mathbf{m}_{t}^{(n)}$ the $\mathrm{MCLD}(\lambda)$ process with initial condition $m^{(n)}$ at time t. For any $t \geq 0$ we have

$$\mathbf{m}_t^{(n)} \xrightarrow{d} \mathbf{m}_t^{(\infty)}, \quad n \to \infty,$$
 (1.3)

where $\stackrel{d}{\longrightarrow}$ denotes convergence in distribution of random variables on the Polish space $(\ell_2^{\downarrow}, d(\cdot, \cdot))$.

We will prove Theorem 1.2 in Section 4 using an argument that involves truncation and coupling.

Remark 1.3. The reason why the proof of the Feller property for $MCLD(\lambda)$ is more involved than the proof of the Feller property for MC (c.f. the proof of [3, Proposition 5] in [3, Section 4.2]) is that the natural graphical construction of $MCLD(\lambda)$ is not *monotone*:

If we obtain $\mathbf{m}'_t, t \ge 0$ from $\mathbf{m}_t, t \ge 0$ by inserting an extra deletion event at time t_1 then it might happen that this deletion prevents later coagulations and deletions, so that \mathbf{m}'_{t_2} has more/bigger components than \mathbf{m}_{t_2} for some $t_2 > t_1$. Similarly, insertion of an extra coagulation event at some time might lead to the deletion of more/bigger components and thus create a state with fewer/smaller components at a later time.

1.1 Motivation, related results

Our reason for developing the theory of $\mathrm{MCLD}(\lambda)$ on the state space ℓ_2^{\downarrow} is that we want to understand the scaling limit of the time evolution of large connected component sizes in the self-organized critical mean field frozen percolation model [22], as we now explain.

The frozen percolation process on the binary tree was defined in [5]: the model is a modification of the dynamical percolation process on the binary tree which makes the following informal description precise: edges appear with rate 1 and if an infinite component appears, we immediately "freeze" it, and we do not allow edges with an end-vertex in a frozen component to appear.

Remark 1.4. I. Benjamini and O. Schramm showed that it is impossible to define a similar modification of the percolation process on \mathbb{Z}^2 , c.f. [9,

Section 3, Remark (i)]. Various modifications of the two-dimensional frozen percolation model where large finite clusters are frozen are further explored in [7, 16, 8, 6]. The result of [17] about the closely related model of two-dimensional self-destructive percolation implies non-existence of the so-called two dimensional forest fire process, c.f. [17, Section 3.2]. However, the result of [2] about self-destructive percolation on the high-dimensional lattice \mathbb{Z}^d indicates that the self-organized critical forest fire process on \mathbb{Z}^d should exist if d is high enough. The existence and uniqueness of the subcritical forest fire process on \mathbb{Z}^d was proved in [14, 15].

Let us now recall the notion of mean-field frozen percolation process from [22] (using slightly different notation).

Definition 1.5 (FP $(n, \lambda(n))$). We start with a graph $F_0^{(n)}$ on n vertices. Between each pair of unconnected vertices an edge appears with rate 1/n; also, every connected component of size k is deleted with rate $\lambda(n) \cdot k$. (When a component is deleted, its vertices as well as its edges are removed from the graph.) Let $F_t^{(n)}$ be the graph at time t. Denote by

$$\mathbf{M}^{(n)}(t) = \left(M_1^{(n)}(t), M_2^{(n)}(t), \dots\right) \in \ell_0^{\downarrow}$$

the sequence of component sizes of $F_t^{(n)}$, arranged in decreasing order.

Then $\mathbf{M}^{(n)}(t), t \ge 0$ is a Markov process – let us call it here the frozen percolation component process on n vertices with lightning rate $\lambda(n)$, or briefly $\mathrm{FP}(n,\lambda(n))$. In fact, up to time-change, $\mathbf{M}^{(n)}(t), t \ge 0$ evolves according to the rules (1.2) of MCLD.

Remark 1.6. We note that $FP(n, \lambda(n))$ is a simplification of the mean field forest fire model [23], the definition of which agrees with Definition 1.5 above, with the only difference that in the forest fire model we only delete the edges of the connected components that are destroyed by fire, i.e., a destroyed component of size k is immediately replaced by k singletons. The mean field forest fire model behaves very similarly to the mean field frozen percolation model (e.g., the self-organized critical behaviour of the two models are quite similar, see also Remark 1.10(iii) below), however the mathematics of the mean field frozen percolation model is simpler than that of the mean field forest fire model, e.g. the solution of the system of differential equations that appears in [22, Theorem 1.2] is fairly explicit, while the system of differential equations that appears in [23, Theorem 2] currently does not have an explicit solution.

Remark 1.7. One studies the asymptotic behaviour of the component size structure of $FP(n, \lambda(n))$ when $1 \ll n$ and $1/n \ll \lambda(n) \ll 1$. Definition 1.5 above is slightly different from the one proposed in [5, Section 5.5] and studied in [21] where connected components are frozen when their size exceeds a threshold $\omega(n)$ satisfying $1 \ll \omega(n) \ll n$. The results [21, Theorem 1.1] and [22, Theorem 1.2] are very similar: indeed, if one is interested in small connected component densities then the two models produce exactly the same (self-organized critical) behaviour. However, if one is interested in the scaling limit of big component dynamics, the exact deletion mechanism does crucially enter the picture.

We are interested in identifying the scaling limit of $\operatorname{FP}(n,\lambda(n))$ as $n\to\infty$. In order to describe the kind of result we are after, let us recall that the large components of the dynamical Erdős-Rényi random graph process in the critical window $\mathcal{G}(n,\frac{1+tn^{-1/3}}{n}), t\in\mathbb{R}$, scaled by $n^{2/3}$, converge in law to the standard multiplicative coalescent process $(\mathcal{M}(t), t\in\mathbb{R})$, see [3, Section 4.3].

Remark 1.8. The family of multiplicative coalescent processes defined for all $t \in \mathbb{R}$ (i.e., the *eternal* MC processes) are characterized in [4]. The class of inhomogeneous random graph models whose scaling limit is the standard MC is explored in [10, 12] (see also references therein). The scaling limits of other classes of inhomogeneous random graph models are related to non-standard eternal MC processes, see [4, 11]. The continuum scaling limit of the metric structure of critical random graphs is studied in [1, 13] (see also references therein).

The next result gives a scaling limit for the frozen percolation process started from a critical Erdős-Rényi graph.

Proposition 1.9. Fix $u \in \mathbb{R}$ and let $F_0^{(n)}$ be an Erdős-Rényi graph $\mathcal{G}(n,p)$ with edge probability $p = \frac{1+un^{-1/3}}{n}$. Let $\lambda > 0$ and let $\mathbf{M}^{(n)}(t), t \geq 0$ be the $\mathrm{FP}(n, \lambda n^{-1/3})$ process with initial state $F_0^{(n)}$. Define $\mathbf{m}^{(n)}(t), t \geq 0$ by

$$\mathbf{m}^{(n)}(t) := \left(n^{-2/3} M_1^{(n)}(n^{-1/3}t), n^{-2/3} M_2^{(n)}(n^{-1/3}t), \dots\right). \tag{1.4}$$

Then as $n \to \infty$ the finite dimensional marginals of the sequence of ℓ_2^{\downarrow} -valued processes $\mathbf{m}^{(n)}(t), t \geq 0$ converge in law to the finite dimensional marginals of the MCLD(λ) process ($\mathbf{m}(t), t \geq 0$) started from an initial state with distribution $\mathbf{m}(0) \sim \mathcal{M}(u)$ (i.e., the state of the standard multiplicative coalescent process at time u), i.e., for every $k \in \mathbb{N}$ and $0 \leq t_1 < t_2 < \cdots < t_k$ we have

$$\left(\mathbf{m}^{(n)}(t_1), \mathbf{m}^{(n)}(t_2), \dots, \mathbf{m}^{(n)}(t_k)\right) \stackrel{\mathrm{d}}{\longrightarrow} \left(\mathbf{m}(t_1), \mathbf{m}(t_2), \dots, \mathbf{m}(t_k)\right), \quad n \to \infty.$$

The proof of Proposition 1.9 follows as an application of Theorem 1.2 (for details of the proof, we refer to [19, Proposition 6.10]).

Remark 1.10.

- (i) Loosely speaking, if $(\mathbf{m}(t), t \geq 0)$ is the MCLD(λ) process started from an initial state with distribution $\mathbf{m}(0) \sim \mathcal{M}(u)$ (this is the limit object that appears in Proposition 1.9), then we have $\mathbf{m}(t) \sim \mathcal{M}(u+t-\Phi(t))$, where $\Phi(t)$ denotes the sum of the sizes of the components deleted up to time t (see [19, Proposition 6.7(ii)] for a precise formulation of this property). In fact, in [19, Proposition 6.7] we give a representation of $(\mathbf{m}(t), t \geq 0)$ on the probability space of a standard Brownian motion using what we call the "rigid" representation of MCLD(λ). We note that Theorem 1.2 is also crucially used when we extend our rigid representation results from ℓ_0^{\downarrow} to ℓ_2^{\downarrow} in [19, Section 5].
- (ii) In [20] we describe the possible scaling limits that can arise from a $FP(n, \lambda n^{-1/3})$ process started from an empty graph. The possible limit objects are eternal $MCLD(\lambda)$ processes (i.e., they are defined for any $t \in \mathbb{R}$). The "arrival at the critical window" gives rise to a non-stationary $MCLD(\lambda)$ scaling limit, while the scaling limit in the "self-organized critical" regime is a stationary $MCLD(\lambda)$ (see also [19, Remark 6.8]).
- (iii) We conjecture that the scaling limit of the coagulation-fragmentation dynamics of big components of the mean field forest fire model (c.f. Remark 1.6 above) with lightning rate $\lambda n^{-1/3}$ is also an MCLD(λ) process.

2 Notation and basic results

The aim of this section is to collect some basic results about the multiplicative coalescent from [3] and [18]. In some cases, we will augment these results to fit our purposes or present them using different notation.

We define

$$\ell_2^+ = \left\{ x = (x_1, x_2, \dots) : \forall i \ x_i \ge 0, \quad \sum_{i \ge 1} x_i^2 < +\infty \right\}.$$

We have $\ell_2^{\downarrow} \subseteq \ell_2^+$. Define the mapping

$$ord: \ell_2^+ \to \ell_2^{\downarrow} \tag{2.1}$$

by letting $\operatorname{ord}(\underline{x})$ be the decreasing rearrangement of $\underline{x} \in \ell_2^+$.

Definition 2.1. If $\underline{m} \in \ell_2^{\downarrow}$ and G is a graph with vertex set $V \subseteq \mathbb{N}_+$, denote by $\operatorname{ord}(\underline{m}, G)$ the ordered sequence of the weights of the connected components of G. More precisely, if C_1, C_2, \ldots is the sequence of the vertex sets of the connected components of G, we define

$$\underline{x}_G = \left(\sum_{i \in \mathcal{C}_1} m_i, \sum_{i \in \mathcal{C}_2} m_i, \dots\right) \quad \text{and} \quad \operatorname{ord}(\underline{m}, G) \stackrel{(2.1)}{=} \operatorname{ord}(\underline{x}_G), \tag{2.2}$$

assuming that $\underline{x}_G \in \ell_2^+$. We also denote

$$S_2^G = \sum_{k=1}^{\infty} \left(\sum_{j \in \mathcal{C}_k} m_j \right)^2 = \|x_G\|_2^2 = \|\operatorname{ord}(\underline{x}_G)\|_2^2.$$
 (2.3)

Let us now state an elementary yet useful result which involves the metric $d(\cdot, \cdot)$ defined in (1.1).

Lemma 2.2. If $\underline{m} \in \ell_2^{\downarrow}$ and G, G' are graphs with vertex sets $V, V' \subseteq \mathbb{N}_+$ such that $V \subseteq V', G \subseteq G'$ and $\operatorname{ord}(\underline{m}, G) \in \ell_2^{\downarrow}$ then we have

$$\mathrm{d}\left(\mathrm{ord}(\underline{m},G),\mathrm{ord}(\underline{m},G')\right)\leqslant\sqrt{\|\mathrm{ord}(\underline{m},G')\|_2^2-\|\mathrm{ord}(\underline{m},G)\|_2^2}.$$

Proof. This is a special case of [3, Lemma 17].

Let us recall the graphical construction used in [3, Section 1.5] to define the multiplicative coalescent process.

Definition 2.3. Let $(\xi_{i,j})_{1 \leq i < j < \infty}$ denote independent random variables with EXP(1) distribution. Given $\underline{x} \in \ell_2^+$ let us define the simple graph G_t with vertex set \mathbb{N}_+ and an edge between i and j if and only if $\xi_{i,j} \leq tx_ix_j$. For $i, j \in \mathbb{N}_+$ we denote by $i \stackrel{G_t}{\longleftrightarrow} j$ the event that i and j are connected by a simple path in the graph G_t .

Given G_t we define the connected components $(\mathcal{C}_k(t))_{k=1}^{\infty}$ of G_t by

$$i_k = \min\{ \mathbb{N}_+ \setminus \bigcup_{l=1}^{k-1} C_l(t) \}, \quad C_k(t) = \{ i \in \mathbb{N}_+ : i \stackrel{G_t}{\longleftrightarrow} i_k \}, \quad k \geqslant 1. \quad (2.4)$$

Note that we have

$$S_2^{G_t} \stackrel{(2.3)}{=} S_2^{G_0} + \sum_{i \neq j} x_i x_j \mathbb{1}[i \stackrel{G_t}{\longleftrightarrow} j]$$
 (2.5)

and $S_2^{G_0} = \sum_{i=1}^{\infty} x_i^2 < +\infty \text{ if } \underline{x} \in \ell_2^+.$

The statement of the next lemma follows from [3, Proposition 5] and shows that Definitions 2.1 and 2.3 give rise to a graphical representation of the ℓ_2^{\downarrow} -valued multiplicative coalescent process with initial state $\underline{m} \in \ell_2^{\downarrow}$ in the form $\operatorname{ord}(m, G_t), t \geq 0$.

Lemma 2.4. For any $t \ge 0$ and $\underline{x} \in \ell_2^+$ we have

$$\mathbf{P}\left(S_2^{G_t} < +\infty\right) = 1. \tag{2.6}$$

In particular, for any $t \in \mathbb{R}_+$ the weights of the connected components of G_t are almost surely finite:

$$\mathbf{P}\left(\forall k \in \mathbb{N}_{+} : \sum_{i \in \mathcal{C}_{k}(t)} x_{i} < +\infty\right) = 1.$$
 (2.7)

The next lemma is an extended version of [18, (2.2)].

Lemma 2.5. For any $\underline{x} \in \ell_2^+$ and $i, j \in \mathbb{N}_+$ and $t < \frac{1}{S_2^{G_0}}$ we have

$$\mathbf{P}\left(i \stackrel{G_t}{\longleftrightarrow} j\right) \leqslant \frac{x_i \cdot x_j \cdot t}{1 - t \cdot S_2^{G_0}}.$$
 (2.8)

Proof.

$$\mathbf{P}\left(i \stackrel{G_t}{\longleftrightarrow} j\right) \leqslant \sum_{k=1}^{\infty} \mathbf{P}\left(\exists i_0, \dots, i_k \in \mathbb{N}_+ : i_0 = i, i_k = j \text{ and } \atop (i_0, i_1, \dots, i_{k-1}, i_k) \text{ is a simple path in } G_t\right) \leqslant \sum_{k=1}^{\infty} \sum_{(i_1, \dots, i_{k-1}) \in \mathbb{N}_+^{k-1}} \prod_{l=1}^k (1 - \exp(-x_{i_{l-1}} x_{i_l} t)) \leqslant \sum_{k=1}^{\infty} \sum_{(i_1, \dots, i_{k-1}) \in \mathbb{N}_+^{k-1}} \prod_{l=1}^k x_{i_{l-1}} x_{i_l} t = x_i x_j t \cdot \sum_{k=1}^{\infty} \sum_{(i_1, \dots, i_{k-1}) \in \mathbb{N}_+^{k-1}} \prod_{l=1}^{k-1} x_{i_l}^2 t = x_i x_j t \cdot \sum_{k=1}^{\infty} (t \cdot S_2^{G_0})^{k-1} = \frac{x_i \cdot x_j \cdot t}{1 - t \cdot S_2^{G_0}}. \quad (2.9)$$

Corollary 2.6. For any $\underline{x} \in \ell_2^+$, $t \ge 0$ and $i, j \in \mathbb{N}_+$, if

$$S_2^{G_0} \leqslant \frac{1}{2t} \tag{2.10}$$

holds then we have

$$\mathbf{E}\left(S_2^{G_t}\right) \leqslant 2S_2^{G_0} \tag{2.11}$$

Proof. Using (2.5), (2.8) and (2.10) we obtain

$$\mathbf{E}\left(S_{2}^{G_{t}}\right) \leqslant S_{2}^{G_{0}} + 2t \sum_{i \neq j} x_{i}^{2} x_{j}^{2} \leqslant S_{2}^{G_{0}} + 2t \cdot \left(S_{2}^{G_{0}}\right)^{2} \stackrel{(2.10)}{\leqslant} 2S_{2}^{G_{0}}. \quad (2.12)$$

The next lemma is based on [3, Lemma 23] and [18, (2.5)]. It will be used in Section 4 to show that the truncated process is close to the original process if the truncation threshold is chosen big enough.

Lemma 2.7. Let $\underline{x}, \underline{y} \in \ell_2^+$ and $t \ge 0$. Denote the index set of \underline{x} by I and the index set of \underline{y} by J. Denote by

$$a = \|\underline{x}\|_2^2 < +\infty$$
 and $b = \|y\|_2^2 < +\infty$.

Consider the bipartite random graph B_t with vertex set $I \cup J$, where $i \in I$ and $j \in J$ are connected with probability $1 - \exp(-tx_iy_j)$. Then we have

$$|I| < +\infty \implies \mathbf{E}\left(S_2^{B_t}\right) < +\infty.$$
 (2.13)

Moreover, if

$$t^2 ab \leqslant \frac{1}{2},\tag{2.14}$$

holds then we have

$$\mathbf{E}\left(S_2^{B_t}\right) - a \leqslant 2b \cdot (1 + ta)^2. \tag{2.15}$$

Proof. First note that, similarly to (2.5), we have

$$\mathbf{E}\left(S_{2}^{B_{t}}\right) = a + b + \sum_{i_{1} \neq i_{2} \in I} x_{i_{1}} x_{i_{2}} \mathbf{P}\left(i_{1} \stackrel{B_{t}}{\longleftrightarrow} i_{2}\right) + \sum_{j_{1} \neq j_{2} \in J} y_{j_{1}} y_{j_{2}} \mathbf{P}\left(j_{1} \stackrel{B_{t}}{\longleftrightarrow} j_{2}\right) + 2 \sum_{i \in I, j \in J} x_{i} y_{j} \mathbf{P}\left(i \stackrel{B_{t}}{\longleftrightarrow} j\right). \quad (2.16)$$

Now note that the number of visits to I of a simple path in B_t is at most |I|. Using this idea and a calculation similar to (2.9), we obtain the inequalities

$$\mathbf{P}\left(i_{1} \overset{B_{t}}{\longleftrightarrow} i_{2}\right) \leqslant (x_{i_{1}}x_{i_{2}} \cdot b \cdot t^{2}) \cdot \sum_{k=1}^{|I|} (t^{2}ab)^{k-1}, \quad i_{1} \neq i_{2}, \ i_{1}, i_{2} \in I$$

$$\mathbf{P}\left(j_{1} \overset{B_{t}}{\longleftrightarrow} j_{2}\right) \leqslant (y_{j_{1}}y_{j_{2}} \cdot a \cdot t^{2}) \cdot \sum_{k=1}^{|I|} (t^{2}ab)^{k-1}, \quad j_{1} \neq j_{2}, \ j_{1}, j_{2} \in J$$

$$\mathbf{P}\left(i \overset{B_{t}}{\longleftrightarrow} j\right) \leqslant (x_{i}y_{j}t) \cdot \sum_{k=1}^{|I|} (t^{2}ab)^{k-1}, \quad i \in I, \ j \in J$$

Combining these inequalities with (2.16) we obtain (2.13) as well as

$$\mathbf{E}\left(S_{2}^{B_{t}}\right) - a \overset{(2.14)}{\leqslant} b + 2\left(a^{2} \cdot b \cdot t^{2} + b^{2} \cdot a \cdot t^{2} + 2a \cdot b \cdot t\right) \overset{(2.14)}{\leqslant} b \cdot \left(1 + 2a^{2}t^{2} + 1 + 4at\right) = 2b\left(1 + at\right)^{2}.$$

This completes the proof of (2.15).

Lemma 2.8. With probability 1, the function $t \mapsto \operatorname{ord}(\underline{m}, G_t)$ (see (2.2)) is càdlàg with respect to the $d(\cdot, \cdot)$ -metric (defined in (1.1)).

Proof. Let us fix some $T \ge 0$. Denote by A the event

$$A = \{S_2^{G_T} < +\infty\} \cap \left\{ \begin{array}{c} \text{for any } i, j \in \mathbb{N} \text{ the number of simple} \\ \text{paths connecting } i \text{ and } j \text{ in } G_T \text{ is finite} \end{array} \right\}$$
 (2.17)

By Lemma 2.4 the event A almost surely holds. Assuming that A holds, we will show that $t \mapsto \operatorname{ord}(\underline{m}, G_t)$ is càdlàg on [0, T).

Since $G_s \subseteq G_t$ if $s \leq t$, we can apply Lemma 2.2 in order to reduce our task to showing that the function $t \mapsto S_2^{G_t}$ is càdlàg on [0,T). If A holds, then for any $i,j \in \mathbb{N}$ the function $t \mapsto \mathbb{1}[i \stackrel{G_t}{\longleftrightarrow} j]$ is càdlàg on [0,T). Using this fact, (2.5) and the dominated convergence theorem, we obtain that indeed $t \mapsto S_2^{G_t}$ is also càdlàg on [0,T).

3 Graphical construction of $MCLD(\lambda)$

Recall the informal definition of the $\mathrm{MCLD}(\lambda)$ process \mathbf{m}_t from (1.2). We now give a graphical construction of the process \mathbf{m}_t with initial state $\underline{m} \in \ell_2^{\downarrow}$ and deletion rate λ . Let

$$(\xi_{i,j})_{1 \leq i < j < \infty}$$
 be random variables with EXP(1) distribution,
 $(\lambda_i)_{1 \leq i < \infty}$ be random variables with EXP(λ) distribution, (3.1)

and let us also assume that all of these random variables are independent.

The heuristic description of our graphical construction is as follows: we increase t continuously and if the event $\xi_{i,j} = tm_im_j$ occurs for some $1 \le i < j < \infty$, we merge the components of the vertices i and j, moreover if $\lambda_i = tm_i$ for some $i \in \mathbb{N}_+$, then we say that a lightning strikes vertex i and delete the connected component of vertex i. Since the total rate of merger and deletion events is infinite if $\sum_i m_i = +\infty$, we need to be careful with the above heuristic definition if we want to make it precise: we will now provide the graphical construction.

In Definition 2.3 we defined the simple graph G_t with vertex set \mathbb{N}_+ . We will define for any $t \in \mathbb{R}_+$

the set of intact vertices
$$\mathcal{V}_t \subseteq \mathbb{N}_+$$
 and the set of burnt vertices $\mathbb{N}_+ \backslash \mathcal{V}_t$. (3.2)

The graph H_t will denote the subgraph of G_t spanned by \mathcal{V}_t and \mathbf{m}_t will denote the ordered sequence of component weights of H_t .

Recall that we enumerated the connected components $C_k(t)$, $k \in \mathbb{N}_+$ of G_t in (2.4). By the properties of exponential random variables, (2.7) and the independence of $(\xi_{i,j})_{1 \leq i < j < \infty}$ and $(\lambda_i)_{i=1}^{\infty}$, we see that for every $t \geq 0$

$$\mathbf{P}\left(\forall k \in \mathbb{N}_{+} : \sum_{i \in \mathcal{C}_{k}(t)} \mathbb{1}[\lambda_{i} \leqslant t m_{i}] < +\infty\right) = 1.$$
(3.3)

This implies that for every $t \ge 0$ and $k \in \mathbb{N}_+$, there exists an almost surely finite \mathbb{N} -valued random variable N (the number of lightnings that hit the component $\mathcal{C}_k(t)$ by time t), indices $i_1, \ldots, i_N \subseteq \mathcal{C}_k(t)$ (the vertices that are hit by lightning) and times $0 < t_1 < \cdots < t_N \le t$ (the ordered sequence of the times of the lightnings) such that

$$\{i \in \mathcal{C}_k(t) : \lambda_i \leqslant t m_i\} = \{i_1, \dots, i_N\} \text{ and } \forall 1 \leqslant l \leqslant N : t_l = \frac{\lambda_{i_l}}{m_{i_l}}.$$

We now define the set of intact vertices $\mathcal{V}_t \subseteq \mathbb{N}_+$ by constructing $\mathcal{V}_t \cap \mathcal{C}_k(t)$ for every $k \in \mathbb{N}_+$.

Let us fix $k \in \mathbb{N}_+$. We recursively define $\mathcal{V}_{t_l} \cap \mathcal{C}_k(t)$ for each $1 \leq l \leq N$ in the following way.

- (i) At $t_0 = 0$ we have $\mathcal{V}_{t_0} \cap \mathcal{C}_k(t) = \mathcal{C}_k(t)$.
- (ii) Assume that we have already constructed $\mathcal{V}_{t_{l-1}} \cap \mathcal{C}_k(t)$ for some $1 \leq l \leq N$. We define $\mathcal{V}_{t_l} \cap \mathcal{C}_k(t)$ by deleting the connected component of i_l in the restriction of the graph G_{t_l} to the vertex set $\mathcal{V}_{t_{l-1}} \cap \mathcal{C}_k(t)$.
- (iii) With this recursion we define $\mathcal{V}_{t_N} \cap \mathcal{C}_k(t)$. Since there are no lightnings hitting $\mathcal{C}_k(t)$ between t_N and t, let $\mathcal{V}_t \cap \mathcal{C}_k(t) = \mathcal{V}_{t_N} \cap \mathcal{C}_k(t)$.

Since $C_k(t)$, $k \in \mathbb{N}_+$ is a partition of \mathbb{N}_+ , we define

$$\mathcal{V}_t = \bigcup_{k \geqslant 1} (\mathcal{V}_t \cap \mathcal{C}_k(t))$$
 and H_t to be the subgraph of G_t spanned by \mathcal{V}_t . (3.4)

Recalling Definition 2.1 we let

$$\mathbf{m}_t = \operatorname{ord}(\underline{m}, H_t). \tag{3.5}$$

Lemma 3.1. For any $\underline{m} \in \ell_2^{\downarrow}$ the graphical construction (3.5) of the process \mathbf{m}_t gives an MCLD(λ) process with initial condition \underline{m} , i.e., an ℓ_2^{\downarrow} -valued Markov process whose dynamics satisfy the informal definition given in (1.2).

Proof. \mathbf{m}_t is a random element of ℓ_2^{\downarrow} , because we have

$$\|\mathbf{m}_t\|_2^2 = S_2^{H_t} \leqslant S_2^{G_t} \stackrel{(2.6)}{<} + \infty.$$

The fact that \mathbf{m}_t is a Markov process with the prescribed transition rates follows from the memoryless property and independence of the random variables $(\xi_{i,j})_{1 \leq i < j < \infty}$ and $(\lambda_i)_{i=1}^{\infty}$. We omit further details.

Proof of Proposition 1.1. We will show that with probability 1, the function $t \mapsto \operatorname{ord}(\underline{m}, H_t)$ is càdlàg with respect to the $d(\cdot, \cdot)$ -metric, see (1.1).

Let us fix some $T \ge 0$. We know that the event A defined in (2.17) almost surely holds. Denote by B the event that every connected component of G_T is exposed to only finitely many lightning strikes on [0,T]. By (3.3), the event B occurs almost surely. Assuming that $A \cap B$ holds, we will show that $t \mapsto \operatorname{ord}(\underline{m}, H_t)$ is càdlàg on [0,T). For any $t \ge 0$, define

- $\hat{H}_{t+\Delta t}$ to be the subgraph of G_t spanned by $\mathcal{V}_{t+\Delta t}$,
- $\check{H}_{t+\Delta t}$ to be the subgraph of $G_{t+\Delta t}$ spanned by \mathcal{V}_t .

Recalling (3.4) and the inclusions $G_t \subseteq G_{t+\Delta t}$ and $\mathcal{V}_{t+\Delta t} \subseteq \mathcal{V}_t$ we see that

$$\hat{H}_{t+\Delta t} \subseteq H_t \subseteq \check{H}_{t+\Delta t}$$
 and $\hat{H}_{t+\Delta t} \subseteq H_{t+\Delta t} \subseteq \check{H}_{t+\Delta t}$

so we can apply Lemma 2.2 and the triangle inequality in order to reduce our task of proving right-continuity of $t \mapsto \operatorname{ord}(\underline{m}, H_t)$ at t to showing that

(a)
$$\lim_{\Delta t \to 0_+} S_2^{\check{H}_{t+\Delta t}} - S_2^{H_t} = 0$$
, (b) $\lim_{\Delta t \to 0_+} S_2^{H_t} - S_2^{\hat{H}_{t+\Delta t}} = 0$.

Now (a) follows from the fact that the graphical representation of the multiplicative coalescent possesses the càdlàg property (see Lemma 2.8).

In order to show (b) we observe that on the event B, for every connected component C of G_T , we have

$$\lim_{\Delta t \to 0} \mathbb{1} \left[\exists i \in \mathcal{C} : t m_i < \lambda_i \leqslant (t + \Delta t) m_i \right] = 0.$$

Given this observation, we see that for every connected component \mathcal{C} of H_t we have $\lim_{\Delta t \to 0} \mathbb{1}[\mathcal{C} \subseteq \mathcal{V}_{t+\Delta t}] = 1$. Using this fact, $S_2^{H_t} < \infty$ and the dominated convergence theorem, we obtain (b).

The proof of the existence of left limits is similar and we omit it. \Box

4 Feller property of $MCLD(\lambda)$

Definition 4.1. The graphical construction of Section 3 gives a joint realization of all of the $\mathrm{MCLD}(\lambda)$ processes with different initial conditions by using the same collection of random variables $(\xi_{i,j})_{1 \leq i < j < \infty}$ and $(\lambda_i)_{1 \leq i < \infty}$ (see (3.1)). We call this coupling the (ξ, λ) -coupling.

Theorem 4.2. Let $\underline{m}^{(n)}$, $n \in \mathbb{N}$ be a convergent sequence of elements of ℓ_2^{\downarrow} and let $\underline{m}^{(\infty)}$ denote their limit, i.e., $\lim_{n\to\infty} d(\underline{m}^{(n)}, \underline{m}^{(\infty)}) = 0$. For any $t \in \mathbb{R}_+$ and $n \in \mathbb{N}_+ \cup {\infty}$, denote by $\mathbf{m}_t^{(n)}$ the $\mathrm{MCLD}(\lambda)$ process with initial condition $\underline{m}^{(n)}$ at time t. Under the (ξ, λ) -coupling, we have

$$d(\mathbf{m}_t^{(n)}, \mathbf{m}_t^{(\infty)}) \xrightarrow{p} 0, \quad n \to \infty.$$
 (4.1)

Theorem 4.2 implies that the $MCLD(\lambda)$ Markov process indeed possesses the Feller property, i.e., Theorem 1.2 holds.

We want to prove Theorem 4.2 using truncation, because (4.1) trivially holds for the truncated process. However, we cannot directly apply Lemma 2.2 to compare the original with the truncated process, because we cannot upper bound the state of the truncated process at time t by the state of the original process at time t (c.f. Remark 1.3).

In Section 4.1 we overcome this problem by introducing two auxiliary objects that upper/lower bound both the original and the truncated object, but yet these auxiliary objects can be shown to be close to each other if we only throw away a small part of the original when we truncate.

In Section 4.2 we prove Theorem 4.2 using the results of Section 4.1 and variant of the $\varepsilon/3$ -argument.

4.1 Bounding the effect of truncation

In this subsection, we will fix $t \ge 0$ as well as an initial state $\underline{m} \in \ell_2^{\downarrow}$, and omit the dependence of random variables on t and \underline{m} . We also fix a truncation threshold $m \in \mathbb{N}$.

Definition 4.3. Recall Definition 2.3. Denote by G, $G^{m\downarrow}$ and $G^{m\uparrow}$ the graphs with adjacency matrix $\mathbb{1}[\xi_{i,j} \leq tm_im_j]$ on the vertex set \mathbb{N}_+ , $\{1,\ldots,m\}$, and $\{m+1,m+2,\ldots\}$, respectively.

Let $\underline{m}^{(m)}$ denote the vector \underline{m} truncated at index m:

$$\underline{m}^{(m)} = (m_1, \dots, m_m, 0, 0, \dots), \text{ where } \underline{m} = (m_1, m_2, \dots).$$

Let \mathbf{m} (resp. $\mathbf{m}^{(m)}$) denote the state at time t of the realization under the (ξ, λ) -coupling of the $\mathrm{MCLD}(\lambda)$ process with initial state \underline{m} (resp. $\underline{m}^{(m)}$).

Denote by \mathcal{V} and $\mathcal{V}^{(m)}$ the corresponding sets of intact vertices, see (3.4). Denote by H and $H^{(m)}$ the subgraphs of G spanned by \mathcal{V} and $\mathcal{V}^{(m)}$.

In order to compare \mathbf{m} with $\mathbf{m}^{(m)}$, we need the following result.

Lemma 4.4. If $\hat{G}^{(m)}$ and $\check{G}^{(m)}$ are random graphs with vertex sets

$$V(\hat{G}^{(m)}), V(\check{G}^{(m)}) \subseteq \mathbb{N}_+$$

and under the (ξ, λ) -coupling we have

$$\widehat{G}^{(m)} \subseteq H^{(m)} \subseteq \widecheck{G}^{(m)}, \quad \widehat{G}^{(m)} \subseteq H \subseteq \widecheck{G}^{(m)} \tag{4.2}$$

then almost surely we have

$$d(\mathbf{m}, \mathbf{m}^{(m)}) \leq 3 \cdot \sqrt{S_2^{\check{G}^{(m)}} - S_2^{\hat{G}^{(m)}}}.$$
 (4.3)

Proof. First note that it follows from (4.2) that

$$S_2^{\hat{G}^{(m)}} \leqslant S_2^{H^{(m)}} \leqslant S_2^{\check{G}^{(m)}}, \quad S_2^{\hat{G}^{(m)}} \leqslant S_2^{H} \leqslant S_2^{\check{G}^{(m)}}.$$
 (4.4)

Thus we have

$$d(\mathbf{m}, \mathbf{m}^{(m)}) \overset{(2.2)}{\leqslant} d(\mathbf{m}, \operatorname{ord}(\underline{m}, \check{G}^{(m)})) + \\ d(\operatorname{ord}(\underline{m}, \check{G}^{(m)}), \operatorname{ord}(\underline{m}^{(m)}, \widehat{G}^{(m)})) + d(\operatorname{ord}(\underline{m}^{(m)}, \widehat{G}^{(m)}), \mathbf{m}^{(m)}) \overset{(*)}{\leqslant} \\ \sqrt{S_2^{\check{G}^{(m)}} - S_2^H} + \sqrt{S_2^{\check{G}^{(m)}} - S_2^{\widehat{G}^{(m)}}} + \sqrt{S_2^{H^{(m)}} - S_2^{\widehat{G}^{(m)}}} \overset{(4.4)}{\leqslant} 3 \cdot \sqrt{S_2^{\check{G}^{(m)}} - S_2^{\widehat{G}^{(m)}}},$$
where (*) follows from (3.5), the inclusions (4.2) and Lemma 2.2.

In Definition 4.8 below we will construct auxiliary graphs $\hat{G}^{(m)}$ and $\check{G}^{(m)}$ in such a way that (4.2) holds. Recall Definition 4.3. Note that $H^{(m)}$ is the subgraph of $G^{m\downarrow}$ spanned by the vertex set $\mathcal{V}^{(m)}$. In particular, every connected component of $H^{(m)}$ is a subset of a connected component of $G^{m\downarrow}$.

The next definition only involves the random variables $(\xi_{i,j})_{1 \leq i < j < \infty}$ (i.e., we don't have to look at $(\lambda_i)_{i=1}^{\infty}$).

Definition 4.5. Given $G^{m\downarrow}$ and $G^{m\uparrow}$, denote the connected components of $G^{m\downarrow}$ by $C_k^{m\downarrow}$, $k \in K$ and the connected components of $G^{m\uparrow}$ by $C_l^{m\uparrow}$, $l \in L$.

Let us define an auxiliary bipartite multigraph \mathcal{B} with vertex set $K \cup L$. Declare $k \in K$ and $l \in L$ connected in \mathcal{B} if $\mathcal{C}_k^{m\downarrow}$ is connected to $\mathcal{C}_l^{m\uparrow}$ in G. We allow parallel edges to be present in \mathcal{B} : if $\mathcal{C}_k^{m\downarrow}$ is connected to $\mathcal{C}_l^{m\uparrow}$ by more than one edge in G, then we put an equal number of parallel edges between $k \in K$ and $l \in L$ in \mathcal{B} .

Now we define a subset $K^* \subseteq K$ indexing "bad" components of $G^{m\downarrow}$. This definition involves the random variables $(\xi_{i,j})_{1 \le i < j < \infty}$ as well as $(\lambda_i)_{i=1}^{\infty}$. The components indexed by $k \in K \backslash K^*$ are "good". The key property of good components will be stated in Lemma 4.7 below.

Definition 4.6. Recall the definition of \mathcal{B} from Definition 4.5.

- (i) An edge-simple path in \mathcal{B} is a path with no repeated edges.
- (ii) We say that $k \in K$ (resp. $l \in L$) is *intact* if no lightning hit any vertex of $C_k^{m\downarrow}$ (resp. $C_l^{m\uparrow}$) before time t. If a vertex of \mathcal{B} is not intact, then we say that it is *damaged*.
- (iii) We say that $k \in K^*$ if $k \in K$ and there is a edge-simple path in \mathcal{B} which consists of at least one edge and connects k to a damaged vertex of \mathcal{B} .

For an illustration of Definition 4.6, see Figure 1.




Figure 1: An illustration of Definition 4.6. The blobs marked with a lightning are damaged connected components of $G^{m\downarrow}$ and $G^{m\uparrow}$. The grey blobs are the "bad" components of $G^{m\downarrow}$. The set of indices of "bad" components is denoted by K^* . Note that intact connected components of $G^{m\downarrow}$ can be "bad" and damaged connected components of $G^{m\downarrow}$ can be "good".

Lemma 4.7. Recalling Definition 4.3, we have

$$\forall k \in K \backslash K^* : \mathcal{C}_k^{m\downarrow} \cap \mathcal{V}^{(m)} = \mathcal{C}_k^{m\downarrow} \cap \mathcal{V}. \tag{4.5}$$

Proof. Let $k \in K \setminus K^*$. Denote by C' the connected component of k in B. We prove (4.5) by considering two cases separately.

First case: k is intact.

Denote by $K' = \mathcal{C}' \cap K$ and $L' = \mathcal{C}' \cap L$. Then

$$\mathcal{C} = \left(igcup_{k' \in K'} \mathcal{C}_{k'}^{m\downarrow}
ight) \cup \left(igcup_{l' \in L'} \mathcal{C}_{l'}^{m\uparrow}
ight)$$

is a connected component of G which contains $C_k^{m\downarrow}$ (c.f. Definition 4.5), moreover our assumption that k is intact together with $k \in K \backslash K^*$ imply that $\mathcal C$ is intact (c.f. Definition 4.6), thus we have $C_k^{m\downarrow} \cap \mathcal V^{(m)} = C_k^{m\downarrow}$ and $C_k^{m\downarrow} \cap \mathcal V = C_k^{m\downarrow}$, therefore (4.5) holds.

Second case: k is damaged.

 $\mathcal{C}'\setminus\{k\}$ is the disjoint union of some connected components $\mathcal{C}'_N, N \in \mathbb{N}$ of $\mathcal{B}\setminus\{k\}$. Our assumption that k is damaged, Definition 4.6 and the fact that $k \in K\setminus K^*$ together imply that there are no parallel edges connected to k in \mathcal{B} and no edge-simple circle of the graph \mathcal{B} contains k as a vertex. Therefore for each $N \in \mathbb{N}$, the cluster \mathcal{C}'_N is connected to k by one single edge e_N of \mathcal{B} . Note that $k \in K\setminus K^*$ implies that \mathcal{C}'_N is intact for all $N \in \mathbb{N}$. Therefore, the fires caused by lightnings can only spread "away" from k on the edges $e_N, N \in \mathbb{N}$, so by the graphical construction given in Section 3 and Definition 4.3 we obtain (4.5).

Now we define auxiliary random graphs $\hat{G}^{(m)}$ and $\check{G}^{(m)}$ (c.f. Lemma 4.4).

Definition 4.8. Let $\check{G}^{(m)}$ be the subgraph of G spanned by the vertices

$$V(\check{G}^{(m)}) = \left(\bigcup_{k \in K \setminus K^*} \mathcal{C}_k^{m\downarrow} \cap \mathcal{V}^{(m)}\right) \cup \left(\bigcup_{k \in K^*} \mathcal{C}_k^{m\downarrow}\right) \cup \{m+1, m+2, \dots\}.$$
(4.6)

Define $\widehat{G}^{(m)}$ to be the subgraph of G spanned by the vertices

$$V(\widehat{G}^{(m)}) = \bigcup_{k \in K \setminus K^*} \mathcal{C}_k^{\downarrow m} \cap \mathcal{V}^{(m)}. \tag{4.7}$$

Lemma 4.9. With the above definitions the inclusions (4.2) hold.

Proof. The inclusions $V(\hat{G}^{(m)}) \subseteq \mathcal{V}^{(m)} \subseteq V(\check{G}^{(m)})$ follow from the definitions (4.6), (4.7). Thus $\hat{G}^{(m)} \subseteq H^{(m)} \subseteq \check{G}^{(m)}$ follows from the fact that $H^{(m)}$ is the subgraph of G spanned by the vertex set $\mathcal{V}^{(m)}$.

The inclusions $\hat{G}^{(m)} \subseteq H \subseteq \check{G}^{(m)}$ follow from Lemma 4.7 and the fact that H is the subgraph of G spanned by the vertex set \mathcal{V} .

The next lemma is similar to Lemma 2.7.

Lemma 4.10. Given the above set-up let us condition on the graphs $G^{m\downarrow}$ and $G^{m\uparrow}$ and denote by

$$\alpha = S_2^{G^{m\downarrow}}, \quad \beta = S_2^{G^{m\uparrow}}.$$

There exists a constant $C = C(\lambda, t)$ such that if

$$t^2 \alpha \beta \leqslant \frac{1}{2} \tag{4.8}$$

holds then we have

$$\mathbf{E}\left(S_2^{\check{G}^{(m)}} - S_2^{\widehat{G}^{(m)}} \middle| G^{m\downarrow}, G^{m\uparrow}\right) \leqslant C \cdot \beta \cdot \left((1 + t\alpha)^2 + (1 + t\alpha) \cdot \alpha^{3/2}\right). \tag{4.9}$$

4.1.1 Proof of Lemma 4.10

For any subset \mathcal{C} of \mathbb{N} , denote by

$$w(\mathcal{C}) = \sum_{i \in \mathcal{C}} m_i$$

the weight of the subset, where $\underline{m} = (m_1, m_2, ...)$.

Definition 4.11. Define a bipartite weighted graph $\widetilde{\mathcal{B}}$ whose "left" vertices correspond to the connected components of the restriction of G to the vertex set

$$\widetilde{V}^{(m)} := V(\widecheck{G}^{(m)}) \cap \{1, \dots, m\} \stackrel{(4.6)}{=} \left(\bigcup_{k \in K \setminus K^*} \mathcal{C}_k^{m\downarrow} \cap \mathcal{V}^{(m)} \right) \cup \left(\bigcup_{k \in K^*} \mathcal{C}_k^{m\downarrow} \right),$$

and the "right" vertices correspond to the components of $G^{m\uparrow}$. Define the weights of the vertices of $\widetilde{\mathcal{B}}$ to be the $w(\cdot)$ -weight of the corresponding connected components. We declare two vertices in $\widetilde{\mathcal{B}}$ to be connected if the corresponding subsets are connected in $\check{G}^{(m)}$. Denote by $\widetilde{G}^{(m)}$ the subgraph of G spanned by $\widetilde{V}^{(m)}$.

With the above notation we have

$$S_2^{\check{G}^{(m)}} \stackrel{(4.6)}{=} S_2^{\tilde{\mathcal{B}}}, \qquad S_2^{\tilde{G}^{(m)}} \stackrel{(4.7)}{=} S_2^{\hat{G}^{(m)}} + \sum_{k \in K^*} w(\mathcal{C}_k^{m\downarrow})^2.$$

Thus we can start to rewrite the left-hand side of (4.9):

$$\mathbf{E}\left(S_{2}^{\tilde{G}^{(m)}} - S_{2}^{\hat{G}^{(m)}} \middle| G^{m\downarrow}, G^{m\uparrow}\right) = \\ \mathbf{E}\left(S_{2}^{\tilde{\mathcal{B}}} - S_{2}^{\tilde{G}^{(m)}} \middle| G^{m\downarrow}, G^{m\uparrow}\right) + \mathbf{E}\left(\sum_{k \in K^{*}} w(\mathcal{C}_{k}^{m\downarrow})^{2} \middle| G^{m\downarrow}, G^{m\uparrow}\right).$$

In order to show (4.9), it is enough to prove that (4.8) implies

$$\mathbf{E}\left(S_2^{\widetilde{\mathcal{B}}} - S_2^{\widetilde{G}^{(m)}} \middle| G^{m\downarrow}, G^{m\uparrow}\right) \leqslant 2\beta \cdot (1 + t\alpha)^2, \tag{4.10}$$

$$\mathbf{E}\left(\sum_{k\in K^*} w(\mathcal{C}_k^{m\downarrow})^2 \left| G^{m\downarrow}, G^{m\uparrow} \right.\right) \leqslant 2t^2 \lambda \beta \cdot (1+t\alpha) \cdot \alpha^{3/2}. \tag{4.11}$$

First we deduce (4.10) from Lemma 2.7, with the underlying bipartite graph being $\widetilde{\mathcal{B}}$. Note that the condition (2.14) holds, because $a=S_2^{\widetilde{G}^{(m)}}\leqslant S_2^{G^{m\downarrow}}=\alpha$ and $b=S_2^{G^{m\uparrow}}=\beta$. Thus we have

$$\mathbf{E}\left(S_2^{\widetilde{\mathcal{B}}} - S_2^{\widetilde{G}^{(m)}} \middle| G^{m\downarrow}, G^{m\uparrow}, (\lambda_i)_{i=1}^m\right) \overset{(2.15)}{\leqslant}$$

$$2S_2^{G^{m\uparrow}} \cdot (1 + tS_2^{\widetilde{G}^{(m)}})^2 \leqslant 2\beta \cdot (1 + t\alpha)^2.$$

Now (4.10) follows by averaging over the values of $(\lambda_i)_{i=1}^m$.

In order to prove (4.11), we first give an upper bound on the probability of the event $\{k \in K^*\}$.

For $k \in K$, denote by $x'_k = w(\mathcal{C}_k^{m\downarrow})$ and for $l \in L$, denote $y'_l = w(\mathcal{C}_l^{m\uparrow})$. Note that we have

$$\alpha = \sum_{k \in K} (x'_k)^2, \qquad \beta = \sum_{l \in L} (y'_l)^2.$$

Recall the definition of K^* from Definition 4.6. The next calculation is similar to (2.9), so we omit the first few steps.

$$\mathbf{P}\left(k \in K^* \mid G^{m\downarrow}, G^{m\uparrow}\right) \leqslant \sum_{l_1 \in L} \left(x'_k y'_{l_1} t\right) \left(\lambda y'_{l_1} t\right) + \sum_{l_1 \in L} \sum_{k_1 \in K} \left(x'_k y'_{l_1} t\right) \left(y'_{l_1} x'_{k_1} t\right) \left(\lambda x'_{k_1} t\right) + \sum_{l_1 \in L} \sum_{k_1 \in K} \sum_{l_2 \in L} \left(x'_k y'_{l_1} t\right) \left(y'_{l_1} x'_{k_1} t\right) \left(x'_{k_1} y'_{l_2} t\right) \left(\lambda y'_{l_2} t\right) + \dots = x'_k t^2 \lambda \beta + x'_k t^3 \lambda \alpha \beta + x'_k t^4 \lambda \alpha \beta^2 + \dots = x'_k t^2 \lambda \beta \cdot (1 + t\alpha) \cdot \sum_{n=0}^{\infty} \left(t^2 \alpha \beta\right)^n \overset{(4.8)}{\leqslant} 2x'_k t^2 \lambda \beta \cdot (1 + t\alpha).$$

Now we are ready to prove (4.11):

$$\mathbf{E}\left(\sum_{k\in K^*} (x_k')^2 \left| G^{m\downarrow}, G^{m\uparrow} \right) \leqslant \sum_{k\in K} 2(x_k')^3 t^2 \lambda \beta \cdot (1+t\alpha) \stackrel{(*)}{\leqslant} 2t^2 \lambda \beta \cdot (1+t\alpha) \cdot \alpha^{3/2},$$

where in (*) we used the fact that $x'_k \leq \sqrt{\alpha}$ for any $k \in K$. This completes the proof of (4.9) and Lemma 4.10.

4.2 Proof of Theorem 4.2

Let us fix $t, \lambda \in \mathbb{R}_+$, the sequence $\underline{m}^{(n)}, n \in \mathbb{N}$ and the limit $\underline{m}^{(\infty)}$. For any $n \in \mathbb{N}_+ \cup \{\infty\}$, let $\mathbf{m}_t^{(n,m)}$ denote the realization under the (ξ, λ) -coupling of the $\mathrm{MCLD}(\lambda)$ with initial state

$$\underline{m}^{(n,m)} = (m_1^{(n)}, \dots, m_m^{(n)}, 0, 0, \dots), \text{ where } \underline{m}^{(n)} = (m_1^{(n)}, m_2^{(n)}, \dots).$$
 (4.12)

We also define $\mathcal{V}_t^{(n,m)}$ to be the set of intact vertices of the graph $H_t^{(n,m)}$ of the MCLD(λ) with initial state $\underline{m}^{(n,m)}$ under the (ξ, λ) -coupling.

In order to prove (4.1) we only need to show that for every $\varepsilon > 0$ there exists $m, n_0 \in \mathbb{N}$ such that for all $n \ge n_0$ we have

$$\mathbf{P}\left(d(\mathbf{m}_t^{(n)}, \mathbf{m}_t^{(n,m)}) \geqslant \varepsilon\right) \leqslant 4\varepsilon, \tag{4.13}$$

$$\mathbf{P}\left(d(\mathbf{m}_{t}^{(n,m)}, \mathbf{m}_{t}^{(\infty,m)}) \geqslant \varepsilon\right) \leqslant \varepsilon, \tag{4.14}$$

$$\mathbf{P}\left(d(\mathbf{m}_{t}^{(\infty,m)},\mathbf{m}_{t}^{(\infty)}) \geqslant \varepsilon\right) \leqslant 4\varepsilon. \tag{4.15}$$

Let us fix $\varepsilon > 0$. We know from Lemma 2.4 that

$$\mathbf{P}\left(S_2^{G_t^{(\infty)}} < +\infty\right) = 1,$$

where $G_t^{(\infty)}$ denotes the random graph constructed from the exponential variables $(\xi_{i,j})_{1\leqslant i< j<\infty}$ and the initial state $\underline{m}^{(\infty)}\in\ell_2^\downarrow$ according to the rules described in Definition 2.3. Given $\varepsilon>0$, we can find $M\in\mathbb{R}_+$ such that

$$\mathbf{P}\left(S_2^{G_t^{(\infty)}} \geqslant M - 1\right) \leqslant \varepsilon. \tag{4.16}$$

Recall the notion of the constant $C=C(t,\lambda)$ from Lemma 4.10. Let us choose $\delta>0$ such that

$$t^2 M \delta \leqslant \frac{1}{2}$$
 and $9C \cdot \delta \cdot ((1 + tM)^2 + (1 + tM) \cdot M^{3/2}) \leqslant \varepsilon^3$. (4.17)

Now we choose the truncation threshold m. Since $\underline{m}^{(n)} \to \underline{m}^{(\infty)}$ in l_2 , we can make

$$\sup_{n \in \mathbb{N} \cup \{\infty\}} \|\underline{m}^{(n)} - \underline{m}^{(n,m)}\|_2$$

(where $\underline{m}^{(n,m)}$ is defined in (4.12)) as small as we wish by making m large. Thus by (2.11) and the Markov inequality we can choose m such that

$$\sup_{n \in \mathbb{N} \cup \{\infty\}} \mathbf{P}\left(S_2^{G_t^{(n,m)\uparrow}} \geqslant \delta\right) \leqslant \varepsilon. \tag{4.18}$$

Having fixed m, we note that under the (ξ, λ) -coupling we have

$$d(\mathbf{m}_t^{(n,m)}, \mathbf{m}_t^{(\infty,m)}) \xrightarrow{p} 0, \quad n \to \infty.$$

We also have

$$S_2^{G_t^{(n,m)\downarrow}} \xrightarrow{p} S_2^{G_t^{(\infty,m)\downarrow}} \leqslant S_2^{G_t^{(\infty)}}, \tag{4.19}$$

thus we can choose n_0 such that for all $n \ge n_0$ we have (4.14) and

$$\forall n \in \{n_0, n_0 + 1, \dots\} \cup \{\infty\} : \mathbf{P}\left(S_2^{G_t^{(n,m)\downarrow}} \geqslant M\right) \stackrel{(4.16), (4.19)}{\leqslant} 2\varepsilon. \quad (4.20)$$

We are ready to show (4.13) and (4.15) for the above choice of m and n_0 . For any $n \in \{n_0, n_0 + 1, ...\} \cup \{\infty\}$ we have

$$\mathbf{P}\left(\mathbf{d}(\mathbf{m}_{t}^{(n)}, \mathbf{m}_{t}^{(n,m)}) \geqslant \varepsilon\right) \leqslant \mathbf{P}\left(S_{2}^{G_{t}^{(n,m)\uparrow}} \geqslant \delta\right) + \mathbf{P}\left(S_{2}^{G_{t}^{(n,m)\downarrow}} \geqslant M\right) + \mathbf{P}\left(\mathbf{d}(\mathbf{m}_{t}^{(n)}, \mathbf{m}_{t}^{(n,m)}) \geqslant \varepsilon, S_{2}^{G_{t}^{(n,m)\uparrow}} \leqslant \delta, S_{2}^{G_{t}^{(n,m)\downarrow}} \leqslant M\right) \stackrel{(4.18),(4.20)}{\leqslant} \varepsilon + 2\varepsilon + \mathbf{P}\left(\mathbf{d}(\mathbf{m}_{t}^{(n)}, \mathbf{m}_{t}^{(n,m)}) \geqslant \varepsilon, A\right),$$

where $A = \{S_2^{G_t^{(n,m)\uparrow}} \leq \delta, S_2^{G_t^{(n,m)\downarrow}} \leq M\}$. We bound

$$\mathbf{P}\left(\mathbf{d}(\mathbf{m}_{t}^{(n)}, \mathbf{m}_{t}^{(n,m)}) \geqslant \varepsilon, A\right) \overset{(4.3)}{\leqslant} \mathbf{P}\left(9 \cdot \left(S_{2}^{\check{G}^{(n,m)}} - S_{2}^{\hat{G}^{(n,m)}}\right) \geqslant \varepsilon^{2}, A\right) = \mathbf{E}\left(\mathbf{P}\left(9 \cdot \left(S_{2}^{\check{G}^{(n,m)}} - S_{2}^{\hat{G}^{(n,m)}}\right) \geqslant \varepsilon^{2} \mid G^{(n,m)\downarrow}, G^{(n,m)\uparrow}\right); A\right)$$

$$\overset{(*)}{\leqslant} \frac{9C \cdot \delta \cdot \left((1 + tM)^{2} + (1 + tM) \cdot M^{3/2}\right)}{\varepsilon^{2}} \overset{(4.17)}{\leqslant} \varepsilon,$$

where in the equation marked by (*) we used Lemma 4.10 and the Markov inequality. This concludes the proof of (4.13),(4.14),(4.15) and Theorem 4.2.

Acknowledgements: I thank James Martin for collaborating with me on [19], which inspired this work. I also thank an anonymous referee for useful comments on the manuscript. This work is partially supported by OTKA (Hungarian National Research Fund) grant K100473, the Postdoctoral Fellowship of the Hungarian Academy of Sciences and the Bolyai Research Scholarship of the Hungarian Academy of Sciences.

References

- [1] L. Addario-Berry, N. Broutin and C. Goldschmidt. The continuum limit of critical random graphs. *Probab. Theo. and Rel. Fields* 152.3-4, 367-406, 2012.
- [2] D. Ahlberg, H. Duminil-Copin, G. Kozma and V. Sidoravicius. Seven-dimensional forest fires. Ann. de l'Institut Henri Poincaré, Probabilités et Statistiques, 51(3), 862-866, 2015.
- [3] D.J. Aldous. Brownian Excursions, Critical Random Graphs and the Multiplicative Coalescent. *Annals of Probability* **25**: 812–854, 1997.
- [4] D.J. Aldous and V. Limic. The entrance boundary of the multiplicative coalescent. *Electronic Journal of Probability* 3, paper 3, 59 pp. (1998)
- [5] D. J. Aldous. The percolation process on a tree where infinite clusters are frozen. *Math. Proc. Cambridge Philos. Soc.*, 128(3):465–477, 2000.
- [6] J. van den Berg, D. Kiss and P. Nolin. Two-dimensional volume-frozen percolation: deconcentration and prevalence of mesoscopic clusters. arXiv:1512.05335, 2015.
- [7] J. van den Berg, B. N. B. de Lima, P. Nolin. A percolation process on the square lattice where large finite clusters are frozen. Rand. Struct. and Alg. 40.2, 220-226, 2012.
- [8] J. van den Berg, P. Nolin. Two-dimensional volume-frozen percolation: exceptional scales. (to appear in Ann. Appl. Probab.), arXiv:1501.05011, 2015.
- [9] J. van den Berg, B. Tóth. A signal-recovery system: asymptotic properties, and construction of an infinite-volume process. Stoch. Proc. and Appl. 96.2, 177-190, 2001.

- [10] S. Bhamidi, R. van der Hofstad, and J. van Leeuwaarden. Scaling limits for critical inhomogeneous random graphs with finite third moments. Electron. J. Probab. **15**(54), 1682-1702, 2010.
- [11] S. Bhamidi, R. van der Hofstad, and J. van Leeuwaarden. Novel scaling limits for critical inhomogeneous random graphs. *Ann. of Probab.* 40.6, 2299-2361, 2012.
- [12] S. Bhamidi, N. Broutin, S. Sen, and X. Wang. Scaling limits of random graph models at criticality: Universality and the basin of attraction of the Erdős-Rényi random graph. arXiv:1411.3417, 2014.
- [13] S. Bhamidi, R. van der Hofstad, S. Sen. The multiplicative coalescent, inhomogeneous continuum random trees, and new universality classes for critical random graphs. arXiv:1508.04645, 2015.
- [14] M. Dürre. Existence of multi-dimensional infinite volume self-organized critical forest-fire models. Electr. J. of Probab. 11.513-539, (2006): 2242654.
- [15] M. Dürre. Uniqueness of multi-dimensional infinite-volume self-organized critical forest fire models. Electr. Comm. in Probab. 11.304-315, (2006): 2266720.
- [16] D. Kiss. Frozen percolation in two dimensions. Prob. The. and Rel. Fields. 163.3-4: 713-768, 2015.
- [17] D. Kiss, I. Manolescu and V. Sidoravicius. *Planar lattices do not recover from forest fires*. Ann. of Probab. 43.6: 3216-3238, 2015.
- [18] V. Limic. Properties of the Multiplicative Coalescent. *PhD thesis*, UC Berkeley, 1998.
- [19] J. B. Martin, B. Ráth. Rigid representations of the multiplicative coalescent with linear deletion. *Electron. J. Probab.* Vol. 22, paper no. 83, 2017.
- [20] J.B. Martin, B. Ráth. Scaling limits of mean-field frozen percolation and forest fire processes (in preparation)
- [21] M. Merle, R. Normand. Self-organized criticality in a discrete model for Smoluchowski's equation. arXiv:1410.8338, 2014.
- [22] B. Ráth. Mean field frozen percolation. J. of Stat. Phys. **137(3)**: 459-499, 2009.

[23] B. Ráth, B. Tóth. Erdős-Rényi Random graphs + forest fires = Self-Organized Criticality Electr. J. of Probab. 14:1290-1327, 2009.