THE FORMAL THEORY OF MULTIMONOIDAL MONADS

GABRIELLA BÖHM

ABSTRACT. Certain aspects of Street's formal theory of monads in 2-categories are extended to multimonoidal monads in symmetric strict monoidal 2-categories. Namely, any symmetric strict monoidal 2-category \mathcal{M} admits a symmetric strict monoidal 2-category of pseudomonoids, monoidal 1-cells and monoidal 2-cells in \mathcal{M} . Dually, there is a symmetric strict monoidal 2-category of pseudomonoids, opmonoidal 1-cells and opmonoidal 2-cells in \mathcal{M} . Extending a construction due to Aguiar and Mahajan for $\mathcal{M} = \mathsf{Cat}$, we may apply the first construction *p*-times and the second one q-times (in any order). It yields a 2-category \mathcal{M}_{pq} . A 0-cell therein is an object A of \mathcal{M} together with p + q compatible pseudomonoid structures; it is termed a (p+q)-oidal object in \mathcal{M} . A monad in \mathcal{M}_{pq} is called a (p,q)-oidal monad in \mathcal{M} ; it is a monad t on A in \mathcal{M} together with p monoidal, and q opmonoidal structures in a compatible way. If \mathcal{M} has monoidal Eilenberg-Moore construction, and certain (Linton type) stable coequalizers exist, then a (p+q)-oidal structure on the Eilenberg-Moore object A^t of a (p,q)-oidal monad (A,t) is shown to arise via a symmetric strict monoidal double functor to Ehresmann's double category $Sqr(\mathcal{M})$ of squares in \mathcal{M} , from the double category of monads in $Sqr(\mathcal{M})$ in the sense of Fiore, Gambino and Kock. While q ones of the pseudomonoid structures of A^t are lifted along the 'forgetful' 1-cell $A^t \to A$, the other p ones are lifted along its left adjoint. In the particular example when \mathcal{M} is an appropriate 2-subcategory of Cat, this yields a conceptually different proof of some recent results due to Aguiar, Haim and López Franco.

INTRODUCTION

Classically, a *monad* on a category A is a monoid in the category of endofunctors on A; that is, a functor $t: A \to A$ together with natural transformations from the twofold iterate t.t and from the identity functor, respectively, to t, regarded as an associative multiplication with a unit. A popular example is the monad $T \otimes -$ induced by an associative unital algebra T on the category of vector spaces.

Any adjunction $l \dashv r : B \to A$ induces a monad r.l on A, with unit provided by the unit of the adjunction, and multiplication induced by the counit. Conversely, any monad is induced by some adjunction in this sense. There is no unique adjunction in general, but a terminal one can easily be described. An *Eilenberg-Moore algebra* of a monad t on a category A consists of an object X of A and an associative and unital action $tX \to X$. A morphism of Eilenberg-Moore algebras is a morphism in Awhich commutes with the actions. (For the monad $T \otimes -$ induced by an algebra Ton the category of vector spaces, these are just left T-modules and their morphisms.) The evident forgetful functor u^t (forgetting the actions) from the so defined *Eilenberg-Moore category* A^t to A has a left adjoint (sending an object X to tX with action provided by the multiplication of the monad) and this adjunction induces the monad

Date: April 2019.

t. Moreover, any other adjunction $l \dashv r : B \to A$ inducing the same monad factorizes through a unique functor $B \to A^t$.

Having functors $f: A \to B$ between the base categories of some respective monads t and s, and natural transformations between them, it is often a relevant question if they lift to the Eilenberg-Moore categories in the sense of the commutative diagram

where u^t and u^s are the respective forgetful functors. It is not hard to see that such liftings of a functor f correspond bijectively to natural transformations $\varphi : s.f \to f.t$ which are compatible with both monad structures. Such a pair (f, φ) is called a *monad functor*. A natural transformation between functors $A \to B$ has at most one lifting to a natural transformation between the lifted functors, and the condition for its existence is a compatibility with the natural transformations φ . A natural transformation satisfying this compatibility condition is termed a *monad transformation*.

The formal theory of monads [28] due to Ross Street provides a wide generalization of the above picture and gives it a conceptual interpretation. The 2-category **Cat** of categories, functors and natural transformations is replaced by an arbitrary 2category \mathcal{M} . A monad in \mathcal{M} on a 0-cell A is defined as a monoid in the hom category $\mathcal{M}(A, A)$. Monads are the 0-cells in a 2-category $\mathsf{Mnd}(\mathcal{M})$ whose 1-cells and 2-cells are the analogues of monad functors, and monad transformations, respectively. Regarding any 0-cell of \mathcal{M} as a trivial monad (with identity 1, and 2-cell parts), regarding any 1-cell of \mathcal{M} as their monad morphism (with identity 2-cell part), and regarding any 2cell of \mathcal{M} as a monad transformation, there is an inclusion 2-functor $\mathcal{M} \to \mathsf{Mnd}(\mathcal{M})$. For $\mathcal{M} = \mathsf{Cat}$ it has a right 2-adjoint. The right 2-adjoint sends a monad to its Eilenberg-Moore category, a monad functor to the corresponding lifted functor, and a monad transformation to the lifted natural transformation. When for some 2-category \mathcal{M} the above inclusion 2-functor $\mathcal{M} \to \mathsf{Mnd}(\mathcal{M})$ possesses a right 2-adjoint, \mathcal{M} is said to admit Eilenberg-Moore construction. By the theory worked out in [28], the right adjoint describes an analogous lifting theory. (More will be said in Section 3.)

For a monad t on a monoidal category A, in [25] the additional structure was described which is equivalent to a monoidal structure on the Eilenberg-Moore category A^t such that the forgetful functor $A^t \to A$ is strict monoidal. The explicit computations of [25] in **Cat** (considered with the Cartesian product of categories as the monoidal structure) were replaced in [24], [9] by abstract arguments about more general monoidal bicategories. Beyond a wide generalization, thereby also a conceptually different proof was obtained. Namely, the structure described in [25] was interpreted as an opmonoidal monad; that is, a monad in the 2-category **Cat**₀₁ of monoidal categories, opmonoidal functors and opmonoidal natural transformations. Now **Cat**₀₁ admits Eilenberg-Moore construction in the sense of [28], see [22] and [31] for conceptually different proofs. Hence there is a 2-functor from Street's 2-category of monoidal **Mnd**(**Cat**₀₁) above to **Cat**₀₁ whose object map sends an opmonoidal monad to its monoidal Eilenberg-Moore category. Recently in [1] a similar analysis to that of [25] was carried out for multimonoidal monads on multimonoidal categories. (In [1] the term higher monoidal was used. However, we prefer to call the same thing multimonoidal and reserve the term higher to be used only for dimensionality of categorical structures.) The aim of this paper is to extend the results of [1] to multimonoidal monads in symmetric strict monoidal 2-categories and, more importantly, derive them from a suitable 'formal theory'. With this we not only gain a deeper understanding of the origin of the formulae in [1] but, as a byproduct, also open a way to some practical applications. For example, we obtain sufficient conditions under which the 2-category of pseudomonoids, monoidal 1-cells and monoidal 2-cells in a strict monoidal 2-category admits Eilenberg–Moore construction.

The development of this formal theory will require a move away from 2-categories to symmetric strict monoidal double categories. By a strict monoidal 2-category we mean a monoid in the category of 2-categories and 2-functors considered with the Cartesian product of 2-categories as the monoidal product. This definition occurred e.g. on page 69 of [20] where also the notion of symmetry was introduced as a suitable 2-natural isomorphism between the monoidal product 2-functor and its reversed mate. Similarly restrictively, we adopt the definition of symmetric strict monoidal double category in [8]. Here again, a strict monoidal double category means a monoid in the category of double categories as the monoidal product. A symmetry in [8] can be interpreted then as a suitable double natural isomorphism between the monoidal product. A symmetry in [12] to yield a symmetric strict monoidal double category in this sense, we need to apply it to a symmetric strict monoidal 2-category.)

The notions recalled in the previous paragraph are very restrictive (by being so strict). One can ask about various levels of generalization whether they are possible. Instead of monoids, one may consider pseudomonoids in the 2-category of 2-categories, 2-functors and 2-natural transformations; and correspondingly, pseudomonoids in the 2-category of double categories, double functors and double natural transformations — considered in both cases with the monoidal structure provided by the Cartesian product. Although we expect that it should be possible, it is not motivated by our examples. Also, the technical complexity resulting from the tedious checking of all coherence conditions could divert attention from the key ideas. It looks more challenging to extend our considerations to monoidal bicategories in the sense of [21, 3, 26] — or at least to their semistrict version known as *Gray monoids* [16, 10]. These are monoids in the category of 2-categories and 2-functors considered with the monoidal structure provided by the *Gray tensor product* [19]. Since the corresponding Gray tensor product of double categories seems not yet available in the literature, this problem does not look to be within reach. We plan to address it elsewhere [5].

For any symmetric strict monoidal 2-category \mathcal{M} , there is a symmetric strict monoidal 2-category \mathcal{M}_{10} whose 0-cells are the pseudomonoids (also called *monoidal objects* e.g. in [31] or *monoidales* e.g. in [9]), the 1-cells are the monoidal 1-cells, and the 2-cells are the monoidal 2-cells in \mathcal{M} . Symmetrically, there is a symmetric strict monoidal 2-category \mathcal{M}_{01} whose 0-cells are again the pseudomonoids, but the 1-cells are the opmonoidal 1-cells, and the 2-cells are the opmonoidal 2-cells in \mathcal{M} . Moreover, these constructions commute with each other (see Section 1). Thus applying p

times the first construction and q times the second one (in an arbitrary order), we get a 2-category \mathcal{M}_{pq} . For a fixed non-negative integer n and every $0 \leq p \leq n$, the 0-cells of $\mathcal{M}_{p,n-p}$ are the same gadgets. They consist of a 0-cell of \mathcal{M} together with n pseudomonoid structures and compatibility morphisms between them (constituting suitable monoidal structures on the structure morphisms of the pseudomonoids). We call a 0-cell of $\mathcal{M}_{p,n-p}$ an *n*-oidal object of \mathcal{M} . (In the particular case of $\mathcal{M} = \mathsf{Cat}$, in [1, 2] it was called an *n*-monoidal category.) A 1-oidal object is a pseudomonoid, in particular a monoidal category for $\mathcal{M} = \mathsf{Cat}$. So we re-obtain the classical terminology if "1" is pronounced as "mono". A 2-oidal object of Cat is a duoidal category (in the sense of [29], termed 2-monoidal in [2]). Again we agree with the established terminology if "2" is pronounced as "duo". A 1-cell in \mathcal{M}_{pq} consists of a 1-cell of \mathcal{M} together with monoidal structures with respect to p ones of the pseudomonoid structures; and opmonoidal structures with respect to the remaining q ones of the pseudomonoid structures of the domain and the codomain. They are subject to suitable compatibility conditions. We term a 1-cell of \mathcal{M}_{pq} a (p,q)-oidal 1-cell in \mathcal{M} (rather than (p, q)-monoidal as in [1, 2]; where in particular a (2, 0)-oidal 1-cell of Cat was called a *double monoidal functor*, a (0,2)-oidal 1-cell was called *double comonoidal* and a (1,1)-oidal 1-cell was called *bimonoidal*). A 2-cell in \mathcal{M}_{pq} — called a (p,q)-oidal 2-cell in \mathcal{M} — is a 2-cell in \mathcal{M} which is compatible with all of the (op)monoidal structures of the domain and the codomain. A monad in \mathcal{M}_{pq} is termed a (p,q)-oidal monad.

As recalled above from [24], the monoidal structure of the base category A of any opmonoidal (or (0, 1)-oidal) monad t in Cat lifts to the Eilenberg-Moore category A^t along the forgetful functor $u^t: A^t \to A$. (This means that the functors and natural transformations constituting the monoidal structures of A and A^t fit in commutative diagrams as in (EM)). Furthermore, if reflexive coequalizers exist in A and they are preserved by the monoidal product of A and by the functor t, then also the monoidal structure of the base category A of a monoidal (or (1,0)-oidal) monad t lifts to the Eilenberg-Moore category A^t . However, at this time it is a lifting along the left adjoint f^t of the forgetful functor $u^t: A^t \to A$. This means that the functors and natural transformations constituting the monoidal structures of A and A^t fit in commutative diagrams obtained from that in (EM) replacing the forgetful functors with their left adjoints in the opposite direction. This is a result of [27]; see also [1]. In [1] it was proven, moreover, that under the same assumptions also the (p+q)-oidal structure of the base category A of any (p,q)-oidal monad t in Cat lifts to the Eilenberg-Moore category A^t . This is a lifting of mixed kind, though. While q ones of the monoidal structures are lifted along the forgetful functor $u^t: A^t \to A$; the remaining p ones are lifted along its left adjoint f^t .

Because of this mixed nature of lifting; that is, since the different ingredients are lifted along different functors u^t and f^t , we do not expect it to be described by some 2-functor (as in the situations of [28] and [9]). Instead, in this paper we deal with symmetric strict monoidal double categories and define their (p, q)-oidal objects (see Section 6). These (p, q)-oidal objects are shown to be preserved by symmetric strict monoidal double functors. In Ehresmann's double category Sqr (\mathcal{M}) of squares (or quintets [12]) in a symmetric strict monoidal 2-category \mathcal{M} , the (p, q)-oidal objects are the same as the (p + q)-oidal objects in \mathcal{M} . Taking the double category of monads [14] in the particular double category $Sqr(\mathcal{M})$, we obtain a symmetric strict monoidal double category which we denote by $Mnd(\mathcal{M})$. Its horizontal 2-category is $Mnd(\mathcal{M})$ and its vertical 2-category is $Mnd(\mathcal{M}_{op})_{op}$ (where $(-)_{op}$ refers to the horizontally opposite 2-category, see [28, Section 4]). The (p, q)oidal objects in $Mnd(\mathcal{M})$ are identified with the (p, q)-oidal monads in \mathcal{M} (see Section 7). Consequently, any symmetric strict monoidal double functor $Mnd(\mathcal{M}) \to Sqr(\mathcal{M})$ takes (p, q)-oidal monads in \mathcal{M} to (p + q)-oidal objects in \mathcal{M} .

Whenever a symmetric strict monoidal 2-category \mathcal{M} admits monoidal Eilenberg-Moore construction (in the sense that the comparison 1-cells $I^1 \to I$ and $A^t B^s \to (AB)^{ts}$ are identities for the identity monad on the monoidal unit I and all monads t on A and s on B, see Definition 3.1) and furthermore some Linton type coequalizers (see (5.1)) exist and are preserved by the horizontal composition, we construct a symmetric strict monoidal double functor $\operatorname{Mnd}(\mathcal{M}) \to \operatorname{Sqr}(\mathcal{M})$ (see Section 8). In the terminology of [15] it provides Eilenberg-Moore construction for the double category $\operatorname{Sqr}(\mathcal{M})$. Its horizontal 2-functor is the Eilenberg-Moore 2-functor $\operatorname{Mnd}(\mathcal{M}) \to \mathcal{M}$ and its vertical 2-functor $\operatorname{Mnd}(\mathcal{M}_{op})_{op} \to \mathcal{M}$ is obtained via Linton type coequalizers. Its taking (p, q)-oidal monads in \mathcal{M} to their (p + q)-oidal Eilenberg-Moore objects provides a 'formal theory' in the background of the liftings in [1].

Acknowledgement. It is a pleasure to thank Ross Street and two anonymous referees for helpful comments, insightful questions and mentioning some relevant references. Financial support by the Hungarian National Research, Development and Innovation Office NKFIH (grant K124138) is gratefully acknowledged.

1. Multimonoidal structures in 2-categories

We begin with a brief review of some definitions and basic constructions for later use. A more detailed introduction can be found e.g. in Chapter 7 of [6].

A 2-category is a category enriched in the category of small categories and functors considered with the Cartesian product of categories as the monoidal product. The explicit definition can be found in [6, Definition 7.1.1]. Throughout, horizontal composition in a 2-category (i.e. the composition functor) will be denoted by a lower dot . and vertical composition (that is, the compositions of the hom categories) will be denoted by an upper dot \cdot . All identity (1- or 2-) cells will be denoted by 1.

Example 1.1. The most well-known 2-category is, perhaps, Cat [6, Example 7.1.4.a]. Its 0-cells (or objects) are small categories, the 1-cells (i.e. objects of the hom categories) are the functors and the 2-cells (i.e. morphisms of the hom categories) are the natural transformations.

For any 2-category \mathcal{M} we denote the *horizontally opposite* 2-category in [28, Section 4] by \mathcal{M}_{op} and we denote the *vertically opposite* 2-category in [28, Section 4] by \mathcal{M}^{\diamond} .

A 2-functor is a functor enriched in the category of small categories and functors considered with the Cartesian product of categories as the monoidal product. The explicit definition can be found in [6, Definition 7.2.1]. Any 2-functor $F : \mathcal{M} \to \mathcal{N}$ can be seen as a 2-functor $F_{op} : \mathcal{M}_{op} \to \mathcal{N}_{op}$ and also as a 2-functor $F^{\mathfrak{d}} : \mathcal{M}^{\mathfrak{d}} \to \mathcal{N}^{\mathfrak{d}}$.

A 2-natural transformation is a natural transformation enriched in the category of small categories and functors considered with the Cartesian product of categories as the monoidal product. The explicit definition can be found in [6, Definition 7.2.2].

Any 2-natural transformation $\Theta : F \to G$ can be seen as a 2-natural transformation $\Theta_{op} : F_{op} \to G_{op}$ and also as a 2-natural transformation $\Theta^{\circ} : F^{\circ} \to G^{\circ}$.

Example 1.2. Small 2-categories are the 0-cells, 2-functors are the 1-cells, and 2-natural transformations are the 2-cells of a 2-category 2Cat, see [6, Proposition 7.2.3].

Definition 1.3. [20, page 69] A *strict monoidal 2-category* is a monoid in the category whose objects are the 2-categories, whose morphisms are the 2-functors and whose monoidal product is the Cartesian product of 2-categories. Explicitly, a strict monoidal 2-category consists of the following data.

- A 2-category \mathcal{M} .
- A 2-functor I from the singleton 2-category to \mathcal{M} . The image of the only object of the singleton category under it is called the *monoidal unit* and it is denoted by the same symbol I.
- A 2-functor $\otimes : \mathcal{M} \times \mathcal{M} \to \mathcal{M}$ called the *monoidal product*. It must be strictly associative with the strict unit *I*. The action of the 2-functor \otimes on (0-, 1- and 2-) cells will be denoted by juxtaposition.

The 2-category Cat of Example 1.1 is strict monoidal via the Cartesian product of categories.

Definition 1.4. A strict monoidal 2-functor is a monoid morphism in the category whose objects are the 2-categories, whose morphisms are the 2-functors and whose monoidal product is the Cartesian product of 2-categories. Explicitly, a 2-functor $F : \mathcal{M} \to \mathcal{M}'$ is strict monoidal whenever FI = I' and $F \otimes = \otimes' \cdot (F \times F)$.


Definition 1.5. A 2-natural transformation $\Theta : \mathsf{F} \to \mathsf{G}$ between strict monoidal 2-functors is said to be *monoidal* if the following diagrams of 2-natural transformations commute.

$$\begin{array}{c} \otimes.(\mathsf{F}\times\mathsf{F}) = & \mathsf{F}.\otimes & I = & \mathsf{F}.I \\ 1.(\Theta\times\Theta) \downarrow & \downarrow \Theta.1 & \downarrow & \downarrow \Theta.1 \\ \otimes.(\mathsf{G}\times\mathsf{G}) = & \mathsf{G}.\otimes & I = & \mathsf{G}.I \end{array}$$

That is, for all objects A and B, $\Theta_{AB} = \Theta_A \Theta_B$ and $\Theta_I = 1$.

For any 2-category \mathcal{M} , there is a 2-functor flip : $\mathcal{M} \times \mathcal{M} \to \mathcal{M} \times \mathcal{M}$, sending a pair of 2-cells (ω, ϑ) to (ϑ, ω) . It occurs in the following definition.

Definition 1.6. [20, Page 69] A symmetric strict monoidal 2-category consists of a strict monoidal 2-category $(\mathcal{M}, \otimes, I)$ together with a 2-natural transformation σ : $\otimes \to \otimes$.flip rendering commutative the following diagrams for all objects A, B and C.



The subscripts p and q of σ_{pq} indicate that it is a morphism $A_1 \dots A_p B_1 \dots B_q \rightarrow B_1 \dots B_q A_1 \dots A_p$.

Definition 1.7. A strict monoidal 2-functor F between symmetric strict monoidal 2-categories is said to be *symmetric* if $\mathsf{F}(AB \xrightarrow{\sigma} BA) = (\mathsf{F}A)(\mathsf{F}B) \xrightarrow{\sigma'} (\mathsf{F}B)(\mathsf{F}A)$ for all objects A and B of the domain 2-category of F .

Example 1.8. Symmetric strict monoidal small 2-categories are the 0-cells, symmetric strict monoidal 2-functors are the 1-cells and monoidal 2-natural transformations are the 2-cells of a 2-category sm-2Cat. Both vertical and horizontal compositions are given by the same formulae as in 2Cat of [6, Proposition 7.2.3].

Example 1.9. Below we describe a 2-functor $\mathsf{Mnd} : \mathsf{sm-2Cat} \to \mathsf{sm-2Cat}$.

For any 2-category \mathcal{M} , there is a 2-category $\mathsf{Mnd}(\mathcal{M})$ as in [28]. Its 0-cells are the monads in \mathcal{M} . That is, quadruples consisting of a 0-cell A of \mathcal{M} , a 1-cell $t: A \to A$ and 2-cells $\mu : t.t \to t$ and $\eta : 1 \to t$. They are required to satisfy the associativity condition $\mu^{\cdot}(\mu,1) = \mu^{\cdot}(1,\mu)$ and the unitality conditions $\mu^{\cdot}(\eta,1) = 1 = \mu^{\cdot}(1,\eta)$. A 1-cell in $\mathsf{Mnd}(\mathcal{M})$ from (A, t, μ, η) to (A', t', μ', η') is a pair consisting of a 1-cell $f: A \to A'$ in \mathcal{M} and a 2-cell $\varphi: t' f \to f t$ satisfying the multiplicativity condition $(1,\mu)^{\cdot}(\varphi,1)^{\cdot}(1,\varphi) = \varphi^{\cdot}(\mu',1)$ and the unitality condition $1,\eta = \varphi^{\cdot}(\eta',1)$. Such a 1-cell is called a monad morphism in \mathcal{M} . The 2-cells $(f, \varphi) \to (g, \gamma)$ in $\mathsf{Mnd}(\mathcal{M})$ are those 2-cells $\omega : f \to g$ in \mathcal{M} for which $(\omega.1)^{\cdot}\varphi = \gamma^{\cdot}(1.\omega)$. They are called monad transformations in \mathcal{M} . The vertical composite of monad transformations is their vertical composite as 2-cells in \mathcal{M} . The horizontal composite of composable 1-cells (f, φ) and (q, γ) is $(q, f, (1, \varphi); (\gamma, 1))$. The horizontal composite of monad transformations is their horizontal composite as 2-cells in \mathcal{M} . Whenever \mathcal{M} is equipped with a strict monoidal structure (\otimes , I), there is an induced strict monoidal structure on $\mathsf{Mnd}(\mathcal{M})$. The monoidal unit is (I, 1, 1, 1), the identity monad on I. The monoidal product of monads (A, t, μ, η) and (A', t', μ', η') is

$$(AA', AA' \xrightarrow{tt'} AA', tt'.tt' = (t.t)(t.'t') \xrightarrow{\mu\mu'} tt', I = II \xrightarrow{\eta\eta'} tt').$$

The monoidal product of monad morphisms $(f, \varphi) : (A, t) \to (B, s)$ and $(f', \varphi') : (A', t') \to (B', s')$ is the monad morphism

$$(AA' \xrightarrow{ff'} BB', ss'.ff' = (s.f)(s.'f') \xrightarrow{\varphi\varphi'} (f.t)(f'.t') = ff'.tt').$$

The monoidal product of monad transformations is their monoidal product as 2cells of \mathcal{M} . If in addition a strict monoidal 2-category $(\mathcal{M}, \otimes, I)$ is equipped with a symmetry σ , then the induced symmetry on $\mathsf{Mnd}(\mathcal{M})$ has the components $(\sigma, 1)$. So far we recalled the 0-cell part of the desired 2-functor Mnd .

For any 2-functor $\mathsf{F} : \mathcal{M} \to \mathcal{N}$ there is a 2-functor $\mathsf{Mnd}(\mathsf{F}) : \mathsf{Mnd}(\mathcal{M}) \to \mathsf{Mnd}(\mathcal{N})$. It sends a monad (A, t, μ, η) to

$$(FA, FA \xrightarrow{\mathsf{F}t} \mathsf{F}A, (\mathsf{F}t).(\mathsf{F}t) = \mathsf{F}(t.t) \xrightarrow{\mathsf{F}\mu} \mathsf{F}t, I = \mathsf{F}I \xrightarrow{\mathsf{F}\eta} \mathsf{F}t);$$

it sends a monad morphism $(f, \varphi) : (A, t) \to (B, s)$ to

$$(\mathsf{F}A \xrightarrow{\mathsf{F}f} \mathsf{F}B, (\mathsf{F}s).(\mathsf{F}f) = \mathsf{F}(s.f) \xrightarrow{\mathsf{F}\varphi} \mathsf{F}(f.t) = (\mathsf{F}f).(\mathsf{F}t)$$

and it sends a monad transformation ω to $F\omega$. Whenever F is strict monoidal then so is Mnd(F) and if in addition F is symmetric then so is Mnd(F). This yields the 1-cell part of the desired 2-functor Mnd.

For any 2-natural transformation Θ there is a 2-natural transformation $\mathsf{Mnd}(\Theta)$ with components $(\Theta, 1)$. It is strict monoidal whenever Θ is so. This finishes the description of the stated 2-functor Mnd. (It would be straightforward to extend it also to modifications but it is not needed for the purposes of the present paper.)

Symmetrically to the above considerations, we introduce another 2-functor $\mathsf{Mnd}_{op} := \mathsf{Mnd}((-)_{op})_{op} : \mathsf{sm-2Cat} \to \mathsf{sm-2Cat}.$

Example 1.10. (1) Below a 2-functor $(-)_{01}$: sm-2Cat \rightarrow sm-2Cat is described.

For any strict monoidal 2-category $(\mathcal{M}, \otimes, I)$, there is a 2-category \mathcal{M}_{01} . Its 0-cells are the *pseudomonoids* in \mathcal{M} , also called *monoidal objects* or *monoidales* by other authors, see [31] and [9], respectively. A pseudomonoid in \mathcal{M} consists of a 0-cell A, 1-cells $m : AA \to A$ and $u : I \to A$ together with invertible 2-cells $\alpha : m.m1 \to m.1m$, $\lambda : m.u1 \to 1$ and $\rho : m.1u \to 1$ satisfying MacLane's pentagon and triangle conditions

$$\begin{array}{c|c} m.m1.m11 \xrightarrow{\alpha.1} m.1m.m11 = m.m1.11m \\ 1.\alpha 1 \\ m.m1.1m1 \xrightarrow{\alpha.1} m.1m.1m1 \xrightarrow{1.1\alpha} m.1m.11m \end{array} \qquad \begin{array}{c} m.m1.1u1 \xrightarrow{\alpha.1} m.1m.1u1 \\ 1.\varrho 1 \\ m.m1.1m1 \xrightarrow{\alpha.1} m.1m.1m1 \xrightarrow{1.1\alpha} m.1m.11m \end{array}$$

A 1-cell in \mathcal{M}_{01} is a so-called *opmonoidal 1-cell*. An opmonoidal 1-cell from $(A, m, u, \alpha, \lambda, \varrho)$ to $(A', m', u', \alpha', \lambda', \varrho')$ consists of a 1-cell $f : A \to A'$ in \mathcal{M} together with 2-cells $\varphi^2 : f.m \to m'.ff$ (called the *binary part*) and $\varphi^0 : f.u \to u'$ (called the *nullary part*) subject to the coassociativity and counitality conditions

The 2-cells $\omega : (f, \varphi^2, \varphi^0) \to (g, \gamma^2, \gamma^0)$ in \mathcal{M}_{01} are the *opmonoidal 2-cells* in \mathcal{M} ; that is, those 2-cells $\omega : f \to g$ in \mathcal{M} for which the following diagrams commute.

The vertical composite of opmonoidal 2-cells is their vertical composite as 2-cells in \mathcal{M} . The horizontal composite of 1-cells $(f, \varphi^2, \varphi^0) : (A, m, u, \alpha, \lambda, \varrho) \to (A', m', u', \alpha', \lambda', \varrho')$ and $(g, \gamma^2, \gamma^0) : (A', m', u', \alpha', \lambda', \varrho') \to (A'', m'', u'', \alpha'', \lambda'', \varrho'')$ is

$$(g.f, \ g.f.m \xrightarrow{1.\varphi^2} g.m'.ff \xrightarrow{\gamma^2.1} m''.gg.ff = m''.(g.f)(g.f) \ , \ g.f.u \xrightarrow{1.\varphi^0} g.u' \xrightarrow{\gamma^0} u'' \)$$

The horizontal composite of opmonoidal 2-cells is their horizontal composite as 2-cells in \mathcal{M} .

If the strict monoidal 2-category $(\mathcal{M}, \otimes, I)$ is equipped with a symmetry σ , then there is an induced symmetric strict monoidal structure on \mathcal{M}_{01} . The monoidal product of pseudomonoids $(A, m, u, \alpha, \lambda, \varrho)$ and $(A', m', u', \alpha', \lambda', \varrho')$ is

$$(AA', AA'AA' \xrightarrow{1\sigma_1} AAA'A' \xrightarrow{mm'} AA', I = II \xrightarrow{uu'} AA',$$
$$mm'.1\sigma_1.mm'_{11.1\sigma_{111}} \qquad mm'.1\sigma_{1.11}mm'_{.111\sigma_{111}} \\ \square \\ mm'.m_1m'_{1.11\sigma_{21}} 1.1\sigma_{111} \xrightarrow{\alpha\alpha'.1.1} mm'.1m_1m'_{.1\sigma_{12}} 11.111\sigma_1,$$
$$mm'.1\sigma_{1.uu'_{11}} = mm'.u_1u'_{11} \xrightarrow{\lambda\lambda'} 1, mm'.1\sigma_{1.11uu'} = mm'.1u_1u' \xrightarrow{\varrho\varrho'} 1).$$

The monoidal product of 1-cells $(f, \varphi^2, \varphi^0) : (A, m, u, \alpha, \lambda, \varrho) \to (A', m', u', \alpha', \lambda', \varrho')$ and $(g, \gamma^2, \gamma^0) : (B, m, u, \alpha, \lambda, \varrho) \to (B', m', u', \alpha', \lambda', \varrho')$ is

$$(AB \xrightarrow{fg} A'B', fg.mm.1\sigma1 \xrightarrow{\varphi^2 \gamma^2.1} m'm'.ffgg.1\sigma1 = m'm'.1\sigma1.fgfg, fg.uu \xrightarrow{\varphi^0 \gamma^0} u'u').$$

The monoidal product of opmonoidal 2-cells is their monoidal product as 2-cells in \mathcal{M} . The symmetry has the components $(\sigma, 1, 1)$. This gives the object map of a 2-functor $(-)_{01}$: sm-2Cat \rightarrow sm-2Cat.

A strict monoidal 2-functor $\mathsf{F} : \mathcal{M} \to \mathcal{N}$ induces a 2-functor $\mathsf{F}_{01} : \mathcal{M}_{01} \to \mathcal{N}_{01}$. It takes a 0-cell of \mathcal{M}_{01} — that is, a pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$ in \mathcal{M} — to the pseudomonoid in \mathcal{N} ,

$$(FA, (FA)(FA) = F(AA) \xrightarrow{Fm} FA, I = FI \xrightarrow{Fu} FA,$$

$$Fm.(Fm)1 = F(m.m1) \xrightarrow{F\alpha} F(m.1m) = Fm.1(Fm),$$

$$Fm.(Fu)1 = F(m.u1) \xrightarrow{F\lambda} F1 = 1, Fm.1(Fu) = F(m.1u) \xrightarrow{F\varrho} F1 = 1).$$

The 2-functor F_{01} sends an opmonoidal 1-cell $(f, \varphi^2, \varphi^0)$ to

$$(\mathsf{F}f, \ \mathsf{F}m'.(\mathsf{F}f)(\mathsf{F}f) = \mathsf{F}(m'.ff) \xrightarrow{\mathsf{F}\varphi^2} \mathsf{F}(f.m) = \mathsf{F}f.\mathsf{F}m \ , \ \mathsf{F}u' \xrightarrow{\mathsf{F}\varphi^0} \mathsf{F}(f.u) = \mathsf{F}f.\mathsf{F}u \) = \mathsf$$

Finally, F_{01} sends an opmonoidal 2-cell ω to $F\omega$. If the strict monoidal 2-functor F is also symmetric, then F_{01} is strict monoidal and symmetric. This gives the 1-cell part of a 2-functor $(-)_{01}$: sm-2Cat \rightarrow sm-2Cat.

For a monoidal 2-natural transformation $\Theta : \mathsf{F} \to \mathsf{G}$, there is a 2-natural transformation $\Theta_{01} : \mathsf{F}_{01} \to \mathsf{G}_{01}$ with the component at a pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$ in \mathcal{M}

$$(\Theta_A, \Theta_A, \mathsf{F}m = \mathsf{G}m.\Theta_{AA} = \mathsf{G}m.\Theta_A\Theta_A, \Theta_A, \Theta_A, \mathsf{F}u = \mathsf{G}u.\Theta_I = \mathsf{G}u).$$

It is strict monoidal thanks to the strict monoidality of Θ . These maps define

- (i) a 2-functor $(-)_{01}$ from the 2-category of strict monoidal 2-categories, strict monoidal 2-functors and monoidal 2-natural transformations to 2Cat,
- (ii) a 2-functor $(-)_{01}$: sm-2Cat \rightarrow sm-2Cat.

(It would be easy to extend them to monoidal modifications — defined in the evident way — but this is not needed for the purposes of the present paper.)

(2) There are symmetrically constructed 2-functors $(-)_{10} := (((-)^{\diamond})_{01})^{\diamond}$ of both kinds in items (i) and (ii) of the list in part (1) above. They send a strict monoidal

2-category \mathcal{M} to the 2-category $\mathcal{M}_{10} = ((\mathcal{M}^{\mathfrak{S}})_{01})^{\mathfrak{S}}$ whose 0-cells are again the pseudomonoids in \mathcal{M} . Its 1-cells and 2-cells are known as monoidal 1-cells and monoidal 2-cells in \mathcal{M} , respectively.

(3) Next we show that the 2-endofunctors on sm-2Cat in parts (1) and (2) commute up-to an irrelevant 2-natural isomorphism (this extends a claim in [2, Proposition 6.75]).

First we compare the symmetric strict monoidal 2-categories $(\mathcal{M}_{10})_{01}$ and $(\mathcal{M}_{01})_{10}$ for a symmetric strict monoidal 2-category \mathcal{M} . A 0-cell of $(\mathcal{M}_{10})_{01}$ is a pseudomonoid in \mathcal{M}_{10} . As such, it consists of the following data.

- A 0-cell of \mathcal{M}_{10} ; that is, a pseudomonoid $(A, m^{!}, u^{!}, \alpha^{!}, \lambda^{!}, \varrho^{!})$ in \mathcal{M} .
- Multiplication and unit 1-cells in \mathcal{M}_{10} . That is, monoidal 1-cells in $\mathcal{M}_{(m^-,\xi,\xi^0)}: (AA, m^!m^!.1\sigma 1, u^!u^!, \alpha^!\alpha^!.1.1, \lambda^!\lambda^!, \varrho^!\varrho^!) \to (A, m^!, u^!, \alpha^!, \lambda^!, \varrho^!)$ and $(u^-,\xi_0,\xi_0^0): (I, 1, 1, 1, 1, 1) \to (A, m^!, u^!, \alpha^!, \lambda^!, \varrho^!).$
- Associativity and unitality 2-cells in \mathcal{M}_{10} . They are invertible monoidal 2-cells in $\mathcal{M}, \alpha^-: m^-.m^-1 \to m^-.1m^-, \lambda^-: m^-.u^-1 \to 1$ and $\varrho^-: m^-.1u^- \to 1$ satisfying MacLane's pentagon and triangle conditions.

Symmetrically, a 0-cell of $(\mathcal{M}_{01})_{10}$ is a pseudomonoid in \mathcal{M}_{01} , which consists of the following data.

- A 0-cell of \mathcal{M}_{01} ; that is, a pseudomonoid $(A, m^{-}, u^{-}, \alpha^{-}, \lambda^{-}, \varrho^{-})$ in \mathcal{M} .
- Multiplication and unit 1-cells in \mathcal{M}_{01} . That is, opmonoidal 1-cells in \mathcal{M}_{01} . $(m^{|}, \xi.1, \xi_0) : (AA, m^-m^-.1\sigma 1, u^-u^-, \alpha^-\alpha^-.1.1, \lambda^-\lambda^-, \varrho^-\varrho^-) \to (A, m^-, u^-, \alpha^-, \lambda^-, \varrho^-),$ $(u^{|}, \xi^0, \xi^0_0) : (I, 1, 1, 1, 1, 1) \to (A, m^-, u^-, \alpha^-, \lambda^-, \varrho^-).$
- Associativity and unitality 2-cells in \mathcal{M}_{01} . They are invertible opmonoidal 2cells in $\mathcal{M}, \alpha^{\scriptscriptstyle |}: m^{\scriptscriptstyle |}.m^{\scriptscriptstyle |}1 \rightarrow m^{\scriptscriptstyle |}.1m^{\scriptscriptstyle |}, \lambda^{\scriptscriptstyle |}: m^{\scriptscriptstyle |}.u^{\scriptscriptstyle |}1 \rightarrow 1$ and $\varrho^{\scriptscriptstyle |}: m^{\scriptscriptstyle |}.1u^{\scriptscriptstyle |} \rightarrow 1$ satisfying MacLane's pentagon and triangle conditions.

Both of these sets of data amount to two pseudomonoid structures $(A, m', u', \alpha', \lambda', \varrho')$ and $(A, m^-, u^-, \alpha^-, \lambda^-, \varrho^-)$ on the same object A together with 2-cells in \mathcal{M}

$$m^{!}.m^{-}m^{-} \xrightarrow{\xi} m^{-}.m^{|}m^{|}.1\sigma 1 \qquad u^{|} \xrightarrow{\xi^{0}} m^{-}.u^{|}u^{|} \qquad m^{|}.u^{-}u^{-} \xrightarrow{\xi_{0}} u^{-} \qquad u^{|} \xrightarrow{\xi^{0}_{0}} u^{-}$$

(giving rise to the 2-cell $\xi.1: m^{!}.m^{-}m^{-}.1\sigma 1 \rightarrow m^{-}.m^{!}m^{!}.1\sigma 1.1\sigma 1 = m^{-}.m^{!}m^{!}$). They are subject to conditions as follows.

- Associativity and unitality of the monoidal 1-cell (m^-, ξ, ξ^0) ; equivalently, compatibility of the opmonoidal 2-cells $\alpha^{|}, \lambda^{|}$ and $\varrho^{|}$ with the binary parts of the opmonoidal structures of their source and target 1-cells.
- Associativity and unitality of the monoidal 1-cell (u^-, ξ_0, ξ_0^0) ; equivalently, compatibility of the opmonoidal 2-cells $\alpha^{\scriptscriptstyle |}$, $\lambda^{\scriptscriptstyle |}$ and $\varrho^{\scriptscriptstyle |}$ with the nullary parts of the opmonoidal structures of their source and target 1-cells.
- Compatibility of the monoidal 2-cells α^- , λ^- and ρ^- with the binary parts of the monoidal structures of their source and target 1-cells; equivalently, coassociativity and counitality of the opmonoidal 1-cell $(m^{\scriptscriptstyle |}, \xi_{\cdot}, 1, \xi_{0})$.
- Compatibility of the monoidal 2-cells α^- , λ^- and ϱ^- with the nullary parts of the monoidal structures of their source and target 1-cells; equivalently, coassociativity and counitality of the opmonoidal 1-cell $(u^{\scriptscriptstyle |}, \xi^0, \xi^0_0)$.

Such a datum is called a *duoidal* or 2-oidal object of \mathcal{M} .

A 1-cell in either one of the 2-categories $(\mathcal{M}_{10})_{01}$ and $(\mathcal{M}_{01})_{10}$ consists of a 1-cell $f: A \to A'$ in \mathcal{M} together with a monoidal structure $(\varphi_2: m'.ff \to f.m', \varphi_0: u' \to f.u')$ and an opmonoidal structure $(\varphi^2: f.m^- \to m^{-\prime}.ff, \varphi^0: f.u^- \to u^{-\prime})$ subject to the following compatibility conditions.

- The monoidal 2-cell φ^2 : $(f, \varphi_2, \varphi_0).(m^-, \xi, \xi^0) \rightarrow (m^{-\prime}, \xi', \xi^{0\prime}).(ff, \varphi_2\varphi_2.1, \varphi_0\varphi_0)$ is compatible with the binary parts of the monoidal structures of its source and target 1-cells; equivalently, the opmonoidal 2-cell $\varphi_2 : (m^{\prime\prime}, \xi^{\prime}.1, \xi_0').(ff, \varphi^2\varphi^2.1, \varphi^0\varphi^0) \rightarrow (f, \varphi^2, \varphi^0).(m^{\prime}, \xi.1, \xi_0)$ is compatible with the binary parts of the opmonoidal structures of its source and target 1-cells.
- The monoidal 2-cell φ^2 is compatible with the nullary parts of the monoidal structures of its source and target 1-cells; equivalently, the opmonoidal 2-cell $\varphi_0 : (u^{\prime}, \xi^{0\prime}, \xi^{0\prime}_0) \rightarrow (f, \varphi^2, \varphi^0) \cdot (u^{\prime}, \xi^0, \xi^0_0)$ is compatible with the binary parts of the opmonoidal structures of its source and target 1-cells.
- The monoidal 2-cell φ^0 : $(f, \varphi_2, \varphi_0).(u^-, \xi_0, \xi_0^0) \rightarrow (u^{-\prime}, \xi_0', \xi_0^{0\prime})$ is compatible with the binary parts of the monoidal structures of its source and target 1cells; equivalently, the opmonoidal 2-cell φ_2 is compatible with the nullary parts of the opmonoidal structures of its source and target 1-cells.
- The monoidal 2-cell φ^0 is compatible with the nullary parts of the monoidal structures of its source and target 1-cells; equivalently, the opmonoidal 2-cell φ_0 is compatible with the nullary parts of the opmonoidal structures of its source and target 1-cells.

Such a datum is termed a (1, 1)-oidal 1-cell. In the particular case of $\mathcal{M} = \mathsf{Cat}$, in [1] it was called a *bimonoidal functor*.

Finally, a 2-cell $(f, \varphi_2, \varphi_0, \varphi^2, \varphi^0) \to (g, \gamma_2, \gamma_0, \gamma^2, \gamma^0)$ in either one of the 2-categories $(\mathcal{M}_{10})_{01}$ and $(\mathcal{M}_{01})_{10}$ is a 2-cell in \mathcal{M} which is both

- a monoidal 2-cell $(f, \varphi_2, \varphi_0) \to (g, \gamma_2, \gamma_0)$ and
- an opmonoidal 2-cell $(f, \varphi^2, \varphi^0) \to (g, \gamma^2, \gamma^0)$.

We say that it is a (1, 1)-oidal 2-cell in \mathcal{M} .

This proves the isomorphism of the 2-categories $(\mathcal{M}_{10})_{01}$ and $(\mathcal{M}_{01})_{10}$. The obtained isomorphism is clearly symmetric strict monoidal and 2-natural.

(4) Thanks to their commuting verified in item (3) above, we may apply the 2functor in part (1) q times and the 2-functor in part (2) p times in an arbitrary order, for any non-negative integers q and p. Thereby we obtain a 2-functor $(-)_{pq}$: sm-2Cat \rightarrow sm-2Cat (so that $(-)_{00}$ is the identity 2-functor). In particular, it takes any symmetric strict monoidal 2-category \mathcal{M} to a symmetric strict monoidal 2-category \mathcal{M}_{pq} . Recall that a 0-cell in $\mathcal{M}_{p+1,q} \cong (\mathcal{M}_{pq})_{10}$ is the same as a 0-cell in $\mathcal{M}_{p,q+1} \cong$ $(\mathcal{M}_{pq})_{01}$; namely, a pseudomonoid in \mathcal{M}_{pq} . Thus the notion of 0-cell in $\mathcal{M}_{p,n-p}$ only depends on the non-negative integer n but not on the integer $0 \leq p \leq n$. (For the particular 2-category $\mathcal{M} =$ Cat this was discussed in Proposition 7.49 and Remark 7.51 of [2].)

2. (0, q)-OIDAL MONADS

The aim of this section is to prove the commutativity (up-to 2-natural isomorphism) of the diagram

$$sm-2Cat \xrightarrow{(-)_{01}} sm-2Cat \qquad (2.1)$$

$$Mnd \downarrow \qquad \qquad \downarrow Mnd$$

$$sm-2Cat \xrightarrow{(-)_{01}} sm-2Cat$$

of the 2-functors of Example 1.9 and Example 1.10 (1). Its iteration yields 2-natural isomorphisms $\mathsf{Mnd}(\mathcal{M}_{0q}) \cong \mathsf{Mnd}(\mathcal{M})_{0q}$ for any symmetric strict monoidal 2-category \mathcal{M} and any non-negative integer q. A 0-cell in $\mathsf{Mnd}(\mathcal{M}_{0q}) \cong \mathsf{Mnd}(\mathcal{M})_{0q}$ is called a (0,q)-oidal monad.

For 2-endofunctors on the 2-category $2Cat_{\times}$ of *Cartesian monoidal* 2-categories (instead of our sm-2Cat), commutativity of the analogous diagram was proved (by actually the same steps) in [31, Lemma 3.1]. A possible generalization, discussed here practically without any additional technical difficulty, is proposed in the introduction and the concluding remarks of [31].

In order to prove the commutativity of (2.1) (up-to 2-natural isomorphism), we need to compare first the actions of the 2-functors around it on an arbitrary 0-cell; that is, symmetric strict monoidal 2-category \mathcal{M} . A 0-cell in $\mathsf{Mnd}(\mathcal{M}_{01})$ is a monad in \mathcal{M}_{01} thus it consists of the following data.

- A 0-cell of \mathcal{M}_{01} ; that is, a pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$ in \mathcal{M} .
- A 1-cell $(A, m, u, \alpha, \lambda, \varrho) \to (A, m, u, \alpha, \lambda, \varrho)$ in \mathcal{M}_{01} ; that is, an opmonoidal 1-cell $(t: A \to A, \tau^2: t.m \to m.tt, \tau^0: t.u \to u)$ in \mathcal{M} .
- Multiplication and unit 2-cells in \mathcal{M}_{01} ; that is, opmonoidal 2-cells $\mu : (t, \tau^2, \tau^0)$. $(t, \tau^2, \tau^0) \rightarrow (t, \tau^2, \tau^0)$ and $\eta : (1, 1, 1) \rightarrow (t, \tau^2, \tau^0)$ in \mathcal{M} satisfying the associativity and unit conditions.

On the other hand, a 0-cell in $\mathsf{Mnd}(\mathcal{M})_{01}$ is a pseudomonoid in $\mathsf{Mnd}(\mathcal{M})$ thus it consists of the following data.

- A 0-cell in $\mathsf{Mnd}(\mathcal{M})$; that is, a monad (A, t, μ, η) in \mathcal{M} .
- Multiplication and unit 1-cells in $\mathsf{Mnd}(\mathcal{M})$; that is, monad morphisms $(m : AA \to A, \tau^2 : t.m \to m.tt)$ and $(u : I \to A, \tau^0 : t.u \to u)$ in \mathcal{M} .
- Associativity and unitality 2-cells in $\mathsf{Mnd}(\mathcal{M})$; that is, invertible monad transformations $\alpha : (m.m1, (1.\tau^21)^{\cdot}(\tau^2.1)) \to (m.1m, (1.1\tau^2)^{\cdot}(\tau^2.1)), \lambda : (m.u1, (1.\tau^01)^{\cdot}(\tau^2.1)) \to (1, 1)$ and $\varrho : (m.1u, (1.1\tau^0)^{\cdot}(\tau^2.1)) \to (1, 1)$ satisfying MacLane's pentagon and triangle conditions.

Both sets of data above amount to a pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$, an opmonoidal 1-cell $(t, \tau^2, \tau^0) : (A, m, u, \alpha, \lambda, \varrho) \to (A, m, u, \alpha, \lambda, \varrho)$ and a monad (A, t, μ, η) in \mathcal{M} with common 1-cell part t; subject to the following compatibility conditions.

The first two diagrams express the opmonoidality of the 2-cell μ ; equivalently, the multiplicativity of the monad morphisms (m, τ^2) and (u, τ^0) , respectively. The last two diagrams express the opmonoidality of the 2-cell η ; equivalently, the unitality of the monad morphisms (m, τ^2) and (u, τ^0) , respectively. This structure is termed a (0, 1)-oidal or opmonoidal monad.

Both in $\mathsf{Mnd}(\mathcal{M}_{01})$ and $\mathsf{Mnd}(\mathcal{M})_{01}$ a 1-cell is an opmonoidal 1-cell $(f, \varphi^2, \varphi^0)$: $(A, m, u, \alpha, \lambda, \varrho) \to (A', m', u', \alpha', \lambda', \varrho')$ and a monad morphism $(f, \Phi) : (A, t, \mu, \eta) \to (A', t', \mu', \eta')$ in \mathcal{M} with common 1-cell part f such that the following compatibility conditions hold.

The first diagram expresses the compatibility of the opmonoidal 2-cell $\Phi : (t', \tau'^2, \tau'^0)$. $(f, \varphi^2, \varphi^0) \rightarrow (f, \varphi^2, \varphi^0).(t, \tau^2, \tau^0)$ with the binary parts of the opmonoidal structures of its source and target 1-cells; equivalently, the requirement that $\varphi^2 : (f, \Phi).(m, \tau^2) \rightarrow (m', \tau'^2).(ff, \Phi\Phi)$ is a monad transformation. The second diagram expresses the compatibility of the opmonoidal 2-cell Φ with the nullary parts of the opmonoidal structures of its source and target 1-cells; equivalently, the requirement that $\varphi^0 : (f, \Phi).(u, \tau^0) \rightarrow (u', \tau'^0)$ is a monad transformation.

Finally, both in $\mathsf{Mnd}(\mathcal{M}_{01})$ and $\mathsf{Mnd}(\mathcal{M})_{01}$ a 2-cell is a 2-cell in \mathcal{M} which is both opmonoidal and a monad transformation.

With this we established an isomorphism between the 2-categories $\mathsf{Mnd}(\mathcal{M}_{01})$ and $\mathsf{Mnd}(\mathcal{M})_{01}$. It is clearly symmetric strict monoidal and it is straightforward to see its 2-naturality.

3. q-oidal Eilenberg-Moore objects

For a symmetric strict monoidal 2-category \mathcal{M} , any symmetric strict monoidal 2-functor $\mathsf{H} : \mathsf{Mnd}(\mathcal{M}) \to \mathcal{M}$ induces a symmetric strict monoidal 2-functor

$$\mathsf{Mnd}(\mathcal{M}_{0q}) \cong \mathsf{Mnd}(\mathcal{M})_{0q} \xrightarrow{\mathsf{H}_{0q}} \mathcal{M}_{0q}$$
(3.1)

whose object map sends a (0, q)-oidal monad to a q-oidal object of \mathcal{M} . The aim of this section is to investigate when the Eilenberg-Moore construction in \mathcal{M} yields such a symmetric strict monoidal 2-functor $\mathsf{H} : \mathsf{Mnd}(\mathcal{M}) \to \mathcal{M}$; hence the q-oidal structure of the base object of a (0, q)-oidal monad in \mathcal{M} lifts to the Eilenberg-Moore object along the 'forgetful' 1-cells (see (EM)). The results of this section extend [1, Theorem 8.2] and place it in a broader context. We apply analogous ideas to those in [31]. In the particular case when the strict monoidal structure of \mathcal{M} is Cartesian, a stronger result — proving also that H_{0q} provides Eilenberg-Moore construction on \mathcal{M}_{0q} ; cf. Remark 8.3 — was obtained in [31, Theorem 5.1].

The Eilenberg-Moore construction for 2-categories was shortly recalled from [28] in the Introduction. Namely, a 2-category \mathcal{M} is said to admit Eilenberg-Moore construction if the inclusion 2-functor $\mathcal{M} \to \mathsf{Mnd}(\mathcal{M})$ possesses a right 2-adjoint H. The image of a monad (A, t) under H is denoted by A^t and it is called the Eilenberg-Moore

object of the monad. The component of the counit of this 2-adjunction at any monad (A, t) is necessarily of the form $(u^t, 1.\epsilon^t : t.u^t = u^t.f^t.u^t \to u^t) : (A^t, 1) \to (A, t)$, where the 1-cell $u^t : A^t \to A$ has a left adjoint f^t in \mathcal{M} and $\epsilon^t : f^t.u^t \to 1$ is the counit of this adjunction, see [28, pages 151-153]. The adjunction $f^t \dashv u^t : A^t \to A$ generates the monad (A, t) in the sense that the 1-cell $u^t.f^t$ is equal to t, the 2-cell $1.\epsilon^t.1 : f^t.u^t.f^t.u^t \to f^t.u^t$ is equal to the multiplication of the monad while the unit $\eta^t : 1 \to u^t.f^t$ of the adjunction is equal to the unit of the monad, see [28, Theorem 2]. By [28, Theorem 3], for any other adjunction $l \dashv r : B \to A$ generating the same monad (A, t), there is a unique comparison 1-cell k rendering commutative the first diagram of

With its help, the counit ϵ of the adjunction $l \dashv r$ fits in the second commutative diagram of (3.2).

Consider now a strict monoidal 2-category which admits Eilenberg-Moore construction. The adjunction $1 \dashv 1 : I \to I$ of identity functors on the monoidal unit I generates the monad (I, 1). Hence there is a comparison 1-cell $I \to I^1$. For any monads (A, t) and (B, s), the adjunction $f^t f^s \dashv u^t u^s : A^t B^s \to AB$ generates the monad (AB, ts). Hence there is a unique comparison 1-cell $A^t B^s \to (AB)^{ts}$.

Definition 3.1. We say that a strict monoidal 2-category admits monoidal Eilenberg– Moore construction if it admits Eilenberg–Moore construction and the comparison 1cells $I \to I^1$ and $A^t B^s \to (AB)^{ts}$ are identities for the monoidal unit I and all monads (A, t) and (B, s).

For example, the strict monoidal 2-category **Cat** of Example 1.1 admits monoidal Eilenberg-Moore construction. More generally, monoidality of the Eilenberg-Moore construction in a Cartesian monoidal 2-category is a key assumption also in [31, Theorem 5.1].

Proposition 3.2. If a strict monoidal 2-category \mathcal{M} admits monoidal Eilenberg-Moore construction, then the Eilenberg-Moore 2-functor $H : Mnd(\mathcal{M}) \to \mathcal{M}$ is strict monoidal. If in addition \mathcal{M} is symmetric then also H is symmetric.

Proof. The monoidal unit I is preserved by the assumption that the comparison 1-cell $I \to I^1 = \mathsf{H}(I,1)$ is the identity. The equality of 2-functors $\mathsf{H}(-\otimes -) = \mathsf{H}(-) \otimes \mathsf{H}(-)$ holds on any 0-cells (A,t) and (B,s) by the assumption that the comparison 1-cell $A^t B^s \to (AB)^{ts}$ is the identity. By the second diagram of (3.2), $\epsilon^{ts} = \epsilon^t \epsilon^s$. Therefore for any 1-cells $(h,\chi) : (A,t) \to (A',t')$ and $(g,\gamma) : (B,s) \to (B',s')$, the 2-natural isomorphism $\mathcal{M}((AB)^{ts}, (A'B')^{t's'}) \cong \mathsf{Mnd}(\mathcal{M})(((AB)^{ts}, 1), (A'B', t's'))$ sends both objects $\mathsf{H}(hg,\chi\gamma)$ and $\mathsf{H}(h,\chi)\mathsf{H}(g,\gamma)$ to the same object

$$(hg.u^{ts}, t's'.hg.u^{ts} \xrightarrow{\chi\gamma.1} hg.ts.u^{ts} = hg.u^{ts}.f^{ts}.u^{ts} \xrightarrow{11.\epsilon^{ts}} hg.u^{ts})$$

which proves their equality. For any 2-cells ω and ϑ , both 2-cells $H(\omega\vartheta)$ and $H(\omega)H(\vartheta)$ are the liftings of the same monad transformation $\omega\vartheta$ hence they are equal.

Assume now the existence of a symmetry σ of \mathcal{M} . For any monads (A, t) and (B, s), the 2-natural isomorphism $\mathcal{M}((AB)^{ts}, (BA)^{st}) \cong \mathsf{Mnd}(\mathcal{M})(((AB)^{ts}, 1), (BA, st))$ sends both objects $\mathsf{H}(\sigma_{A,B}, 1)$ and σ_{A^t,B^s} to the same object

$$(\sigma_{A,B}.u^{ts}, st.\sigma_{A,B}.u^{ts} = \sigma_{A,B}.ts.u^{ts} = \sigma_{A,B}.u^{ts}.f^{ts}.u^{ts} \xrightarrow{1.1.\epsilon^{ts}} \sigma_{A,B}.u^{ts})$$

which proves their equality.

From (3.1) we obtain the following generalization of [1, Theorem 8.2].

Corollary 3.3. In a symmetric strict monoidal 2-category which admits monoidal Eilenberg-Moore construction, the q-oidal structure of the base object A of any (0,q)-oidal monad (A,t) lifts to the Eilenberg-Moore object A^t along the 1-cell $u^t : A^t \to A$.

4. (p, 0)-OIDAL MONADS

Without entering the details, in this section we sketch the dual of the situation in Section 2. Although the omitted proofs are analogous to those in Section 2, they do not seem to follow by any kind of abstract duality.

The 2-functors $\mathsf{Mnd}_{\mathsf{op}}$ of Example 1.9 and $(-)_{10}$ of part (2) of Example 1.10 constitute a diagram

$$\begin{array}{c} \text{sm-2Cat} \xrightarrow{(-)_{10}} \text{sm-2Cat} \\ \xrightarrow{\text{Mnd}_{op}} & & & \\ \text{sm-2Cat} \xrightarrow{(-)_{10}} \text{sm-2Cat} \end{array}$$

which is commutative up-to 2-natural isomorphism. So whenever there is a symmetric strict monoidal 2-functor V from $Mnd_{op}(\mathcal{M}) = Mnd(\mathcal{M}_{op})_{op}$ to some symmetric strict monoidal 2-category \mathcal{M} , it induces a symmetric strict monoidal 2-functor

$$\operatorname{Mnd}_{\operatorname{op}}(\mathcal{M}_{p0}) \cong \operatorname{Mnd}_{\operatorname{op}}(\mathcal{M})_{p0} \xrightarrow{\operatorname{V}_{p0}} \mathcal{M}_{p0}$$

$$(4.1)$$

for any non-negative integer p. Its object map sends a (p, 0)-oidal monad to a p-oidal object in \mathcal{M} .

5. *p*-OIDAL EILENBERG-MOORE OBJECTS

In this section we consider a symmetric strict monoidal 2-category \mathcal{M} and a nonnegative integer p. Then we describe a setting in which the p-oidal structure of the base object A of any (p, 0)-oidal monad (A, t) lifts to the Eilenberg–Moore object A^t along the 1-cell $f^t : A \to A^t$. (A 1-cell $\overline{h} : A^t \to B^s$ is said to be a lifting of 1-cell $h : A \to B$ along the left adjoint 1-cells $f^t : A \to A^t$ and $f^s : B \to B^s$ if $f^s \cdot h = \overline{h} \cdot f^t$.) The results of this section extend [1, Theorem 7.1] and give it a different explanation.

Let \mathcal{M} be a 2-category which admits Eilenberg–Moore construction. For any 1-cell $(h, \chi) : (A, t) \to (B, s)$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$, consider the reflexive pair of morphisms in $\mathcal{M}(A^t, B^s)$,

$$f^{s}.h.t.u^{t} \underbrace{f^{s}.h.u^{t}.f^{t}.u^{t} - \frac{1.1.1.\epsilon^{t}}{1.\chi.1}}_{1.\chi.1} f^{s}.s.h.u^{t} = f^{s}.u^{s}.f^{s}.h.u^{t} - \frac{\pi(h,\chi)}{\epsilon^{s}.1.1.1}} F^{s}.h.u^{t} - \frac{\pi(h,\chi)}{1.\chi.1} \to \mathsf{V}(h,\chi).$$
(5.1)

(A common section of the parallel morphisms is $1.1.\eta^t.1$ in terms of the unit η^t of the monad (A,t).) Assuming that their coequalizer exists for all 1-cells (h,χ) in $\mathsf{Mnd_{op}}(\mathcal{M})$, and it is preserved by the horizontal composition on either side with any 1-cell in \mathcal{M} , we construct a 2-functor $\mathsf{V} : \mathsf{Mnd_{op}}(\mathcal{M}) \to \mathcal{M}$ whose action on any 1-cell (h,χ) is given by the coequalizer in (5.1). Coequalizers of this form were studied in [23].

Lemma 5.1. Consider a 2-category \mathcal{M} which admits Eilenberg–Moore construction and take any monads (A, t) and (B, s) in \mathcal{M} . If the coequalizer (5.1) exists for all objects of $\mathsf{Mnd_{op}}(\mathcal{M})((A, t), (B, s))$, then there is a functor $\mathsf{Mnd_{op}}(\mathcal{M})((A, t), (B, s)) \to \mathcal{M}(A^t, B^s)$ whose object map is given by the coequalizer in (5.1).

Proof. Take a morphism $\omega : (h, \chi) \to (k, \kappa)$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})((A, t), (B, s))$. By the universality of the coequalizer in the top row of the serially commutative diagram

$$\begin{array}{c|c} f^{s}.h.t.u^{t} & f^{s}.h.u^{t}.f^{t}.u^{t} & 1.1.1.\epsilon^{t} \\ f^{s}.h.t.u^{t} & f^{s}.s.h.u^{t} = f^{s}.u^{s}.f^{s}.h.u^{t} & \epsilon^{s}.1.1.1 \\ 1.\omega.1 & f^{s}.s.h.u^{t} = f^{s}.u^{s}.f^{s}.h.u^{t} & \epsilon^{s}.1.1.1 \\ f^{s}.k.t.u^{t} & f^{s}.k.u^{t}.f^{t}.u^{t} & 1.1.1.\epsilon^{t} \\ f^{s}.k.t.u^{t} & f^{s}.k.u^{t} = f^{s}.u^{s}.f^{s}.k.u^{t} & \epsilon^{s}.1.1.1 \end{array}$$

the image of ω under the desired functor V occurs in the right vertical. Functoriality follows by construction.

In order to interpret the functors of Lemma 5.1 as the hom functors of a 2-functor, we need to see their compatibility with the horizontal composition and the identity 1-cells. The proof of this rests on the following.


Lemma 5.2. Consider a 2-category \mathcal{M} which admits Eilenberg–Moore construction. Assume that the coequalizer (5.1) exists for some 1-cell $(h, \chi) : (A, t) \to (B, s)$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$, and it is preserved by the horizontal composition on either side with any 1-cell in \mathcal{M} . Then

$$(f^s.h.u^t.f^t \xrightarrow{\pi(h,\chi).1} \mathsf{V}(h,\chi).f^t) = (f^s.h.t \xrightarrow{1.\chi} f^s.s.h = f^s.u^s.f^s.h \xrightarrow{\epsilon^s.1.1} f^s.h).$$

Proof. By assumption $V(h, \chi)$. f^t appears in the coequalizer

$$f^{s}.h.t.t \xrightarrow{1.1,\mu^{t}} f^{s}.h.t = f^{s}.u^{s}.f^{s}.h.u^{t}.f^{t} \xrightarrow{\pi(h,\chi).1} \mathsf{V}(h,\chi).f^{t}$$

(where μ^t denotes the multiplication of the monad (A, t)). So the claim follows by the observation that



is a split coequalizer (where η^t is the unit of the monad t).

Proposition 5.3. Consider a 2-category \mathcal{M} which admits Eilenberg–Moore construction. If the coequalizer (5.1) exists for all 1-cells in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$, and it is preserved by the horizontal composition on either side with any 1-cell in \mathcal{M} , then the following hold.

(1) There is a 2-functor $V : \mathsf{Mnd}_{\mathsf{op}}(\mathcal{M}) \to \mathcal{M}$ which sends a monad (A, t) to the Eilenberg-Moore object A^t and whose hom functors are defined as in Lemma 5.1.

(2) If moreover \mathcal{M} is a strict monoidal 2-category which admits monoidal Eilenberg-Moore construction, then the 2-functor V of part (1) is strict monoidal.

(3) If in addition \mathcal{M} has a symmetry then the 2-functor V of part (1) is symmetric as well.

Proof. (1) There is a monad t.(-) on the category $\mathcal{M}(A^t, A)$ whose Eilenberg–Moore category is isomorphic to $\mathsf{Mnd}(\mathcal{M})((A^t, 1), (A, t)) \cong \mathcal{M}(A^t, A^t)$; and whose forgetful functor differs by this isomorphism from

$$u^{t}.(-): \mathcal{M}(A^{t}, A^{t}) \to \mathcal{M}(A^{t}, A).$$
 (5.2)

Thus (5.2) is monadic and therefore the coequalizers of those forks exist and are preserved by (5.2) which are sent by (5.2) to split coequalizers, see Proposition 3.5 on page 95 of [4] or Theorem 4.4.4 in [7].

Now the functor (5.2) sends the fork

$$f^{t}.t.u^{t} = f^{t}.u^{t}.f^{t}.u^{t} \xrightarrow{1.1.\epsilon^{t}} f^{t}.u^{t} \xrightarrow{\epsilon^{t}} 1$$

$$(5.3)$$

to the split coequalizer

$$u^{t}.f^{t}.t.u^{t} = u^{t}.f^{t}.u^{t}.f^{t}.u^{t} \xrightarrow{1.1.1.\epsilon^{t}} u^{t}.f^{t}.u^{t} \xrightarrow{1.\epsilon^{t}} u^{t}$$

so that (5.3) is a coequalizer and thus V preserves identity 1-cells.

For the preservation of the horizontal composition recall that for arbitrary 1-cells $(h, \chi) : (A, t) \to (B, s)$ and $(k, \kappa) : (B, s) \to (C, z)$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$, $\mathsf{V}(k, \kappa).\mathsf{V}(h, \chi)$ and $\mathsf{V}((k, \kappa).(h, \chi))$ are defined as the coequalizers of the respective pairs of parallel

morphisms in

The vertical equalities hold by Lemma 5.2. The square with the lower ones of the parallel arrows commutes since the following diagram commutes and its dotted arrow denotes an epimorphism.


$$\begin{array}{c|c} \mathsf{V}(k,\kappa).f^s.u^s.f^s & \xrightarrow{1.\epsilon^{s}.1} & \mathsf{V}(k,\kappa).f^s \\ \hline f^z.k.u^s.f^s.u^s.f^s & \xrightarrow{1.1.\epsilon^{s}.1} & f^z.k.u^s.f^s \\ \hline f^z.k.s.s & \xrightarrow{1.1.\mu^s} & f^z.k.s \\ \text{Lemma 5.2} & 1.\kappa.1 & 1.\kappa & \text{Lemma 5.2} \\ f^z.z.k.s & \xrightarrow{1.1.\kappa} & f^z.z.z.k & \xrightarrow{1.\mu^{z}.1} & f^z.z.k \\ \hline f^z.u^z.f^z.k.s & \xrightarrow{1.1.\kappa} & f^z.u^z.f^z.k & \xrightarrow{f^z}.u^z.f^z.k \\ \hline f^z.u^z.f^z.k.s & \xrightarrow{f^z}.u^z.f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1.1} & f^z.k & \xrightarrow{\epsilon^z.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z.1} & f^z.k.s & \xrightarrow{\epsilon^z.1} & \xrightarrow{\epsilon^z.1} & \xrightarrow{\epsilon^z.1} & f^z.k \\ \hline f^z.k.s & \xrightarrow{\epsilon^z$$

This also proves the equality

$$(\mathsf{V}(k,\kappa).f^{s}.h.u^{t} \xrightarrow{1.\pi(h,\chi)} \mathsf{V}(k,\kappa).\mathsf{V}(h,\chi)) = (f^{z}.k.h.u^{t} \xrightarrow{\pi((k,\kappa).(h,\chi))} \mathsf{V}((k,\kappa).(h,\chi))).$$
(5.4)

Part (2) is immediate. For part (3) recall that for the symmetry σ of \mathcal{M} , the action of V on the symmetry $(\sigma, 1) : (AB, ts) \to (BA, st)$ of $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$ is defined as the

coequalizer of the morphisms in the top row of



Both the vertical equalities and serial commutativity of the diagram follow by the 2-naturality of σ . The bottom row is the image under σ .(-) of parallel morphisms as in (5.3) hence their coequalizer is σ .1 = σ as stated.

In view of Lemma 5.2, the action of V can be interpreted as a lifting along the left adjoint 1-cells of the kind $f^t : A \to A^t$. So from (4.1) we obtain the following.

Corollary 5.4. In the setting of Proposition 5.3 (3), the p-oidal structure of the base object A of any (p, 0)-oidal monad (A, t) in \mathcal{M} lifts to the Eilenberg-Moore object A^t along the 1-cell $f^t : A \to A^t$.

Remark 5.5. The premises of Proposition 5.3 (3) hold in a suitable 2-subcategory of the 2-category Cat of Example 1.1 which we describe next.

Let \mathcal{M} be the symmetric strict monoidal 2-subcategory of Cat whose 0-cells are those categories in which the coequalizers of reflexive pairs exist; whose 1-cells are those functors which preserve reflexive coequalizers; and whose 2-cells are all natural transformations between them.

For any monad (A, t) in \mathcal{M} , the coequalizer of any reflexive pair in A^t exists and it is preserved by the forgetful functor $A^t \to A$, see [23, Corollary 3] or [7, Proposition 4.3.2]. Since the forgetful functor is conservative, it also reflects the coequalizers of reflexive pairs. Then if some functor $h : A \to B$ preserves reflexive coequalizers, then so does $h.u^t = u^s.\mathsf{H}(h,\chi)$ for any monad functor $(h,\chi) : (A,t) \to (B,s)$ and the Eilenberg–Moore 2-functor H . So since u^s reflects reflexive coequalizers, $\mathsf{H}(h,\chi) : A^t \to A^s$ preserves them. Because in this way \mathcal{M} is closed under the monoidal Eilenberg–Moore construction of Cat , it admits monoidal Eilenberg–Moore construction itself.

Evaluating the parallel natural transformations of (5.1) at an arbitrary *t*-algebra we get a reflexive pair in B^s . Their coequalizer exists by the considerations in the previous paragraph and it is evidently preserved by any 1-cell in \mathcal{M} . Then also the coequalizer (5.1) exists and it is preserved by the horizontal composition.


Applying Corollary 5.4 to the 2-category of Remark 5.5, we obtain an alternative proof of related [27, Theorem 2.6.4] and [1, Theorem 4.14]; as well as of [1, Theorem 7.1].

6. Multimonoidal structures in double categories

For a study of the lifting of the (p+q)-oidal structure of the base object of a (p,q)oidal monad in a symmetric strict monoidal 2-category, in the case when both p and q are strictly positive integers, we leave the realm of 2-categories and operate, more generally, with double categories. Therefore this section begins with the introduction of a 2-category sm-DblCat of symmetric strict monoidal double categories. Its choice is motivated by two requirements. On one hand, it should allow for the interpretation of the symmetry of a symmetric strict monoidal double category in [8, Definition 5] as a suitable 2-cell in the 2-category of double categories. On the other hand, even more importantly, a definition of sm-DblCat would serve our purposes if it admits several 2-functors constructed in this section and the next one.

In Example 6.12 we introduce 2-endofunctors $(-)_{01}$ and $(-)_{10}$ on sm-DblCat; and show that they commute up-to 2-natural isomorphism. Then iterating them in an arbitrary order q and p times, respectively, we obtain a 2-endofunctor $(-)_{pq}$ on sm-DblCat. The 0-cells of \mathbb{D}_{pq} , for any symmetric strict monoidal double category \mathbb{D} , can be interpreted as (p, q)-oidal objects in \mathbb{D} .

Ehresmann's square or quintet construction [12] yields a 2-functor $Sqr : sm-2Cat \rightarrow sm-DblCat$. Together with the 2-functor $(-)_{pq} : sm-2Cat \rightarrow sm-2Cat$ of Example 1.10 (4) for any non-negative integers p and q, it fits in the commutative diagram



The 2-functors around it take a symmetric strict monoidal 2-category \mathcal{M} to the symmetric strict monoidal double category $\mathbb{Sqr}(\mathcal{M})_{pq} \cong \mathbb{Sqr}(\mathcal{M}_{pq})$. Thus the (p,q)-oidal objects in $\mathbb{Sqr}(\mathcal{M})$ are the same as the (p+q)-oidal objects of \mathcal{M} .

We begin with quickly recalling some basic definitions. More details can be found e.g. in [8].

Definition 6.1. A *double category* is an internal category in the category of small categories and functors. Thus a double category \mathbb{D} consists of a category of objects V, a category of morphisms \mathcal{V} and structure functors

$$V \xrightarrow[t^-]{s^-} \mathcal{V} \xleftarrow{c^-} \mathcal{V} \times_V \mathcal{V}$$

which are subject to the usual axioms of internal category.

The objects of V are called the 0-cells or objects of \mathbb{D} and the morphisms of V are called the *vertical 1-cells* of \mathbb{D} . The objects of \mathcal{V} are called the *horizontal 1-cells* of \mathbb{D} and the morphisms of \mathcal{V} are called the 2-cells of \mathbb{D} . The compositions of both categories V and \mathcal{V} are called the *vertical composition* (of 1-cells and 2-cells, respectively). The functor c^- is called the *horizontal composition*. A general 2-cell is

depicted as

where s^{\dagger} denotes the source maps, and t^{\dagger} denotes the target maps of both categories V and \mathcal{V} . In this graphical notation, vertical composition is denoted by vertical juxtaposition and horizontal composition is denoted by horizontal juxtaposition.

In Definition 6.1 the roles of the vertical and horizontal structures turn out to be symmetric. This leads to a symmetric description of the same double category \mathbb{D} . It can be seen equivalently as an internal category in the category of small categories and functors with another category of objects H and another category of morphisms \mathcal{H} . In H the objects are again the 0-cells of \mathbb{D} but now the morphisms are the horizontal 1-cells. The objects of \mathcal{H} are the vertical 1-cells and the morphisms are the 2-cells again. The compositions in H and \mathcal{H} are provided by the horizontal composition of \mathbb{D} . In this description the structure functors of the internal category are

$$H \xrightarrow[]{s^{|}}_{\underset{t^{|}}{\underbrace{\overset{s^{|}}{\longleftarrow}}}} \mathcal{H} \xleftarrow{c^{|}} \mathcal{H} \times_{H} \mathcal{H}.$$

Here the functor i sends any object A of H (equivalently, of V) to the corresponding identity morphism $1 : A \to A$ in V (regarded now as an object of \mathcal{H}) and it sends a horizontal morphism h to the identity morphism $h \to h$ in \mathcal{V} (regarded now as a non-identity morphism in \mathcal{H}).

Any double category \mathbb{D} contains two 2-categories. In the *horizontal 2-category* $\mathsf{Hor}(\mathbb{D})$ the 0-cells are the 0-cells, and the 1-cells are the horizontal 1-cells of \mathbb{D} . The 2-cells are those 2-cells in \mathbb{D} which are taken by s^- and t^- to identity vertical 1-cells. Both horizontal and vertical compositions are the restrictions of the respective composition in \mathbb{D} . Symmetrically, in the *vertical 2-category* $\mathsf{Ver}(\mathbb{D})$ the 0-cells are the 0-cells are the vertical 1-cells of \mathbb{D} . The 2-cells are those 2-cells in \mathbb{D} which are taken by $s^{|}$ and $t^{|}$ to identity horizontal 1-cells. In this 2-category the horizontal composition is the restriction of the vertical composition of \mathbb{D} and vertical composition is the restriction of the vertical composition of \mathbb{D} .

Definition 6.2. A *double functor* is an internal functor in the category of small categories and functors. Thus a double functor $\mathbf{F} : \mathbb{D} \to \mathbb{D}'$ consists of compatible functors $(V \to V', \mathcal{V} \to \mathcal{V}')$. Equivalently, it consists of compatible functors $(H \to H', \mathcal{H} \to \mathcal{H}')$.

The object maps of the functors $V \to V'$ and $H \to H'$ in Definition 6.2 are equal. The morphism map of the functor $V \to V'$ is equal to the object map of the functor $\mathcal{H} \to \mathcal{H}'$ and the morphism map of the functor $H \to H'$ is equal to the object map of the functor $\mathcal{V} \to \mathcal{V}'$. The morphism maps of the functors $\mathcal{V} \to \mathcal{V}'$ and $\mathcal{H} \to \mathcal{H}'$ are equal again. Any double functor \mathbf{F} induces 2-functors $\mathsf{Hor}(\mathbf{F})$ and $\mathsf{Ver}(\mathbf{F})$ between the horizontal and vertical 2-categories, respectively, which we call the horizontal and the vertical 2-functors, respectively.

Between double functors $\mathbf{F}, \mathbf{G} : \mathbb{D} \to \mathbb{D}'$, there are two symmetric variants of internal natural transformations. A *horizontal transformation* $x : \mathbf{F} \to \mathbf{G}$ is an internal natural transformation in the category of small categories and functors from $(\mathbf{F}: V \to V', \mathbf{F}: \mathcal{V} \to \mathcal{V}')$ to $(\mathbf{G}: V \to V', \mathbf{G}: \mathcal{V} \to \mathcal{V}')$. Thus it is a functor $V \to \mathcal{V}'$ sending any vertical 1-cell $f: A \to B$ to a 2-cell

$$\begin{array}{c} \mathbf{F}A \xrightarrow{x_A} \mathbf{G}A \\ \mathbf{F}f & \downarrow x_f & \downarrow \mathbf{G}f \\ \mathbf{F}B \xrightarrow{x_B} \mathbf{G}B \end{array}$$

subject to the naturality condition

For any horizontal transformation $x : \mathbf{F} \to \mathbf{G}$, the components $x_A : \mathbf{F}A \to \mathbf{G}A$ constitute a 2-natural transformation $\mathsf{Hor}(x) : \mathsf{Hor}(\mathbf{F}) \to \mathsf{Hor}(\mathbf{G})$.

There is then a 2-category in which the 0-cells are the double categories, the 1cells are the double functors and the 2-cells are the horizontal transformations. The horizontal composition is induced by the consecutive application of double functors and the vertical composition is induced by the horizontal composition in the target double category. We may see Hor as a 2-functor from this 2-category to 2Cat.

Symmetrically, a vertical transformation $y : \mathbf{F} \to \mathbf{G}$ is an internal natural transformation in the category of small categories and functors from $(\mathbf{F} : H \to H', \mathbf{F} : \mathcal{H} \to \mathcal{H}')$ to $(\mathbf{G} : H \to H', \mathbf{G} : \mathcal{H} \to \mathcal{H}')$. Thus it is a functor $H \to \mathcal{H}'$ sending any horizontal 1-cell $l : A \to C$ to a 2-cell

$$\begin{array}{c} \mathbf{F}A \xrightarrow{\mathbf{F}l} \mathbf{F}C \\ y_A & \downarrow y_l & \downarrow y_c \\ \mathbf{G}A \xrightarrow{\mathbf{G}l} \mathbf{G}C \end{array}$$

subject to the naturality condition

For any vertical transformation $y : \mathbf{F} \to \mathbf{G}$, the components $y_A : \mathbf{F}A \to \mathbf{G}A$ constitute a 2-natural transformation $\mathsf{Ver}(y) : \mathsf{Ver}(\mathbf{F}) \to \mathsf{Ver}(\mathbf{G})$.

There is then a 2-category of double categories, double functors and vertical transformations, and a 2-functor Ver from it to 2Cat.

The notion of double natural transformation, which is relevant for our purposes, symmetrically combines vertical and horizontal transformations. It is a particular case of *generalized natural transformations* in [8].

Definition 6.3. For double functors $\mathbf{F}, \mathbf{G} : \mathbb{D} \to \mathbb{D}'$, a double natural transformation $\mathbf{F} \rightarrow \mathbf{G}$ consists of

- a horizontal transformation $x : \mathbf{F} \to \mathbf{G}$ a vertical transformation $y : \mathbf{F} \to \mathbf{G}$ a map θ associating to each 0-cell A of \mathbb{D} a 2-cell $\mathbf{G}A = \mathbf{G}A$

such that θ gives rise to natural transformation from the functor $x: V \to \mathcal{V}'$ to the functor sending a vertical 1-cell $f: A \to B$ to the identity morphism $1: \mathbf{G}f \to \mathbf{G}f$ in \mathcal{H}' (regarded as a non-identity morphism of \mathcal{V}'); as well as a natural transformation from the functor $H \to \mathcal{H}'$ sending a horizontal 1-cell $l: A \to C$ to the identity morphism 1 : $\mathbf{G}l \to \mathbf{G}l$ in \mathcal{V}' (regarded as a non-identity morphism of \mathcal{H}') to y. Equivalently, θ is a *modification* in the sense of [17, Section 1.6]. That is, in addition to the conditions (6.1) and (6.2), also the further naturality conditions

hold, for all vertical 1-cells $f: A \to B$ and horizontal 1-cells $l: A \to C$.

Double categories are the 0-cells, double functors are the 1-cells and double natural transformations are the 2-cells of the 2-category DblCat. For double functors \mathbf{F} , \mathbf{G} and $\mathbf{H}: \mathbb{D} \to \mathbb{D}'$, the vertical composite of double natural transformations (x, θ, y) : $\mathbf{F} \to \mathbf{G}$ and $(x', \theta', y') : \mathbf{G} \to \mathbf{H}$ has the component

$$\begin{array}{c} \mathbf{F}A \xrightarrow{x_A} \mathbf{G}A \xrightarrow{x'_A} \mathbf{H}A \\ y_A \downarrow & \| \theta_A & \| & \| 1 \\ \mathbf{G}A = \mathbf{G}A \xrightarrow{x'_A} \mathbf{H}A \\ y'_A \downarrow & \| 1 & y'_A & \| \theta'_A \\ \mathbf{H}A = \mathbf{H}A = \mathbf{H}A \end{array}$$
(6.4)

for any 0-cell A of \mathbb{D} . This is the *diagonal composition* of [8]. Double functors \mathbf{F} : $\mathbb{D} \to \mathbb{D}'$ and $\mathbf{F}' : \mathbb{D}' \to \mathbb{D}''$ are composed as any internal functors. For another pair of double functors $\mathbf{G} : \mathbb{D} \to \mathbb{D}'$ and $\mathbf{G}' : \mathbb{D}' \to \mathbb{D}''$, and for double natural transformations $(x, \theta, y) : \mathbf{F} \to \mathbf{G}$ and $(x', \theta', y') : \mathbf{F}' \to \mathbf{G}'$, the horizontal composite has the component

for any 0-cell A of \mathbb{D} . We leave it to the reader to check that DblCat is a 2-category indeed; in particular, to derive the middle four interchange law from (6.1), (6.2) and (6.3). There are evident forgetful 2-functors from DblCat to the 2-category of double categories, double functors and horizontal or vertical transformations.

Although double natural transformations as in Definition 6.3 were introduced in [8] (without using this name for them), we did not manage to find an explicit description of the above 2-category DblCat anywhere in the literature.

The 2-category DblCat can be made monoidal with the Cartesian product of double categories as the monoidal product. Then it has a symmetry **X** as well, provided by the flip double functors $\mathbb{D} \times \mathbb{C} \to \mathbb{C} \times \mathbb{D}$, sending a pair of 2-cells (ω, ϑ) to (ϑ, ω) .

Remark 6.4. The author is grateful to Ross Street for kindly pointing out the following derivation of the 2-category DblCat via the 'change of enriching category construction' in [13].

The category dblcat of double categories and double functors is Cartesian closed. (In the internal hom double category $[\mathbb{A}, \mathbb{B}]$ the 0-cells are the double functors $\mathbb{A} \to \mathbb{B}$, the horizontal/vertical 1-cells are the horizontal/vertical transformations and the 2-cells are the modifications in [17, Section 1.6].) Hence it can be seen as a dblcat-enriched category.


Now let us consider the following monoidal functor F from dblcat to the category cat of small categories and functors. It sends a double category \mathbb{D} to the category whose objects are the 0-cells of \mathbb{D} ; whose morphisms $P \to Q$ are the triples consisting of a horizontal 1-cell x, a vertical 1-cell y and a 2-cell

in \mathbb{D} ; and whose composition is the diagonal composition as in (6.4). A double functor **H** is taken by F to the functor which acts on the objects as **H** does; and acts on the morphisms componentwise.

Following [13], the monoidal functor F induces a 2-functor from the 2-category of dblcat-enriched categories to the 2-category 2Cat of cat-enriched categories. This induced 2-functor sends the dblcat-enriched category dblcat precisely to the 2-category DblCat above.

Example 6.5. There are several notions of duality for double categories (see e.g. [17, Section 1.2]). In addition to the horizontal and vertical opposites as for 2-categories,

there is a further possibility to take the *diagonal dual* or *transpose* \mathbb{D}^* of a double category \mathbb{D} . Its 0-cells are the 0-cells of \mathbb{D} , its horizontal 1-cells are the vertical 1-cells of \mathbb{D} and vice versa, its vertical 1-cells are the horizontal 1-cells of \mathbb{D} . A 2-cell in \mathbb{D}^* as on the left,



is a 2-cell in \mathbb{D} on the right. The horizontal composition in \mathbb{D}^* is the vertical composition in \mathbb{D} and the vertical composition in \mathbb{D}^* is the horizontal composition in \mathbb{D} .

Any double functor $\mathbf{F} : \mathbb{D} \to \mathbb{C}$ induces a double functor $\mathbf{F}^* : \mathbb{D}^* \to \mathbb{C}^*$. On the 0-cells and on the 2-cells it acts as \mathbf{F} . Its action on the horizontal and vertical 1-cells is the action of \mathbf{F} on the vertical and horizontal 1-cells, respectively.

The components of a double natural transformation $(x, \theta, y) : \mathbf{F} \to \mathbf{G}$ on the left



can be seen as the components on the right of a double natural transformation $(x, \theta, y)^* : \mathbf{F}^* \to \mathbf{G}^*$.

This defines 2-functors $(-)^*$ from the 2-category of double categories, double functors and horizontal/vertical transformations to the 2-category of double categories, double functors and vertical/horizontal transformations; as well as a 2-endofunctor $(-)^*$ on the 2-category DblCat of double categories, double functors and double natural transformations.

Definition 6.6. [8, Definition 4] A strict monoidal double category is a monoid in the category of double categories and double functors considered with the Cartesian product of double categories as the monoidal product. That is, a strict monoidal double category consists of a double category \mathbb{D} together with double functors \otimes the monoidal product — from the Cartesian product double category $\mathbb{D} \times \mathbb{D}$ to \mathbb{D} and I — the monoidal unit — from the singleton double category (with a single 0-cell and only identity higher cells) to \mathbb{D} such that \otimes is strictly associative with the strict unit I.

The action of the monoidal product double functor \otimes on any cells will be denoted by juxtaposition. Note that in a strict monoidal double category \mathbb{D} all constituent categories V, H, \mathcal{V} and \mathcal{H} are strict monoidal, and so are the horizontal and vertical 2-categories Hor(\mathbb{D}) and Ver(\mathbb{D}).

Definition 6.7. [8, Definition 4] A strict monoidal double functor is a monoid morphism in the category of double categories and double functors considered with the Cartesian product of double categories as the monoidal product. That is, a double functor $\mathbf{F} : \mathbb{D} \to \mathbb{D}'$ such that $\mathbf{F}(-\otimes -) = \mathbf{F}(-) \otimes' \mathbf{F}(-)$ and $\mathbf{F}I = I'$.

Definition 6.8. A monoidal horizontal transformation between strict monoidal double functors $(\mathbb{D}, \otimes, I) \to (\mathbb{D}', \otimes', I')$ is a horizontal transformation z such that $z : V \to \mathcal{V}'$ is a strict monoidal functor. Symmetrically, a monoidal vertical transformation between strict monoidal double functors is a vertical transformation v such that $v : H \to \mathcal{H}'$ is a strict monoidal functor. A monoidal double natural transformation is a double natural transformation (z, θ, v) between strict monoidal double functors **F** and $\mathbf{G} : (\mathbb{D}, \otimes, I) \to (\mathbb{D}', \otimes', I')$ such that $z : V \to \mathcal{V}'$ and $v : H \to \mathcal{H}'$ are strict monoidal functors and θ , seen as a natural transformation in either way, is monoidal. That is to say, the equalities of 2-cells

$$(\mathbf{F}A)(\mathbf{F}B) \xrightarrow{z_A z_B} (\mathbf{G}A)(\mathbf{G}B) \qquad \mathbf{F}(AB) \xrightarrow{z_{AB}} \mathbf{G}(AB) \qquad \mathbf{F}I \xrightarrow{z_I} \mathbf{G}I \qquad I' = I' \\ v_A v_B \downarrow \qquad \downarrow \theta_A \theta_B \qquad = v_{AB} \downarrow \qquad \downarrow \theta_{AB} \qquad v_I \downarrow \qquad \downarrow \theta_I \qquad = \parallel \qquad \downarrow 1 \qquad \parallel \\ (\mathbf{G}A)(\mathbf{G}B) = (\mathbf{G}A)(\mathbf{G}B) \qquad \mathbf{G}(AB) = \mathbf{G}(AB) \qquad \mathbf{G}I = \mathbf{G}I \qquad I' = I'$$

hold for all 0-cells A and B in \mathbb{D} .

Strict monoidal double categories are the 0-cells and strict monoidal double functors are the 1-cells in various 2-categories. The 2-cells can be chosen to be monoidal horizontal transformations, monoidal vertical transformations, or monoidal double natural transformations.

Definition 6.9. [8, Definition 5] A symmetric strict monoidal double category consists of a strict monoidal double category (\mathbb{D}, \otimes, I) together with a double natural transformation $(x, \sigma, y) : \otimes \to \otimes \mathbf{X}$ (where **X** denotes the relevant component of the symmetry of the 2-category DblCat, that is, the flip double functor $\mathbb{D} \times \mathbb{D} \to \mathbb{D} \times \mathbb{D}$). It is required to be involutive in the sense that


$$\begin{array}{c|c} \otimes & \xrightarrow{x} & \otimes .\mathbf{X} \xrightarrow{x.1} & \otimes \\ y \downarrow & \downarrow \sigma & \parallel & \downarrow 1 \\ \otimes .\mathbf{X} & \xrightarrow{ \otimes .\mathbf{X} \xrightarrow{x.1} & \otimes } & = & \parallel & \downarrow 1 \\ y.1 \downarrow & \downarrow 1 & y_{1}^{1} \downarrow & \sigma.1 & \parallel & \otimes & = & \otimes \\ \otimes & \xrightarrow{ \otimes . \infty & \otimes & = & \otimes } & \otimes & & & & \\ \end{array}$$

and the *hexagon condition* is required to hold, which says that the 2-cells of Figure 1 are equal.

The symmetry (x, σ, y) of a symmetric strict monoidal double category \mathbb{D} induces symmetries on all constituent categories V, H, \mathcal{V} and \mathcal{H} .

It follows from Definition 6.9 that for any 0-cell A,

Recall from [8, Section 4] that in a strict monoidal double category (\mathbb{D}, \otimes, I) with symmetry (x, σ, y) , the 2-cell σ has a horizontal inverse σ^h on the left, and a vertical






FIGURE 1. Hexagon condition for a symmetric strict monoidal double category

inverse σ^v on the right of

$$AB \xrightarrow{x} BA \qquad AB \xrightarrow{x} BA \xrightarrow{x} AB$$

$$y \downarrow \downarrow \sigma \parallel \qquad y \downarrow \downarrow \sigma \parallel \downarrow 1 \parallel \qquad BA \xrightarrow{x} BA \xrightarrow{x} AB$$

$$y \downarrow \downarrow \sigma \parallel \downarrow 1 \parallel \qquad BA \xrightarrow{x} AB.$$

$$y \downarrow \downarrow 1 \downarrow y$$

$$AB \xrightarrow{x} AB$$

From these explicit expressions it follows for any vertical 1-cells $f: A \to A'$ and $g: B \to B'$ that

$$AB \xrightarrow{x} BA = BA = BA \qquad (6.5)$$

$$\| \qquad \downarrow \sigma^{h} \qquad \downarrow y \qquad gf \qquad \downarrow 1 \qquad \downarrow gf \qquad AB \xrightarrow{x} BA$$

$$AB = AB \qquad \downarrow 1 \qquad B'A' = B'A' = fg \qquad \downarrow x \qquad \downarrow gf$$

$$fg \qquad \downarrow 1 \qquad \downarrow fg \qquad y \qquad \downarrow \sigma^{v} \qquad A'B' \xrightarrow{x} B'A'$$

$$A'B' = A'B' \xrightarrow{x} B'A'$$

and it also follows for any horizontal 1-cells $f: A \to A'$ and $g: B \to B'$ that

$$AB \xrightarrow{x} BA \xrightarrow{gf} B'A' = B'A' \qquad AB \xrightarrow{x} BA \xrightarrow{gf} B'A' \qquad (6.6)$$

$$\| \qquad \downarrow \sigma^{h} \qquad \downarrow y \qquad \downarrow y \qquad \downarrow y \qquad \downarrow \sigma^{v} \qquad = \qquad \| \qquad \downarrow 1 \qquad \| \qquad AB \xrightarrow{x} B'A' \qquad AB \xrightarrow{x} B'A' \qquad (6.6)$$

$$AB = AB \xrightarrow{fg} A'B' \xrightarrow{x} B'A' \qquad AB \xrightarrow{fg} A'B' \xrightarrow{x} B'A'.$$

Definition 6.10. A symmetric strict monoidal double functor is a strict monoidal double functor $\mathbf{F} : (\mathbb{D}, \otimes, I) \to (\mathbb{D}', \otimes', I')$ which preserves the symmetry in the sense of the equalities

$$\mathbf{F}.x = x'.(\mathbf{F} \times \mathbf{F}) \qquad \mathbf{F}.y = y'.(\mathbf{F} \times \mathbf{F}) \qquad \mathbf{F}.\sigma = \sigma'.(\mathbf{F} \times \mathbf{F}).$$

(of functors $V \times V \to \mathcal{V}'$ and $H \times H \to \mathcal{H}'$, and of natural transformations, respectively).

Definition 6.11. Consider some symmetric strict monoidal double functors \mathbf{F}, \mathbf{G} : $\mathbb{D} \to \mathbb{D}'$. A monoidal horizontal transformation $h: \mathbf{F} \to \mathbf{G}$ is said to be symmetric if the strict monoidal functor $h: V \to \mathcal{V}'$ is symmetric. Symmetrically, a monoidal vertical transformation $v: \mathbf{F} \to \mathbf{G}$ is said to be symmetric if the strict monoidal functor $v: H \to \mathcal{H}'$ is symmetric. A monoidal double natural transformation $(h, \theta, v):$ $\mathbf{F} \to \mathbf{G}$ is said to be symmetric if h and v are symmetric.

Symmetric strict monoidal double categories are the 0-cells and symmetric strict monoidal double functors are the 1-cells in various 2-categories. The 2-cells can be chosen to be symmetric monoidal horizontal transformations, symmetric monoidal vertical transformations or symmetric monoidal double natural transformations. We denote this latter 2-category by sm-DblCat. Its vertical and horizontal compositions are given by the same expressions as in DblCat.

This finishes our review of the basic theory of double categories.

Example 6.12. (1) For any strict monoidal double category \mathbb{D} there is a double category \mathbb{D}_{01} which we describe next. Its 0-cells are the pseudomonoids

$$(A, AA \xrightarrow{m} A, I \xrightarrow{u} A, A \xrightarrow{m1} AA \xrightarrow{m} A \xrightarrow{m} A \xrightarrow{u1} AA \xrightarrow{m} A \xrightarrow{h} A \xrightarrow{1u} AA \xrightarrow{m} A \xrightarrow{$$

in the horizontal 2-category $Hor(\mathbb{D})$ of \mathbb{D} (which we call the *horizontal pseudomonoids* in \mathbb{D}), and the horizontal 1-cells are the opmonoidal 1-cells

$$(A \xrightarrow{h} A', \| \underset{AA \xrightarrow{m}}{\longrightarrow} A \xrightarrow{h} A', I \xrightarrow{u} A \xrightarrow{h} A' \\ \downarrow \chi^2 \| , \| \underset{AA \xrightarrow{h}}{\longrightarrow} A'A' \xrightarrow{m'} A', I \xrightarrow{u'} A'$$

in the horizontal 2-category of \mathbb{D} . A vertical 1-cell consists of a vertical 1-cell on the left, together with 2-cells on the right of

$$\begin{array}{cccc} A & & AA \xrightarrow{m} A & & I \xrightarrow{u} A \\ \downarrow f & & ff \downarrow & \downarrow \varphi^2 & \downarrow f & & \\ A' & & A'A' \xrightarrow{m'} A' & & I \xrightarrow{u'} A' \end{array}$$

which are required to satisfy the following coassociativity and counitality conditions.

A 2-cell in \mathbb{D}_{01} with boundaries on the left of

$$\begin{array}{cccc} A \xrightarrow{(h,\chi^{2},\chi^{0})} C & & A \xrightarrow{h} C \\ (f,\varphi^{2},\varphi^{0}) & & & \downarrow (g,\gamma^{2},\gamma^{0}) \\ B \xrightarrow{(k,\kappa^{2},\kappa^{0})} D & & B \xrightarrow{k} D \end{array}$$

is a 2-cell of \mathbb{D} on the right, subject to the opmonoidality conditions

(where all occurring pseudomonoid structures are denoted by the same symbols $(m, u, \alpha, \lambda, \varrho)$). Horizontal 1-cells are composed by the composition rule of opmonoidal 1-cells in a 2-category, see Example 1.10. Vertical 1-cells are vertically composed according to the rule

Horizontal and vertical compositions of 2-cells are computed in \mathbb{D} . The horizontal 2-category of the resulting double category \mathbb{D}_{01} can be obtained by applying the 2-functor $(-)_{01}$ of Example 1.10 to the horizontal 2-category of \mathbb{D} . That is, $Hor(\mathbb{D})_{01} \cong Hor(\mathbb{D}_{01})$. (The analogous statement fails to hold for the vertical 2-category.)

If in addition the strict monoidal double category (\mathbb{D}, \otimes, I) has a symmetry (x, σ, y) , then the above double category \mathbb{D}_{01} can be equipped with a strict monoidal structure. The monoidal unit is the monoidal unit I of \mathbb{D} seen as a trivial pseudomonoid, see Example 1.10. The monoidal product of 0-cells and of horizontal 1-cells is their monoidal product as pseudomonoids and opmonoidal 1-cells, respectively, in the horizontal 2-category, see again Example 1.10. The monoidal product of vertical 1-cells

$$\begin{array}{ccc} (f,\varphi^2,\varphi^0):A\to A' \text{ and } (g,\gamma^2,\gamma^0):B\to B' \text{ is} \\ & AB & ABAB \xrightarrow{1x1} AABB \xrightarrow{mm} AB & I \xrightarrow{uu} AB \\ & \downarrow fg & fgfg \downarrow & \downarrow 1x1 & ffgg \downarrow \varphi^2\gamma^2 & \downarrow fg & & \parallel \downarrow \varphi^0\gamma^0 & \downarrow fg \\ & A'B' & A'B'A'B' \xrightarrow{1x1} A'A'B'B' \xrightarrow{m'm'} A'B' & I' \xrightarrow{u'u'} A'B'. \end{array}$$

The monoidal product of 2-cells is their monoidal product as 2-cells of \mathbb{D} . What is more, the above strict monoidal double category \mathbb{D}_{01} has a symmetry with components

$$\begin{array}{c|c} AB \xrightarrow{(x,1,1)} BA \\ \xrightarrow{(y,\nu^2,\nu^0)} & & & \\ BA \xrightarrow{} BA \end{array}$$

where the binary part ν^2 and the nullary part ν^0 of the opmonoidal 1-cell y are the 2-cells

respectively, (where σ^h stands for the horizontal inverse of σ in \mathbb{D}).

For any strict monoidal double functor $\mathbf{F} : \mathbb{D} \to \mathbb{D}'$ there is a double functor $\mathbf{F}_{01} : \mathbb{D}_{01} \to \mathbb{D}'_{01}$. On the horizontal 2-category it acts as the 2-functor obtained by applying the 2-functor $(-)_{01}$ of Example 1.10 to the 2-functor induced by \mathbf{F} between the horizontal 2-categories. It takes a vertical 1-cell $(f, \varphi^2, \varphi^0)$ to

$$\begin{array}{cccc} \mathbf{F}A & (\mathbf{F}A)(\mathbf{F}A) == \mathbf{F}(AA) \xrightarrow{\mathbf{F}m} \mathbf{F}A & I == \mathbf{F}I \xrightarrow{\mathbf{F}u} \mathbf{F}A \\ \downarrow \mathbf{F}f & (\mathbf{F}f)(\mathbf{F}f) & \downarrow 1 & \mathbf{F}(f) & \downarrow \mathbf{F}\varphi^2 & \downarrow \mathbf{F}f & \parallel \downarrow 1 & \parallel \downarrow \mathbf{F}\varphi^0 & \downarrow \mathbf{F}f \\ \mathbf{F}B & (\mathbf{F}B)(\mathbf{F}B) == \mathbf{F}(BB) \xrightarrow{\mathbf{F}m} \mathbf{F}B & I == \mathbf{F}I \xrightarrow{\mathbf{F}u} \mathbf{F}B \end{array}$$

and it acts on the 2-cells as \mathbf{F} .

If the strict monoidal double categories (\mathbb{D}, \otimes, I) and $(\mathbb{D}', \otimes', I')$ are symmetric and **F** preserves the symmetry, then \mathbf{F}_{01} is symmetric strict monoidal with respect to the symmetric strict monoidal structures of \mathbb{D}_{01} and \mathbb{D}'_{01} in the previous paragraphs.

Finally, take any strict monoidal double functors \mathbf{F} and \mathbf{G} . For any monoidal horizontal transformation $z : \mathbf{F} \to \mathbf{G}$, there is a horizontal transformation $z_{01} : \mathbf{F}_{01} \to \mathbf{G}_{01}$ with the components $(z_A, 1, 1)$ at any pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$ in \mathbb{D} , and x_f at any vertical 1-cell f in \mathbb{D} . Whenever \mathbf{F} and \mathbf{G} are symmetric as well — so that \mathbf{F}_{01} and \mathbf{G}_{01} are strict monoidal too — z_{01} is monoidal too; and it is symmetric if z is so.

For any monoidal vertical transformation $v : \mathbf{F} \to \mathbf{G}$ there is a vertical transformation $v_{01} : \mathbf{F}_{01} \to \mathbf{G}_{01}$ with the components (v_A, v_m, v_u) at any pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$ in \mathbb{D} ; and v_h at any horizontal 1-cell h in \mathbb{D} . It is symmetric monoidal whenever v is a symmetric monoidal vertical transformation between symmetric strict monoidal double functors.

For any monoidal double natural transformation $(z, \theta, v) : \mathbf{F} \to \mathbf{G}$ there is a double natural transformation $\mathbf{F}_{01} \to \mathbf{G}_{01}$ with the components z_{01}, v_{01} and

$$\begin{array}{c|c} \mathbf{F}A \xrightarrow{(z,1,1)} \mathbf{G}A \\ \xrightarrow{(v,v_m,v_u)} & & & & \\ \mathbf{G}A \xrightarrow{\qquad} \mathbf{G}A \end{array} \end{array}$$

at any pseudomonoid $(A, m, u, \alpha, \lambda, \varrho)$ in \mathbb{D} . It is monoidal whenever \mathbf{F} , \mathbf{G} and v are symmetric as well; and symmetric if so is z in addition.

These maps define various 2-functors $(-)_{01}$:

- (i) from the 2-category of strict monoidal double categories, strict monoidal double functors and monoidal horizontal transformations to the 2-category of double categories, double functors and horizontal transformations,
- (ii) from the 2-category of strict monoidal double categories, strict monoidal double functors and monoidal vertical transformations to the 2-category of double categories, double functors and vertical transformations,
- (iii) from the 2-category of strict monoidal double categories, strict monoidal double functors and monoidal double natural transformations to the 2-category DblCat of double categories, double functors and double natural transformations,
- (iv) from the 2-category of symmetric strict monoidal double categories, symmetric strict monoidal double functors and symmetric monoidal horizontal transformations to itself,
- (v) from the 2-category of symmetric strict monoidal double categories, symmetric strict monoidal double functors and symmetric monoidal vertical transformations to itself,
- (vi) from the 2-category sm-DblCat of symmetric strict monoidal double categories, symmetric strict monoidal double functors and symmetric monoidal double natural transformations to itself.

(2) The 'diagonal duality' 2-functors $(-)^*$ (see Example 6.5) induce 2-functors between the (domain, respectively, codomain) 2-categories in items (i) and (ii), as well as those in items (iv) and (v); and they induce 2-endofunctors on the 2-categories in items (iii) and (vi) of the list of part (1) above. Hence symmetrically to part (1), we define 2-functors $(-)_{10} := (((-)^*)_{01})^*$ of all kinds in this list. They send a strict monoidal double category \mathbb{D} to the double category \mathbb{D}_{10} whose 0-cells are the pseudomonoids in the vertical 2-category of \mathbb{D} (i.e. the *vertical pseudomonoids* in \mathbb{D}), and the vertical 1-cells are the monoidal morphisms in the vertical 2-category of \mathbb{D} . The horizontal 1-cells are monoidal horizontal 1-cells with respect to the vertical pseudomonoids and the 2-cells are the 2-cells of \mathbb{D} which are monoidal in a suitable sense. If in addition the strict monoidal double category \mathbb{D} is symmetric then also \mathbb{D}_{10} is symmetric strict monoidal. (3) Next we show that the diagram

sm-DblCat
$$\xrightarrow{(-)_{01}}$$
 sm-DblCat
 $(-)_{10} \downarrow \qquad \qquad \downarrow (-)_{10}$
sm-DblCat $\xrightarrow{(-)_{01}}$ sm-DblCat

of the 2-functors in parts (1) and (2) commutes up-to 2-natural isomorphism. (The same proof shows, in fact, the commutativity of the 2-endofunctors $(-)_{01}$ and $(-)_{10}$ also in items (iv) and (v) of the list of part (1).)

For any symmetric strict monoidal double category \mathbb{D} , a 0-cell of $(\mathbb{D}_{01})_{10}$ is a vertical pseudomonoid in \mathbb{D}_{01} ,

$$((A, m^-, u^-, \alpha^-, \lambda^-, \varrho^-), (m^{\scriptscriptstyle |}, \zeta, \zeta_0), (u^{\scriptscriptstyle |}, \zeta^0, \zeta_0^0), \alpha^{\scriptscriptstyle |}, \lambda^{\scriptscriptstyle |}, \varrho^{\scriptscriptstyle |}).$$

Symmetrically, a 0-cell of $(\mathbb{D}_{10})_{01}$ is a horizontal pseudomonoid in \mathbb{D}_{10} ,

 $((A,m^{\scriptscriptstyle |},u^{\scriptscriptstyle |},\alpha^{\scriptscriptstyle |},\lambda^{\scriptscriptstyle |},\varrho^{\scriptscriptstyle |}),(m^-\!\!\!,\xi,\xi^0),(u^-\!\!\!,\xi_0,\xi_0^0),\alpha^-\!\!\!,\lambda^-\!\!\!,\varrho^-).$

A bijective correspondence between them is given by $\zeta_0^0 = \xi_0^0$, $\zeta_0 = \xi_0$, $\zeta^0 = \xi^0$ and ζ equal to

$$\begin{array}{c|c} AAAA \xrightarrow{1x1} AAAA \xrightarrow{m^-m^-} AA \\ \downarrow y_1 & \downarrow 1\sigma1 & \downarrow \downarrow 1 & \downarrow \\ AAAA \xrightarrow{m^-m^-} AAA \\ \downarrow y_1 & \downarrow 1 & \downarrow \\ AAAA & \downarrow \xi & \downarrow m^{|} \\ AAAAA & \downarrow \xi & \downarrow m^{|} \\ AA \xrightarrow{m^-} AA & \downarrow A \end{array}$$

Similarly to Example 1.10 (3), the following properties are pairwise equivalent (for checking it use also (6.5)).

- Associativity and unitality of the monoidal horizontal 1-cell (m^-, ξ, ξ^0) are equivalent to the compatibilities of the opmonoidal 2-cells α^{\downarrow} , λ^{\downarrow} and ρ^{\downarrow} with the binary parts of their source and target opmonoidal vertical 1-cells.
- Associativity and unitality of the monoidal horizontal 1-cell (u^-, ξ_0, ξ_0^0) are equivalent to the compatibilities of the opmonoidal 2-cells $\alpha^{\scriptscriptstyle |}$, $\lambda^{\scriptscriptstyle |}$ and $\varrho^{\scriptscriptstyle |}$ with the nullary parts of their source and target opmonoidal vertical 1-cells.
- Compatibilities of the monoidal 2-cells α^- , λ^- and ϱ^- with the binary parts of their source and target monoidal horizontal 1-cells are equivalent to the coassociativity and counitality of the opmonoidal vertical 1-cell $(m^{\downarrow}, \zeta, \zeta_0)$.
- Compatibilities of the monoidal 2-cells α^- , λ^- and ϱ^- with the nullary parts of their source and target monoidal horizontal 1-cells are equivalent to the coassociativity and counitality of the opmonoidal vertical 1-cell $(u^{\downarrow}, \zeta^0, \zeta_0^0)$.

A horizontal 1-cell in $(\mathbb{D}_{01})_{10}$ is a monoidal horizontal 1-cell in \mathbb{D}_{01} ,

$$((f, \varphi^2, \varphi^0), \varphi_2, \varphi_0)$$

while a horizontal 1-cell in $(\mathbb{D}_{10})_{01}$ is an opmonoidal horizontal 1-cell in \mathbb{D}_{10} ,

$$((f, \varphi_2, \varphi_0), \varphi^2, \varphi^0).$$

Both of them amount to a monoidal structure (φ_2, φ_0) on f with respect to the vertical pseudomonoids and an opmonoidal structure (φ^2, φ^0) on f with respect to the horizontal pseudomonoids, which are subject to four compatibility conditions.

- Compatibility of the opmonoidal 2-cell φ_0 with the nullary parts of its surrounding opmonoidal 1-cells coincides with the compatibility of the monoidal 2-cell φ^0 with the nullary parts of its surrounding monoidal 1-cells.
- Compatibility of the opmonoidal 2-cell φ_0 with the binary parts of its surrounding opmonoidal 1-cells is the same condition as the compatibility of the monoidal 2-cell φ^2 with the nullary parts of surrounding monoidal 1-cells.
- Symmetrically, compatibility of the opmonoidal 2-cell φ_2 with the nullary parts of its surrounding opmonoidal 1-cells coincides with the compatibility of the monoidal 2-cell φ^0 and the binary parts of its surrounding monoidal 1-cells.
- Compatibility of the opmonoidal 2-cell φ_2 with the binary parts of its surrounding opmonoidal 1-cells, and compatibility of the monoidal 2-cell φ^2 with the binary parts of its surrounding monoidal 1-cells are equivalent to each other (for its proof also (6.6) is used).

So a bijective correspondence between the horizontal 1-cells of $(\mathbb{D}_{01})_{10}$ and $(\mathbb{D}_{10})_{01}$ is obtained by re-ordering the constituent 2-cells.

A bijection between the vertical 1-cells of $(\mathbb{D}_{01})_{10}$ and $(\mathbb{D}_{10})_{01}$ is obtained symmetrically.

Finally, the 2-cells both in $(\mathbb{D}_{01})_{10}$ and $(\mathbb{D}_{10})_{01}$ are those 2-cells of \mathbb{D} which are both

- monoidal (with respect to the monoidal structures of the surrounding 1-cells),
- opmonoidal (with respect to the opmonoidal structures of these 1-cells).

So there is a trivial (identity) bijection between them.

The above bijections combine into an iso double functor $(\mathbb{D}_{01})_{10} \to (\mathbb{D}_{10})_{01}$ which is symmetric strict monoidal and 2-natural.

(4) By the commutativity of the first diagram of part (3), we may apply in any order the 2-functor in its columns p times and the 2-functor in its rows q times. This yields a 2-functor $(-)_{pq}$: sm-DblCat \rightarrow sm-DblCat. Analogously we obtain 2-endofunctors $(-)_{pq}$ on the 2-categories in items (iv) and (v) of the list of part (1). For any symmetric strict monoidal double category \mathbb{D} , we term the 0-cells of \mathbb{D}_{pq} as the (p,q)-oidal objects of \mathbb{D} . In particular, the (0, 1)-oidal objects are the horizontal pseudomonoids in \mathbb{D} while the (1, 0)-oidal objects are the vertical pseudomonoids. Thus in contrast to part (4) of Example 1.10, the structure of (p,q)-oidal objects in \mathbb{D} depends both on p and q not only their sum.

Example 6.13. We may regard Ehresmann's square or quintet construction [12] as a 2-functor $\text{Sqr} : \text{sm-2Cat} \rightarrow \text{sm-DblCat}$.

Recall that for any 2-category \mathcal{M} , the 0-cells of the double category $\operatorname{Sqr}(\mathcal{M})$ are the 0-cells of \mathcal{M} . Both the horizontal and the vertical 1-cells in $\operatorname{Sqr}(\mathcal{M})$ are the 1-cells

of \mathcal{M} . The 2-cells of the form

$$\begin{array}{c} A \xrightarrow{h} C \\ f \downarrow & \downarrow \omega & \downarrow g \\ B \xrightarrow{k} D \end{array}$$

in $\operatorname{Sqr}(\mathcal{M})$ are the 2-cells $\omega : g.h \to k.f$ in \mathcal{M} . Both the horizontal composition of horizontal 1-cells, and the vertical composition of vertical 1-cells in $\operatorname{Sqr}(\mathcal{M})$ are given by the composition of 1-cells in \mathcal{M} . The horizontal composition results in

$$t.n.h \xrightarrow{\vartheta.1} p.g.h \xrightarrow{1.\omega} p.k.f \quad \text{for 2-cells} \qquad \begin{array}{c} A \xrightarrow{h} C & C \xrightarrow{n} E \\ f \downarrow & \downarrow \omega & \downarrow g & g \downarrow & \downarrow \vartheta & \downarrow t \\ B \xrightarrow{k} D & D \xrightarrow{p} F \end{array}$$
(6.7)

while the vertical composition results in

$$p.g.h \xrightarrow{1.\omega} p.k.f \xrightarrow{\vartheta.1} t.n.f \quad \text{for 2-cells} \quad \begin{cases} A \xrightarrow{h} C & B \xrightarrow{k} D \\ f \downarrow \downarrow \omega \downarrow g & n \downarrow \downarrow \vartheta \downarrow p \\ B \xrightarrow{k} D & E \xrightarrow{k} F. \end{cases}$$
(6.8)

Both the horizontal and the vertical 2-categories of $Sqr(\mathcal{M})$ are isomorphic to \mathcal{M} .

Whenever \mathcal{M} possesses a strict monoidal structure, it induces an evident strict monoidal structure on $Sqr(\mathcal{M})$. If moreover \mathcal{M} has a symmetry x then a symmetry on $Sqr(\mathcal{M})$ is given by the horizontal and vertical transformations in the first two diagrams, and 2-cell part in the third diagram of

for any 1-cells $f : A \to A'$ and $g : B \to B'$ in \mathcal{M} .


For any 2-functor $F : \mathcal{M} \to \mathcal{N}$, there is a double functor $Sqr(F) : Sqr(\mathcal{M}) \to Sqr(\mathcal{N})$. Its action on the 0-cells is equal to the action of F on the 0-cells. Its action both on the horizontal and vertical 1-cells is equal to the action of F on the 1-cells. Its action on the 2-cells is equal to the action of F on the 2-cells. Whenever F is strict monoidal, clearly Sqr(F) is strict monoidal; and whenever F is symmetric, so is Sqr(F).

Finally, a 2-natural transformation $\omega : \mathsf{F} \to \mathsf{G}$ induces a double natural transformation $\operatorname{Sqr}(\omega) : \operatorname{Sqr}(\mathsf{F}) \to \operatorname{Sqr}(\mathsf{G})$ with the horizontal and vertical transformations in the first two diagrams, and 2-cell part in the third diagram of

$$\begin{array}{cccc} \mathsf{F}A & \stackrel{\omega_A}{\longrightarrow} \mathsf{G}A & \mathsf{F}A & \stackrel{\mathsf{F}f}{\longrightarrow} \mathsf{F}B & \mathsf{F}A & \stackrel{\omega_A}{\longrightarrow} \mathsf{G}A \\ \mathsf{F}f & & \downarrow 1 & \downarrow \mathsf{G}f & & & \\ \mathsf{F}B & \stackrel{\omega_A}{\longrightarrow} \mathsf{G}B & & & \\ \mathsf{G}A & \stackrel{\omega_A}{\longrightarrow} \mathsf{G}B & & & \\ \mathsf{G}A & \stackrel{\mathsf{G}f}{\longrightarrow} \mathsf{G}B & & & \\ \end{array}$$

for any 1-cell $f: A \to B$ in \mathcal{M} . It is symmetric monoidal whenever ω is monoidal.

The so constructed 2-functor $Sqr : sm-2Cat \rightarrow sm-DblCat$, together with the 2-functors of Example 1.10 and Example 6.12 in the columns, render strictly commutative the diagrams of 2-functors



Then Sqr commutes also with the 2-functors $(-)_{pq}$ of Example 1.10 (4) and Example 6.12 (4) for any non-negative integers p and q.

7. (p,q)-OIDAL MONADS

In this section we describe a 2-functor Mnd : sm-2Cat \rightarrow sm-DblCat. Its object map sends a symmetric strict monoidal 2-category \mathcal{M} to the double category of monads in Sqr(\mathcal{M}) in the sense of [14] (so the horizontal 2-category of Mnd(\mathcal{M}) is Mnd(\mathcal{M}) and its vertical 2-category is Mnd_{op}(\mathcal{M})). Together with the 2-functor $(-)_{pq}$: sm-2Cat \rightarrow sm-2Cat of Example 1.10 (4) and the 2-functor $(-)_{pq}$: sm-DblCat \rightarrow sm-DblCat of Example 6.12 (4) for any non-negative integers p and q, it fits in the diagram

$$\begin{array}{c|c} \text{sm-2Cat} & \xrightarrow{\mathbb{M}\text{nd}} \text{sm-DblCat} \\ (-)_{pq} & & & \downarrow (-)_{pq} \\ \text{sm-2Cat} & \xrightarrow{\mathbb{M}\text{nd}} \text{sm-DblCat} \end{array}$$

which commutes up-to 2-natural isomorphism. The 2-functors around it take a symmetric strict monoidal 2-category \mathcal{M} to a symmetric strict monoidal double category $\mathbb{M}nd(\mathcal{M})_{pq} \cong \mathbb{M}nd(\mathcal{M}_{pq})$. Thus the (p,q)-oidal objects in $\mathbb{M}nd(\mathcal{M})$ are the same as the (p,q)-oidal monads in \mathcal{M} . Consequently, any symmetric strict monoidal double functor $\mathbf{K} : \mathbb{M}nd(\mathcal{M}) \to \mathbb{S}qr(\mathcal{M})$ induces a symmetric strict monoidal double functor

$$\mathbb{M}\mathrm{nd}(\mathcal{M}_{pq}) \cong \mathbb{M}\mathrm{nd}(\mathcal{M})_{pq} \xrightarrow{\mathbf{K}_{pq}} \mathbb{S}\mathrm{qr}(\mathcal{M})_{pq} \cong \mathbb{S}\mathrm{qr}(\mathcal{M}_{pq})$$

whose object map sends a (p,q)-oidal monad in \mathcal{M} to a (p+q)-oidal object in \mathcal{M} .

As anticipated above, for any 2-category \mathcal{M} we take the double category $Mnd(\mathcal{M})$ of monads in the sense of [14] in the double category $Sqr(\mathcal{M})$ of Example 6.13. Thus a 0-cell in $Mnd(\mathcal{M})$ is a monad in \mathcal{M} . Below it will be written as a pair (A, t) where A is the 0-cell part and $t : A \to A$ is the 1-cell part. Whenever needed, the multiplication and unit 2-cells will be denoted by $\mu : t.t \to t$ and $\eta : 1 \to t$, respectively, but in most cases they will not be explicitly written. A horizontal 1-cell in $Mnd(\mathcal{M})$ is a 1-cell in $Mnd(\mathcal{M})$, see Example 1.9. A vertical 1-cell in $Mnd(\mathcal{M})$ is a 1-cell in $Mnd_{op}(\mathcal{M})$, see again Example 1.9. A 2-cell in $Mnd(\mathcal{M})$ with boundaries as in the first diagram of

36

is a 2-cell in \mathcal{M} as in the second diagram, such that the third diagram of 2-cells in \mathcal{M} commutes. The horizontal composition of horizontal 1-cells is the composition of 1-cells in $\mathsf{Mnd}(\mathcal{M})$ and the vertical composition of vertical 1-cells is the composition of 1-cells in $\mathsf{Mnd}_{op}(\mathcal{M})$. The horizontal and vertical compositions of 2-cells are given by the same formulae as in (6.7) and (6.8), respectively.

A strict monoidal structure (\otimes, I) of \mathcal{M} induces a strict monoidal structure on $\mathbb{M}nd(\mathcal{M})$ analogously to Example 1.9, and a symmetry x on $(\mathcal{M}, \otimes, I)$ induces a symmetry of $\mathbb{M}nd(\mathcal{M})$ with horizontal and vertical transformations in the first two diagrams, and 2-cell part in the third diagram of

$$\begin{array}{cccc} (AB,ts) \xrightarrow{(x,1)} (BA,st) & (AB,ts) \xrightarrow{(hk,\Xi\Theta)} (A'B',t's') & (AB,ts) \xrightarrow{(x,1)} (BA,st) \\ (fg,\Phi\Gamma) & & \downarrow 1 & \downarrow (gf,\Gamma\Phi) & (x,1) & \downarrow 1 & \downarrow (x,1) & (x,1) & \downarrow 1 & \downarrow \\ (A'B',t's') \xrightarrow{(x,1)} (B'A',s't') & (BA,st) \xrightarrow{(kh,\Theta\Xi)} (B'A',s't') & (BA,st) \xrightarrow{(BA,st)} (BA,st) \end{array}$$

for any 1-cells $(h, \Xi) : (A, t) \to (A', t')$ and $(k, \Theta) : (B, s) \to (B', s')$ in $\mathsf{Mnd}(\mathcal{M})$ and $(f, \Phi) : (A, t) \to (A', t')$ and $(g, \Gamma) : (B, s) \to (B', s')$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$.

Any 2-functor $F : \mathcal{M} \to \mathcal{N}$ gives rise to a double functor $Mnd(F) : Mnd(\mathcal{M}) \to Mnd(\mathcal{N})$. On the 0-cells it acts as Mnd(F) of Example 1.9; equivalently, as $Mnd_{op}(F)$ of Example 1.9. On the horizontal and vertical 1-cells, Mnd(F) acts as Mnd(F) and $Mnd_{op}(F)$, respectively. On the 2-cells Mnd(F) acts as F. The double functor Mnd(F) is strict monoidal whenever F is strict monoidal and it is symmetric whenever F is so.

A 2-natural transformation $\omega : F \to G$ induces a double natural transformation $Mnd(\omega) : Mnd(F) \to Mnd(G)$ with horizontal and vertical transformations in the first two diagrams, and 2-cell part in the third diagram of

$$\begin{array}{c|c} (\mathsf{F}A,\mathsf{F}t) \xrightarrow{(\omega_{A},1)} (\mathsf{G}A,\mathsf{G}t) & (\mathsf{F}A,\mathsf{F}t) \xrightarrow{(\mathsf{F}h,\mathsf{F}\Xi)} (\mathsf{F}B,\mathsf{F}s) & (\mathsf{F}A,\mathsf{F}t) \xrightarrow{(\omega_{A},1)} (\mathsf{G}A,\mathsf{G}t) \\ \hline (\mathsf{F}f,\mathsf{F}\Phi) & & \downarrow 1 & \downarrow (\mathsf{G}f,\mathsf{G}\Phi) & (\omega_{A},1) & \downarrow 1 & \downarrow (\omega_{B},1) & (\omega_{A},1) & \downarrow 1 & \downarrow 1 \\ (\mathsf{F}B,\mathsf{F}s) \xrightarrow{(\omega_{B},1)} (\mathsf{G}B,\mathsf{G}s) & (\mathsf{G}A,\mathsf{G}t) \xrightarrow{(\mathsf{G}A,\mathsf{G}t)} (\mathsf{G}B,\mathsf{G}s) & (\mathsf{G}A,\mathsf{G}t) \xrightarrow{(\mathsf{G}A,\mathsf{G}t)} (\mathsf{G}A,\mathsf{G}t) \end{array}$$

for any 1-cells $(h, \Xi) : (A, t) \to (B, s)$ in $\mathsf{Mnd}(\mathcal{M})$ and $(f, \Phi) : (A, t) \to (B, s)$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$. It is symmetric monoidal whenever ω is monoidal.

With this we constructed a 2-functor $Mnd : sm-2Cat \rightarrow sm-DblCat$ which we want to relate next to the 2-functors $(-)_{01}$ of Example 1.10 and Example 6.12.

For any symmetric strict monoidal 2-category \mathcal{M} , a 0-cell of $\mathbb{M}nd(\mathcal{M})_{01}$ is a horizontal pseudomonoid in $\mathbb{M}nd(\mathcal{M})$. Since the horizontal 2-category of $\mathbb{M}nd(\mathcal{M})$ is $\mathbb{M}nd(\mathcal{M})$, this is the same as a pseudomonoid $((A, t, \mu, \eta), (m, \tau^2), (u, \tau^0), \alpha, \lambda, \varrho)$ in $\mathbb{M}nd(\mathcal{M})$; equivalently, a 0-cell of $\mathbb{M}nd(\mathcal{M})_{01}$. Thus the isomorphism of Section 2 takes it to the 0-cell $((A, m, u, \alpha, \lambda, \varrho), (t, \tau^2, \tau^0), \mu, \eta)$ of $\mathbb{M}nd(\mathcal{M}_{01})$; that is, a 0-cell of $\mathbb{M}nd(\mathcal{M}_{01})$ (which was termed an opmonoidal monad in \mathcal{M}).

In the same way, a horizontal 1-cell of $Mnd(\mathcal{M})_{01}$ is a 1-cell of $Mnd(\mathcal{M})_{01}$, taken by the isomorphism of Section 2 to a 1-cell of $Mnd(\mathcal{M}_{01})$; that is, a horizontal 1-cell of $Mnd(\mathcal{M}_{01})$.

GABRIELLA BÖHM

A vertical 1-cell of $Mnd(\mathcal{M})_{01}$ is an opmonoidal vertical 1-cell of $Mnd(\mathcal{M})$,

$$\begin{array}{cccc} (A,t) & (AA,tt) \xrightarrow{(m,\tau^2)} (A,t) & (I,1) \xrightarrow{(u,\tau^0)} (A,t) \\ & \downarrow^{(f,\Phi)} & (ff,\Phi\Phi) \downarrow & \Downarrow \varphi^2 & \downarrow^{(f,\Phi)} & \parallel & \Downarrow \varphi^0 & \downarrow^{(f,\Phi)} \\ (A',t') & (A'A',t't') \xrightarrow{(m',\tau'^2)} (A',t') & (I,1) \xrightarrow{(u',\tau'^0)} (A',t'). \end{array}$$

It means a 1-cell $(f, \Phi) : (A, t) \to (A', t')$ in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$ and a 1-cell $(f, \varphi^2, \varphi^0) : (A, m, u, \alpha, \lambda, \varrho) \to (A', m', u', \alpha', \lambda', \varrho')$ in \mathcal{M}_{01} such that in addition φ^2 and φ^0 are 2-cells of $\mathsf{Mnd}(\mathcal{M})$; that is, the following diagrams of 2-cells in \mathcal{M} commute.

Now these conditions can be interpreted, equivalently, as the opmonoidality of the 2cell $\Phi : (f, \varphi^2, \varphi^0).(t, \tau^2, \tau^0) \to (t', \tau'^2, \tau'^0).(f, \varphi^2, \varphi^0)$ in \mathcal{M}_{01} . This amounts to saying that $((f, \Phi), \varphi^2, \varphi^0)$ is a vertical 1-cell of $\mathbb{M}nd(\mathcal{M})_{01}$ if and only if $((f, \varphi^2, \varphi^0), \Phi)$ is a vertical 1-cell of $\mathbb{M}nd(\mathcal{M}_{01})$.

Finally, a 2-cell of $Mnd(\mathcal{M})_{01}$ is an opmonoidal 2-cell in $Mnd(\mathcal{M})$; that is, a 2-cell of \mathcal{M} as in the first diagram of

$$\begin{array}{cccc} A & \stackrel{h}{\longrightarrow} C & (A,t) & \stackrel{(h,\Xi)}{\longrightarrow} (C,z) & A & \stackrel{(h,\chi^{2},\chi^{0})}{\longrightarrow} C \\ f & & \downarrow \omega & \downarrow g & (f,\Phi) & \downarrow \omega & \downarrow (g,\Gamma) & (f,\varphi^{2},\varphi^{0}) & \downarrow \omega & \downarrow (g,\gamma^{2},\gamma^{0}) \\ B & \stackrel{}{\longrightarrow} D & (B,s) & \stackrel{}{\longrightarrow} (D,v) & B & \stackrel{}{\longrightarrow} D \end{array}$$

which is both a 2-cell in $\mathbb{M}nd(\mathcal{M})$ as in the second diagram and a 2-cell in \mathcal{M}_{01} as in the third diagram. Now these are the same conditions characterizing a 2-cell of $\mathbb{M}nd(\mathcal{M}_{01})$.

With this we constructed an iso double functor $Mnd(\mathcal{M})_{01} \to Mnd(\mathcal{M}_{01})$ which is symmetric strict monoidal and 2-natural. This proves the commutativity, up-to the above 2-natural isomorphism, of the first diagram of 2-functors in

$$\begin{array}{c|c} \mathsf{sm-2Cat} & \stackrel{\mathbb{M}\mathsf{nd}}{\longrightarrow} \mathsf{sm-DblCat} & \mathsf{sm-2Cat} & \stackrel{\mathbb{M}\mathsf{nd}}{\longrightarrow} \mathsf{sm-DblCat} \\ (-)_{01} & & & & & & & & \\ \mathsf{sm-2Cat} & & & & & & & & \\ \hline \mathbb{M}\mathsf{nd}} \mathsf{sm-DblCat} & & & & & & & \\ \mathsf{sm-2Cat} & & & & & & \\ \hline \mathbb{M}\mathsf{nd}} \mathsf{sm-DblCat}. \end{array}$$

Commutativity of the second diagram follows by symmetric steps (although it does not seem to result from any kind of abstract duality). Then Mnd commutes also with the 2-functors $(-)_{pq}$ of Example 1.10 (4) and Example 6.12 (4) for any non-negative integers p and q.

8. (p+q)-OIDAL EILENBERG-MOORE OBJECTS

In this final section we consider a symmetric strict monoidal 2-category \mathcal{M} which satisfies the assumptions in all parts of Proposition 5.3. Then we construct a symmetric strict monoidal double functor $\mathbf{K} : \mathbb{M}nd(\mathcal{M}) \to \mathbb{S}qr(\mathcal{M})$. Its horizontal 2-functor is the Eilenberg–Moore 2-functor $\mathsf{H} : \mathbb{M}nd(\mathcal{M}) \to \mathcal{M}$ and its vertical 2-functor is $\mathsf{V} : \mathbb{M}nd_{op}(\mathcal{M}) \to \mathcal{M}$ of Proposition 5.3. Throughout, the hypotheses of Proposition 5.3 (including those in parts (2) and (3)) are assumed to hold and the notation of Proposition 5.3 (and its proof) is used.

A 0-cell of $Mnd(\mathcal{M})$ is a monad (A, t) in \mathcal{M} hence it can be seen as a 0-cell in either 2-category $Mnd(\mathcal{M})$ or $Mnd_{op}(\mathcal{M})$. Then we may apply to it either one of the Eilenberg-Moore 2-functor $H : Mnd(\mathcal{M}) \to \mathcal{M}$ or the 2-functor $V : Mnd_{op}(\mathcal{M}) \to \mathcal{M}$ of Proposition 5.3. Both of them yield the Eilenberg-Moore object $\mathbf{K}(A, t) := A^t$.

A horizontal 1-cell of $Mnd(\mathcal{M})$ is a 1-cell (h, Ξ) of $Mnd(\mathcal{M})$ so it makes sense to put $\mathbf{K}(h, \Xi) := \mathsf{H}(h, \Xi)$. Similarly, a vertical 1-cell of $Mnd(\mathcal{M})$ is a 1-cell (f, Φ) of $Mnd_{op}(\mathcal{M})$ so we may put $\mathbf{K}(f, \Phi) := \mathsf{V}(f, \Phi)$.

For the definition of the action of the desired double functor \mathbf{K} on a 2-cell

universality of the coequalizer in the top row of

$$f^{v}.g.z.u^{z}.\mathsf{H}(h, \Xi) \xrightarrow{f^{v}.g.u^{z}.f^{z}.u^{z}.\mathsf{H}(h, \Xi)} \xrightarrow{f^{v}.g.u^{z}.\mathsf{H}(h, \Xi)} \xrightarrow{\pi(g,\Gamma).1} \mathsf{V}(g,\Gamma).\mathsf{H}(h, \Xi)$$

$$f^{v}.g.z.h.u^{t} \xrightarrow{f^{v}.u^{v}.f^{v}.g.u^{z}.\mathsf{H}(h, \Xi)} \xrightarrow{\epsilon^{v}.1.1.1} \bigvee \qquad \mathsf{V}(g,\Gamma).\mathsf{H}(h, \Xi)$$

$$f^{v}.g.z.h.u^{t} \xrightarrow{f^{v}.g.u^{z}.\mathsf{H}(h, \Xi)} \xrightarrow{f^{v}.g.h.u^{t}} \xrightarrow{f^{v}.g.h.u^{t}} \xrightarrow{\mathsf{I}(h, \Xi)} \bigvee \mathsf{V}(g,\Gamma).\mathsf{H}(h, \Xi)$$

$$f^{v}.g.z.h.u^{t} \xrightarrow{f^{v}.g.h.u^{v}.f^{v}.g.u^{z}.\mathsf{H}(h, \Xi)} \xrightarrow{f^{v}.g.h.u^{t}} \xrightarrow{\mathsf{I}(h, \Xi)} \bigvee \mathsf{V}(g,\Gamma).\mathsf{H}(h, \Xi)$$

$$f^{v}.g.h.t.u^{t} \xrightarrow{f^{v}.g.h.u^{v}.f^{v}.g.u^{z}.\mathsf{H}(h, \Xi)} \xrightarrow{f^{v}.g.h.u^{t}} \xrightarrow{\mathsf{I}(h, \Xi)} \xrightarrow{\mathsf{I}(h, E)} \bigvee \mathsf{I}(h, E)$$

$$H(h, E) \xrightarrow{\mathsf{I}(h, E)} \xrightarrow{\mathsf{I}(h, E)} \xrightarrow{\mathsf{I}(h, E)} \xrightarrow{\mathsf{I}(h, E)} \xrightarrow{\mathsf{I}(h, E)} \xrightarrow{\mathsf{I}(h, E)} \bigvee \mathsf{I}(h, E)$$

$$H(h, E) \xrightarrow{\mathsf{I}(h, E)} \xrightarrow{$$

is used, see (5.1). Note that for any 1-cell $(k, \Theta) : (B, s) \to (D, v)$ in $\mathsf{Mnd}(\mathcal{M})$ the multiplicativity condition

$$\begin{array}{c} v.v.k \xrightarrow{\mu.1} v.k \\ 1.\Theta \downarrow & \downarrow \Theta \\ v.k.s \xrightarrow{\Theta.1} k.s.s \xrightarrow{1.\mu} k.s \end{array}$$

$$(8.2)$$

holds, where $\mu : v.v \to v$ is the multiplication of the monad (D, v) and $\mu : s.s \to s$ is the multiplication of the monad (B, s). This can be interpreted as Θ being a 2-cell in $\mathsf{Mnd}(\mathcal{M})$ as on the left of

$$\begin{array}{ccc} (B,1) \xrightarrow{(k,1)} (D,1) & B \xrightarrow{k} D \\ (s,\mu) & & \downarrow \Theta & \downarrow (v,\mu) \\ (B,s) \xrightarrow{(k,\Theta)} (D,v) & B^s \xrightarrow{k} D \\ f^s & \downarrow H\Theta & \downarrow f^s \\ B^s \xrightarrow{k} D \\ f^s & \downarrow H\Theta & \downarrow f^s \\ H(k,\Theta) & D^v. \end{array}$$

Thus the Eilenberg-Moore 2-functor H takes it to a 2-cell of \mathcal{M} on the right. The resulting 2-cell H Θ occurs at the bottom of the vertical paths of (8.1).

In order for the 2-cell $\mathbf{K}\omega$ to be well-defined, the diagram of (8.1) should serially commute. The square on its left commutes if we take the upper ones of the parallel arrows: use that for the Eilenberg-Moore 2-functor H the diagram

commutes by the 2-naturality of the counit $(u^t, 1.\epsilon^t)$ of the Eilenberg–Moore 2-adjunction, and use the middle four interchange law in the 2-category \mathcal{M} . Commutativity of the square on the left of (8.1) with the lower ones of the parallel arrows is slightly more involved; it is checked in Figure 2. The region marked by (*) commutes because ω is a 2-cell of $\mathbb{M}nd(\mathcal{M})$. The image of the square marked by (**) under the functor $u^v.(-): \mathcal{M}(A^t, D^v) \to \mathcal{M}(A^t, D)$ commutes by (8.2). Hence this square commutes by the faithfulness of $u^v.(-): \mathcal{M}(A^t, D^v) \to \mathcal{M}(A^t, D)$.

We turn to showing that the so defined maps combine into a double functor \mathbf{K} . The maps in question preserve horizontal and vertical identity 1-cells, as well as the horizontal composition of horizontal 1-cells and the vertical composition of vertical 1-cells since the 2-functors H and V preserve identity 1-cells and compositions of 1-cells (see Proposition 5.3 about V). Preservation of identity 2-cells as on the left of




FIGURE 2. Serial commutativity of (8.1)

is obvious since then also the middle column of (8.1) is an identity 2-cell. For an identity 2-cell as on the right, (8.1) takes the form

$$\begin{array}{cccc} f^{z}.z.u^{z}.\mathsf{H}(h,\Xi) & & & f^{z}.u^{z}.\mathsf{H}(h,\Xi) & \xrightarrow{\epsilon^{z}.1} & \mathsf{H}(h,\Xi) \\ & & & & \\ & & & & \\ f^{z}.z.h.u^{t} & & & \\ f^{z}.h.t.u^{t} & & & \\ & & & \\ \mathsf{H}\Xi.1.1 & & & \\ \mathsf{H}(h,\Xi).f^{t}.t.u^{t} & & & \\ & & \mathsf{H}(h,\Xi).f^{t}.u^{t} & \xrightarrow{1.\epsilon^{t}} & \mathsf{H}(h,\Xi). \end{array}$$

The square on its right is taken by the functor $u^z(-) : \mathcal{M}(A^t, C^z) \to \mathcal{M}(A^t, C)$ to the commutative square of (8.3). Hence it commutes by the faithfulness of $u^z(-) :$ $\mathcal{M}(A^t, C^z) \to \mathcal{M}(A^t, C)$ so that also identity 2-cells of this kind are preserved by **K**. For 2-cells

$$(A, t) \xrightarrow{(h, \Xi)} (C, z) \qquad (C, z) \xrightarrow{(h', \Xi')} (E, y)$$

$$(n, \Phi) \downarrow \qquad \qquad \downarrow \omega \qquad \downarrow (g, \Gamma) \qquad \text{and} \qquad (g, \Gamma) \downarrow \qquad \downarrow \omega' \qquad \downarrow (l, \Lambda)$$

$$(B, s) \xrightarrow{(k, \Theta)} (D, v) \qquad (D, v) \xrightarrow{(k', \Theta')} (F, w)$$

of $Mnd(\mathcal{M})$, the 2-cells $(1.\mathbf{K}\omega)^{\cdot}(\mathbf{K}\omega'.1)$ and $\mathbf{K}((1.\omega)^{\cdot}(\omega'.1))$ are defined as the unique 2-cells making the respective diagrams of Figure 3 commute. The top and bottom rows of the diagrams of Figure 3 are equal epimorphisms. So from the equality of their left columns we infer the equality of their right columns. This proves that \mathbf{K} preserves the horizontal composition of 2-cells.

For 2-cells

$$\begin{array}{cccc} (A,t) & \xrightarrow{(h,\Xi)} (C,z) & (B,s) \xrightarrow{(k,\Theta)} (D,v) \\ (n,\Phi) & & \downarrow \omega & \downarrow (g,\Gamma) & \text{and} & (n',\Phi') & \downarrow \omega' & \downarrow (g',\Gamma') \\ (B,s) & \xrightarrow{(k,\Theta)} (D,v) & (E,y) \xrightarrow{(l,\Lambda)} (F,w) \end{array}$$

of $Mnd(\mathcal{M})$, the 2-cells $\mathbf{K}((\omega'.1)^{\cdot}(1.\omega))$ and $(\mathbf{K}\omega'.1)^{\cdot}(1.\mathbf{K}\omega)$ are defined as the unique 2-cells making commute the diagram of Figure 4 and the diagram of Figure 5, respectively. The top and bottom rows of the diagrams of Figure 4 and Figure 5 are equal epimorphisms. So from the equality of their left columns we infer the equality of their right columns. This proves that \mathbf{K} preserves the vertical composition of 2-cells. Here again, in Figure 4 the region marked by (*) commutes since ω' is a 2-cell of $Mnd(\mathcal{M})$ and the region marked by (**) commutes by the same reason as the region (**) of Figure 2.

Theorem 8.1. Consider a 2-category \mathcal{M} which admits Eilenberg–Moore construction. If the coequalizer (5.1) exists for all 1-cells in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$, and it is preserved by the horizontal composition on either side with any 1-cell in \mathcal{M} , then the following hold.

(1) The maps, constructed preceding the theorem, constitute a double functor \mathbf{K} : $\mathbb{M}nd(\mathcal{M}) \to \mathbb{S}qr(\mathcal{M}).$




FIGURE 3. K preserves the horizontal composition




FIGURE 4. K preserves the vertical composition — first part




FIGURE 5. K preserves the vertical composition — second part

GABRIELLA BÖHM

(2) If moreover \mathcal{M} is a strict monoidal 2-category which admits monoidal Eilenberg– Moore construction, then the double functor \mathbf{K} of part (1) is strict monoidal.

(3) If in addition the strict monoidal 2-category \mathcal{M} has a symmetry then the strict monoidal double functor \mathbf{K} of part (2) is symmetric as well.

Proof. We checked part (1) during the construction. Parts (2) and (3) clearly follow from the strict monoidality and the symmetry, respectively, of the 2-functors H and V under the stated conditions (see Proposition 3.2 and Proposition 5.3). \Box

In the setting of Theorem 8.1 (3), the object map of the induced symmetric strict monoidal double functor

$$\mathbb{M}\mathsf{nd}(\mathcal{M}_{pq}) \cong \mathbb{M}\mathsf{nd}(\mathcal{M})_{pq} \xrightarrow{\mathbf{K}_{pq}} \mathbb{S}\mathsf{qr}(\mathcal{M})_{pq} \cong \mathbb{S}\mathsf{qr}(\mathcal{M}_{pq})$$

sends a (p,q)-oidal monad (A,t) in \mathcal{M} to its (p+q)-oidal Eilenberg–Moore object A^t . While q ones of the pseudomonoid structures of A^t are liftings of the respective pseudomonoid structures of A along the 1-cell $u^t : A^t \to A$, the other p pseudomonoid structures of A along the corresponding pseudomonoid structures of A along the 1-cell $f^t : A \to A^t$.

Applying this to the particular symmetric strict monoidal 2-category of Remark 5.5, we obtain very similar results to [1, Corollary 6.4 and Theorem 9.1]. The only difference is that our assumptions are slightly stronger. Recall that a (p, q)-oidal object in the symmetric strict monoidal 2-category of Remark 5.5 is in fact a (p + q)-odial category with some further properties. They include, in particular, that all monoidal products preserve reflexive coequalizers — not only the first p ones as required in [1, Theorem 9.1].

Remark 8.2. As already discussed in the Introduction, a 2-category \mathcal{M} is said to admit Eilenberg-Moore construction if the inclusion 2-functor $\mathcal{M} \to \mathsf{Mnd}(\mathcal{M})$ possesses a right 2-adjoint H. As it is explained below, under the standing assumptions of Theorem 8.1 (see also Proposition 5.3), there is an analogous interpretation of the double functor $\mathbf{K} : \mathsf{Mnd}(\mathcal{M}) \to \mathsf{Sqr}(\mathcal{M})$ of Theorem 8.1 (1) as the right adjoint of an inclusion type double functor, in the sense of [15]. In the terminology of [15] this means that \mathbf{K} provides Eilenberg-Moore construction for the double category $\mathsf{Sqr}(\mathcal{M})$. We are grateful to an anonymous referee for raising this question.

(1) Any 0-cell of a 2-category \mathcal{M} can be seen as a monad with identity 1-cell part and identity multiplication and unit 2-cells. Any 1-cell in \mathcal{M} can be regarded as a monad morphism between these trivial monads — that is, as a horizontal 1-cell in $\mathbb{M}nd(\mathcal{M})$ — with identity 2-cell part. Symmetrically, it can be regarded as a vertical 1-cell in $\mathbb{M}nd(\mathcal{M})$ with identity 2-cell part. Finally, any 2-cell of $Sqr(\mathcal{M})$ yields a 2-cell in $\mathbb{M}nd(\mathcal{M})$ surrounded by the above trivial 1-cells. These maps constitute the inclusion double functor $\mathbf{I} : Sqr(\mathcal{M}) \to \mathbb{M}nd(\mathcal{M})$.

Assume now that \mathcal{M} admits Eilenberg–Moore construction; and that in \mathcal{M} the coequalizer (5.1) exists for all 1-cells in $\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})$ and it is preserved by the horizontal composition on either side with any 1-cell in \mathcal{M} . Then the double functor \mathbf{K} of Theorem 8.1 (1) is available and it turns out to be the right adjoint of the above inclusion double functor \mathbf{I} in the 2-category of double categories, double functors and *horizontal* transformations. The unit of the adjunction is trivial while the counit has

the horizontal 1-cell part and the 2-cell part

$$(A^{t},1) \xrightarrow{(u^{t},1.\epsilon^{t})} (A,t) \qquad g.u^{t} \xrightarrow{\eta^{s}.1.1} u^{s}.f^{s}.g.u^{t} \xrightarrow{1.\pi(g,\gamma)} u^{s}.\mathsf{V}(g,\gamma) \qquad (8.4)$$

respectively, for any monads (A, t) and (B, s) in \mathcal{M} , and any vertical 1-cell (g, γ) : $(A, t) \to (B, s)$ in $\mathbb{M}\mathsf{nd}(\mathcal{M})$.

However, we were not able to construct vertical transformations for the unit and counit of the adjunction $\mathbf{I} \dashv \mathbf{K}$, let alone double natural transformations.

(2) If in addition \mathcal{M} is a strict monoidal 2-category which admits monoidal Eilenberg– Moore construction then $\mathbf{I} \dashv \mathbf{K}$ of part (1) is an adjunction in the 2-category of strict monoidal double categories, strict monoidal double functors and monoidal horizontal transformations.

(3) If, moreover, the strict monoidal 2-category \mathcal{M} has a symmetry then $\mathbf{I} \dashv \mathbf{K}$ of part (2) is an adjunction in the 2-category of symmetric strict monoidal double categories, symmetric strict monoidal double functors and symmetric monoidal horizontal transformations.

(4) Recall that for any monad (A, t) in a 2-category \mathcal{M} admitting Eilenberg-Moore construction, the forgetful 1-cell $u^t : A^t \to A$ possesses a left adjoint f^t in \mathcal{M} (with unit and counit denoted by $\eta^t : 1 \to u^t f^t$ and $\epsilon^t : f^t . u^t \to 1$, respectively). Extending this observation, for the double functors of part (1) there is a vertical transformation $1 \to \mathbf{I}.\mathbf{K}$ with vertical 1-cell part and 2-cell part

$$\begin{array}{ccc} (A,t) \\ \downarrow (f^t,\epsilon^{t}.1) & f^s.h \xrightarrow{1.1.\eta^t} f^s.h.u^t.f^t = f^s.u^s.\mathsf{H}(h,\chi).f^t \xrightarrow{\epsilon^s.1.1} \mathsf{H}(h,\chi).f^t & (8.5) \\ (A^t,1) & \end{array}$$

respectively, for any monads (A, t) and (B, s) in \mathcal{M} , and any horizontal 1-cell (h, χ) : $(A, t) \to (B, s)$ in $\mathbb{M}nd(\mathcal{M})$. The transformations of (8.4) and (8.5) turn out to be orthogonal adjoints in the sense of [18] — termed conjoints in [11] — in the double category whose 0-cells are the double functors, horizontal and vertical 1-cells are the horizontal and vertical transformations, respectively, and whose 2-cells are the modifications in the sense of [18, Section 1.6]. By [18, Section 1.3] or [11, page 314] this means the existence of modifications

$$(A^{t}, 1) \xrightarrow{(u^{t}, 1, \epsilon^{t})} (A, t) \qquad (A, t) = (A, t)$$

$$\downarrow \downarrow \epsilon^{t} \downarrow (f^{t}, \epsilon^{t}, 1) \qquad (f^{t}, \epsilon^{t}, 1) \downarrow \downarrow \downarrow \eta^{t} \mid$$

$$(A^{t}, 1) = (A^{t}, 1) \qquad (A^{t}, 1) \xrightarrow{(u^{t}, 1, \epsilon^{t})} (A, t)$$

satisfying the triangle conditions requiring that

are identity 2-cells of \mathcal{M} .

In Section 6 we described a 2-functor Hor from the 2-category of double categories, double functors and horizontal transformations to 2Cat (it assigns to a double category its horizontal 2-category). Applying it to the adjunction $\mathbf{I} \dashv \mathbf{K}$ of Remark 8.2 (1) we re-obtain the inclusion 2-functor $\mathcal{M} \to \mathsf{Mnd}(\mathcal{M})$ and its right 2-adjoint H in Section 3; that is, the Eilenberg–Moore construction in \mathcal{M} . However, since from the same 2-category of double categories, double functors, and horizontal transformations there seems to be no analogous 2-functor taking the vertical 2-category of a double category, Remark 8.2 should have no message about the 2-functor V of Proposition 5.3 being a right 2-adjoint or not.

Remark 8.3. Whenever a strict monoidal 2-category \mathcal{M} admits monoidal Eilenberg– Moore construction, the 2-functor H of Section 3 is right adjoint of the inclusion 2-functor $J : \mathcal{M} \to \mathsf{Mnd}(\mathcal{M})$ in the 2-category of strict monoidal 2-categories, strict monoidal 2-functors and monoidal natural transformations. Then we may apply to this adjunction the 2-functor $(-)_{01}$ in item (i) of the list in Example 1.10 (1) so to infer that H_{01} is the right 2-adjoint of the inclusion

$$\mathcal{M}_{01} \xrightarrow{\mathsf{J}_{01}} \mathsf{Mnd}(\mathcal{M})_{01} \xrightarrow{\cong} \mathsf{Mnd}(\mathcal{M}_{01}).$$

This shows that under the stated assumptions also \mathcal{M}_{01} admits Eilenberg–Moore construction described by H_{01} . This argument can be found in [31, Theorem 5.1] for Cartesian monoidal 2-categories \mathcal{M} .

On the contrary, since the 2-functors Mnd and $(-)_{10}$ do not commute, no similarly simple argument seems available showing that under suitable circumstances also \mathcal{M}_{10} admits Eilenberg–Moore construction. However, an application of the adjunction of Remark 8.2 (2) yields an easy proof. The so obtained result extends Theorem 4.1 of [30]. The author is grateful to an anonymous referee for indicating this question.

As in Remark 8.2 (2), consider a strict monoidal 2-category \mathcal{M} which admits monoidal Eilenberg–Moore construction, and in which the coequalizer (5.1) exists for all 1-cells in $\mathsf{Mnd}_{op}(\mathcal{M})$ and it is preserved by the horizontal composition on either side with any 1-cell in \mathcal{M} .

In Example 6.12 (2) we constructed a 2-functor $(-)_{10}$ from the 2-category of strict monoidal double categories, strict monoidal double functors and monoidal horizontal transformations to the 2-category of double categories, double functors and horizontal transformations (see item (ii) of the list in Example 6.12 (1)). It takes the adjunction $\mathbf{I} \dashv \mathbf{K}$ of Remark 8.2 (2) to an adjunction $\mathbf{I}_{10} \dashv \mathbf{K}_{10}$ which induces a further adjunction

$$\mathbb{M}\mathrm{nd}(\mathcal{M}_{10}) \xrightarrow{\cong} \mathbb{M}\mathrm{nd}(\mathcal{M})_{10} \ \top \ \mathbb{S}\mathrm{qr}(\mathcal{M})_{10} \xrightarrow{\cong} \mathbb{S}\mathrm{qr}(\mathcal{M}_{10})$$

in the 2-category of double categories, double functors and horizontal transformations. Applying to it the 2-functor Hor to 2Cat in Section 6, we obtain a right 2-adjoint of the inclusion 2-functor $\mathcal{M}_{10} \to \mathsf{Mnd}(\mathcal{M}_{10})$, proving that under the stated assumptions \mathcal{M}_{10} admits Eilenberg–Moore construction. Even the explicit form of this right 2-adjoint can be read off this proof. Its action on a 0-cell; that is, the Eilenberg–Moore object of a monad $((A, m, u, \alpha, \lambda, \varrho), (t, \tau_2, \tau_0), \mu, \eta)$ in \mathcal{M}_{10} is equal to its image $(A^t = \mathsf{V}(A, t, \mu, \eta), \mathsf{V}(m, \tau_2), \mathsf{V}(u, \tau_0), \mathsf{V}\alpha, \mathsf{V}\lambda, \mathsf{V}\varrho)$ under the 2-functor

$$\mathsf{Mnd}_{\mathsf{op}}(\mathcal{M}_{10}) \xrightarrow{\cong} \mathsf{Mnd}_{\mathsf{op}}(\mathcal{M})_{10} \xrightarrow{\mathsf{V}_{10}} \mathcal{M}_{10}.$$

A 1-cell $((h, \chi_2, \chi_0), \Xi)$ of $\mathsf{Mnd}(\mathcal{M}_{10})$ is sent to the 1-cell $(\mathbf{K}(h, \Xi) = \mathsf{H}(h, \Xi), \mathbf{K}\chi_2, \mathbf{K}\chi_0)$ of \mathcal{M}_{10} and a 2-cell ω is taken to $\mathbf{K}\omega = \mathsf{H}\omega$. Thus it describes a lifting along the forgetful 1-cell $u^t : A^t \to A$.

If, furthermore, the strict monoidal 2-category \mathcal{M} has a symmetry then, based on Remark 8.2 (3), the above arguments can be iterated to obtain Eilenberg-Moore construction in \mathcal{M}_{pq} for any non-negative integers p, q.

References

- Marcelo Aguiar, Mariana Haim and Ignacio López Franco, Monads on higher monoidal categories, Applied Categorical Structures 26 no. 3 (2018) 413–458. 3, 4, 5, 11, 13, 15, 19, 46
- [2] Marcelo Aguiar and Swapneel Mahajan, Monoidal functors, species and Hopf algebras, CRM Monograph Series 29 American Mathematical Society 2010. 4, 10, 11
- [3] John C. Baez and Martin Neuchl, Higher-dimensional algebra. I. Braided monoidal 2-categories, Advances in Mathematics 121 no. 2 (1996) 196–244. 3
- [4] Michael Barr and Charles Wells, Toposes, triples and theories, Reprints in Theory and Applications of Categories 12, 2005. Originally published by Springer-Verlag, NewYork, 1985. 17
- [5] Gabriella Böhm, The Gray monoidal product of double categories, preprint available at arXiv:1901.10707.3
- [6] Francis Borceux, Handbook of Categorical Algebra: Volume 1, Basic category theory, Cambridge University Press, 1994. 5, 6, 7
- [7] Francis Borceux, Handbook of Categorical Algebra: Volume 2, Categories and Structures, Cambridge University Press, 1994. 17, 19
- [8] Roberto Bruni, José Meseguer and Ugo Montanari, Symmetric monoidal and cartesian double categories as a semantic framework for tile logic, Mathematical Structures in Computer Science 12 no. 1 (2002) 53–90. 3, 20, 23, 24, 25, 26
- [9] Dimitri Chikhladze, Stephen Lack and Ross Street, Hopf monoidal comonads, Theory and Applications of Categories 24 no. 19 (2010) 554–563. 2, 3, 4, 8
- [10] Brian Day and Ross Street, Monoidal bicategories and Hopf algebroids, Advances in Mathematics 129 no. 1 (1997) 99–157. 3
- [11] Robert Dawson, Robert Par and Dorette Pronk, The span construction, Theory and Applications of Categories 24 no. 13 (2010) 302–377. 47
- [12] Charles Ehresmann, Catégories structurées III. Quintettes et applications covariantes, in:
 "Séminaire Ehresmann. Topologie et géométrie différentielle" 5 (1963) 1–21. 3, 4, 20, 34
- [13] Samuel Eilenberg and G. Max Kelly, *Closed categories*, in: "Proceedings of the Conference on Categorical Algebra (La Jolla 1965)", Samuel Eilenberg, David Kent Harrison, Helmut Rhrl and Saunders MacLane (eds.), pp 421–562. Springer Verlag 1966. 24
- [14] Thomas M. Fiore, Nicola Gambino and Joachim Kock, Monads in double categories, Journal of Pure and Applied Algebra 215 no. 6 (2011) 1174–1197. 5, 36
- [15] Thomas M. Fiore, Nicola Gambino, and Joachim Kock. Double adjunctions and free monads, Cahiers de Topologie et Géométrie Différentielle Catégoriques, 53 no. 4 (1012) 242–306. 5, 46
- [16] Robert Gordon, Anthony John Power and Ross Street, Coherence for tricategories, Memoirs of the American Mathematical Society 117 no. 558 (1995). 3
- [17] Marco Grandis and Robert Paré, *Limits in double categories*, Cahiers de Topologie et Géométrie Différentielle Catégoriques, 40 no. 3 (1999) 162–220. 23, 24
- [18] Marco Grandis and Robert Paré, Adjoint for double categories, Cahiers de Topologie et Géométrie Différentielle Catégoriques, 45 no. 3 (2004) 193–240. 47

GABRIELLA BÖHM

- [19] John W. Gray, Formal category theory: adjointness for 2-categories, Lecture Notes in Mathematics 391 Springer-Verlag, Berlin-New York, 1974. 3
- [20] C. Barry Jay, Languages for monoidal categories, Journal of Pure and Applied Algebra 59 no. 1 (1989) 61–85. 3, 6
- [21] Mikhail M. Kapranov and Vladimir Voevodsky, 2-categories and Zamolodchikov tetrahedra equations, in: "Algebraic groups and their generalizations: quantum and infinite-dimensional methods" William J. Haboush, Brian J. Parshall (eds.) pp. 177–259 Proceedings of Symposia in Pure Mathematics 56, American Mathematical Society, Providence, RI 1994. 3
- [22] Stephen Lack, Limits for Lax Morphisms, Applied Categorical Structures 13 no. 3 (2005) 189– 203. 2
- [23] Fred E. J. Linton, Coequalizers in categories of algebras, In: "Seminar on Triples and Categorical Homology Theory" (ETH, Zürich, 1966/67) Beno Eckmann and Myles Tierny (eds.) pp 61–72 Reprints in Theory and Applications of Categories 18, 2008. Originally published by Springer, Berlin, 1969 (pp 75–90). 16, 19
- [24] Paddy McCrudden, Opmonoidal monads, Theory and Applications of Categories 10 no. 19 (2002) 469–485. 2, 4
- [25] Ieke Moerdijk, Monads on tensor categories, Journal of Pure and Applied Algebra 168 no. 2-3 (2002) 189–208. 2, 3
- [26] Christopher Schommer-Pries, The Classification of Two-Dimensional Extended Topological Field Theories. Ph. D. Dissertation, University of California, Berkeley, 2009. available at arXiv:1112.1000. 3
- [27] Gavin J. Seal, Tensors, monads and actions, Theory and Applications of Categories 28 no. 15 (2013) 403–434. 4, 19
- [28] Ross H. Street, The formal theory of monads, Journal of Pure and Applied Algebra 2 (1972) 149–168. 2, 4, 5, 7, 13, 14
- [29] Ross Street, Monoidal categories in, and linking, geometry and algebra, Bulletin of the Belgian Mathematical Society Simon Stevin 19 no. 5 (2012) 769–821. 4
- [30] Stanisław Szawiel and Marek Zawadowski, *Polynomial and Analytic Functors and Monads*, revisited, preprint available at arXiv:1506.04317. 48
- [31] Marek Zawadowski, The formal theory of monoidal monads, Journal of Pure and Applied Algebra 216 no. 8-9 (2012) 1932–1942. 2, 3, 8, 12, 13, 14, 48

WIGNER RESEARCH CENTRE FOR PHYSICS, H-1525 BUDAPEST 114, P.O.B. 49, HUNGARY *E-mail address*: bohm.gabriella@wigner.mta.hu