

Article

Brassica oleracea var. italica extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke

Rivan Virlando Suryadinata, Dwi Martha Nur Aditya, Viera Permatasari Wiana, Diana Rahman

*Faculty of Medicine, Universitas Surabaya (UBAYA), Surabaya, Indonesia***Abstract**

Introduction: Herbal extracts are often administered to cigarette smokers to prevent excessive free radicals. These include *Brassica oleracea* var. *Italica*, known to contain high antioxidant flavonoids and selenium micronutrients. Therefore, this study aims to determine the efficacy of *Brassica oleracea* var extract. *italica* in reducing the free radicals and inflammation present in experimental animals exposed to cigarette smoke.

Design and Methods: This research was conducted based on an experimental method using a randomized controlled trial (RCT) for 21 days. The animals used were divided into six groups (negative control, positive control, and four treatments). Particularly, the positive control and treatment groups were exposed to cigarette smoke for 2 minutes, twice a day, at 50 PPM CO levels. The treatment groups were administered the extract at different doses (0.5 ml; 0.75 ml; 1 ml; 1.25 ml), before assessing the blood level of malondialdehyde and C-Reactive Protein.

Result: The results showed the tendency for exposure to smoke to increase the number of free radicals and stimulate inflammation responses in the body ($P < 0,05$). In addition, a strong correlation between variables was established ($p = 0.000$; $r = 0.713$).

Conclusions: Broccoli extracts (*Brassica oleracea* L. var. *Italica*) administration has the potential to cause a decline in the two aspects, including free radicals and inflammation responses resulting from exposure to cigarette smoke.

Introduction

Tobacco cigarette is one of the products consumed through smoking. In addition, there has been an increase in popularity despite the negative health effects. Numerous countries have explored various means to stop the use,¹ but these efforts have not yielded significant results. Based on previous reports, one of the serious potential problems of every smoker is the elevated risk of disease infections, which is implicated in around 6 million deaths, worldwide. Despite the practice of abstinence for several years, smokers are known to have a sustained higher risk of infection with earlier onset than non-smokers. These are strongly associated with 10 years of shorter life expectancy.² The total number of

smokers worldwide is estimated to have reached 1 billion people and consist predominantly of men.³ In addition, the prevalence in developed countries, including America is relatively high, at about 15% of the total population, and consists more of people in the productive age groups.⁴ The number in developing countries, including Armenia, Laos and Indonesia, is comparably more, reaching over 50% of the total population.⁵

The increased mortality rate of smokers is evidence of the dangerous effects. Previous studies showed the development of various diseases, including lung cancer, respiratory tract, heart problems, and problems during pregnancy. Furthermore, about 40% of the total smoking population are known to die prematurely.⁶ The increase in complications from diseases have been attributed to the free radicals present in cigarette smoke and are known to directly stimulate the body's inflammatory response.⁷ Further exposure potentially instigates a decline in antioxidant levels and propagates immunosuppression.⁸ Previous studies also highlighted the tendency for cigarettes to initiate endothelial dysfunction as a result of excessive free radicals and inflammation.⁹

Physiologically, free radicals play an important role in cellular functions related to the enzymatic defence system assumed to prevent further accumulation. The presence of excessive radicals causes cellular changes, and consequently oxidative stress.^{10,11} The production of reactive oxygen species, including superoxide ($O_2 \cdot^-$), peroxynitrite ($ONOO \cdot^-$), and hydroxyl (OH) is possible during metabolism.¹² The enzymatic antioxidants, in form of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), and Catalase (CAT) function to neutralize these free radicals.¹³ Other variants are possibly derived from food intake, including non-enzymatic antioxidants.¹⁴ In addition, inflammation is another expression of the body's natural defence mechanism, which occurs in response to the entry of various foreign objects, including bacteria, viruses, allergens, toxic chemicals, and cigarettes. Pathogens or other hazardous substances continuously perpetuates chronic oxidative stress and consequently trigger protein oxidation, which releases pro-inflammatory cytokines and stimulates inflammatory responses all over the body.¹⁵

The intake of non-enzyme antioxidants is necessary to reduce excessive free radicals. These include polyphenols, a natural compound obtained from numerous fruits and vegetables, including broccoli. This constituent serves as a defence system from ultravi-

Significance for public health

The increasing number of active smokers is expected to cause serious health problems. This habit has been implicated in the increase in free radicals entering the respiratory tract, which stimulate an inflammatory process in the body. This process triggers cell damage and death, eventually causing health problems. Moreover, Brassica oleracea var. italica is known to contain high antioxidants, hence, there is a postulation of potential capacity to neutralize free radicals and reduce body inflammations.

oiet radiation as well as various pathogens and also has anti-inflammatory properties with the potential to confer cellular protection against oxidative stress.¹⁶ The broccoli plant (*Brassica oleracea* var. *italica*) is known to contain one of the polyphenol types termed flavonoids and considered to have higher amounts compared to other vegetables.^{17,18} Therefore, this research aims to determine the effectiveness of broccoli in reducing free radicals and inflammation resulting from exposure to cigarette smoke.

Design and Methods

This is an experimental research with a post-test control group design. This involved the use of male Wistar rats (*Rattus norvegicus*) as samples during a 21 days experiment (No:139/KE/X/2021). The animals were subdivided into 6 groups (one negative control group, one positive control group, and four treatment groups). In addition, basic maintenance and care were provided with reference to the 3R principle (Replacement, Reduction, and Refinement).

Kretek cigarettes, predominantly consumed by Indonesians, was used in this study, and is characterized by 2.4 mg nicotine and 38 mg tar. The smoke exposure was carried out two times daily for 21 days (each for 2 minutes with 50 PPM Carbon Monoxide levels measured using a Carbon Monoxide Meter device).

The broccoli was extracted through a maceration process using 96% ethanol as the solvent. Furthermore, the preparations were obtained by evaporating the extracts with a vacuum rotary evaporator. Based on Laurence and Bacharach's framework, a flavonoid present in Broccoli extract (18 mg/ml) was measured and evaluated by comparing it with the adequacy daily rate of (8 mg/day). Hence, a dose of 0.5 ml; 0.75 ml; 1 ml; 1.25 ml were derived and used for the experiment.

The negative control (Group I) consisted of experimental animals administered a daily intake, while the positive control (II) were only exposed to cigarette smoke. In addition, each of the four treatment groups (Group III, IV, V, VI) comprised of experimental animals exposed to cigarette smoke and the different doses of Broccoli extracts (0.5 ml; 0.75 ml; 1 ml; 1.25 ml). Therefore, the ratio data of malondialdehyde and C-Reactive Protein for each

group were obtained and analysed to ascertain the possible differences and relationships.

Results and Discussions

The results showed the mean levels of malondialdehyde in each group. Table 1 highlights the highest values (11.23 ± 1.45) in Group II, while the lowest (3.36 ± 0.74) were demonstrated in I. Furthermore, all data obtained were subjected to a normality test ($p > 0.05$) to ensure a well-modelled dataset is used, based on a normal distribution. The homogeneity test ($p = 0.018$) identified the dataset as non-homogeneous ($p < 0.05$). Therefore, differences in both results were calculated using the T-Test. Table 2 showed differences in level within each group, although no significant differences were observed between II and III ($p = 0.898$) as well as between Group I and V ($p = 0.465$).

The free radicals contained in cigarette smoke are assumed to instigate tissue damage through various mechanical processes, and consequently trigger the peroxidation of lipids, proteins, and DNA. This damage reportedly reduces the antioxidant defence system and stimulates the release of pro-inflammatory cytokines.¹⁹ In addition, an increase in the predominance of cellular damage from exposure to cigarette smoke that enters the respiratory tract directly elevates malondialdehyde levels in the blood. This compound is not considered a free radical, but the end product of lipid peroxidation.²⁰ The intrinsic characteristics include relatively strong stability and potential for use as an indicator of oxidative stress or body radical content.^{21,22}

Table 3 shows the highest levels of C-reactive protein (6.58 ± 0.58) in Group II, while I had the lowest values (4.44 ± 0.74). Therefore, a normality test ($P > 0.05$) was performed on all the data obtained to determine if the dataset was well-modelled by a normal distribution. The homogeneity test ($p = 0.831$) showed the samples as homogeneous ($p > 0.05$). The differences in both results for each group were then calculated using the ANOVA test, followed by the Least Significance Different (LSD) assessment.

In addition, free radicals entering the body potentially stimulate the phagocytosis process of the body's immune system. This

Table 1. Means of malondialdehyde between all groups.

Groups	Means \pm SD	Normality	Homogeneity
I A group only given daily intake	3.36 ± 0.74	0.754	0.018
II A group only exposed to cigarette smoke	11.23 ± 1.45	0.367	
III A group given daily intake, cigarette smoke, and 0.5 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	11.14 ± 1.62	0.580	
IV A group given daily intake, cigarette smoke, and 0.75 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	5.60 ± 0.94	0.237	
V A group given daily intake, cigarette smoke, and 1 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	3.87 ± 0.83	0.640	
VI A group given daily intake, cigarette smoke, and 1.25 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	1.90 ± 0.70	0.552	

Table 2. The results of T-test on malondialdehyde levels between all groups.

Groups	I	II	III	IV	V	VI
I	-	-	-	-	-	-
II	0.000	-	-	-	-	-
III	0.000	0.898	-	-	-	-
IV	0.003	0.000	0.000	-	-	-
V	0.465	0.000	0.000	0.019	-	-
VI	0.038	0.000	0.000	0.000	0.006	-

process stimulates the secretion of pro-inflammatory cytokines, including interleukin-1, interleukin-6, and tumour necrosis factor- α (TNF- α), known to be responsible for inflammatory responses.^{23,24,25} The reaction serves as a defence mechanism against various microorganisms or foreign objects, and there is a need to control this effect to avoid attacks by autoimmune diseases.²⁶ Furthermore, a gradual change in the cells surrounding the site is observed during chronic attacks, and potentially results in permanent damage.²⁷ These conditions are also possibly created after chronic cigarette exposure, which initiates a plaque build-up in the artery. This formation is consequently implicated in atherosclerosis, which leads to serious diseases, including cardiovascular and stroke.²⁸ The increase in free radicals is signalled by an elevation in malondialdehyde levels, which directly triggers inflammatory responses with negative health impacts.

The ANOVA analysis results showed differences in the levels of C-reactive protein in each group. These were further evaluated using the Least Significance Different (LSD) test. Table 4 showed a significant difference ($p < 0.05$) in Group I compared to others, while II demonstrated no substantial variation in contrast with III ($p = 0.917$) and Group IV ($p = 0.152$). However, no significant difference were observed between Group IV and III ($p = 0.182$), as well as V ($p = 0.139$) and VI ($p = 0.085$). The results also indicate no substantial variations between Group V and VI ($p = 0.794$). Table 5 depicts the relationship between increased malondialdehyde and C-reactive protein levels. This was ascertained using the Pearson test, and the results showed a strong correlation between both parameters ($r = 0.713$).

The external antioxidants or exogen antioxidants obtained from food possess different action mechanisms compared to enzymatic antioxidants, despite the intrinsic neutralizing capacity. These compounds are known to reduce free radicals in various ways, including through the protection of cells from lipid peroxidation (Vitamin E), as strong reducing agents to be potentially reduced by enzymes and glutathione (Vitamin C). Previous reports showed proficiency in boosting the immune system (β carotene) and polyphenol groups, known to act as both antioxidant and anti-inflammatory agents.^{29,30,31} In addition, non-enzymatic antioxidants reportedly reduce muscle pain and physical strain resulting from oxidative stress reactions.³² Broccoli (*Brassica oleracea* var. italica) contains various types of non-enzymatic antioxidants and high polyphenol, including flavonoids, which is estimated to reduce free radicals and inflammation caused by cigarette smoke exposure.^{33,34} This vegetable also contains selenium, and is known to potentially increase various enzymatic antioxidants, including superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px).³⁵

Conclusions

The conclusion of this study is exposure to cigarette smoke increases free radicals and stimulates an inflammatory response in the body. Broccoli (*Brassica oleracea* var. italica) extracts plays an intrinsic role in reducing these negative impacts, and the effect is dose-dependent.

Table 3. Means of C-reactive protein levels between all groups.

Groups	Means \pm SD	Normality	Homogeneity	P
I A group only given daily intake	4.44 \pm 0.74	0.530	0.831	0.000
II A group only exposed to cigarette smoke	6.58 \pm 0.58	0.642		
III A group given daily intake, cigarette smoke, and 0.5 ml of <i>Brassica oleracea</i> var. italica extract	6.54 \pm 0.50	0.384		
IV A group given daily intake, cigarette smoke, and 0.75 ml of <i>Brassica oleracea</i> var. italica extract	6.02 \pm 0.52	0.758		
V A group given daily intake, cigarette smoke, and 1 ml of <i>Brassica oleracea</i> var. italica extract	5.44 \pm 0.45	0.074		
VI A group given daily intake, cigarette smoke, and 1.25 ml of <i>Brassica oleracea</i> var. italica extract	5.34 \pm 0.74	0.248		

Table 4. The results of the Least Significance Different (LSD) test between all groups.

Groups	I	II	III	IV	V	VI
I	-	-	-	-	-	-
II	0.000	-	-	-	-	-
III	0.000	0.917	-	-	-	-
IV	0.000	0.152	0.182	-	-	-
V	0.014	0.006	0.008	0.139	-	-
VI	0.026	0.003	0.004	0.085	0.794	-

Table 5. The results of the Pearson test between all groups.

Groups	P	Pearson Test
Malondialdehyde	0.000	0.713
C-Reactive Protein		

Correspondence: Rivan Virlando Suryadinata, Faculty of Medicine, Universitas Surabaya (UBAYA), Surabaya, Jl. Tenggilis Mejoyo, KaliRungkut, Kec. Rungkut, Kota SBY, Jawa Timur 60293, Indonesia.

Tel.: +62.31.2981000. E-mail: rivan.virlando.s@staff.ubaya.ac.id

Key words: *Brassica oleracea*; C-reactive protein; free radical; inflammation; malondialdehyde.

Acknowledgement: The authors are sincerely grateful to the staff at the Faculty of Medicine, Universitas Surabaya (UBAYA) for the assistance rendered during this research.

Contributions: All authors contributed equally to this research.

Conflict of Interest: The author declare no conflicts of interest.

Funding: This research uses independent funds from all research members.

Clinical trials: This research was approved by the Health Research Ethics Committee of the University of Surabaya.

Availability of data and materials: All data generated or analyzed during this study are included in this published article.

Informed consent: Written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Conference presentation: Part of this paper was presented at the 2nd International Nursing and Health Sciences Symposium that took place at the Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.

Received for publication: 3 December 2021.

Accepted for publication: 10 May 2022.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Healthcare in Low-resource Settings 2023; 11(s1):11210

doi:10.4081/hls.2023.11210


Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

References

- Lorensia A, Muntu CM, Suryadinata RV, et al. Effect of lung function disorders and physical activity on smoking and non-smoking students. *J Prev Med Hygiene* 2021;62:E89-E96.
- West R. Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychol Health* 2017;32:1018–36.
- Drope J, Schluger N, Cahn Z, et al. The Tobacco Atlas. Atlanta: American Cancer Society and Vital Strategies. *Am Cancer Soc* 2018;26 p.
- Jamal A, King BA, Neff LJ, et al. Current Cigarette Smoking Among Adults — United States, 2005–2015. *MMWR Morb Mortal Wkly Rep* 2016;65:1205–1211.
- Saleheen D, Zhao W, Rasheed A. Epidemiology and Public Health Policy of Tobacco Use and Cardiovascular Disorders in Low- and Middle-Income Countries. *Arterioscler Thromb Vasc Biol* 2014;34:1811–9.
- National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention (US); 2014. PMID: 24455788.
- Suryadinata RV, Wirjatmadi B. Selenium Linked to Increased Antioxidant Levels and Decreased Free Radicals in Lung Tissue of Wistar Rats Exposed to E-Cigarette Smoke. *J Global Pharma Technol* 2020;12:32-39.
- Khanna A, Guo M, Mehra M, et al. Inflammation and Oxidative Stress Induced by Cigarette Smoke in Lewis Rat Brains. *J Neuroimmunol* 2013;254:69–75.
- Winkelmann BR, von Holt K, Unverdorben M. Smoking and atherosclerotic cardiovascular disease: Part I: Atherosclerotic disease process. *Biomarkers in Medicine* 2009;3:411-428.
- Suryadinata RV, Wirjatmadi B, Adriani M, et al. Effects of knowledge of vitamin D on attitudes toward sun exposure among middle-aged and elderly Indonesian adults. *Indian J Public Health Res Develop* 2018;9:11-15.
- Bardaweel SK, Gul M, Alzweiri M, et al. Reactive Oxygen Species: the Dual Role in Physiological and Pathological Conditions of the Human Body. *Eurasian J Med* 2018;50:193–201.
- Weng M, Xie X, Liu C, et al. The Sources of Reactive Oxygen Species and Its Possible Role in the Pathogenesis of Parkinson's Disease. *Parkinsons Dis* 2018;9:163040.
- Wang Y, Chun O, Song W. Plasma and dietary antioxidant status as cardiovascular disease risk factors: A review of human studies. *Nutrients* 2013;5:2969-3004.
- Suryadinata RV, Wirjatmadi B, Adriani M. Efektivitas penurunan malondialdehyde dengan kombinasi suplemen antioksidan superoxide dismutase melon dengan gliadin akibat paparan asap rokok. *Global Medical and Health Communication* 2017;5:79-83
- Biswas S, Das R, Banerjee ER. Role of free radicals in human inflammatory diseases. *AIMS Biophysics* 2017;4:596-614.
- Hussain T, Tan B, Yin Y, et al. Oxidative Stress and Inflammation: What Polyphenols Can Do for Us?. *Oxidative Medicine and Cellular Longevity* 2016;1-9.
- Sami FJ, Rahimah S. Uji aktivitas antioksidan ekstrak metanol bunga brokoli (*Brassica oleracea* L. var. *Italica*) dengan metode DPPH (2,2 diphenyl-1-picrylhydrazyl) dan metode ABTS (2,2 azinobis (3-etilbenzotiazolin)-6-asam sulfonat). [Test di attività antiossidante dell'estratto metanolo di fiori di broccoli (*Brassica oleracea* L. var. *Italica*) utilizzando i metodi DPPH (2,2 difenil-1-picrylhydrazyl) e ABTS (2,2 azinobis (3-etilbenzotiazolina)-6-solfonico)] *Jurnal Fitofarmaka Indonesia* 2015;2:107-110.
- Suryadinata RV, Sukarno DA, Sardjono SC, et al. Antioxidant activity in red mulberries on sperm development exposed by cigarette smoke. *Bali Med J* 2021;10:583-586.
- Suryadinata RV, Wirjatmadi B. The Molecular Pathways of Lung Damage by E-Cigarette in Experimental Mice. *Sultan Qaboos University Med J* 2021;1(1).
- Cherian DA, Peter T, Narayanan A, et al. Malondialdehyde as a Marker of Oxidative Stress in Periodontitis Patients. *J Pharm Bioallied Sci* 2019;11: S297–S300.
- Cui X, Gang J, Han H, et al. Relationship between free and total malondialdehyde, a well-established marker of oxidative stress, in various types of human biospecimens. *J Thorac Dis* 2018;10:3088–3097.
- Ito F, Sono Y, Ito T. Measurement and Clinical Significance of Lipid Peroxidation as a Biomarker of Oxidative Stress:

- Oxidative Stress in Diabetes, Atherosclerosis, and Chronic Inflammation. *Antioxidants* 2019;8:72.
23. Mantovani A, Dinarello C, Molgora M, et al. Interleukin-1 and Related Cytokines in the Regulation of Inflammation and Immunity. *Immunity* 2019;50:778-795.
 24. Borsini A, Benedetto M, Giacobbe J, et al. Pro- and Anti-Inflammatory Properties of Interleukin in Vitro: Relevance for Major Depression and Human Hippocampal Neurogenesis. *Int J Neuropsychopharmacol* 2020;23:738-750.
 25. Phaniendra A, Jestadi DB. Free Radicals: Properties, Sources, Targets, and Their Implication in Various Diseases. *Ind J Clin Biochem* 2015;30:11-26.
 26. Chen L, Deng H, Cui H, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget* 2018;9:7204-7218.
 27. Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010;140:805-20.
 28. Jee Y, Jung KJ, Lee S, et al. Smoking and atherosclerotic cardiovascular disease risk in young men: the Korean Life Course Health Study. *BMJ Open* 2019;9: e024453.
 29. Azlina MFN, Qodriyah MS, Kamisah Y. Tocopherol and Tocotrienol: Therapeutic Potential in Animal Models of Stress. *Curr Drug Targets* 2018;19:1456-1462.
 30. Peternej TT, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med* 2011;41:1043-69.
 31. Lohan SB, Vitt K, Scholz P, et al. ROS production and glutathione response in keratinocytes after application of β -carotene and VIS/NIR irradiation. *Chem Biol Interact* 2018;280:1-7.
 32. Simioni C, Zauli G, Martelli AM, et al. Oxidative stress: role of physical exercise and antioxidant nutraceuticals in adulthood and aging. *Oncotarget* 2018;9:17181-17198.
 33. Teixeira J, Chavarria D, Borges F, et al. Dietary Polyphenols and Mitochondrial Function: Role in Health and Disease. *Curr Med Chem* 2019;26:3376-3406.
 34. Suryadinata RV, Lorensia A, Sefania K. Effectiveness of Lime Peel Extract (*Citrus aurantifolia* Swingle) against C-Reactive Protein Levels in Alloxan-Induced Wistar Rats. *Global Medical and Health Communication* 2021;9:23-28.
 35. Suryadinata RV, Wirjatmadi G, Lorensia A. The time pattern of selenomethionine administration in preventing free radicals due to exposure to electric cigarette smoke. *J Public Health Res* 2021;10:2232.

HLS



Healthcare in Low-resource Settings

Volume 10 Number 1

January 2013

Editorial Board

Editor-in-Chief

Luigi Barberini

University of Cagliari and University Hospital of Cagliari, Italy | [Curriculum Vitae](#)

Editorial Board

Saurav Basu

Indian Institute of Public Health (Public Health Foundation of India), New Delhi, India

Mauro Carta

Department of Medical Sciences and Public Health, University of Cagliari, Italy

Vittorio Colizzi

University of Rome Tor Vergata, Italy and Faculty of Science & Technology, Evangelic University of Cameroon

Claudia Fattuoni

Department of Chemical and Geological Sciences, University of Cagliari, Italy

Simona Gurzu

George Emil Palade University of Medicine, Pharmacy, Sciences and Technology, Targu-Mures, Romania (UMFST)

Uzzam Ahmed Khawaja

Division of Pulmonary Medicine and Critical Care, Medicare Cardiac and General Hospital (MCGH), Karachi, Pakistan

Mohsen Khosravi

Department of Psychiatry and Clinical Psychology, Zahedan University of Medical Sciences, Zahedan, Iran

Roy Rillera Marzo

Department Chair of Public Health, International Medical School, Management and Science University, Malaysia

ert Nyenswah

Johns Hopkins University Bloomberg School of Public Health, United States

Salvatore Sardo

Dipartimento di Scienze Mediche e Sanità Pubblica, Università degli Studi di Cagliari, Italy

FOR AUTHORS

SUBMIT YOUR PAPER

[Guide for Authors](#)

[Benefits for Authors](#)

[How to write a scientific paper](#)

[How to write a Review article](#)

[Article Processing Charge](#)

FOR REVIEWERS

[Benefits for Reviewers](#)

[How to review](#)

[Thanks to Reviewers](#)

INDEXING

[DOAJ](#)

[ESCI](#)

MOST READ LAST MONTH

Efficacy and safety of Camosunate for the treatment of uncomplicated malaria in the University of Benin Teaching Hospital, Benin City, Nigeria

 85

Vol. 11 No. s1 (2023): 2nd International Nursing and Health Sciences Symposium

Published: 30-12-2022



ORIGINAL ARTICLES



The effects of TeamSTEPPS implementation by nurses on situation monitoring in hospital

Ahsan Ahsan, Imeldha Monitasari, Evi Harwiati Ningrum, Ike Nesdia Rahmawati, Linda Wieke Noviyanti, Kuswanto Rusca Putra

<https://doi.org/10.4081/hls.2023.11163>

0 0 0 0

471 PDF: 10

PDF



Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

Amelia Lorensia, Rivan Virlando Suryadinata

<https://doi.org/10.4081/hls.2023.11164>

0 0 0 0

430 PDF: 6

PDF



β -(1,3)-D-glucan from *Pleurotus ostreatus* correlates with lower plasma IL-6, IL-1 β , HOMA-IR, and higher pancreatic beta cell count in High-Fat and High-Fructose Diet (HFFD) rats

Alma Maghfrotun Innayah, Elvira Nur Sa'idah Hariani, Husnul Khotimah, Inggita Kusumastuty,

Ema Pristi Yunita, Dian Handayani

 <https://doi.org/10.4081/hls.2023.11165>

 0  0  0  0

 442  PDF: 11

 PDF



The effect of community of practice in improving the role of clinical instructor in patient safety implementation by prelicensure nursing students

Dyah Wiji Puspita Sari, Hanny Handiyani, Kuntarti Kuntarti, Yuni Sufyanti Arief

 <https://doi.org/10.4081/hls.2023.11166>

 0  0  0  0

 141  PDF: 6

 PDF



Identification of new cases of tuberculosis during the COVID-19 pandemic using model strategic management

Eppy Setiyowati, Umi Hanik, Ni Njoman Juliasih, Priyo Susilo

 <https://doi.org/10.4081/hls.2023.11167>

 0  0  0  0

 121  PDF: 8

 PDF



The effects of a 'COVID Nurse Assistant' application on patient satisfaction in COVID isolation rooms

Evi Harwiati Ningrum, Annisa Wuri Kartika, Ahmad Hasyim Wibisono, Ike Nesda Rahmawati, Linda Wieke Noviyanti, Ahsan Ahsan, Kuswantoro Rusca

Putra, Ungky Agus Setyawan, Lusia Titik
Andayani, Ririn Widayanti, Arif Jati Purnanto, Gatot
Subroto, Nurul Laili, Judith Anderson

 <https://doi.org/10.4081/hls.2023.11168>

 0  0  0  0

 111  PDF: 8

 PDF



The three month's dietary brown rice intervention has not significantly decreased levels of CRP, TNF- α , and IL6 of type 2 diabetes mellitus patients

Fajar Ari Nugroho, Inggita Kusumastuty, Anggun Rindang Cempaka, Atifa Nafia Hasantie Latif, Dian Handayani

 <https://doi.org/10.4081/hls.2023.11172>

 0  0  0  0

 114  PDF: 17

 PDF



Correlation of leg pain responses with ankle-brachial index and peripheral sensory responses in foot of type 2 diabetes mellitus

Heri Kristianto, Tina Handayani Nasution, Endah Panca Lidya Fatma, Efris Kartika Sari, Ahmad Hasyim Wibisono, Haryadi Kurniawan, Firdausy Ratna, Ica Cristiningtyas, Endang Listyowati

 <https://doi.org/10.4081/hls.2023.11173>

 0  0  0  0

 139  PDF: 14

 PDF



Sciences Symposium

The use of web-based digital media to enhance admission orientation for patients in the hospital

Ike Nesdia Rahmawati, Imeldha Monitasari, Debby Hamsa Putri, Delfira Arizda, Febry Pricila, Fitri Rosyidawati, Bagus Lanang Sejati

<https://doi.org/10.4081/hls.2023.11174>

0 0 0 0

94 PDF: 8

 PDF



The ethical principles and caring behavior of Indonesian nurses

Ilkafah Ilkafah, Anestasia Pangestu Mei Tyas, Rini Rachmawaty

<https://doi.org/10.4081/hls.2023.11176>

0 0 0 0

128 PDF: 8

 PDF



A cross-sectional study of the knowledge, skills, and 6 rights on medication administration by nurses at emergency department

Linda Wieke Noviyanti, Andri Junianto, Ahsan Ahsan

<https://doi.org/10.4081/hls.2023.11177>

0 0 0 0

120 PDF: 7

 PDF



The efficacy of Raspberry Pi-based automatic voice message education on knowledge level and prevention behavior of high-risk population

Mifetika Lukitasari, Wahyu Indah Windarti, Endah Panca Lydia Fatma, Tony Suharsono, Dwi Adi Nugroho

 <https://doi.org/10.4081/hls.2023.11178>

 0  0  0  0

 112  PDF: 6

 PDF



Patient's knowledge, gender, and physical activity level as the predictors of self-care in heart failure patients

Mifetika Lukitasari, Ulfa Fitriani Nafista

 <https://doi.org/10.4081/hls.2023.11179>

 0  0  0  0

 97  PDF: 6

 PDF



The correlation between family role and anemia prevention in pregnancy

Mira Triharini, Ni Ketut Alit Armini, Sekar Ayu Pitaloka

 <https://doi.org/10.4081/hls.2023.11180>

 0  0  0  0

 102  PDF: 4

 PDF



Relationship between postural stability and fall risk in young adult after lower limb muscle fatigue

Mohd Khairuddin Mohd Safee, Noor Azuan Abu Osman

 <https://doi.org/10.4081/hls.2023.11182>

 0  0  0  0

 94  PDF: 7

 PDF



Knowledge and peer support for increase Menstrual Hygiene Management (MHM) in adolescents

Ni Ketut Alit Armini, Arik Setyani, Aria Aulia Nastiti, Mira Triharini

 <https://doi.org/10.4081/hls.2023.11193>

 0  0  0  0

 101  PDF: 6

 PDF



Family health task implementation and the health status of diabetes mellitus patients: A correlational study

Niko Dima Kristianingrum Kristianingrum, Delfira Arizda, Setyoadi Setyoadi, Yati Sri Hayati, Michael Leo Parchman

 <https://doi.org/10.4081/hls.2023.11195>

 0  0  0  0

 104  PDF: 5

 PDF



Physical activity, exercise habits, and body mass index of adults

Ratna Candra Dewi, Bambang Wirjatmadi

 <https://doi.org/10.4081/hls.2023.11197>

 0  0  0  0

👁 90 📄 PDF: 8

📄 PDF



Correlation between family support with work productivity of people with severe mental disorders (PSMD) in Bantur Community Health Center, Malang, Indonesia

Renny Nova, Heni Dwi Windarwati, Erik Meidianto

🌐 <https://doi.org/10.4081/hls.2023.11207>

📄 0 ✅ 0 🔄 0 ❓ 0

👁 85 📄 PDF: 6

📄 PDF



A societal adaptation model as a novel approach toward the recovery of people with schizophrenia

Retno Lestari, Ah Yusuf, Febri Endra Budi Setyawan, Ahsan Ahsan, Rachmat Hargono

🌐 <https://doi.org/10.4081/hls.2023.11208>

📄 0 ✅ 0 🔄 0 ❓ 0

👁 97 📄 PDF: 6

📄 PDF



Mother's knowledge as a dominant factor for the success of exclusive breastfeeding in Indonesia

Rinik Eko Kapti, Yuni Sufyanti Arief, Nurona Azizah

🌐 <https://doi.org/10.4081/hls.2023.11209>

📄 0 ✅ 0 🔄 0 ❓ 0

👁 107 📄 PDF: 9

📄 PDF



Brassica oleracea var. italica extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke

Rivan Virlando Suryadinata, Dwi Martha Nur Aditya, Viera Permatasari Wiana, Diana Rahman

<https://doi.org/10.4081/hls.2023.11210>

0 0 0 0

71 PDF: 4

PDF



The effects of exclusive breastfeeding booklets on mothers' knowledge in providing exclusive breastfeeding in Mataram City, Indonesia

Septa Katmawanti, Farah Paramita, Agung Kurniawan, Siti Khuzaimah A Sharoni, Rosuzeita Fauzi, Intan Gumilang Pratiwi, Dea Aflah Samah, Yassinda Thasia Audina, Oktavia Sri Wahyuni, Meyralda Dara Adisa, Alinda Rahmani

<https://doi.org/10.4081/hls.2023.11211>

0 0 0 0

94 PDF: 9

PDF



The correlation between family support and health status in patients with diabetes mellitus

Setyoadi Setyoadi, Ah Yusuf, Niko Dima Kristianingrum, Yati Sri Hayati, Linda Wieke Noviyanti, Nurmalia Filda Syafiky

<https://doi.org/10.4081/hls.2023.11212>

0 0 0 0

85 PDF: 5

PDF



Experience of nurses using the basic nursing science in performing nursing care for patients at RSUD Dr. Soedomo Trenggalek, Indonesia

Rizki Bahtiyar Ardyansah, Shila Wisnasari, Titin Andri Wihastuti, Dina Dewi Sartika Lestari Ismail

<https://doi.org/10.4081/hls.2023.11213>

0 0 0 0

100 PDF: 8

PDF



The effectiveness of nursing psychoeducation toward family burden and quality life on caregiver of people with schizophrenia in the community

Suharsono Suharsono, Noor Faidah, Moh Hanafi

<https://doi.org/10.4081/hls.2023.11215>

0 0 0 0

90 PDF: 21

PDF



The implementation of code blue by nurses as first responders in outpatient and inpatient rooms at Malang Indonesia Hospital

Tony Suharsono, Sunarmi Sunarmi, Nur Ida, Bella Nove Khirria, Nazla Asrin, Ikhda Ulya

<https://doi.org/10.4081/hls.2023.11217>

0 0 0 0

83 PDF: 5

PDF

REVIEWS





Strategies to improving patient safety in hospitals

Mohamad Taji, Kuswantoro Rusca Putra, Dina Dewi Sartika Lestari Ismail

<https://doi.org/10.4081/hls.2023.11181>

0 0 0 0

98 PDF: 14

PDF



A systematic review on telenursing as a solution in improving the treatment compliance of tuberculosis patients in the COVID-19 pandemic

Ni Nyoman Elfiyunai, Nursalam Nursalam, Tintin Sukartini, Ferry Efendi

<https://doi.org/10.4081/hls.2023.11194>

0 0 0 0

118 PDF: 7

PDF



The effectiveness of self-care management in treating heart failure: A scoping review

Alfrina Hany, Ratih Arum Vatmasari

<https://doi.org/10.4081/hls.2023.11196>

0 0 0 0

82 PDF: 6 Supplementary Materials: 2

PDF

SUPPLEMENTARY MATERIALS



The application of the public health model in pregnancy coaching for preventing high-risk pregnancy

Sri Wahyuni, Apriliani Yulianti Wuriningsih,
Nursalam Nursalam, Yuni Sufyanti Arief

 <https://doi.org/10.4081/hls.2023.11214>

 0  0  0  0

FOR AUTHORS

 93  PDF: 7  Sup

SUBMIT YOUR PAPER

 **PDF**

Guide for Authors

 **SUPPLEMENTARY**

Benefits for Authors

How to write a scientific paper

How to write a Review article

Article Processing Charge

FOR REVIEWERS

Benefits for Reviewers

How to review

Thanks to Reviewers

INDEXING

DOAJ

ESCI

MOST READ LAST MONTH

Efficacy and safety of Camosunate for the treatment of uncomplicated malaria in the University of Benin Teaching Hospital, Benin City, Nigeria

 85

Color coding: a tool to enhance the quality of health care in low resource settings

64

Knowledge, attitude and practice of contraceptive use among female students of Dilla secondary and preparatory school, Dilla town, South Ethiopia, 2014

61

KEYWORDS



Healthcare in Low-resource Settings

is an Open Access, peer-reviewed journal published by PAGEPress®. Pavia, Italy. All credits and honors to PKP for their OJS.

eISSN: 2281-7824

About

About The Journal

Editorial Board

Info Submission

Submissions

Guidelines For Authors

Our App Is Free For Anyone!

