

Pace University

DigitalCommons@Pace

Faculty Papers

Dyson College of Arts & Sciences

6-2019

Crystal structure of hexa- μ -chlorido- μ 4-oxido-tetrakis{[1-(2-hydroxyethyl)-2-methyl-5-nitro-1H-imidazole- κ N3]copper(II)} containing short NO₂ · · · NO₂ contacts

Ja-Shin Wu

Pace University

Daniel G. Shlian

Columbia University

Joshua H. Palmer

Columbia University

Rita K. Upmacis

Pace University

Follow this and additional works at: https://digitalcommons.pace.edu/dyson_fac

Recommended Citation

Wu, Ja-Shin; Shlian, Daniel G.; Palmer, Joshua H.; and Upmacis, Rita K., "Crystal structure of hexa- μ -chlorido- μ 4-oxido-tetrakis{[1-(2-hydroxyethyl)-2-methyl-5-nitro-1H-imidazole- κ N3]copper(II)} containing short NO₂ · · · NO₂ contacts" (2019). *Faculty Papers*. 1.

https://digitalcommons.pace.edu/dyson_fac/1

This Article is brought to you for free and open access by the Dyson College of Arts & Sciences at DigitalCommons@Pace. It has been accepted for inclusion in Faculty Papers by an authorized administrator of DigitalCommons@Pace. For more information, please contact nmcguire@pace.edu.



Crystal structure of hexa- μ -chlorido- μ_4 -oxido-tetrakis{[1-(2-hydroxyethyl)-2- methyl-5-nitro-1*H*-imidazole- κN^3]copper(II)} containing short NO₂· · · NO₂ contacts

Ja-Shin Wu, Daniel G. Shlian, Joshua H. Palmer and Rita K. Upmacis

Acta Cryst. (2019). E75, 1057–1060



IUCr Journals

CRYSTALLOGRAPHY JOURNALS ONLINE

This open-access article is distributed under the terms of the Creative Commons Attribution Licence
<http://creativecommons.org/licenses/by/4.0/legalcode>, which permits unrestricted use, distribution, and
reproduction in any medium, provided the original authors and source are cited.



Crystal structure of hexa- μ -chlorido- μ_4 -oxido-tetra-kis{[1-(2-hydroxyethyl)-2-methyl-5-nitro-1H-imidazole- κN^3]copper(II)} containing short NO₂ \cdots NO₂ contacts

Ja-Shin Wu,^a Daniel G. Shlian,^b Joshua H. Palmer^b and Rita K. Upmacis^{c*}

Received 12 February 2019

Accepted 16 June 2019

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

Keywords: crystal structure; tetranuclear copper; metronidazole; bridging chloride; NO₂ interactions.

CCDC reference: 1923275


Supporting information: this article has supporting information at journals.iucr.org/e

^aDepartment of Chemistry & Physical Sciences, Pace University, New York, NY 10038, USA, ^bDepartment of Chemistry, Columbia University, New York, NY 10027, USA, and ^cDept. of Chemistry & Physical Sciences, Pace University, New York, NY 10038, USA. *Correspondence e-mail: rupmacis@pace.edu


The title tetranuclear copper complex, [Cu₄Cl₆O(C₆H₉N₃O₃)₄] or [Cu₄Cl₆O(MET)₄] [MET is 1-(2-hydroxyethyl)-2-methyl-5-nitro-1H-imidazole or metronidazole], contains a tetrahedral arrangement of copper(II) ions. Each copper atom is also linked to the other three copper atoms in the tetrahedron *via* bridging chloride ions. A fifth coordination position on each metal atom is occupied by a nitrogen atom of the monodentate MET ligand. The result is a distorted CuCl₃NO trigonal–bipyramidal coordination polyhedron with the axial positions occupied by oxygen and nitrogen atoms. The extended structure displays O–H \cdots O hydrogen bonding, as well as unusual short O \cdots N interactions [2.775 (4) Å] between the nitro groups of adjacent clusters that are oriented perpendicular to each other. The scattering contribution of disordered water and methanol solvent molecules was removed using the SQUEEZE procedure [Spek (2015). *Acta Cryst. C*71, 9–16] in PLATON [Spek (2009). *Acta Cryst. D*65, 148–155].

1. Chemical context

Metronidazole (C₆H₉N₃O₃; MET) is a medication that was discovered to be effective against both bacteria and parasites more than 50 years ago (Samuelson, 1999). MET is currently incorporated in the World Health Organization (WHO) list of essential medicines, *i.e.* medications that are considered to be effective and safe to meet the most important needs in a health system (WHO, 2015). Despite the widespread use of MET as a drug, relatively little structural data concerning its interactions with metal ions exist, and there are few structurally characterized copper compounds of MET (Galván-Tejada *et al.*, 2002; Barba-Behrens *et al.*, 1991; Athar *et al.*, 2005; Ratajczak-Sitarz *et al.*, 1998; Bharti *et al.*, 2002). Our recent work has sought to develop further metal–MET chemistry and we have reported structures containing Cu (Palmer *et al.*, 2015; Quinlivan & Upmacis, 2016), as well as Ag (Palmer & Upmacis, 2015) and Au (Quinlivan *et al.*, 2015). Tetranuclear copper(II) compounds of the form [Cu₄O_xL₄] are relatively well known, with the first example described in 1996 (Bertrand & Kelley, 1966). In this regard, although the structure of a [Cu₄O_xL₄] structure, where L = imidazole, has been previously described (Atria *et al.*, 1999), a counterpart containing L = MET has not been reported. Herein, we describe the structure of a tetranuclear Cu–MET complex [Cu₄Cl₆O(MET)₄] that is obtained by the reaction of anhydrous copper(I) chloride with MET in MeOH under aerobic conditions.



OPEN ACCESS



2. Structural commentary

The structure of the $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$ complex is shown in Fig. 1. Four copper atoms are arranged around an oxygen atom in a tetrahedral fashion, with $\text{Cu}–\text{O}$ distances ranging from 1.8960 (18) to 1.913 (2) Å. The $\text{Cu}–\text{O}–\text{Cu}$ angles range from 108.36 (10) to 110.80 (9)°, indicating a fairly uniform tetrahedron with little distortion. In fact, the degree of distortion from a tetrahedral arrangement can be readily quantified by the τ_4 four-coordinate geometry index that is reported and discussed elsewhere (Yang *et al.*, 2007; Palmer *et al.*, 2015; Brescia *et al.*, 2018). Briefly, τ_4 is obtained from the expression, $\tau_4 = [360 - (\alpha + \beta)]/141$, where α and β represent the two largest angles; a τ_4 value of 1.00 indicates an idealized tetrahedral geometry, whereas a value of 0.00 indicates an idealized square-planar geometry. In the title complex, $\alpha = 110.80$ (9)° and $\beta = 109.55$ (9)°, such that τ_4 is 0.990, which indicates negligible deviation from a tetrahedral geometry for oxygen (Yang *et al.*, 2007).




Figure 1

The molecular structure of $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$. For clarity, hydrogen atoms have been omitted. The ethoxy group of the MET ligand attached to Cu3 (comprising C34, C35 and O31) is disordered over two sets of sites in a 0.515 (19):0.485 (19) ratio.

Each of the four copper atoms is linked to the other three copper atoms *via* three chloride bridges, with the $\text{Cu}–\text{Cl}$ bridging distances varying from 2.3579 (10) to 2.4435 (9) Å (for $\text{Cu}2–\text{Cl}6$ and $\text{Cu}1–\text{Cl}2$, respectively). Each copper atom is also bound to a nitrogen atom of a MET ligand. The $\text{Cu}–\text{N}$ lengths range from 1.949 (2) to 1.972 (3) Å (for $\text{Cu}1–\text{N}11$ and $\text{Cu}4–\text{N}41$, respectively). Thus, each copper atom sits within a trigonal-bipyramidal arrangement, with the oxygen and nitrogen atoms forming the axial coordination points, and the bridging chloride ligands occupying the equatorial plane. The trigonal-bipyramidal structure is somewhat distorted, as indicated by the fact that the $\text{O}–\text{Cu}–\text{N}$ angles are less than 180°, ranging from 173.12 (10) to 176.91 (10)° (for $\text{O}1–\text{Cu}1–\text{N}11$ and $\text{O}1–\text{Cu}2–\text{N}21$, respectively), and the $\text{Cl}–\text{Cu}–\text{Cl}$ angles differ significantly from 120°, ranging from 109.97 (3) to 134.02 (3)° (for $\text{Cl}2–\text{Cu}2–\text{Cl}4$ and $\text{Cl}3–\text{Cu}1–\text{Cl}2$, respectively). Furthermore, the $\text{O}–\text{Cu}–\text{Cl}$ angles are all less than 90°, ranging from 83.33 (6) to 86.13 (6)° (for $\text{O}1–\text{Cu}1–\text{Cl}2$ and $\text{O}1–\text{Cu}–\text{Cl}1$, respectively), indicating that the equatorial chloride ligands are displaced slightly more towards the axial oxygen atom in the center of the molecule, than towards the nitrogen-containing ligand in the opposite axial position.

The τ_5 geometry index is a general descriptor of five-coordinate molecules and provides a way to determine the extent of distortion of a molecule from trigonal bipyramidal to square pyramidal (Addison *et al.*, 1984). The τ_5 geometry index is calculated by using the equation: $\tau_5 = (\beta - \alpha)/60$, where $\beta - \alpha$ is the difference between the two largest angles (Addison *et al.*, 1984; Palmer & Parkin, 2014). The values for τ_5 are calculated to be 0.65 (Cu1), 0.74 (Cu2), 0.84 (Cu3) and 0.73 (Cu4) for the five-coordinate copper centers, giving an average τ_5 value of 0.74. The τ_5 values obtained indicate that the copper-centered structures are closer to an idealized trigonal-bipyramidal (1.00) than a square-pyramidal (0.00).




Figure 2

Unit-cell packing of $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$ viewed down [100].

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O41—H41A···O31 ⁱ	0.89 (2)	2.13 (3)	2.738 (8)	125 (2)

Symmetry code: (i) $x + \frac{1}{2}, y - \frac{1}{2}, z$.

3. Supramolecular features

Fig. 2 shows the packing in the unit cell. As well as the O—H···O hydrogen bonds shown in Table 1, O11—H11A and O21—H21A probably form links to the disordered solvent molecules removed with SQUEEZE (see *Experimental*). The most interesting observation is the existence of short O···N interactions between the N13/O12/O13 and N33/O32/O33 nitro groups of adjacent clusters that are oriented perpendicular to each other, as illustrated in Fig. 3 with $\text{O}12\cdots\text{N}33 = 2.775 (4)$ \AA . This type of contact has previously been described as an $\text{O}_{\text{NO}_2}\cdots\pi(\text{N})_{\text{NO}_2}$ interaction (Daszkiewicz, 2013); such contacts are typically shorter than 3 \AA .

4. Database survey

The tetranuclear copper motif, $L_4\text{Cu}_4\text{Cl}_6\text{O}$, where L is a nitrogen-containing Lewis base ligand, is common. For instance, several structures have been reported in which L contains either an imidazole or substituted imidazole moiety (Clegg *et al.*, 1988; Norman *et al.*, 1989; Erdonmez *et al.*, 1990; Atria *et al.*, 1999; Cortés *et al.*, 2006; Chiarella *et al.*, 2009, 2010; She *et al.*, 2010) or a benzimidazole moiety (Tosik *et al.*, 1991; Zhang *et al.*, 2003; Jian *et al.*, 2004; Li *et al.*, 2011).

The title compound $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$ contains Cu—X distances that are similar to those in $[\text{Cu}_4\text{Cl}_6\text{O}(\text{imidazole})_4]$ (Atria *et al.*, 1999). For example, the Cu—O distances in $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$ are 1.8960 (18)–1.913 (2) \AA , compared to 1.903 (4)–1.924 (4) \AA for $[\text{Cu}_4\text{Cl}_6\text{O}(\text{imidazole})_4]$. Likewise, the Cu—Cl distances in $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$ are 2.3579 (10)–

Table 2
Experimental details.

Crystal data	
Chemical formula	$[\text{Cu}_4\text{Cl}_6\text{O}(\text{C}_6\text{H}_9\text{N}_3\text{O}_3)_4]$
M_r	1167.51
Crystal system, space group	Monoclinic, $C2/c$
Temperature (K)	130
a, b, c (\AA)	22.125 (3), 13.361 (2), 32.633 (5)
β ($^\circ$)	94.752 (2)
V (\AA^3)	9613 (3)
Z	8
Radiation type	Mo $K\alpha$
μ (mm^{-1})	2.14
Crystal size (mm)	0.36 × 0.20 × 0.10
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2008)
T_{\min}, T_{\max}	0.586, 0.746
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	78050, 15003, 11100
R_{int}	0.048
$(\sin \theta/\lambda)_{\text{max}}$ (\AA^{-1})	0.720
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.045, 0.118, 1.03
No. of reflections	15003
No. of parameters	579
No. of restraints	120
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ ($e \text{\AA}^{-3}$)	1.55, −1.09

Computer programs: *APEX2* and, *SAINT* (Bruker, 2008), *SHELXS97* (Sheldrick 2008), *SHELXL2014* (Sheldrick, 2015) and *SHELXTL* (Sheldrick, 2008).

2.4435 (9) \AA , compared to 2.374 (2)–2.564 (2) \AA for $[\text{Cu}_4\text{Cl}_6\text{O}(\text{imidazole})_4]$. Moreover, the Cu—N distances in $[\text{Cu}_4\text{Cl}_6\text{O}(\text{MET})_4]$ are 1.949 (2)–1.972 (3) \AA , compared to 1.934 (6)–1.961 (6) \AA .

5. Synthesis and crystallization

Anhydrous copper(I) chloride (0.015 g, 0.00015 mol) was mixed with MET (0.05075 g, 0.00030 mol) in methanol (2 ml) in a glass vial, forming a dark olive-colored solution. After allowing the solution to evaporate for eight days, gold-colored plates, suitable for X-ray diffraction, were obtained.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. Hydrogen atoms on carbon were placed in calculated positions (C—H = 0.95–1.00 \AA) and included as riding contributions with isotropic displacement parameters $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{Csp}^2)$ or $1.5U_{\text{eq}}(\text{Csp}^3)$. Atoms C34, C35 and O31 and their attached H atoms were modeled as disordered over two sets of sites in a 0.515 (19):0.485 (19) ratio. The structure contains two methanol molecules and one water molecule, but they are disordered and were removed by the SQUEEZE procedure in *PLATON* (Spek, 2015); the stated crystal data (M_r , μ , etc.) only refer to the main molecule.




Figure 3
Detail of the O···N interaction between the nitro groups of adjacent clusters.

Acknowledgements

RKU thanks Pace University for research support. Gerard Parkin (Columbia University) is thanked for helpful discussions.

References

- Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
- Athar, F., Husain, K., Abid, M., Agarwal, S. M., Coles, S. J., Hursthouse, M. B., Maurya, M. R. & Azam, A. (2005). *Chem. Biodivers.* **2**, 1320–1330.
- Atria, A. M., Vega, A., Contreras, M., Valenzuela, J. & Spodine, E. (1999). *Inorg. Chem.* **38**, 5681–5685.
- Barba-Behrens, N., Mutio-Rico, A. M., Joseph-Nathan, P. & Contreras, R. (1991). *Polyhedron*, **10**, 1333–1341.
- Bertrand, J. A. & Kelley, J. A. (1966). *J. Am. Chem. Soc.* **88**, 4746–4747.
- Bharti, N., Shailendra, Coles, S. J., Hursthouse, M. B., Mayer, T. A., Garza, M. G., Cruz-Vega, D. E., Mata-Cardenas, B. D., Naqvi, F., Maurya, M. R. & Azam, A. (2002). *Helv. Chim. Acta*, **85**, 2704–2712.
- Brescia, T. K., Mulosmani, K., Gulati, S., Athanasopoulos, D. & Upmacis, R. K. (2018). *Acta Cryst. E***74**, 309–312.
- Bruker (2008). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chiarella, G. M., Melgarejo, D. Y. & Fackler, J. P. Jr (2009). *Acta Cryst. C***65**, m228–m230.
- Chiarella, G. M., Melgarejo, D. Y., Prosvirin, A. V., Dunbar, K. R. & Fackler, J. P. (2010). *J. Clust Sci.* **21**, 551–565.
- Clegg, W., Nicholson, J. R., Collison, D. & Garner, C. D. (1988). *Acta Cryst. C***44**, 453–461.
- Cortés, P., Atria, A. M., Garland, M. T. & Baggio, R. (2006). *Acta Cryst. C***62**, m311–m314.
- Daszkiewicz, M. (2013). *CrystEngComm*, **15**, 10427–10430.
- Erdonmez, A., van Diemen, J. H., de Graaff, R. A. G. & Reedijk, J. (1990). *Acta Cryst. C***46**, 402–404.
- Galván-Tejada, N., Bernès, S., Castillo-Blum, S. E., Nöth, H., Vicente, R. & Barba-Behrens, N. (2002). *J. Inorg. Biochem.* **91**, 339–348.
- Jian, F. F., Zhao, P. S., Wang, H. X. & Lu, L. D. (2004). *Bull. Kor. Chem. Soc.* **25**, 673–675.
- Li, H., Jiang, H. & Sun, H. (2011). *Acta Cryst. E***67**, m1372.
- Norman, R. E., Rose, N. J. & Stenkamp, R. E. (1989). *Acta Cryst. C***45**, 1707–1713.
- Palmer, J. H. & Parkin, G. (2014). *Dalton Trans.* **43**, 13874–13882.
- Palmer, J. H. & Upmacis, R. K. (2015). *Acta Cryst. E***71**, 284–287.
- Palmer, J. H., Wu, J. S. & Upmacis, R. K. (2015). *J. Mol. Struct.* **1091**, 177–182.
- Quinlivan, P. J. & Upmacis, R. K. (2016). *Acta Cryst. E***72**, 1633–1636.
- Quinlivan, P. J., Wu, J.-S. & Upmacis, R. K. (2015). *Acta Cryst. E***71**, 810–812.
- Ratajczak-Sitarz, M., Katrusiak, A., Wojakowska, H., Januszczak, M., Krzyminiewski, R. & Pietrzak, J. (1998). *Inorg. Chim. Acta*, **269**, 326–331.
- Samuelson, J. (1999). *Antimicrob. Agents Chemother.* **43**, 1533–1541.
- She, G., Liu, S. Y., Wu, X. M., Guo, J. H., Wang, X. G. & Liu, Q. X. (2010). *Chin. J. Inorg. Chem.* **26**, 515–520.
- Sheldrick, G. M. (2008). *Acta Cryst. A***64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst. C***71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst. D***65**, 148–155.
- Spek, A. L. (2015). *Acta Cryst. C***71**, 9–18.
- Tosik, A., Bukowska-Strzyzewska, M. & Mrozinski, J. (1991). *J. Coord. Chem.* **24**, 113–125.
- WHO (2015). *Who. Tech. Rep. Ser.* **994**, 1–546.
- Yang, L., Powell, D. R. & Houser, R. P. (2007). *Dalton Trans.* pp. 955–964.
- Zhang, Y.-Q., Xu, D.-J. & Su, J.-R. (2003). *Acta Cryst. E***59**, m919–m920.

supporting information

Acta Cryst. (2019). E75, 1057-1060 [https://doi.org/10.1107/S2056989019008570]

Crystal structure of hexa- μ -chlorido- μ_4 -oxido-tetrakis{[1-(2-hydroxyethyl)-2-methyl-5-nitro-1*H*-imidazole- κN^3]copper(II)} containing short NO₂···NO₂ contacts


Ja-Shin Wu, Daniel G. Shlian, Joshua H. Palmer and Rita K. Upmacis

Computing details

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Hexa- μ -chlorido- μ_4 -oxido-tetrakis{[1-(2-hydroxyethyl)-2-methyl-5-nitro-1*H*-imidazole- κN^3]copper(II)}

Crystal data



$M_r = 1167.51$

Monoclinic, $C2/c$

$a = 22.125$ (3) Å

$b = 13.361$ (2) Å

$c = 32.633$ (5) Å

$\beta = 94.752$ (2)°

$V = 9613$ (3) Å³

$Z = 8$

$F(000) = 4688$

$D_x = 1.613 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9836 reflections

$\theta = 2.2\text{--}29.8^\circ$

$\mu = 2.14 \text{ mm}^{-1}$

$T = 130$ K

Plate, gold

0.36 × 0.20 × 0.10 mm

Data collection

Bruker APEXII CCD

diffractometer

φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2008)

$T_{\min} = 0.586$, $T_{\max} = 0.746$

78050 measured reflections

15003 independent reflections

11100 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

$\theta_{\max} = 30.8^\circ$, $\theta_{\min} = 1.3^\circ$

$h = -31 \rightarrow 31$

$k = -19 \rightarrow 19$

$l = -46 \rightarrow 46$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.118$

$S = 1.03$

15003 reflections

579 parameters

120 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0497P)^2 + 31.4385P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.002$

$\Delta\rho_{\max} = 1.55 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -1.09 \text{ e \AA}^{-3}$ *Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.80290 (2)	-0.01300 (3)	0.39169 (2)	0.02214 (8)	
Cu2	0.70660 (2)	-0.01915 (3)	0.31768 (2)	0.02614 (8)	
Cu3	0.66594 (2)	0.02107 (3)	0.40449 (2)	0.03036 (9)	
Cu4	0.71326 (2)	-0.19124 (3)	0.38380 (2)	0.02342 (8)	
Cl1	0.81567 (3)	-0.18214 (5)	0.41707 (2)	0.02840 (14)	
Cl2	0.81362 (3)	-0.00755 (7)	0.31780 (2)	0.03451 (17)	
Cl3	0.75591 (3)	0.09683 (6)	0.43943 (2)	0.03046 (16)	
Cl4	0.67511 (3)	-0.19355 (6)	0.31175 (2)	0.03082 (15)	
Cl5	0.63812 (3)	-0.13927 (7)	0.42920 (2)	0.03596 (17)	
Cl6	0.63258 (5)	0.09252 (9)	0.33861 (3)	0.0547 (3)	
N11	0.88549 (10)	0.03600 (18)	0.40341 (7)	0.0233 (5)	
N12	0.96498 (10)	0.13435 (19)	0.39975 (9)	0.0302 (5)	
N13	1.04472 (14)	0.0232 (3)	0.43287 (15)	0.0646 (12)	
N21	0.68926 (12)	0.0204 (2)	0.26024 (8)	0.0307 (5)	
N22	0.67466 (12)	0.1147 (2)	0.20463 (8)	0.0312 (6)	
N23	0.62700 (16)	-0.0128 (3)	0.15652 (10)	0.0497 (9)	
N31	0.60566 (11)	0.0980 (2)	0.43173 (8)	0.0316 (6)	
N32	0.51725 (10)	0.15805 (18)	0.44639 (7)	0.0243 (5)	
N33	0.55277 (14)	0.2806 (3)	0.50054 (11)	0.0513 (9)	
N41	0.70835 (10)	-0.33784 (19)	0.38980 (7)	0.0254 (5)	
N42	0.71449 (13)	-0.4914 (2)	0.41434 (9)	0.0355 (6)	
N43	0.71274 (19)	-0.5839 (3)	0.34682 (11)	0.0591 (10)	
O1	0.72212 (8)	-0.05076 (15)	0.37460 (6)	0.0222 (4)	
O11	0.9926 (2)	0.1888 (4)	0.31860 (12)	0.0965 (15)	
H11A	0.991 (3)	0.228 (4)	0.2983 (14)	0.145*	
O12	1.05456 (14)	-0.0584 (3)	0.44694 (19)	0.125 (2)	
O13	1.08424 (12)	0.0857 (3)	0.43043 (14)	0.0822 (12)	
O21	0.55815 (16)	0.1846 (3)	0.17440 (17)	0.0943 (14)	
H21A	0.5252 (8)	0.206 (4)	0.1839 (17)	0.141*	
O22	0.59231 (17)	-0.0862 (2)	0.15657 (10)	0.0714 (11)	
O23	0.64160 (13)	0.0312 (3)	0.12553 (8)	0.0606 (9)	
O32	0.59897 (13)	0.3128 (3)	0.52030 (10)	0.0717 (11)	
O33	0.50093 (13)	0.3067 (3)	0.50600 (10)	0.0683 (10)	
O41	0.81469 (16)	-0.5364 (3)	0.47601 (15)	0.0891 (14)	
H41A	0.8525 (9)	-0.5590 (19)	0.479 (2)	0.134*	
O42	0.70514 (17)	-0.5744 (2)	0.30998 (9)	0.0643 (9)	
O43	0.7238 (3)	-0.6628 (3)	0.36399 (13)	0.139 (2)	
C11	0.90440 (12)	0.1254 (2)	0.39148 (9)	0.0261 (6)	

C12	0.93477 (13)	-0.0145 (2)	0.42001 (11)	0.0333 (7)	
H12A	0.9349	-0.0801	0.4312	0.040*	
C13	0.98391 (13)	0.0457 (3)	0.41783 (12)	0.0380 (8)	
C14	1.00057 (14)	0.2241 (3)	0.39029 (12)	0.0403 (8)	
H14A	1.0302	0.2388	0.4139	0.048*	
H14B	0.9729	0.2822	0.3864	0.048*	
C15	1.03387 (19)	0.2108 (4)	0.35240 (16)	0.0627 (13)	
H15A	1.0563	0.2729	0.3469	0.075*	
H15B	1.0636	0.1557	0.3567	0.075*	
C16	0.86509 (14)	0.2053 (3)	0.37248 (12)	0.0369 (7)	
H16A	0.8229	0.1822	0.3698	0.055*	
H16B	0.8780	0.2213	0.3452	0.055*	
H16C	0.8684	0.2652	0.3899	0.055*	
C21	0.69669 (14)	0.1114 (3)	0.24465 (9)	0.0311 (6)	
C22	0.66075 (16)	-0.0364 (3)	0.22997 (11)	0.0382 (8)	
H22A	0.6493	-0.1046	0.2323	0.046*	
C23	0.65153 (15)	0.0209 (3)	0.19587 (10)	0.0355 (7)	
C24	0.66535 (15)	0.2064 (3)	0.18002 (11)	0.0398 (8)	
H24A	0.6692	0.1905	0.1507	0.048*	
H24B	0.6972	0.2557	0.1889	0.048*	
C25	0.60375 (18)	0.2517 (3)	0.18455 (15)	0.0514 (10)	
H25A	0.6014	0.2739	0.2133	0.062*	
H25B	0.5983	0.3113	0.1666	0.062*	
C26	0.7245 (2)	0.1984 (3)	0.26728 (12)	0.0515 (10)	
H26A	0.7510	0.1746	0.2908	0.077*	
H26B	0.6925	0.2407	0.2771	0.077*	
H26C	0.7483	0.2373	0.2489	0.077*	
C31	0.54558 (12)	0.0912 (2)	0.42365 (9)	0.0247 (5)	
C32	0.61649 (13)	0.1716 (2)	0.46056 (9)	0.0299 (6)	
H32A	0.6552	0.1934	0.4719	0.036*	
C33	0.56247 (14)	0.2077 (2)	0.47002 (10)	0.0308 (6)	
C34	0.4518 (4)	0.1791 (10)	0.4409 (4)	0.027 (2)	0.515 (19)
H34A	0.4346	0.1503	0.4145	0.033*	0.515 (19)
H34B	0.4451	0.2523	0.4399	0.033*	0.515 (19)
C35	0.4205 (4)	0.1352 (8)	0.4754 (3)	0.034 (2)	0.515 (19)
H35A	0.4351	0.1686	0.5014	0.041*	0.515 (19)
H35B	0.3763	0.1469	0.4706	0.041*	0.515 (19)
O31	0.4317 (4)	0.0314 (7)	0.4788 (3)	0.040 (2)	0.515 (19)
H31A	0.4316 (19)	0.012 (3)	0.5031 (8)	0.060*	0.515 (19)
C34A	0.4496 (4)	0.1552 (12)	0.4507 (5)	0.036 (3)	0.485 (19)
H34D	0.4353	0.2234	0.4568	0.044*	0.485 (19)
H34E	0.4283	0.1335	0.4243	0.044*	0.485 (19)
C35A	0.4336 (5)	0.0858 (14)	0.4841 (4)	0.053 (4)	0.485 (19)
H35D	0.4523	0.1103	0.5108	0.063*	0.485 (19)
H35E	0.3890	0.0857	0.4854	0.063*	0.485 (19)
O31A	0.4535 (6)	-0.0129 (10)	0.4775 (2)	0.054 (3)	0.485 (19)
H31D	0.483 (6)	-0.012 (11)	0.489 (4)	0.081*	0.485 (19)
C36	0.51388 (13)	0.0216 (3)	0.39382 (11)	0.0357 (7)	

H36A	0.5428	-0.0279	0.3850	0.053*
H36B	0.4966	0.0594	0.3699	0.053*
H36C	0.4813	-0.0126	0.4068	0.053*
C41	0.71482 (13)	-0.3938 (2)	0.42409 (9)	0.0287 (6)
C42	0.70446 (13)	-0.4025 (2)	0.35718 (9)	0.0281 (6)
H42A	0.6992	-0.3840	0.3290	0.034*
C43	0.70926 (16)	-0.4966 (2)	0.37164 (10)	0.0352 (7)
C44	0.71595 (19)	-0.5751 (3)	0.44430 (12)	0.0480 (9)
H44A	0.7000	-0.5519	0.4701	0.058*
H44B	0.6897	-0.6302	0.4330	0.058*
C45	0.7785 (2)	-0.6121 (4)	0.45304 (15)	0.0602 (11)
H45A	0.7781	-0.6746	0.4693	0.072*
H45B	0.7964	-0.6269	0.4269	0.072*
C46	0.71969 (17)	-0.3551 (3)	0.46674 (10)	0.0393 (8)
H46A	0.7583	-0.3765	0.4809	0.059*
H46B	0.6861	-0.3813	0.4813	0.059*
H46C	0.7179	-0.2818	0.4662	0.059*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.01369 (14)	0.02604 (17)	0.02712 (17)	-0.00543 (12)	0.00426 (12)	-0.00550 (13)
Cu2	0.02511 (17)	0.03088 (19)	0.02237 (16)	-0.00373 (14)	0.00162 (13)	-0.00551 (14)
Cu3	0.01595 (15)	0.0409 (2)	0.0351 (2)	-0.00491 (14)	0.00700 (13)	-0.01941 (16)
Cu4	0.01913 (15)	0.02818 (18)	0.02352 (16)	-0.00774 (13)	0.00518 (12)	-0.00593 (13)
Cl1	0.0216 (3)	0.0303 (4)	0.0323 (3)	-0.0077 (3)	-0.0032 (3)	0.0011 (3)
Cl2	0.0254 (3)	0.0527 (5)	0.0265 (3)	-0.0088 (3)	0.0084 (3)	-0.0048 (3)
Cl3	0.0178 (3)	0.0377 (4)	0.0364 (4)	-0.0069 (3)	0.0055 (3)	-0.0146 (3)
Cl4	0.0320 (3)	0.0325 (4)	0.0270 (3)	-0.0100 (3)	-0.0035 (3)	-0.0038 (3)
Cl5	0.0238 (3)	0.0510 (5)	0.0351 (4)	-0.0049 (3)	0.0145 (3)	-0.0015 (3)
Cl6	0.0554 (6)	0.0713 (7)	0.0395 (5)	0.0348 (5)	0.0163 (4)	0.0098 (4)
N11	0.0136 (9)	0.0250 (12)	0.0317 (12)	-0.0037 (8)	0.0049 (9)	-0.0061 (9)
N12	0.0161 (10)	0.0266 (13)	0.0485 (16)	-0.0053 (9)	0.0071 (10)	-0.0078 (11)
N13	0.0175 (13)	0.049 (2)	0.127 (4)	0.0027 (13)	-0.0002 (17)	0.006 (2)
N21	0.0349 (13)	0.0325 (14)	0.0240 (12)	-0.0034 (11)	-0.0015 (10)	-0.0082 (10)
N22	0.0273 (12)	0.0421 (15)	0.0240 (12)	-0.0001 (11)	0.0020 (10)	-0.0039 (11)
N23	0.0530 (19)	0.051 (2)	0.0406 (17)	0.0300 (16)	-0.0228 (15)	-0.0193 (15)
N31	0.0183 (11)	0.0410 (15)	0.0358 (14)	-0.0054 (10)	0.0040 (10)	-0.0198 (12)
N32	0.0215 (11)	0.0276 (12)	0.0237 (11)	0.0048 (9)	0.0007 (9)	-0.0023 (9)
N33	0.0396 (16)	0.055 (2)	0.057 (2)	0.0174 (15)	-0.0098 (14)	-0.0336 (16)
N41	0.0189 (10)	0.0317 (13)	0.0259 (12)	-0.0082 (9)	0.0033 (9)	-0.0044 (10)
N42	0.0404 (15)	0.0300 (14)	0.0340 (14)	-0.0092 (11)	-0.0094 (12)	-0.0012 (11)
N43	0.090 (3)	0.0332 (17)	0.050 (2)	0.0034 (17)	-0.0176 (19)	-0.0105 (15)
O1	0.0163 (8)	0.0279 (10)	0.0227 (9)	-0.0040 (7)	0.0036 (7)	-0.0080 (8)
O11	0.085 (3)	0.148 (4)	0.061 (2)	-0.048 (3)	0.031 (2)	-0.016 (2)
O12	0.0294 (16)	0.068 (2)	0.273 (7)	0.0062 (16)	-0.022 (3)	0.061 (3)
O13	0.0176 (12)	0.066 (2)	0.161 (4)	-0.0098 (13)	-0.0059 (17)	0.005 (2)
O21	0.0401 (18)	0.073 (3)	0.168 (4)	-0.0018 (17)	-0.004 (2)	-0.021 (3)

O22	0.097 (3)	0.0370 (16)	0.070 (2)	0.0127 (16)	-0.0530 (19)	-0.0170 (14)
O23	0.0467 (16)	0.104 (3)	0.0293 (13)	0.0229 (16)	-0.0096 (11)	-0.0139 (15)
O32	0.0461 (16)	0.088 (2)	0.077 (2)	0.0221 (16)	-0.0221 (15)	-0.0596 (19)
O33	0.0407 (15)	0.080 (2)	0.082 (2)	0.0242 (14)	-0.0056 (14)	-0.0527 (18)
O41	0.055 (2)	0.070 (2)	0.134 (4)	-0.0152 (17)	-0.038 (2)	0.037 (2)
O42	0.108 (3)	0.0457 (17)	0.0396 (15)	-0.0042 (17)	0.0097 (16)	-0.0170 (13)
O43	0.290 (7)	0.045 (2)	0.071 (3)	0.046 (3)	-0.061 (4)	-0.0152 (19)
C11	0.0193 (12)	0.0275 (14)	0.0320 (14)	-0.0053 (10)	0.0064 (10)	-0.0070 (11)
C12	0.0186 (13)	0.0289 (15)	0.053 (2)	0.0019 (11)	0.0050 (13)	-0.0024 (14)
C13	0.0141 (12)	0.0352 (17)	0.065 (2)	-0.0015 (11)	0.0042 (13)	-0.0069 (16)
C14	0.0221 (14)	0.0362 (18)	0.063 (2)	-0.0153 (13)	0.0074 (14)	-0.0067 (16)
C15	0.037 (2)	0.071 (3)	0.084 (3)	-0.022 (2)	0.026 (2)	-0.004 (2)
C16	0.0244 (14)	0.0344 (17)	0.052 (2)	-0.0068 (12)	0.0012 (13)	0.0044 (15)
C21	0.0313 (15)	0.0392 (17)	0.0229 (14)	-0.0076 (13)	0.0035 (11)	-0.0058 (12)
C22	0.0420 (18)	0.0303 (16)	0.0394 (18)	0.0068 (14)	-0.0137 (14)	-0.0113 (14)
C23	0.0357 (16)	0.0406 (18)	0.0284 (15)	0.0114 (14)	-0.0086 (12)	-0.0134 (13)
C24	0.0319 (16)	0.051 (2)	0.0359 (17)	-0.0074 (15)	-0.0003 (13)	0.0097 (15)
C25	0.042 (2)	0.043 (2)	0.069 (3)	0.0006 (17)	0.0065 (19)	0.0059 (19)
C26	0.069 (3)	0.048 (2)	0.0355 (19)	-0.025 (2)	-0.0065 (18)	-0.0030 (16)
C31	0.0176 (12)	0.0325 (15)	0.0245 (13)	-0.0023 (10)	0.0040 (10)	-0.0050 (11)
C32	0.0244 (13)	0.0346 (16)	0.0305 (15)	-0.0031 (12)	0.0001 (11)	-0.0095 (12)
C33	0.0273 (14)	0.0327 (16)	0.0310 (15)	0.0078 (12)	-0.0053 (11)	-0.0106 (12)
C34	0.017 (3)	0.035 (5)	0.030 (5)	0.010 (3)	0.003 (3)	0.000 (3)
C35	0.027 (3)	0.043 (5)	0.035 (4)	0.001 (3)	0.013 (3)	0.003 (3)
O31	0.033 (3)	0.037 (4)	0.052 (4)	0.004 (3)	0.013 (3)	0.011 (3)
C34A	0.022 (4)	0.045 (7)	0.040 (7)	0.019 (4)	-0.010 (4)	-0.005 (5)
C35A	0.023 (4)	0.091 (11)	0.045 (5)	-0.001 (7)	0.010 (4)	0.014 (8)
O31A	0.055 (6)	0.065 (7)	0.041 (4)	-0.023 (5)	-0.007 (3)	0.020 (4)
C36	0.0197 (13)	0.0441 (19)	0.0427 (18)	-0.0038 (12)	-0.0005 (12)	-0.0198 (15)
C41	0.0229 (13)	0.0331 (16)	0.0294 (14)	-0.0129 (11)	-0.0008 (11)	-0.0043 (12)
C42	0.0250 (13)	0.0332 (16)	0.0265 (14)	-0.0074 (11)	0.0038 (11)	-0.0078 (12)
C43	0.0394 (17)	0.0300 (16)	0.0348 (16)	-0.0058 (13)	-0.0052 (13)	-0.0074 (13)
C44	0.059 (2)	0.039 (2)	0.044 (2)	-0.0141 (17)	-0.0106 (18)	0.0024 (16)
C45	0.060 (3)	0.058 (3)	0.060 (3)	-0.004 (2)	-0.009 (2)	0.011 (2)
C46	0.049 (2)	0.0420 (19)	0.0265 (15)	-0.0144 (16)	0.0005 (14)	-0.0057 (14)

Geometric parameters (\AA , $^\circ$)

Cu1—O1	1.8960 (18)	N31—C31	1.337 (3)
Cu1—N11	1.949 (2)	N31—C32	1.368 (4)
Cu1—Cl1	2.4152 (9)	N32—C31	1.348 (4)
Cu1—Cl3	2.4351 (8)	N32—C33	1.381 (4)
Cu1—Cl2	2.4435 (9)	N32—C34	1.472 (9)
Cu2—O1	1.908 (2)	N32—C34A	1.516 (10)
Cu2—N21	1.955 (3)	N33—O33	1.226 (4)
Cu2—Cl6	2.3579 (10)	N33—O32	1.240 (4)
Cu2—Cl2	2.3726 (9)	N33—C33	1.423 (4)
Cu2—Cl4	2.4351 (9)	N41—C41	1.343 (4)

Cu3—O1	1.9022 (19)	N41—C42	1.368 (4)
Cu3—N31	1.955 (2)	N42—C41	1.342 (4)
Cu3—Cl5	2.3877 (10)	N42—C43	1.390 (4)
Cu3—Cl6	2.4113 (11)	N42—C44	1.484 (5)
Cu3—Cl3	2.4312 (8)	N43—O42	1.207 (4)
Cu4—O1	1.913 (2)	N43—O43	1.209 (5)
Cu4—N41	1.972 (3)	N43—C43	1.426 (5)
Cu4—Cl5	2.4186 (8)	O11—C15	1.404 (6)
Cu4—Cl4	2.4314 (9)	O21—C25	1.370 (5)
Cu4—Cl1	2.4332 (8)	O41—C45	1.458 (6)
N11—C11	1.335 (4)	C11—C16	1.480 (4)
N11—C12	1.356 (4)	C12—C13	1.359 (4)
N12—C11	1.350 (4)	C14—C15	1.501 (6)
N12—C13	1.373 (4)	C21—C26	1.482 (5)
N12—C14	1.481 (4)	C22—C23	1.352 (5)
N13—O12	1.195 (5)	C24—C25	1.510 (5)
N13—O13	1.217 (4)	C31—C36	1.480 (4)
N13—C13	1.426 (4)	C32—C33	1.348 (4)
N21—C21	1.334 (4)	C34—C35	1.490 (10)
N21—C22	1.359 (4)	C35—O31	1.411 (9)
N22—C21	1.356 (4)	C34A—C35A	1.495 (13)
N22—C23	1.376 (4)	C35A—O31A	1.413 (13)
N22—C24	1.469 (4)	C41—C46	1.480 (4)
N23—O23	1.236 (5)	C42—C43	1.343 (5)
N23—O22	1.246 (5)	C44—C45	1.474 (6)
N23—C23	1.425 (4)		
O1—Cu1—N11	173.12 (10)	C31—N31—C32	107.4 (2)
O1—Cu1—Cl11	86.13 (6)	C31—N31—Cu3	125.4 (2)
N11—Cu1—Cl11	99.55 (7)	C32—N31—Cu3	127.1 (2)
O1—Cu1—Cl13	84.63 (6)	C31—N32—C33	106.1 (2)
N11—Cu1—Cl13	96.61 (7)	C31—N32—C34	123.8 (6)
Cl1—Cu1—Cl13	112.86 (3)	C33—N32—C34	129.5 (6)
O1—Cu1—Cl12	83.33 (6)	C31—N32—C34A	122.8 (7)
N11—Cu1—Cl12	91.01 (7)	C33—N32—C34A	129.4 (7)
Cl1—Cu1—Cl12	110.37 (3)	O33—N33—O32	124.6 (3)
Cl3—Cu1—Cl12	134.02 (3)	O33—N33—C33	119.6 (3)
O1—Cu2—N21	176.91 (10)	O32—N33—C33	115.9 (3)
O1—Cu2—Cl6	86.06 (6)	C41—N41—C42	107.0 (3)
N21—Cu2—Cl6	91.20 (8)	C41—N41—Cu4	129.2 (2)
O1—Cu2—Cl12	85.06 (6)	C42—N41—Cu4	123.4 (2)
N21—Cu2—Cl12	95.77 (8)	C41—N42—C43	106.5 (3)
Cl6—Cu2—Cl12	132.47 (4)	C41—N42—C44	125.2 (3)
O1—Cu2—Cl4	83.85 (6)	C43—N42—C44	128.2 (3)
N21—Cu2—Cl4	98.64 (8)	O42—N43—O43	124.0 (4)
Cl6—Cu2—Cl4	115.31 (4)	O42—N43—C43	118.0 (3)
Cl2—Cu2—Cl4	109.97 (3)	O43—N43—C43	117.9 (4)
O1—Cu3—N31	176.21 (10)	Cu1—O1—Cu3	110.80 (9)

O1—Cu3—Cl5	85.31 (6)	Cu1—O1—Cu2	108.46 (9)
N31—Cu3—Cl5	96.51 (9)	Cu3—O1—Cu2	108.36 (10)
O1—Cu3—Cl6	84.68 (6)	Cu1—O1—Cu4	108.74 (10)
N31—Cu3—Cl6	91.57 (9)	Cu3—O1—Cu4	109.55 (9)
Cl5—Cu3—Cl6	126.01 (4)	Cu2—O1—Cu4	110.93 (9)
O1—Cu3—Cl3	84.61 (6)	N11—C11—N12	110.5 (3)
N31—Cu3—Cl3	97.52 (7)	N11—C11—C16	125.5 (3)
Cl5—Cu3—Cl3	116.05 (3)	N12—C11—C16	124.0 (3)
Cl6—Cu3—Cl3	115.56 (4)	N11—C12—C13	107.8 (3)
O1—Cu4—N41	175.50 (9)	C12—C13—N12	108.4 (3)
O1—Cu4—Cl5	84.21 (6)	C12—C13—N13	126.4 (3)
N41—Cu4—Cl5	100.26 (7)	N12—C13—N13	125.2 (3)
O1—Cu4—Cl4	83.84 (6)	N12—C14—C15	112.4 (3)
N41—Cu4—Cl4	93.78 (7)	O11—C15—C14	109.9 (3)
Cl5—Cu4—Cl4	113.26 (3)	N21—C21—N22	110.5 (3)
O1—Cu4—Cl1	85.25 (6)	N21—C21—C26	125.7 (3)
N41—Cu4—Cl1	93.57 (7)	N22—C21—C26	123.7 (3)
Cl5—Cu4—Cl1	111.98 (3)	C23—C22—N21	108.1 (3)
Cl4—Cu4—Cl1	131.90 (3)	C22—C23—N22	108.5 (3)
Cu1—Cl1—Cu4	79.38 (2)	C22—C23—N23	125.7 (3)
Cu2—Cl2—Cu1	79.70 (2)	N22—C23—N23	125.6 (3)
Cu3—Cl3—Cu1	79.95 (3)	N22—C24—C25	111.6 (3)
Cu4—Cl4—Cu2	80.61 (2)	O21—C25—C24	111.5 (4)
Cu3—Cl5—Cu4	80.86 (3)	N31—C31—N32	110.3 (2)
Cu2—Cl6—Cu3	80.74 (3)	N31—C31—C36	125.6 (3)
C11—N11—C12	107.5 (2)	N32—C31—C36	124.2 (2)
C11—N11—Cu1	123.7 (2)	C33—C32—N31	107.8 (3)
C12—N11—Cu1	128.4 (2)	C32—C33—N32	108.4 (3)
C11—N12—C13	105.8 (2)	C32—C33—N33	126.4 (3)
C11—N12—C14	124.5 (3)	N32—C33—N33	125.1 (3)
C13—N12—C14	129.7 (3)	N32—C34—C35	110.2 (7)
O12—N13—O13	122.9 (3)	O31—C35—C34	110.9 (8)
O12—N13—C13	117.5 (3)	C35A—C34A—N32	112.3 (8)
O13—N13—C13	119.7 (4)	O31A—C35A—C34A	111.8 (9)
C21—N21—C22	107.3 (3)	N42—C41—N41	110.2 (3)
C21—N21—Cu2	126.3 (2)	N42—C41—C46	124.1 (3)
C22—N21—Cu2	126.1 (2)	N41—C41—C46	125.7 (3)
C21—N22—C23	105.6 (3)	C43—C42—N41	108.6 (3)
C21—N22—C24	125.2 (3)	C42—C43—N42	107.6 (3)
C23—N22—C24	127.8 (3)	C42—C43—N43	124.9 (3)
O23—N23—O22	125.4 (3)	N42—C43—N43	127.3 (3)
O23—N23—C23	118.8 (4)	C45—C44—N42	110.4 (3)
O22—N23—C23	115.9 (4)	O41—C45—C44	109.5 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
-------------	-------------	-------------	---------------------

supporting information

O41—H41A···O31 ⁱ	0.89 (2)	2.13 (3)	2.738 (8)	125 (2)
-----------------------------	----------	----------	-----------	---------

Symmetry code: (i) $x+1/2, y-1/2, z$.