

Pace University

DigitalCommons@Pace

Faculty Papers

Dyson College of Arts & Sciences

2015

Crystal structure of bis[1-(2-hydroxyethyl)-2-methyl-5-nitro- 1H-imidazole-κN3]silver(I) tetrafluoridoborate methanol monosolvate

Joshua H. Palmer

Rita K. Upmacis

Follow this and additional works at: https://digitalcommons.pace.edu/dyson_fac



Part of the Chemistry Commons

STRUCTURE
REPORTSISSN 1600-5368
journals.iucr.org/e

Crystal structure of bis[1-(2-hydroxyethyl)-2-methyl-5-nitro- $1H$ -imidazole- κN^3]silver(I) tetrafluoridoborate methanol monosolvate

Joshua H. Palmer and Rita K. Upmacis*Acta Cryst.* (2015). **E71**, 284–287**IUCr Journals**

CRYSTALLOGRAPHY JOURNALS ONLINE

This open-access article is distributed under the terms of the Creative Commons Attribution Licence
<http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and
reproduction in any medium, provided the original authors and source are cited.



Received 19 January 2015
Accepted 9 February 2015

Edited by A. J. Lough, University of Toronto,
Canada

Keywords: crystal structure; silver; metronidazole; Flagyl; tetrafluoridoborate; hydrogen bonding

CCDC reference: 1048516

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of bis[1-(2-hydroxyethyl)-2-methyl-5-nitro-1H-imidazole- κN^3]silver(I) tetrafluoridoborate methanol monosolvate

Joshua H. Palmer^a and Rita K. Upmacis^{b*}

^aDepartment of Chemistry, Columbia University, New York, NY 10027, USA, and ^bHaskins Laboratories, Dept. of Chemistry, Pace University, New York, NY 10038, USA. *Correspondence e-mail: rupmacis@pace.edu


1-(2-Hydroxyethyl)-2-methyl-5-nitro-1H-imidazole (metronidazole, MET) is a medication that is used to treat infections by a variety of anaerobic organisms, but there are relatively few reports of the structures of metal compounds that exhibit coordination of metronidazole. We have demonstrated that MET reacts with AgBF_4 to give $[\text{Ag}(\text{MET})_2]\text{BF}_4 \cdot \text{CH}_3\text{OH}$, in which the Ag^{I} cation is coordinated by two MET ligands with a *trans* arrangement. The structure of $[\text{Ag}(\text{MET})_2]\text{BF}_4$ exhibits some interesting differences from its nitrate counterpart, $[\text{Ag}(\text{MET})_2]\text{NO}_3$ [Fun *et al.* (2008). *Acta Cryst. E* **64**, m668]. For instance, although the two MET ligands of both $[\text{Ag}(\text{MET})_2]\text{BF}_4$ and $[\text{Ag}(\text{MET})_2]\text{NO}_3$ are almost coplanar, the former compound has an *anti*-like geometry with a molecular inversion center, but the latter has a *syn*-like arrangement. In the crystal, the BF_4^- anion is linked by an $\text{O}-\text{H}\cdots\text{F}$ hydrogen bond to the methanol solvent molecule, which is, in turn, linked to the cation by an $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond; the components of the structure are linked by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming chains along [001]. One of the MET ligands and the BF_4^- anion are disordered over two sets of sites with ratios of refined occupancies 0.501 (17):0.499 (17) and 0.539 (19):0.461 (19), respectively.

1. Chemical context


1-(2-Hydroxyethyl)-2-methyl-5-nitro-1H-imidazole, also known as metronidazole (MET) or Flagyl, is a medication used particularly for treatment of parasitic infections, such as trichomoniasis, amoebiasis and giardiasis, but is also effective against anaerobic bacteria (Freeman *et al.*, 1997; Miljkovic *et al.*, 2014; Soares *et al.*, 2012; Samuelson, 1999; Lofmark *et al.*, 2010). There are relatively few reports of the structures of metal compounds that exhibit coordination of MET. For example, with respect to silver, only the nitrate compound, $[\text{Ag}(\text{MET})_2]\text{NO}_3$, has been structurally characterized by X-ray diffraction (Fun *et al.*, 2008). Herein, we describe the structure of the tetrafluoridoborate derivative, $[\text{Ag}(\text{MET})_2]\text{BF}_4$, which is obtained by addition of MET to AgBF_4 in methanol (see Scheme).

2. Structural commentary

Crystals of composition $[\text{Ag}(\text{MET})_2]\text{BF}_4 \cdot \text{MeOH}$ were obtained from a solution in methanol. The asymmetric unit consists of a silver cation, $[\text{Ag}(\text{MET})_2]^+$, a tetrafluoridoborate anion, BF_4^- , and a solvent methanol molecule. The silver atom of $[\text{Ag}(\text{MET})_2]^+$ is coordinated by two MET ligands in a *trans* manner by their N^3 nitrogen atoms, as illustrated in Fig. 1.



OPEN ACCESS



One of the MET ligands exhibits disorder resulting from rotation about the Ag—N bond [the dihedral angle between the planes of the disordered 5-membered rings is 11.0 (9) $^\circ$]. The Ag—N bond lengths [Ag—N11 = 2.082 (15), Ag—N11A = 2.163 (16) and Ag—N21 = 2.1193 (15) Å] are comparable to those values in the nitrate derivative, [2.1489 (11) and 2.1475 (11) Å; Fun *et al.*, 2008]. There are, however, some interesting differences between the two compounds.

First, while the two MET ligands of both $[\text{Ag}(\text{MET})_2]\text{BF}_4$ and $[\text{Ag}(\text{MET})_2]\text{NO}_3$ are almost coplanar, the former compound has an *anti*-like geometry, and the latter has a *syn*-like arrangement. Thus, the C13—N11···N21—C23 torsion angle for $[\text{Ag}(\text{MET})_2]\text{BF}_4$ is 160.8 (9) $^\circ$ [148.6 (11) $^\circ$ for the minor component of disorder], while the value for $[\text{Ag}(\text{MET})_2]\text{NO}_3$ is 24.10 $^\circ$ (Fun *et al.*, 2008). These differences are illustrated in Fig. 2, which shows that the $[\text{Ag}(\text{MET})_2]^+$ unit of $[\text{Ag}(\text{MET})_2]\text{BF}_4$ has an approximate inversion center at the Ag^I ion, whereas $[\text{Ag}(\text{MET})_2]\text{NO}_3$ does not.




Figure 1

The molecular structure of the cation of the title compound, with displacement ellipsoids drawn at the 30% probability level. The disorder is not shown.




Figure 2

Comparison of the $[\text{Ag}(\text{MET})_2]^+$ units in $[\text{Ag}(\text{MET})_2]\text{BF}_4$ (top) and $[\text{Ag}(\text{MET})_2]\text{NO}_3$ (bottom).

A second interesting difference is that the N11—Ag—N21 angle of 175.7 (5) $^\circ$ for $[\text{Ag}(\text{MET})_2]\text{BF}_4$ is much closer to 180 $^\circ$ than is the corresponding value for $[\text{Ag}(\text{MET})_2]\text{NO}_3$ [165.34 (4) $^\circ$; Fun *et al.*, 2008]. It is possible that this could be attributed to the tetrafluoridoborate ligand being considered a non-coordinating ion relative to nitrate, and this is reflected by the fact that $[\text{Ag}(\text{MET})_2]\text{NO}_3$ exhibits Ag···O contacts of 2.63 and 2.67 Å, which are comparable to distances in other silver nitrate compounds (Wu *et al.*, 2012).

3. Supramolecular features

The hydroxyethyl group of one of the MET ligands [O21—H] serves as a donor in an intermolecular hydrogen-bonding interaction with the other hydroxyethyl group [O11—H] of an adjacent molecule. In turn, the latter hydroxyethyl group serves as a hydrogen-bond donor to a methanol molecule, which also hydrogen bonds to a tetrafluoridoborate anion. In




Figure 3

Part of a hydrogen-bonded chain along [001]. The disorder is not shown and hydrogen bonds are shown as dashed lines.

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots F2	0.84	1.84	2.673 (9)	173
O11—H11A \cdots O1	0.84	1.86	2.697 (10)	175
O21—H21A \cdots O11 ⁱ	0.84	1.91	2.726 (11)	164
O21—H21A \cdots O11A ⁱ	0.84	1.87	2.712 (11)	176

Symmetry code: (i) $x, y, z - 1$.

the crystal, the components of the structure are linked into chains along [001] by the O—H \cdots O hydrogen bonds (Table 1 and Fig. 3).

4. Database survey

In addition to coordination to silver, metronidazole has also been shown to coordinate to other metals, and structurally characterized compounds have been reported for Co (Galván-Tejada *et al.*, 2002), Cu (Galván-Tejada *et al.*, 2002; Barba-Behrens *et al.*, 1991; Athar *et al.*, 2005; Ratajczak-Sitarz *et al.*, 1998; Bharti *et al.*, 2002), Zn (Galván-Tejada *et al.*, 2002), Ru (Wu *et al.*, 2003; Kennedy *et al.*, 2006), Rh (Dyson *et al.*, 1990; Nothenberg *et al.*, 1994), Pd (Bharti *et al.*, 2002; De Bondt *et al.*, 1994; Rochon *et al.*, 1993) and Pt (Bharti *et al.*, 2002; Bales *et al.*, 1983). In these compounds, the coordination number of the central atom ranges from four for Cu, Zn, Pd and Pt to six for Ru and Rh.

5. Synthesis and crystallization

Crystals of composition $[\text{Ag}(\text{MET})_2]\text{BF}_4 \cdot \text{MeOH}$ were obtained by combining AgBF_4 with MET in a 1:2 molar ratio in methanol and allowing the solution to evaporate slowly at room temperature.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. Hydrogen atoms were refined using a riding-model approximation with C—H = 0.95–0.99 \AA , O—H = 0.84 \AA and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{O})$. One of the MET ligands was refined as rotationally disordered with occupancies of 0.501 (17) and 0.499 (17) and the configurations were modeled using the SAME command in *SHELXL2013* (Sheldrick, 2015). The tetrafluoroborate counter-ion was also refined as disordered and was modeled with two site occupancies, 0.539 (19) and 0.461 (19).

Acknowledgements

RKU would like to thank Pace University for research support. Gerard Parkin (Columbia University) is thanked for helpful discussions.

Table 2
Experimental details.

Crystal data	
Chemical formula	$[\text{Ag}(\text{C}_6\text{H}_9\text{N}_3\text{O}_3)_2]\text{BF}_4 \cdot \text{CH}_4\text{O}$
M_r	569.04
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	130
a, b, c (\AA)	9.2592 (10), 10.5339 (10), 12.3995 (12)
α, β, γ ($^\circ$)	106.940 (11), 92.788 (9), 112.439 (10)
V (\AA^3)	1051.7 (2)
Z	2
Radiation type	Mo $K\alpha$
μ (mm^{-1})	1.04
Crystal size (mm)	1.00 \times 0.51 \times 0.31
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2013)
T_{\min}, T_{\max}	0.551, 0.747
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	17164, 6401, 6107
R_{int}	0.023
(sin θ/λ) _{max} (\AA^{-1})	0.714
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.028, 0.067, 1.12
No. of reflections	6401
No. of parameters	440
No. of restraints	144
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ ($e \text{\AA}^{-3}$)	1.00, -1.03

Computer programs: *APEX2* and *SAINT* (Bruker, 2013), *SHELXS97* and *SHELXTL* (Sheldrick, 2008), *SHELXL2013* (Sheldrick, 2015).

References

- Athar, F., Husain, K., Abid, M., Agarwal, S. M., Coles, S. J., Hursthouse, M. B., Maurya, M. R. & Azam, A. (2005). *Chem. Biodivers.*, **2**, 1320–1330.
- Bales, J. R., Coulson, C. J., Gilmour, D. W., Mazid, M. A., Neidle, S., Kuroda, R., Pearse, B. J., Ramsden, C. A. & Sadler, P. J. (1983). *J. Chem. Soc. Chem. Commun.* pp. 432–433.
- Barba-Behrens, N., María Mutio-Rico, A., Joseph-Nathan, P. & Contreras, R. (1991). *Polyhedron*, **10**, 1333–1341.
- Bharti, N., Shailendra, Coles, S. J., Hursthouse, M. B., Mayer, T. A., Gonzalez Garza, M. T., Cruz-Vega, D. E., Mata-Cardenas, B. D., Naqvi, F., Maurya, M. R. & Azam, A. (2002). *Helv. Chim. Acta*, **85**, 2704–2712.
- Bruker (2013). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- De Bondt, H. L., Blaton, N. M., Peeters, O. M. & De Ranter, C. J. (1994). *Acta Cryst. C* **50**, 180–181.
- Dyson, T. M., Morrison, E. C., Tocher, D. A., Dale, L. D. & Edwards, D. I. (1990). *Inorg. Chim. Acta*, **169**, 127–131.
- Freeman, C. D., Klutman, N. E. & Lamp, K. C. (1997). *Drugs*, **54**, 679–708.
- Fun, H.-K., Jebas, S. R. & Balasubramanian, T. (2008). *Acta Cryst. E* **64**, m668–m669.
- Galván-Tejada, N., Bernès, S., Castillo-Blum, S. E., Nöth, H., Vicente, R. & Barba-Behrens, N. (2002). *J. Inorg. Biochem.*, **91**, 339–348.
- Kennedy, D. C., Wu, A., Patrick, B. O. & James, B. R. (2006). *J. Inorg. Biochem.*, **100**, 1974–1982.
- Lofmark, S., Edlund, C. & Nord, C. E. (2010). *Clin. Infect. Dis. Suppl. 1*, S16–S23.
- Miljkovic, V., Arsic, B., Bojanic, Z., Nikolic, G., Nikolic, Lj., Kalicanin, B. & Savic, V. (2014). *Pharmazie*, **69**, 571–577.

- Nothenberg, M. S., Zyngier, S. B., Giesbrecht, A. M., Gambardella, M. T. P., Santos, R. H. A., Kimura, E. & Najjar, R. (1994). *J. Braz. Chem. Soc.* **5**, 23–29.
- Ratajczak-Sitarz, M., Katrusiak, A., Wojakowska, H., Januszczyk, M., Krzymiński, R. & Pietrzak, J. (1998). *Inorg. Chim. Acta*, **269**, 326–331.
- Rochon, F. D., Melanson, R. & Farrell, N. (1993). *Acta Cryst. C* **49**, 1703–1706.
- Samuelson, J. (1999). *Antimicrob. Agents Chemother.* **43**, 1533–1541.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Soares, G. M. S., Figueiredo, L. C., Faveri, M., Cortelli, S. C., Duarte, P. M. & Feres, M. (2012). *J. Appl. Oral Sci.* **20**, 295–309.
- Wu, J., Huang, C., Li, G.-Q., Tian, H.-Y. & Jiang, R.-W. (2012). *Acta Cryst. E* **68**, m185–m186.
- Wu, A., Kennedy, D. C., Patrick, B. O. & James, B. R. (2003). *Inorg. Chem.* **42**, 7579–7586.

supporting information

Acta Cryst. (2015). E71, 284-287 [doi:10.1107/S2056989015002819]

Crystal structure of bis[1-(2-hydroxyethyl)-2-methyl-5-nitro-1*H*-imidazole- κN^3]silver(I) tetrafluoridoborate methanol monosolvate

Joshua H. Palmer and Rita K. Upmacis

Computing details

Data collection: *APEX2* (Bruker, 2013); cell refinement: *SAINT* (Bruker, 2013); data reduction: *SAINT* (Bruker, 2013); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2015); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Bis[1-(2-hydroxyethyl)-2-methyl-5-nitro-1*H*-imidazole- κN^3]silver(I) tetrafluoridoborate methanol monosolvate

Crystal data

$[\text{Ag}(\text{C}_6\text{H}_9\text{N}_3\text{O}_3)_2](\text{BF}_4)\cdot\text{CH}_4\text{O}$	$Z = 2$
$M_r = 569.04$	$F(000) = 572$
Triclinic, $P\bar{1}$	$D_x = 1.797 \text{ Mg m}^{-3}$
$a = 9.2592 (10) \text{ \AA}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$b = 10.5339 (10) \text{ \AA}$	Cell parameters from 9065 reflections
$c = 12.3995 (12) \text{ \AA}$	$\theta = 2.3\text{--}32.9^\circ$
$\alpha = 106.940 (11)^\circ$	$\mu = 1.04 \text{ mm}^{-1}$
$\beta = 92.788 (9)^\circ$	$T = 130 \text{ K}$
$\gamma = 112.439 (10)^\circ$	Block, colourless
$V = 1051.7 (2) \text{ \AA}^3$	$1.00 \times 0.51 \times 0.31 \text{ mm}$

Data collection

Bruker APEXII CCD	6401 independent reflections
diffractometer	6107 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.023$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2013)	$\theta_{\text{max}} = 30.5^\circ$, $\theta_{\text{min}} = 1.8^\circ$
$T_{\text{min}} = 0.551$, $T_{\text{max}} = 0.747$	$h = -13 \rightarrow 13$
17164 measured reflections	$k = -15 \rightarrow 15$
	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0161P)^2 + 1.0163P]$
$R[F^2 > 2\sigma(F^2)] = 0.028$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.067$	$(\Delta/\sigma)_{\text{max}} = 0.007$
$S = 1.12$	$\Delta\rho_{\text{max}} = 1.00 \text{ e \AA}^{-3}$
6401 reflections	$\Delta\rho_{\text{min}} = -1.03 \text{ e \AA}^{-3}$
440 parameters	Extinction correction: <i>SHELXL2013</i> (Sheldrick, 2015), $\text{Fc}^* = k\text{Fc}[1 + 0.001 \times \text{Fc}^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
144 restraints	Extinction coefficient: 0.0027 (4)
Hydrogen site location: inferred from neighbouring sites	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ag	0.34662 (2)	0.82628 (2)	-0.00390 (2)	0.02854 (5)	
N11	0.4710 (18)	0.786 (2)	0.1171 (7)	0.018 (2)	0.501 (17)
N12	0.5696 (15)	0.7754 (15)	0.2748 (8)	0.0211 (16)	0.501 (17)
N13	0.7178 (8)	0.6262 (7)	0.1956 (9)	0.0273 (14)	0.501 (17)
O11	0.3317 (11)	0.6459 (13)	0.4129 (9)	0.0404 (16)	0.501 (17)
H11A	0.2863	0.5926	0.3453	0.048*	0.501 (17)
O12	0.7349 (8)	0.5491 (6)	0.1064 (9)	0.0406 (13)	0.501 (17)
O13	0.7868 (7)	0.6518 (5)	0.2921 (10)	0.0368 (14)	0.501 (17)
C11	0.5471 (13)	0.6968 (12)	0.0851 (8)	0.0227 (15)	0.501 (17)
H11B	0.5535	0.6475	0.0092	0.027*	0.501 (17)
C12	0.6111 (12)	0.6925 (11)	0.1829 (8)	0.0197 (15)	0.501 (17)
C13	0.4855 (12)	0.8352 (11)	0.2301 (9)	0.0149 (15)	0.501 (17)
C14	0.3914 (18)	0.9109 (16)	0.2896 (10)	0.039 (4)	0.501 (17)
H14A	0.3477	0.9470	0.2375	0.047*	0.501 (17)
H14B	0.4599	0.9932	0.3572	0.047*	0.501 (17)
H14C	0.3043	0.8428	0.3136	0.047*	0.501 (17)
C15	0.5998 (12)	0.7981 (8)	0.3981 (7)	0.0282 (14)	0.501 (17)
H15A	0.5816	0.8841	0.4416	0.034*	0.501 (17)
H15B	0.7124	0.8190	0.4220	0.034*	0.501 (17)
C16	0.4946 (13)	0.6673 (12)	0.4271 (9)	0.0346 (19)	0.501 (17)
H16A	0.5035	0.5792	0.3772	0.042*	0.501 (17)
H16B	0.5316	0.6806	0.5076	0.042*	0.501 (17)
N11A	0.470 (2)	0.778 (2)	0.1207 (9)	0.028 (3)	0.499 (17)
N12A	0.5515 (16)	0.7755 (15)	0.2949 (8)	0.0179 (12)	0.499 (17)
N13A	0.7308 (8)	0.6471 (7)	0.2389 (8)	0.0224 (12)	0.499 (17)
O11A	0.2839 (10)	0.6195 (10)	0.4105 (9)	0.0349 (13)	0.499 (17)
H11C	0.2496	0.5780	0.3398	0.042*	0.499 (17)
O12A	0.7814 (9)	0.5918 (10)	0.1564 (8)	0.0368 (16)	0.499 (17)
O13A	0.7663 (7)	0.6551 (5)	0.3379 (7)	0.0314 (11)	0.499 (17)
C11A	0.5764 (13)	0.7137 (13)	0.1118 (8)	0.0212 (15)	0.499 (17)
H11D	0.6121	0.6788	0.0434	0.025*	0.499 (17)
C12A	0.6218 (12)	0.7076 (11)	0.2150 (7)	0.0170 (13)	0.499 (17)
C13A	0.4578 (12)	0.8126 (11)	0.2331 (8)	0.0140 (15)	0.499 (17)
C14A	0.3830 (17)	0.9117 (14)	0.2874 (9)	0.033 (4)	0.499 (17)
H14D	0.4602	1.0132	0.3073	0.039*	0.499 (17)
H14E	0.3490	0.8932	0.3571	0.039*	0.499 (17)
H14F	0.2906	0.8940	0.2338	0.039*	0.499 (17)
C15A	0.5567 (9)	0.7874 (8)	0.4170 (6)	0.0213 (11)	0.499 (17)
H15C	0.5264	0.8670	0.4571	0.026*	0.499 (17)

H15D	0.6671	0.8140	0.4521	0.026*	0.499 (17)
C16A	0.4482 (11)	0.6486 (10)	0.4345 (9)	0.0275 (16)	0.499 (17)
H16C	0.4661	0.5659	0.3840	0.033*	0.499 (17)
H16D	0.4758	0.6557	0.5148	0.033*	0.499 (17)
N21	0.20394 (19)	0.85591 (17)	-0.12498 (13)	0.0280 (3)	
N22	0.06590 (19)	0.82730 (17)	-0.28744 (13)	0.0270 (3)	
N23	-0.0132 (2)	1.0346 (2)	-0.2186 (2)	0.0433 (5)	
O21	0.25596 (17)	0.86179 (17)	-0.45811 (12)	0.0340 (3)	
H21A	0.2608	0.7842	-0.4975	0.051*	
O22	-0.0234 (2)	1.1214 (2)	-0.1326 (2)	0.0737 (7)	
O23	-0.0629 (3)	1.0220 (3)	-0.3151 (2)	0.0775 (8)	
C21	0.1481 (2)	0.9618 (2)	-0.10391 (19)	0.0350 (4)	
H21B	0.1652	1.0344	-0.0319	0.042*	
C22	0.0644 (2)	0.94554 (19)	-0.20292 (19)	0.0306 (4)	
C23	0.1525 (2)	0.77595 (19)	-0.23615 (14)	0.0250 (3)	
C24	0.1834 (3)	0.6463 (2)	-0.29385 (16)	0.0349 (4)	
H24A	0.2455	0.6290	-0.2379	0.042*	
H24B	0.2429	0.6636	-0.3552	0.042*	
H24C	0.0821	0.5608	-0.3265	0.042*	
C25	-0.0068 (2)	0.7652 (3)	-0.40921 (17)	0.0375 (5)	
H25A	-0.1117	0.7691	-0.4181	0.045*	
H25B	-0.0236	0.6617	-0.4390	0.045*	
C26	0.0986 (3)	0.8488 (3)	-0.4785 (2)	0.0456 (6)	
H26A	0.0535	0.7976	-0.5612	0.055*	
H26B	0.0997	0.9473	-0.4580	0.055*	
C1	0.0949 (3)	0.4974 (3)	0.1155 (3)	0.0598 (7)	
H1A	0.0573	0.5721	0.1520	0.072*	
H1B	0.0045	0.4081	0.0682	0.072*	
H1C	0.1706	0.5328	0.0672	0.072*	
O1	0.1702 (3)	0.4678 (3)	0.20073 (19)	0.0787 (8)	
H1	0.1856	0.3926	0.1706	0.094*	
B1	0.3719 (2)	0.2457 (2)	0.19180 (16)	0.0263 (4)	
F1	0.4132 (11)	0.1347 (10)	0.1303 (10)	0.0303 (13)	0.461 (19)
F2	0.2428 (8)	0.2394 (9)	0.1193 (6)	0.0313 (12)	0.461 (19)
F3	0.4916 (8)	0.3783 (7)	0.2162 (11)	0.0487 (18)	0.461 (19)
F4	0.3177 (7)	0.2190 (9)	0.2882 (5)	0.0395 (15)	0.461 (19)
F1A	0.4089 (12)	0.1282 (10)	0.1456 (10)	0.045 (2)	0.539 (19)
F2A	0.2240 (8)	0.2191 (10)	0.1397 (9)	0.0575 (19)	0.539 (19)
F3A	0.4837 (8)	0.3671 (8)	0.1725 (8)	0.0457 (13)	0.539 (19)
F4A	0.3783 (17)	0.2765 (12)	0.3073 (4)	0.080 (2)	0.539 (19)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag	0.03598 (8)	0.02979 (7)	0.02076 (7)	0.01318 (6)	0.00356 (5)	0.01063 (5)
N11	0.020 (4)	0.016 (3)	0.025 (4)	0.013 (3)	0.012 (3)	0.008 (3)
N12	0.021 (3)	0.022 (2)	0.020 (3)	0.012 (2)	-0.003 (2)	0.002 (2)
N13	0.0202 (18)	0.018 (2)	0.045 (4)	0.0087 (15)	0.004 (3)	0.012 (2)

O11	0.040 (4)	0.050 (5)	0.0295 (18)	0.023 (3)	0.007 (3)	0.004 (3)
O12	0.040 (2)	0.034 (2)	0.055 (4)	0.0256 (18)	0.018 (2)	0.010 (2)
O13	0.0310 (19)	0.0325 (17)	0.051 (4)	0.0180 (14)	-0.004 (2)	0.015 (2)
C11	0.026 (4)	0.020 (3)	0.024 (3)	0.010 (3)	0.008 (3)	0.010 (3)
C12	0.019 (2)	0.015 (2)	0.024 (4)	0.0092 (18)	0.004 (3)	0.002 (3)
C13	0.007 (3)	0.009 (3)	0.025 (2)	-0.002 (3)	0.0010 (18)	0.0077 (17)
C14	0.054 (6)	0.047 (7)	0.028 (6)	0.033 (5)	-0.004 (4)	0.016 (5)
C15	0.035 (4)	0.025 (2)	0.024 (3)	0.018 (3)	-0.005 (2)	0.0024 (18)
C16	0.042 (5)	0.047 (4)	0.024 (3)	0.028 (4)	0.002 (3)	0.011 (2)
N11A	0.033 (5)	0.027 (6)	0.027 (5)	0.011 (4)	0.007 (3)	0.014 (4)
N12A	0.020 (2)	0.0159 (18)	0.019 (3)	0.0107 (16)	0.0043 (19)	0.004 (2)
N13A	0.019 (2)	0.020 (2)	0.032 (3)	0.0085 (16)	0.006 (2)	0.012 (2)
O11A	0.031 (3)	0.036 (3)	0.0304 (18)	0.008 (2)	0.010 (3)	0.0074 (17)
O12A	0.041 (3)	0.049 (3)	0.050 (3)	0.035 (3)	0.030 (3)	0.032 (3)
O13A	0.034 (2)	0.0296 (15)	0.033 (3)	0.0163 (13)	-0.0055 (17)	0.0111 (16)
C11A	0.025 (4)	0.024 (3)	0.021 (3)	0.015 (3)	0.009 (3)	0.010 (3)
C12A	0.0166 (19)	0.018 (3)	0.016 (3)	0.0060 (19)	0.001 (2)	0.007 (3)
C13A	0.006 (3)	0.008 (3)	0.022 (2)	-0.004 (3)	0.0011 (17)	0.0072 (17)
C14A	0.057 (6)	0.042 (6)	0.023 (5)	0.045 (6)	0.018 (4)	0.009 (4)
C15A	0.024 (3)	0.0262 (19)	0.015 (2)	0.012 (2)	0.0007 (16)	0.0060 (14)
C16A	0.037 (4)	0.025 (2)	0.022 (2)	0.012 (3)	0.009 (3)	0.0104 (16)
N21	0.0342 (8)	0.0243 (7)	0.0263 (7)	0.0143 (6)	0.0021 (6)	0.0072 (6)
N22	0.0333 (8)	0.0331 (8)	0.0280 (7)	0.0192 (6)	0.0108 (6)	0.0207 (6)
N23	0.0287 (8)	0.0314 (9)	0.0875 (16)	0.0166 (7)	0.0213 (9)	0.0373 (10)
O21	0.0335 (7)	0.0498 (8)	0.0312 (7)	0.0258 (6)	0.0119 (5)	0.0185 (6)
O22	0.0565 (11)	0.0321 (9)	0.1166 (19)	0.0296 (9)	-0.0178 (12)	-0.0075 (10)
O23	0.1030 (17)	0.131 (2)	0.0963 (16)	0.1008 (17)	0.0702 (14)	0.0957 (17)
C21	0.0299 (9)	0.0222 (8)	0.0454 (11)	0.0116 (7)	0.0000 (8)	0.0007 (7)
C22	0.0243 (8)	0.0201 (7)	0.0522 (11)	0.0102 (6)	0.0080 (7)	0.0173 (8)
C23	0.0348 (9)	0.0254 (8)	0.0225 (7)	0.0163 (7)	0.0063 (6)	0.0135 (6)
C24	0.0541 (12)	0.0358 (10)	0.0248 (8)	0.0301 (9)	0.0044 (8)	0.0089 (7)
C25	0.0317 (9)	0.0647 (14)	0.0302 (9)	0.0243 (9)	0.0083 (7)	0.0295 (9)
C26	0.0383 (11)	0.0888 (18)	0.0413 (11)	0.0383 (12)	0.0188 (9)	0.0481 (13)
C1	0.0312 (11)	0.0628 (17)	0.0691 (18)	0.0079 (11)	0.0019 (11)	0.0160 (14)
O1	0.0709 (13)	0.0848 (15)	0.0653 (13)	0.0613 (13)	-0.0276 (11)	-0.0290 (11)
B1	0.0270 (9)	0.0233 (8)	0.0235 (8)	0.0066 (7)	0.0073 (7)	0.0053 (7)
F1	0.037 (3)	0.028 (2)	0.031 (2)	0.0183 (18)	0.0130 (18)	0.0090 (19)
F2	0.037 (3)	0.029 (2)	0.0255 (16)	0.015 (2)	0.0004 (15)	0.0061 (13)
F3	0.0250 (17)	0.0217 (15)	0.083 (5)	0.0004 (12)	0.007 (3)	0.007 (3)
F4	0.039 (2)	0.055 (3)	0.0244 (17)	0.0204 (19)	0.0125 (14)	0.0110 (17)
F1A	0.070 (4)	0.031 (2)	0.039 (3)	0.025 (2)	0.010 (2)	0.0146 (19)
F2A	0.0192 (14)	0.047 (3)	0.074 (4)	0.0064 (16)	-0.001 (2)	-0.015 (3)
F3A	0.0341 (16)	0.0302 (19)	0.068 (3)	0.0038 (13)	0.006 (2)	0.023 (2)
F4A	0.125 (6)	0.081 (4)	0.0201 (16)	0.033 (5)	0.019 (2)	0.008 (2)

Geometric parameters (\AA , $^{\circ}$)

Ag—N11	2.082 (15)	C15A—C16A	1.510 (9)
Ag—N21	2.1193 (15)	C15A—H15C	0.9900
Ag—N11A	2.163 (16)	C15A—H15D	0.9900
N11—C13	1.325 (9)	C16A—H16C	0.9900
N11—C11	1.365 (9)	C16A—H16D	0.9900
N12—C13	1.356 (9)	N21—C23	1.338 (2)
N12—C12	1.389 (9)	N21—C21	1.365 (2)
N12—C15	1.467 (8)	N22—C23	1.350 (2)
N13—O12	1.224 (6)	N22—C22	1.380 (3)
N13—O13	1.233 (6)	N22—C25	1.466 (3)
N13—C12	1.436 (8)	N23—O23	1.212 (3)
O11—C16	1.431 (9)	N23—O22	1.221 (3)
O11—H11A	0.8400	N23—C22	1.429 (2)
C11—C12	1.345 (9)	O21—C26	1.413 (2)
C11—H11B	0.9500	O21—H21A	0.8400
C13—C14	1.479 (9)	C21—C22	1.349 (3)
C14—H14A	0.9800	C21—H21B	0.9500
C14—H14B	0.9800	C23—C24	1.486 (2)
C14—H14C	0.9800	C24—H24A	0.9800
C15—C16	1.509 (9)	C24—H24B	0.9800
C15—H15A	0.9900	C24—H24C	0.9800
C15—H15B	0.9900	C25—C26	1.527 (3)
C16—H16A	0.9900	C25—H25A	0.9900
C16—H16B	0.9900	C25—H25B	0.9900
N11A—C13A	1.365 (8)	C26—H26A	0.9900
N11A—C11A	1.379 (9)	C26—H26B	0.9900
N12A—C13A	1.369 (9)	C1—O1	1.410 (4)
N12A—C12A	1.389 (9)	C1—H1A	0.9800
N12A—C15A	1.483 (8)	C1—H1B	0.9800
N13A—O13A	1.226 (5)	C1—H1C	0.9800
N13A—O12A	1.231 (6)	O1—H1	0.8400
N13A—C12A	1.445 (8)	B1—F3	1.346 (6)
O11A—C16A	1.430 (9)	B1—F4A	1.366 (4)
O11A—H11C	0.8400	B1—F2A	1.373 (6)
C11A—C12A	1.355 (9)	B1—F4	1.381 (4)
C11A—H11D	0.9500	B1—F1A	1.382 (7)
C13A—C14A	1.491 (9)	B1—F1	1.395 (6)
C14A—H14D	0.9800	B1—F3A	1.399 (5)
C14A—H14E	0.9800	B1—F2	1.428 (6)
C14A—H14F	0.9800		
N11—Ag—N21	175.7 (5)	N12A—C15A—H15D	108.9
C13—N11—C11	111.5 (7)	C16A—C15A—H15D	108.9
C13—N11—Ag	128.4 (6)	H15C—C15A—H15D	107.7
C11—N11—Ag	121.4 (7)	O11A—C16A—C15A	112.9 (7)
C13—N12—C12	106.6 (7)	O11A—C16A—H16C	109.0

C13—N12—C15	122.1 (7)	C15A—C16A—H16C	109.0
C12—N12—C15	131.2 (7)	O11A—C16A—H16D	109.0
O12—N13—O13	125.0 (6)	C15A—C16A—H16D	109.0
O12—N13—C12	115.7 (5)	H16C—C16A—H16D	107.8
O13—N13—C12	119.2 (5)	C23—N21—C21	106.99 (16)
C16—O11—H11A	109.5	C23—N21—Ag	126.88 (12)
C12—C11—N11	105.1 (7)	C21—N21—Ag	126.14 (13)
C12—C11—H11B	127.5	C23—N22—C22	105.84 (15)
N11—C11—H11B	127.5	C23—N22—C25	124.53 (16)
C11—C12—N12	109.3 (7)	C22—N22—C25	129.62 (16)
C11—C12—N13	127.3 (6)	O23—N23—O22	123.6 (2)
N12—C12—N13	123.2 (6)	O23—N23—C22	119.0 (2)
N11—C13—N12	107.3 (7)	O22—N23—C22	117.3 (2)
N11—C13—C14	122.3 (8)	C26—O21—H21A	109.5
N12—C13—C14	127.2 (10)	C22—C21—N21	108.23 (17)
C13—C14—H14A	109.5	C22—C21—H21B	125.9
C13—C14—H14B	109.5	N21—C21—H21B	125.9
H14A—C14—H14B	109.5	C21—C22—N22	108.31 (15)
C13—C14—H14C	109.5	C21—C22—N23	126.3 (2)
H14A—C14—H14C	109.5	N22—C22—N23	125.40 (19)
H14B—C14—H14C	109.5	N21—C23—N22	110.63 (15)
N12—C15—C16	112.0 (8)	N21—C23—C24	124.38 (15)
N12—C15—H15A	109.2	N22—C23—C24	124.97 (16)
C16—C15—H15A	109.2	C23—C24—H24A	109.5
N12—C15—H15B	109.2	C23—C24—H24B	109.5
C16—C15—H15B	109.2	H24A—C24—H24B	109.5
H15A—C15—H15B	107.9	C23—C24—H24C	109.5
O11—C16—C15	112.1 (8)	H24A—C24—H24C	109.5
O11—C16—H16A	109.2	H24B—C24—H24C	109.5
C15—C16—H16A	109.2	N22—C25—C26	110.98 (19)
O11—C16—H16B	109.2	N22—C25—H25A	109.4
C15—C16—H16B	109.2	C26—C25—H25A	109.4
H16A—C16—H16B	107.9	N22—C25—H25B	109.4
C13A—N11A—C11A	103.4 (6)	C26—C25—H25B	109.4
C13A—N11A—Ag	122.6 (6)	H25A—C25—H25B	108.0
C11A—N11A—Ag	132.1 (8)	O21—C26—C25	111.84 (15)
C13A—N12A—C12A	104.6 (7)	O21—C26—H26A	109.2
C13A—N12A—C15A	127.3 (7)	C25—C26—H26A	109.2
C12A—N12A—C15A	127.5 (7)	O21—C26—H26B	109.2
O13A—N13A—O12A	125.3 (5)	C25—C26—H26B	109.2
O13A—N13A—C12A	118.6 (5)	H26A—C26—H26B	107.9
O12A—N13A—C12A	116.1 (5)	O1—C1—H1A	109.5
C16A—O11A—H11C	109.5	O1—C1—H1B	109.5
C12A—C11A—N11A	110.0 (10)	H1A—C1—H1B	109.5
C12A—C11A—H11D	124.5	O1—C1—H1C	109.5
N11A—C11A—H11D	124.5	H1A—C1—H1C	109.5
C11A—C12A—N12A	107.9 (7)	H1B—C1—H1C	109.5
C11A—C12A—N13A	125.8 (7)	C1—O1—H1	109.5

N12A—C12A—N13A	126.2 (7)	F4A—B1—F2A	110.2 (4)
N11A—C13A—N12A	112.9 (7)	F3—B1—F4	112.8 (4)
N11A—C13A—C14A	123.6 (8)	F4A—B1—F1A	110.7 (5)
N12A—C13A—C14A	122.9 (9)	F2A—B1—F1A	110.7 (5)
C13A—C14A—H14D	109.5	F3—B1—F1	111.9 (5)
C13A—C14A—H14E	109.5	F4—B1—F1	109.9 (6)
H14D—C14A—H14E	109.5	F4A—B1—F3A	108.5 (4)
C13A—C14A—H14F	109.5	F2A—B1—F3A	108.4 (5)
H14D—C14A—H14F	109.5	F1A—B1—F3A	108.4 (5)
H14E—C14A—H14F	109.5	F3—B1—F2	107.7 (5)
N12A—C15A—C16A	113.4 (8)	F4—B1—F2	107.7 (3)
N12A—C15A—H15C	108.9	F1—B1—F2	106.5 (5)
C16A—C15A—H15C	108.9		
C13—N11—C11—C12	1.2 (14)	Ag—N11A—C13A—N12A	-178.7 (9)
Ag—N11—C11—C12	-179.8 (8)	C11A—N11A—C13A—C14A	166.5 (12)
N11—C11—C12—N12	1.2 (14)	Ag—N11A—C13A—C14A	-12 (2)
N11—C11—C12—N13	-173.5 (11)	C12A—N12A—C13A—N11A	-4.3 (15)
C13—N12—C12—C11	-3.1 (15)	C15A—N12A—C13A—N11A	-176.3 (12)
C15—N12—C12—C11	175.3 (13)	C12A—N12A—C13A—C14A	-168.1 (12)
C13—N12—C12—N13	171.8 (10)	C15A—N12A—C13A—C14A	20 (2)
C15—N12—C12—N13	-10 (2)	C13A—N12A—C15A—C16A	93.2 (15)
O12—N13—C12—C11	-10.3 (15)	C12A—N12A—C15A—C16A	-77.0 (15)
O13—N13—C12—C11	168.2 (11)	N12A—C15A—C16A—O11A	-71.9 (10)
O12—N13—C12—N12	175.6 (11)	C23—N21—C21—C22	-0.5 (2)
O13—N13—C12—N12	-5.8 (16)	Ag—N21—C21—C22	179.02 (13)
C11—N11—C13—N12	-3.2 (14)	N21—C21—C22—N22	0.6 (2)
Ag—N11—C13—N12	178.0 (9)	N21—C21—C22—N23	-179.57 (17)
C11—N11—C13—C14	-168.5 (12)	C23—N22—C22—C21	-0.4 (2)
Ag—N11—C13—C14	10 (2)	C25—N22—C22—C21	-179.75 (19)
C12—N12—C13—N11	3.8 (14)	C23—N22—C22—N23	179.73 (17)
C15—N12—C13—N11	-174.8 (11)	C25—N22—C22—N23	0.4 (3)
C12—N12—C13—C14	167.9 (13)	O23—N23—C22—C21	168.6 (2)
C15—N12—C13—C14	-11 (2)	O22—N23—C22—C21	-10.3 (3)
C13—N12—C15—C16	103.4 (13)	O23—N23—C22—N22	-11.6 (3)
C12—N12—C15—C16	-74.8 (16)	O22—N23—C22—N22	169.5 (2)
N12—C15—C16—O11	-68.9 (10)	C21—N21—C23—N22	0.2 (2)
C13A—N11A—C11A—C12A	1.5 (18)	Ag—N21—C23—N22	-179.28 (12)
Ag—N11A—C11A—C12A	-179.9 (13)	C21—N21—C23—C24	-178.24 (19)
N11A—C11A—C12A—N12A	-2.1 (15)	Ag—N21—C23—C24	2.2 (3)
N11A—C11A—C12A—N13A	-179.3 (10)	C22—N22—C23—N21	0.1 (2)
C13A—N12A—C12A—C11A	3.7 (15)	C25—N22—C23—N21	179.47 (17)
C15A—N12A—C12A—C11A	175.8 (12)	C22—N22—C23—C24	178.57 (18)
C13A—N12A—C12A—N13A	-179.0 (10)	C25—N22—C23—C24	-2.1 (3)
C15A—N12A—C12A—N13A	-7 (2)	C23—N22—C25—C26	-96.7 (2)
O13A—N13A—C12A—C11A	176.7 (10)	C22—N22—C25—C26	82.5 (2)
O12A—N13A—C12A—C11A	-2.9 (15)	N22—C25—C26—O21	52.0 (3)
O13A—N13A—C12A—N12A	-0.1 (16)	C13—N11—N21—C23	-160.8 (9)

O12A—N13A—C12A—N12A	−179.7 (11)	C13A—N11A—N21—C23	−148.6 (11)
C11A—N11A—C13A—N12A	0.8 (18)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···F2	0.84	1.84	2.673 (9)	173
O11—H11 <i>A</i> ···O1	0.84	1.86	2.697 (10)	175
O21—H21 <i>A</i> ···O11 <i>i</i>	0.84	1.91	2.726 (11)	164
O21—H21 <i>A</i> ···O11 <i>A</i> ⁱ	0.84	1.87	2.712 (11)	176

Symmetry code: (i) $x, y, z-1$.