

Pace University

DigitalCommons@Pace

Faculty Papers

Dyson College of Arts & Sciences

2013

Mass Spectrometric Analysis of Oxidized Eicosapentaenoic Acid Sodium Salt

Kelsey D. Jordan

Rita K. Upmacis

Follow this and additional works at: https://digitalcommons.pace.edu/dyson_fac

 Part of the [Chemistry Commons](#)

ORIGINAL RESEARCH

OPEN ACCESS
Full open access to this and
thousands of other papers at
<http://www.la-press.com>.

Mass Spectrometric Analysis of Oxidized Eicosapentaenoic Acid Sodium Salt

Kelsey D. Jordan and Rita K. Upmacis

Haskins Laboratories, Department of Chemistry and Physical Sciences, Pace University, New York, NY.
Corresponding author email: rupmacis@pace.edu

Abstract: Eicosapentaenoic acid (EPA) is an omega-3 polyunsaturated fatty acid (PUFA) with 20 carbon atoms and 5 carbon-carbon double bonds. Mammalian cells cannot synthesize long chain PUFAs such as EPA de novo, and, thus, the most effective way to enrich cells in EPA is by dietary intake of fish oils. EPA supplementation causes an increase in its concentration in plasma lipids and in cell membrane phospholipids. Many beneficial effects of EPA supplementation have been noted, including (1) the potential to sensitize cancerous tumors towards chemotherapy, (2) the promotion of cardiovascular health, and (3) the alleviation of some mental disorders, but results from clinical trials have sometimes been disparate. In this study, we report the use of mass spectrometry to investigate the autoxidation of EPA, thereby demonstrating the formation of a variety of oxidized products. The oxidative stress of the patient may affect the response to EPA and may, in part, explain divergent results from clinical trials.

Keywords: eicosapentaenoic acid, oxidation, mass spectrometry

Lipid Insights 2013:6 21–35

doi: [10.4137/LPI.S10862](https://doi.org/10.4137/LPI.S10862)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article published under the Creative Commons CC-BY-NC 3.0 license.



Introduction

Eicosapentaenoic acid (EPA) is an omega-3 polyunsaturated fatty acid (PUFA) with 20 carbon atoms and 5 carbon-carbon double bonds and is designated as 20:5 (ω -3) (Fig. 1). Mammalian cells cannot synthesize long chain PUFAs *de novo*, although EPA can be synthesized from the essential fatty acid, α -linolenic acid (ALA), 18:3 (ω -3), but conversion is minimal.¹ The most effective way to enrich cells in EPA is by dietary intake of fish oils.²

There is an active interest in understanding the effects of dietary supplementation of EPA as evidenced by the number of trials exploring the use of this substance in improving human health. For instance, with respect to cancer, marine-derived lipids, such as EPA, have the potential to sensitize tumors toward chemotherapy and enhance the cytotoxicity of certain anti-cancer drugs while preserving (or possibly) enhancing the integrity of non-tumor tissue.³ PUFAs are known to increase the presence of reactive oxygen species in cancer cells and may contribute to improving neutrophil function thus enhancing the action of the innate immune system.⁴ Further studies, however, are necessary to understand the mechanism whereby PUFAs selectively sensitize tumor cells but not normal tissue toward chemotherapeutic drugs. Certain cancers also lead to muscle loss or wasting (known as cancer cachexia), with patients exhibiting reduced levels of plasma omega-3 lipids that continue to decline over the course of the disease.⁵ In this case, ongoing studies and trials indicate that fish oil supplements are beneficial.⁶

There is also evidence from clinical trials and randomized controlled trials that omega-3 fatty acid supplementation decreases cardiovascular events and the risk of cardiac death.⁷ The exact mechanisms whereby omega-3 fatty acids contribute to decreasing cardiovascular events are not clear, although effects such as plaque stabilization,⁸ lipid profile alteration,⁹ triglyceride reduction,¹⁰ blood pressure reduction,¹¹ inhibition of inflammation,¹² or reduction in

arrhythmia have been suggested.¹³ While the general trend might be that these oils are beneficial, other data indicate either insufficient evidence^{14,15} or even an increased risk of cardiac death among patients advised to take oily fish.¹⁶ The reasons for these discrepancies are not clear, although potential concerns have been raised with regard to the quality and rigor of some trials and the fact that in some studies, patients were already receiving cardioprotective agents, for example, statins, such that further supplementation with an omega-3 fatty acid may not have had an impact.^{17,18}

The effects of EPA supplementation have also been investigated in patients with schizophrenia and other mental health-related disorders. Certain lipids are integral to the brain and significant membrane abnormalities have been identified in the brains of schizophrenic patients.¹⁹ The enzyme phospholipase A (PLA₂), which releases arachidonic acid, 20:4 (ω -6) from membrane phospholipids is thought to be overactive in schizophrenic brains. EPA supplementation may be beneficial since EPA may compete with arachidonic acid for incorporation into membrane phospholipids and thus inhibit the enzyme.²⁰ While a pilot study indicated that EPA is beneficial in the treatment of schizophrenia,²¹ a more recent meta-analysis of EPA data does not indicate a favorable role in schizophrenia and related psychological disorders.²² However, there may be confounding effects in previous trials since a differential response to EPA may occur depending on the history of pharmacological intervention, and an overall conclusion may not be available as yet.

The role of EPA may be controversial, but the indications are that EPA supplementation has, in some cases, shown promise in ameliorating or preventing several conditions. Therefore, it becomes important to understand the mechanisms of action. However, before biological mechanisms can be fully interpreted, it is essential to appreciate the chemical nature of EPA and the transformations that EPA can undergo. EPA shows reactivity toward oxygen, and is known to autoxidize readily. In this study, we investigated the autoxidation of EPA by mass spectrometry and report a number of oxidized products and their possible assignments. In this regard, we have reviewed some of the schemes involving the autoxidation of EPA.




Figure 1. Structure of eicosapentaenoic acid (EPA), 20:5 (ω -3).
Note: Positions 7, 10, 13, and 16 indicate bis-allylic (methylene) hydrogen locations.

Methods

Materials

The sodium salt of cis-5,8,11,14,17-eicosapentaenoic acid (EPA, $\geq 99\%$ purity) was purchased from Sigma Aldrich (St. Louis, MO). The sodium salt of EPA was stored prior to use as received (under nitrogen) in the solid form at $-80\text{ }^{\circ}\text{C}$. Henceforth, the sodium salt of EPA is referred to as “EPA.” Several portions of EPA (0.5 mg) in glass vials (approximately 2 cm in diameter and 6 cm in height) were allowed to oxidize in air at ambient temperature ($22.2\text{ }^{\circ}\text{C}$ – $25.4\text{ }^{\circ}\text{C}$) over 2 to 4 days under conditions of ~ 12 hours light and ~ 12 hours dark. At the time of analysis, EPA was dissolved in molecular grade water (1 mL) from Cellgro (distilled; deionized; DNase, RNase, and protease tested). While the sodium salt of EPA has limited solubility, we found that the amounts used (i.e. 0.5 mg/mL water) were fully soluble. Stock solutions were further diluted with molecular grade water before introduction to the mass spectrometer. When analyzing fresh EPA (i.e. when recording control mass spectra of the sodium salt of EPA), all manipulations were performed under nitrogen gas. In this regard, EPA was handled under nitrogen gas and dissolved in molecular grade water that had previously been purged with nitrogen.

Mass spectrometry

Native EPA and air-oxidized EPA samples were prepared as indicated above and dissolved in molecular grade water (0.17 mM). Samples were analyzed using an Applied Biosystems MDS SCIEX API 2000 instrument (AB SCIEX, Framingham, MA). The mass spectrometer was operated in negative ion mode with a mass range of m/z 250 to 600 amu, using a declustering potential (DP) of -60.0 V , a focusing potential (FP) of -400.0 V , and an entrance potential (EP) of -100.0 V . Nitrogen was used as both the sheath gas and the auxiliary gas. Data acquisition and analysis were performed using Analyst software, version 1.4. Samples were introduced via syringe injection at a flow rate of $10.00\text{ }\mu\text{L}/\text{min}$. Spectra (100 cycles) were accumulated over a 5-minute period.

Results

Mass spectral analysis of autoxidized EPA

The sodium salt of EPA, a white powder, oxidizes readily in air as evidenced by a change in color to

a yellow solid within a short period of time. When this process was monitored by mass spectrometry, several new species were observed within 0 to 4 days of exposure to air (Fig. 2). Mass spectral analysis of fresh native EPA gives rise to a main peak at $300.9\text{ }m/z$ (exact mass of acidic form of EPA = 302.2 amu), which corresponds to the deprotonated anion, that is, (exact mass-H)/ z . The main peak at $300.9\text{ }m/z$ displays the correct isotopic pattern, giving two shoulders at $302.1\text{ }m/z$ (exact mass = 303.2 amu) and $303.1\text{ }m/z$ (exact mass = 304.2 amu) (Fig. 2, blue spectrum). While care was taken to exclude oxygen, some small peaks mainly centered at $317.0\text{ }m/z$ and $333.1\text{ }m/z$ were also observed, indicating that either the EPA contains some oxidized product, or else a small amount of oxidation occurred during the handling process. However, mass spectrometric analysis of several samples of fresh native EPA consistently revealed small amounts of oxidized product.

Oxidation over several days revealed the formation of species with higher masses (Fig. 2 A–C). For example, after 2 days of air oxidation, small clusters of ions centered at 317.0 , 333.1 , 348.9 , and $365.0\text{ }m/z$ were observed to increase with a corresponding decrease of native EPA at $300.9\text{ }m/z$ (Fig. 2B, red spectrum). After 4 days of air oxidation, new species clustered at $380.7\text{ }m/z$ and $396.7\text{ }m/z$ appeared, accompanied by a decrease of parent EPA ($300.9\text{ }m/z$) together with a decrease in ions centered at 317.0 and $331.1\text{ }m/z$ (Fig. 2B, green spectrum). Notably, the average difference in mass between the peaks of greatest intensity in each cluster group is $15.97 \pm 0.07\text{ amu}$, corresponding to mass of an oxygen atom. Thus, the groups of ions centered at 317.0 , 333.1 , 348.9 , 365.0 , 380.7 , and $396.7\text{ }m/z$ represent, respectively, the addition of 1, 2, 3, 4, 5, and 6 oxygen atoms to native EPA. The facts that (1) the relative intensities of these peaks changed with time and (2) higher masses were observed after longer exposures to air indicate that their presence is real and not an artifact of the mass spectrometric process. In some cases, where the peaks representing oxidized EPA ions are larger, it is possible to discern an isotopic pattern (which is also clearly apparent in non-oxidized EPA at $300.9\text{ }m/z$).

While each cluster group can be ascribed to an ion having 1 to 6 additional oxygen atoms compared with the parent ion, it also appears that species within




Figure 2. Air oxidation of EPA leads to products of increased mass. Mass spectra of (A) native EPA (blue), and EPA oxidized in air for 2 and 4 days (red and green spectra, respectively). Native EPA gives a peak at 300.9 m/z (amu). (B) Magnified view of the same mass range as (a), with the mass of the dominant peak in each cluster group identified as the addition of 1, 2, 3, 4, 5, or 6 oxygen [O] atoms to EPA. (C) Magnified view of the 325–365 m/z range, highlighting the separation of peaks in each cluster of ions by 2.05 ± 0.04 (amu), representing the loss or gain of 2 hydrogen atoms. Peaks may also represent ions formed upon loss of water as, for instance, indicated for a species of mass 365.0 m/z losing H_2O to yield a species of mass 346.8 m/z.



each cluster group are separated by 2.05 ± 0.04 amu (Fig. 2C). This difference represents either an incremental gain or loss of 2 hydrogen atoms. While autoxidation pathways are covered in more detail below, possible mechanisms could include ring-opening of cyclic peroxides resulting in the formation of hydroxy compounds (i.e. gain of 2 hydrogen atoms) or, conversely, keto formation from an alcohol (i.e. loss of

2 hydrogen atoms). Loss of water from compounds could also result (i.e. loss of 18 amu).

The species that result from autoxidation of EPA appeared and disappeared at different rates, as shown by plots of peak intensity versus time (Fig. 3). The peak intensity of the parent EPA ion decreased over 4 days (Fig. 3A). Several species in the cluster groups with 1 or 2 oxygen atom additions




Figure 3. Variation of peak intensity with time. Peak intensities are monitored over time (0–4 days) to indicate the rate at which ions appear or disappear. (A) Parent EPA ion at 300.9 m/z and the cluster group of ions with (B) 1 oxygen [O] atom addition, (C) 2 oxygen atoms, (D) 3 oxygen atoms, (E) 4 oxygen atoms, (F) 5 oxygen atoms, and (G) 6 oxygen atoms.



(designated as EPA + 1[O] and EPA + 2[O] in Fig. 3B and C, respectively) also disappeared over 4 days. Conversely, some species (indicated by peaks at 312.9, 326.8, 328.8, and 330.8 *m/z*) were formed during this time, perhaps from precursors that are being reduced in intensity. The clusters of peaks representing the addition of 3, 4, 5, and 6 oxygen atoms all increased with time (Fig. 3D–G). Notably, the higher-mass species (e.g. EPA + 5[O] and EPA + 6[O]) only started to appear after 2 days, indicating that the higher-mass oxidized species are formed from the lower-mass oxidized species and that oxygen addition is a sequential process.

Comparison of the total intensity of peaks in the 290 to 450 *m/z* region shows that ~30% of the total intensity is lost after 2 days and is further reduced after 4 days (~50% that of the original spectrum). While a comparison of total intensity may not provide a consistent method of comparing spectra, the indication is that there is an overall loss of intensity. Although the loss of intensity is not fully understood, it is possible that some of the species formed are not detected by the mass spectrometric process or else occur at masses outside the region shown. Although oxidized EPA appears soluble giving rise to clear, yellow solutions, it is also possible that the solubility of some of the oxidized species is limited.

Discussion

In this study, we sought to identify the types of products formed when the sodium salt of EPA is autoxidized in air at ambient temperature. To the best of our knowledge, this represents the first study in which the autoxidation process was allowed to occur in the solid phase in the presence of air and light but in the absence of any solvent or exogenously-added peroxide/free-radical initiating species. Previous mass spectrometric studies of *in vitro* oxidized EPA products have involved the addition of free-radical initiating compounds to EPA dissolved in organic solvents.^{23,24} Gas chromatography coupled with mass spectrometry (GC-MS) provides a sensitive method for characterizing products of PUFA oxidation, but often requires the derivatization of products prior to analysis.²⁵ To prevent additional modification that may occur during the GC-MS process, an alternative method may be used that utilizes liquid chromatography to separate compounds before being introduced


to the ion source and characterized by tandem mass spectrometry (LC-MS/MS). These methods have previously been successfully applied to profiling oxidized lipids, for instance, in plasma and urine samples.^{26,27} In the current study, we applied electrospray ionization mass spectrometry to the direct observation of oxidized EPA species.

Our results indicated that species containing up to 6 oxygen atoms result from air oxidation (Fig. 2). In addition to these species, we also observed ions that were separated by 2.05 ± 0.04 amu from the main peak of each cluster group (and, in some cases, by 18 amu between peaks of different cluster groups). In order to make potential assignments, it is necessary to consider schemes of oxygen addition and autoxidation, as described below.

Basic products of EPA autoxidation

Unsaturated fatty acids are susceptible to oxidation due to the fact that the formation of a free radical on a carbon atom located between 2 carbon-carbon double bonds (otherwise known as a bis-allylic position) is remarkably stable.²⁸ Thus, it is relatively easy to remove a hydrogen atom from EPA to produce a radical that can interact with an oxygen molecule. EPA has four bis-allylic moieties, centered at C-7, C-10, C-13, and C-16 (Fig. 1).

Scheme 1 illustrates a mechanism of EPA oxygenation, which is based on the mechanism of arachidonic acid oxygenation.²³ EPA is similar in structure to arachidonic acid in that it has a 20-carbon backbone but, unlike arachidonic acid, EPA has 5 carbon-carbon double bonds (versus 4 in arachidonic acid) and is an omega-3 fatty acid (rather than arachidonic acid which is an omega-6 fatty acid). Due to their similarities in structure, similar mechanisms may apply to oxygenation of both EPA and arachidonic acid. Lipid oxidation can be initiated by certain transition metals, such as iron, but physiologically, lipid oxidation may occur enzymatically in a selective manner. For instance, oxygenation of arachidonic acid by the cyclooxygenase enzymes (COX-1 and COX-2) initiates the cascade leading to prostaglandin (PG) formation. Oxygenation by COX occurs when arachidonic acid lies in the cyclooxygenase channel with the C-13 position adjacent to tyrosine-385 in the polypeptide backbone. The catalytic process involves tyrosine cation radical production,



Scheme 1. Mechanism of EPA oxygenation. Oxygenation of EPA (1) is initiated by hydrogen abstraction of the C-13 position (2), allowing oxygen to add at C-11 (3), followed by endoperoxide formation (4), further cyclization (5) and the addition of a second oxygen to produce a bicyclic endoperoxide (6), which can be reduced to a F₃ isoprostane (7). For simplicity, the stereochemistry of the structures is not shown.


which is responsible for stereoselective abstraction of the 13-pro-S hydrogen from the C-13 position and thus initiating oxygenation.²⁹ EPA can readily replace arachidonic acid in the lipid bilayer and act as a substrate for the COX enzymes.

Physiologically, EPA oxidation is expected to occur enzymatically but, under situations of oxidative stress, nonenzymatic routes may also occur in living systems.³⁰ EPA is prone to peroxidation and can be oxidized nonenzymatically, often resulting in prostaglandin-like substances, called isoprostanes.²⁴ When only lipid radicals and oxygen are involved, the process is known as “autoxidation,” but similar mechanisms may be involved in both enzyme-catalyzed lipid oxidation and autoxidation. Thus, Scheme 1 illustrates a mechanism of EPA oxygenation with initial hydrogen abstraction occurring at the C-13 position. Rearrangement leads to radical formation at the C-11 position with subsequent radical trapping of oxygen, followed by the formation of a 5-membered endoperoxide ring, cyclization, further rearrangement and a second oxygenation at C-15. Reduction results in the production of F₃ isoprostane (also known as PGF_{3α}).

An important difference between enzyme-catalyzed lipid oxygenation and autoxidation is that enzyme-directed oxygenations are stereospecific. Autoxidation of arachidonic acid is well elucidated and can occur at different bis-allylic positions.³¹ In this regard, autoxidation of EPA can, in principle, be initiated at any of the C-7, C-10, C-13, or C-16 sites.²³ Novel oxidation products of EPA have previously been identified using mass spectrometric techniques including Ag⁺ coordination ionspray and atmospheric pressure chemical ionization mass spectrometry.²³ Importantly, these oxidation products were observed both in vitro and in vivo.²³ Scheme 2 illustrates the collection of hydroperoxyeicosapentaenoic acid (HpEPE) and hydroxyeicosapentaenoic (HEPE) isomers that could form following initiation at the 4 different bis-allylic positions. According to Scheme 2, the peaks observed at 317.0 and 333.1 m/z (Fig. 2) represent the addition of 1 and 2 oxygen atoms to EPA, respectively, and can potentially be ascribed to HEPE and HpEPE isomers of EPA (exact masses = 318.2 amu and 334.2 amu, respectively).

Monocyclic and serial cyclic peroxide formation

Monocyclic and serial cyclic peroxide products of EPA have previously been investigated.²³ Scheme 3A represents the type of monocyclic (1,2-dioxolane) and serial cyclic peroxides that can, in principle, form




Scheme 2. Hydrogen abstraction at C-7, C-10, C-13, and C-16 yields a wide variety of hydroperoxyeicosapentaenoic acid (HpEPE) and hydroxyeicosapentaenoic acid (HEPE) products. Hydrogen abstraction of EPA (1) at C-7, leads to the 5- and 9-series of HEPE and HpEPE products (8–11); at C-10 results in the 8- and 12-series of HEPE and HpEPE products (12–15); at C-13 yields the 11- and 15-series of HEPE and HpEPE products (16–19), and at C-16 leads to the 14- and 18-series of HEPE and HpEPE products (20–23).

when initial hydrogen abstraction occurs at C-13, followed by rearrangement to yield a carbon radical at C-11 to allow initial oxygen addition, that is, (1) to (3) in Scheme 1. In this particular example, EPA can potentially gain 4 oxygen atoms (monocyclic peroxide), or

else peroxidation can go further yielding a structure with 6 oxygen atoms (serial cyclic peroxide).

The position of initial hydrogen abstraction and subsequent location of the carbon radical (with respect to its proximity to carbon-carbon double bonds)



Scheme 3. Oxygenation of EPA yields monocyclic and serial cyclic peroxides. **(A)** Following hydrogen abstraction from C-13 of EPA (1) and subsequent endoperoxide radical formation (4), oxygen addition can occur at the C-8 position leading to the formation of a monocyclic peroxide (24), with possible further oxygen addition at C-5 yielding a serial cyclic peroxide (25). **(B)** Hydrogen abstraction from C-7 leading to a radical on C-9 allows addition of 4 oxygen atoms (26), 6 oxygen atoms (27) or 8 oxygen atoms (28).

determine the extent of peroxidation and dioxolane formation. For instance, structures are possible with only 2 oxygen atom additions (i.e. when oxygen first adds to a carbon radical on either C-5 or C-18) or up to 8 oxygen atom additions (i.e. when oxygen first adds to C-9 or C-14 radicals). For example, hydrogen abstraction at C-7 leading to a radical on C-9 allows addition of 4, 6, or 8 oxygen atoms, as shown in Scheme 3B. When hydrogen abstraction at

C-7 instead leads to radical formation on C-5, only 2 oxygen atom additions are possible with the result that 5-HpEPE is formed.

In summary, depending on the position of carbon radical formation, it is possible to form HpEPEs (2 oxygen atoms; exact mass = 334.21 amu), monocyclic peroxides (4 oxygen atoms; exact mass = 366.20 amu) and serial cyclic peroxides containing 2 dioxolane rings (6 oxygen atoms, exact



mass = 398.19 amu) or 3 dioxolane rings (8 oxygen atoms, exact mass = 430.18 amu). As these rearrangements and additions occur, EPA, which is not initially conjugated, becomes a partially conjugated system. Upon reduction of the hydroperoxyl moiety to yield a hydroxyl group, it is possible to form the corresponding dioxolane structures with 1, 3, 5, and 7 oxygen atoms (exact masses = 318.22, 350.21, 382.20, and 414.19 amu, respectively). The mass spectra did not yield evidence of a structure corresponding to 8 oxygen atom additions (exact mass = 430.18 amu), but it was possible to observe a species with $m/z = 413.0$, potentially indicative of a structure with 7 oxygen atoms (exact mass = 414.19 amu; data not shown).

Hydrogen abstraction at multiple bis-allylic carbon positions

Hydrogen abstraction that occurs at multiple bis-allylic carbon positions in EPA can result in structures that contain multiple hydroxyl and/or peroxy adducts. This is illustrated in Scheme 4 for 5-HpEPE that results from initial hydrogen abstraction at C-7. Subsequent removal of a hydrogen atom from C-10 results in diHpEPE, which has another bis-allylic position at C-16 that can undergo further hydrogen abstraction. The resulting structures include triHpEPE and triHEPE. The triHEPE compounds are also known as resolvins and have been found to have anti-inflammatory properties.^{32–34} Notably, in human endothelial cells, aspirin therapy causes COX-2 to switch its catalytic activity from forming prostanoids to producing specific HEPEs and novel trihydroxy-containing resolvins.³² In particular, 5,12,18-triHEPE, also known as RvE1, has been associated with having potent anti-inflammatory and analgesic properties.³⁵


Products of enzymatic oxygenation versus autoxidation

EPA can readily replace arachidonic acid in the lipid bilayer and act as a substrate for the COX enzymes. When arachidonic acid is the COX substrate, the prostaglandin (PG-2) series of products is formed, where the number 2 corresponds to the number of carbon-carbon double bonds in the product. However, when COX metabolizes EPA, the PG-3 series is formed, with products containing 3 double bonds.³⁶ Eicosanoids that are produced with arachidonic acid

as the initiating substrate are often associated with proatherogenic and proinflammatory properties. In contrast, eicosanoids derived from EPA are associated with less biological activity and therefore have less of an impact on the process of inflammation or proliferation.³⁷ For instance, oxidized EPA significantly inhibited human neutrophil and monocyte adhesion to endothelial cells by preventing expression of the endothelial adhesion receptor via a mechanism involving activation of the peroxisome proliferator-activated receptor α (PPAR α) and subsequent inhibition of NF- κ B.^{38,39}

Scheme 5 indicates that following bicyclic endoperoxide (PGG₃) formation, reduction occurs forming PGH₃, which can then be converted to several different types of eicosanoids, depending on the available enzymes. In the presence of PGE synthase (PGES) or PGDS, the products PGE₃ and PGD₃ can form. Interestingly, it has previously been noted that both PGE₂ and PGD₂ (the analogous products of arachidonic acid metabolism) can form independent of COX.³⁰ In general, although the products that result from enzymatic catalysis and autoxidation may be similar, there may be important distinctions in cis-trans orientations of side chains with respect to the prostane ring, with non-enzymatic routes favoring racemic mixtures. However, for COX-independent formation of PGE₂ and PGD₂, the products were found to be identical to COX-derived PGE₂ and PGD₂.³⁰ Thus, it may be possible for PGE₃ and PGD₃ to result from autoxidation. PGA₃ and PGJ₃ (not shown) result from dehydration of PGE₃ and PGD₃, respectively. Interestingly, it has previously been found that incubation of EPA with free-radical generating compounds in ethanol leads to the identification by mass spectrometry of PGA₃ and PGJ₃-like products (termed A₃/J₃ isoprostanes).²⁴ Thus, products similar to PGA₃ and PGJ₃ may also result during autoxidation.

Further products requiring enzymatic catalysis include PGI₃ (catalyzed by prostacyclin synthase), which is likely unstable and converts to 6-keto-PGF_{3 α} (as PGI₂ converts readily to 6-keto-PGF_{2 α}). Importantly, the arachidonate-derived PGI₂ is a potent vasodilator and inhibitor of platelet aggregation. Platelet-derived thromboxane synthase would convert PGH₃ to thromboxane B₃, the stable hydrolysis product of TxA₃. Notably, the arachidonate-derived




Scheme 4. Hydrogen abstraction at multiple bis-allylic carbon positions yields a mix of hydroxy and hydroperoxy products. Initial hydrogen abstraction at C-7 results in 5-HpEPE (8), which can be reduced to 5-HEPE (9), or alternately, be subject to removal of a hydrogen atom at C-10 to form diHpEPE (30). Further hydrogen abstraction at the bis-allylic position at C-16 results in the formation of structures that include triHpEPE (34) and triHEPE (37), as well as other structures (and isomers) containing hydroxyl and peroxy groups.

TxA₂ has properties completely opposed to PGI₂, in that it is a potent inducer of platelet aggregation and vasoconstrictor. A diet rich in EPA has been found to lead to both PGI₃ and TxA₃ production and a concomitant reduction in thrombogenic TxA₂.⁴⁰

Possible assignments of EPA autoxidation products

Possible assignments of EPA autoxidation products observed by mass spectrometry are listed in Table 1. Compounds, outlined in Schemes 1–5, that result from



Scheme 5. EPA metabolism by cyclooxygenase and other enzymes. Following bicyclic endoperoxide (PGG₃) formation (6), reduction occurs forming PGH₃ (38), which can then be converted to several types of eicosanoids, in the presence of different enzymes. PGE synthase (PGES) or PGDS converts PGH₃ to PGE₃ (39) and PGD₃ (40), respectively. PGA₃ (43) and PGI₃ (not shown) result from dehydration of PGE₃ (39) and PGD₃ (40), respectively. Prostacyclin synthase converts PGH₃ to PGI₃ (41), which is likely unstable and dehydrates to 6-keto-PGF_{3α} (44). Platelet-derived thromboxane synthase converts PGH₃ to thromboxane B₃ (42), the stable hydrolysis product of TxA₃.

non-enzymatic oxidation of EPA include HEPEs, HpEPEs, monocyclic peroxides, bicyclic endoperoxides, serial cyclic peroxides, and potentially a series of compounds with several hydroxyl or peroxy moieties are included as possible products. Table 1 shows that several isomers are possible for a single peak (and

not all possible isomers are considered). Other species were also noted that are incrementally 2.05 ± 0.04 amu lower or higher in mass than the main ions in each cluster group representing a gain of 1–6 oxygen atoms (Fig. 2). This mass is consistent with 2 hydrogen atoms. Possible mechanisms that involve a gain of

**Table 1.** Examples of possible assignments of mass spectrometric peaks.

Observed mass (anion) m/z (amu)	Formula	Possible assignments of the neutral form	Exact mass (amu)
300.9	C ₂₀ H ₃₀ O ₂	EPA (1)	302.22
317.0	C ₂₀ H ₃₀ O ₃	1 oxygen addition HEPE (Scheme 2) (9), (11), (13), (15), (17), (19), (21), (23)	318.22
330.8	C ₂₀ H ₂₈ O ₄	2 oxygen additions PGA ₃ (PGJ ₃) (Scheme 5) (43)	332.20
333.1	C ₂₀ H ₃₀ O ₄	HpEPE (Scheme 2) (8), (10), (12), (14), (16), (18), (20), (22) diHEPE (Scheme 4) (32)	334.21
348.9	C ₂₀ H ₃₀ O ₅	3 oxygen additions Hydroxy-HpEPE (Scheme 4) (31) triHEPE (Scheme 4) (37) PGH ₃ (Scheme 5) (38), PGE ₃ (39), PGD ₃ (40)	350.21
351.0	C ₂₀ H ₃₂ O ₅	F ₃ isoprostane (Scheme 1) (7)	352.22
365.0	C ₂₀ H ₃₀ O ₆	4 oxygen additions Bicyclic endoperoxide, PGG ₃ (Scheme 1) (6) diHpEPE (Scheme 4) (30) dihydroxy-HpEPE (Scheme 4) (35) Monocyclic peroxide (Scheme 3) (24), (26)	366.20
380.7	C ₂₀ H ₃₀ O ₇	5 oxygen additions Hydroxy-diHpEPE (Scheme 4) (36)	382.20
396.7	C ₂₀ H ₃₀ O ₈	6 oxygen additions Serial cyclic peroxide (Scheme 3) (25), (27) triHpEPE (Scheme 4) (34)	398.19

2 hydrogen atoms include ring-opening of cyclic peroxides resulting in hydroxyl compounds or, conversely, keto formation from an alcohol involves a loss of 2 hydrogen atoms. Dehydration could also contribute, as this would result in a loss of 18 amu.

Conclusion

Lipid peroxidation can be a function of physiological as well as pathophysiological processes. Dissecting which processes are the result of normal function versus those that are the result of disease and aging might be difficult. Given that enzymatic processes might also be accompanied by lipid peroxidation by non-enzymatic routes, it may be possible that, given their oxidative stress level, patients may differ in their response to EPA and thus give rise to divergent results in clinical trials. Given the propensity of EPA to react with oxygen in the absence of exogenously added free-radical initiators, it is important that patient populations receive the same quality of EPA to ensure that results from clinical trials are comparable. However, it may be possible that differential responses to EPA may still occur given the oxidative stress level of the patient.

Acknowledgements

RKU would like to thank Pace University for research support. RKU and KDJ would like to thank Pace University for an Undergraduate Student-Faculty Research Award.

Author Contributions

Conceived and designed the experiments: RKU. Analyzed the data: RKU, KDJ. Wrote the first draft of the manuscript: RKU. Contributed to the writing of the manuscript: RKU, KDJ. Agree with manuscript results and conclusions: RKU, KDJ. Jointly developed the structure and arguments for the paper: RKU, KDJ. Made critical revisions and approved final version: RKU, KDJ. All authors reviewed and approved of the final manuscript.

Funding

Funding for these studies was provided by Pace University.

Competing Interests

Author(s) disclose no potential conflicts of interest.



Disclosures and Ethics

As a requirement of publication the authors have provided signed confirmation of their compliance with ethical and legal obligations including but not limited to compliance with ICMJE authorship and competing interests guidelines, that the article is neither under consideration for publication nor published elsewhere, of their compliance with legal and ethical guidelines concerning human and animal research participants (if applicable), and that permission has been obtained for reproduction of any copyrighted material. This article was subject to blind, independent, expert peer review. The reviewers reported no competing interests. Provenance: the authors were invited to submit this paper.

References

- Anderson BM, Ma DW. Are all n-3 polyunsaturated fatty acids created equal? *Lipids Health Dis*. 2009;8:33.
- Macintosh BA, Ramsden CE, Fautot KR, et al. Low-n-6 and low-n-3 diets for use in clinical research. *Br J Nutr*. 2013;1–10.
- Hajjaji N, Bougnoux P. Selective sensitization of tumors to chemotherapy by marine-derived lipids: a review. *Cancer Treat Rev*. Jul 29, 2012. [Epub ahead of print.]
- Bonatto SJ, Oliveira HH, Nunes EA, et al. Fish oil supplementation improves neutrophil function during cancer chemotherapy. *Lipids*. 2012;47(4):383–9.
- Zuidgeest-van Leeuwen SD, van der Heijden MS, Rietveld T, et al. Fatty acid composition of plasma lipids in patients with pancreatic, lung and oesophageal cancer in comparison with healthy subjects. *Clin Nutr*. 2002;21(3):225–30.
- Murphy RA, Mourtzakis M, Mazurak VC. n-3 polyunsaturated fatty acids: the potential role for supplementation in cancer. *Curr Opin Clin Nutr Metab Care*. 2012;15(3):246–51.
- Delgado-Lista J, Perez-Martinez P, Lopez-Miranda J, Perez-Jimenez F. Long chain omega-3 fatty acids and cardiovascular disease: a systematic review. *Br J Nutr*. 2012;107(Suppl 2):S201–13.
- Thies F, Garry JM, Yaqoob P, et al. Association of n-3 polyunsaturated fatty acids with stability of atherosclerotic plaques: a randomised controlled trial. *Lancet*. 2003;361(9356):477–85.
- Yamagishi K, Nettleton JA, Folsom AR. Plasma fatty acid composition and incident heart failure in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Am Heart J*. 2008;156(5):965–74.
- Kris-Etherton PM, Harris WS, Appel LJ. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation*. 2002;106(21):2747–57.
- Morris MC, Sacks F, Rosner B. Does fish oil lower blood pressure? A meta-analysis of controlled trials. *Circulation*. 1993;88(2):523–33.
- Serhan CN, Chiang N, Van Dyke TE. Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol*. 2008;8(5):349–61.
- Lombardi F, Terranova P. Anti-arrhythmic properties of N-3 poly-unsaturated fatty acids (n-3 PUFA). *Curr Med Chem*. 2007;14(19):2070–80.
- Kwak SM, Myung SK, Lee YJ, Seo HG. Efficacy of omega-3 fatty acid supplements (eicosapentaenoic acid and docosahexaenoic acid) in the secondary prevention of cardiovascular disease: a meta-analysis of randomized, double-blind, placebo-controlled trials. *Arch Intern Med*. 2012;172(9):686–94.
- Kromhout D, Giltay EJ, Geleijnse JM. n-3 fatty acids and cardiovascular events after myocardial infarction. *N Engl J Med*. 2010;363(21):2015–2026.
- Burr ML, Ashfield-Watt PA, Dunstan FD, et al. Lack of benefit of dietary advice to men with angina: results of a controlled trial. *Eur J Clin Nutr*. 2003;57(2):193–200.
- Lavie CJ, Milani RV, Mehra MR, Ventura HO. Omega-3 polyunsaturated fatty acids and cardiovascular diseases. *J Am Coll Cardiol*. 2009;54(7):585–94.
- Kwak SM, Myung SK. The big ones that got away: omega-3 meta-analysis flawed by excluding the biggest fish oil trials-reply. *Arch Intern Med*. 2012;172(18):1427–8.
- Berger GE, Wood SJ, Pantelis C, Velakoulis D, Wellard RM, McGorry PD. Implications of lipid biology for the pathogenesis of schizophrenia. *Aust N Z J Psychiatry*. 2002;36(3):355–66.
- Law MH, Cotton RG, Berger GE. The role of phospholipases A2 in schizophrenia. *Mol Psychiatry*. 2006;11(6):547–56.
- Peet M, Brind J, Ramchand CN, Shah S, Vankar GK. Two double-blind placebo-controlled pilot studies of eicosapentaenoic acid in the treatment of schizophrenia. *Schizophr Res*. 2001;49(3):243–51.
- Fusar-Poli P, Berger G. Eicosapentaenoic acid interventions in schizophrenia: meta-analysis of randomized, placebo-controlled studies. *J Clin Psychopharmacol*. 2012;32(2):179–85.
- Yin H, Brooks JD, Gao L, Porter NA, Morrow JD. Identification of novel autoxidation products of the omega-3 fatty acid eicosapentaenoic acid in vitro and in vivo. *J Biol Chem*. 2007;282(41):29890–901.
- Brooks JD, Milne GL, Yin H, Sanchez SC, Porter NA, Morrow JD. Formation of highly reactive cyclopentenone isoprostane compounds (A3/J3-isoprostanes) in vivo from eicosapentaenoic acid. *J Biol Chem*. 2008;283(18):12043–55.
- Nourooz-Zadeh J, Liu EH, Anggard E, Halliwell B. F4-isoprostanes: a novel class of prostanoids formed during peroxidation of docosahexaenoic acid (DHA). *Biochem Biophys Res Commun*. 1998;242(2):338–44.
- Strassburg K, Huijbrechts AM, Kortekaas KA, et al. Quantitative profiling of oxylipins through comprehensive LC-MS/MS analysis: application in cardiac surgery. *Anal Bioanal Chem*. 2012;404(5):1413–26.
- Lawson JA, Kim S, Powell WS, FitzGerald GA, Rokach J. Oxidized derivatives of omega-3 fatty acids: identification of IPF3 alpha-VI in human urine. *J Lipid Res*. 2006;47(11):2515–24.
- Sevanian A, Hochstein P. Mechanisms and consequences of lipid peroxidation in biological systems. *Ann Rev Nutr*. 1985;5:365–90.
- Malkowski MG, Ginell SL, Smith WL, Garavito RM. The productive conformation of arachidonic acid bound to prostaglandin synthase. *Science*. 2000;289(5486):1933–7.
- Gao L, Zacker WE, Hasford JJ, et al. Formation of prostaglandins E2 and D2 via the isoprostane pathway: a mechanism for the generation of bioactive prostaglandins independent of cyclooxygenase. *J Biol Chem*. Aug 1, 2003;278(31):28479–89.
- Yin H, Havrilla CM, Gao L, Morrow JD, Porter NA. Mechanisms for the formation of isoprostane endoperoxides from arachidonic acid. “Dioxetane” intermediate versus beta-fragmentation of peroxy radicals. *J Biol Chem*. 2003;278(19):16720–5.
- Serhan CN, Clish CB, Brannon J, Colgan SP, Gronert K, Chiang N. Anti-microinflammatory lipid signals generated from dietary N-3 fatty acids via cyclooxygenase-2 and transcellular processing: a novel mechanism for NSAID and N-3 PUFA therapeutic actions. *J Physiol Pharmacol*. 2000;51(4 Pt 1):643–54.
- Serhan CN, Clish CB, Brannon J, Colgan SP, Chiang N, Gronert K. Novel functional sets of lipid-derived mediators with antiinflammatory actions generated from omega-3 fatty acids via cyclooxygenase 2-nonsteroidal antiinflammatory drugs and transcellular processing. *J Exp Med*. 2000;192(8):1197–204.
- Serhan CN, Hong S, Gronert K, et al. Resolvins: a family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals. *J Exp Med*. 2002;196(8):1025–37.
- Ji RR, Xu ZZ, Strichartz G, Serhan CN. Emerging roles of resolvins in the resolution of inflammation and pain. *Trends Neurosci*. 2011;34(11):599–609.
- Wada M, DeLong CJ, Hong YH, et al. Enzymes and receptors of prostaglandin pathways with arachidonic acid-derived versus eicosapentaenoic acid-derived substrates and products. *J Biol Chem*. 2007;282(31):22254–66.



37. Bagga D, Wang L, Farias-Eisner R, Glaspy JA, Reddy ST. Differential effects of prostaglandin derived from omega-6 and omega-3 polyunsaturated fatty acids on COX-2 expression and IL-6 secretion. *Proc Natl Acad Sci U S A*. 2003;100(4):1751–6.
38. Sethi S. Inhibition of leukocyte-endothelial interactions by oxidized omega-3 fatty acids: a novel mechanism for the anti-inflammatory effects of omega-3 fatty acids in fish oil. *Redox Rep*. 2002;7(6):369–78.
39. Sethi S, Ziouzenkova O, Ni H, Wagner DD, Plutzky J, Mayadas TN. Oxidized omega-3 fatty acids in fish oil inhibit leukocyte-endothelial interactions through activation of PPAR alpha. *Blood*. 2002;100(4):1340–6.
40. Fischer S, Weber PC. Thromboxane (TX)A₃ and prostaglandin (PG)I₃ are formed in man after dietary eicosapentaenoic acid: identification and quantification by capillary gas chromatography-electron impact mass spectrometry. *Biomed Mass Spectrom*. 1985;12(9):470–6.