



*Citation for published version:*

Airdrie, J, Lambe, S & Cooper, K 2023, 'Treating trauma-driven OCD with narrative exposure therapy alongside cognitive behavioural therapy', *The Cognitive Behaviour Therapist*, vol. 16, e2.  
<https://doi.org/10.1017/S1754470X22000605>

*DOI:*

[10.1017/S1754470X22000605](https://doi.org/10.1017/S1754470X22000605)

*Publication date:*

2023

*Document Version*

Peer reviewed version

[Link to publication](#)

This article has been published in the *Cognitive Behaviour Therapist*  
<https://doi.org/10.1017/S1754470X22000605> This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © The Author(s), 2023

**University of Bath**

**Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# **Treating Trauma-Driven OCD with Narrative Exposure Therapy alongside Cognitive Behavioural Therapy**

**Jac Airdrie, Sinéad Lambe and Kate Cooper**

## **Acknowledgements**

The author would like to thank Arwyn (pseudonym) for his consent to share his case study.

## **Conflict of Interest**

The authors declare no conflicts of interest. The authors received no specific financial support in relation to this study. All authors were employed by an NHS trust during the completion of the case study.

## **Treating Trauma-Driven OCD with Narrative Exposure Therapy alongside Cognitive Behavioural Therapy**

When Post Traumatic Stress Disorder (PTSD) co-occurs with Obsessive Compulsive Disorder (OCD), symptoms of the former can interfere with evidence-based treatment of the latter. As a result, exposure-based treatments are recommended for both OCD and PTSD, potentially facilitating a concurrent treatment approach. This case study describes the application of concurrent Cognitive Behaviour Therapy (CBT including Exposure and Response Prevention; ERP) for OCD and Narrative Exposure Therapy to treat a patient whose PTSD symptoms of intrusive images of memories and hyperarousal were interfering with standard CBT (including ERP) treatment for OCD. Following this concurrent approach, the patient's symptoms of OCD reduced to non-clinical levels and showed reliable improvement in PTSD symptoms. Whilst further, methodologically robust research is required, this case study highlights that this approach may be beneficial to the treatment of OCD where PTSD symptoms are impacting on treatment.

**Keywords:** OCD; PTSD; Trauma; ERP; Narrative Exposure Therapy; Case Study

### **Key Learning Objectives**

- To explore the literature considering explanations of the co-occurrence of OCD and PTSD symptomology
- To consider how symptoms of two mental health conditions can maintain one another and attenuate the effectiveness of evidence-based treatment for the other mental health condition
- Consider the use of concurrent therapeutic approaches to treat co-occurring mental health conditions

## **Treating Trauma-Driven OCD with Narrative Exposure Therapy alongside Cognitive Behavioural Therapy**

Obsessive Compulsive Disorder (OCD) is characterised by recurrent obsessions (intrusive, persistent, unwanted thoughts or images) and/or compulsions accompanied by marked distress and impairment in daily living (American Psychiatric Association; APA, 2013). Cognitive behavioural explanations of OCD (e.g. Salkovskis, 1985) postulate normal intrusive thoughts can become clinical obsessions when the thought is interpreted by an individual as having personal significance and are ego-dystonic (Salkovskis, 1985). These thoughts bring with them significant distress, which results in the individual completing some action (compulsion) to reduce this i.e. a neutralising behaviour. This reduction in distress is reinforcing, leading to an increase in occurrence of the neutralising behaviour. Central components of this model, such as unwanted intrusive thoughts and avoidance, are shared by a number of anxiety conditions, including Post Traumatic Stress Disorder (PTSD; APA, 2013).

PTSD consists of the co-existence of the re-experiencing of elements of previously experienced traumatic event(s), avoidance of stimuli associated with the event(s), hyperarousal and negative alterations in cognitions and mood persisting for at least one month (APA, 2013). The cognitive model of PTSD (Ehlers & Clark, 2000) suggests individuals process a traumatic event in a way that leads to a sense current threat. This arises due to negative appraisals of the trauma and/or its consequences which go beyond how the trauma might be generally appraised as horrific (e.g. the fact the trauma happened to them being significant) and due to the autobiographical memory lacking contextualisation. Instead it is characterised by strong associative memory and perceptual (sensory) priming leading to the memory having a sense of “nowness”. This problem becomes enduring due to

problematic behavioural and cognitive strategies (i.e. SSBs) preventing new interpretations and further memory (cognitive, sensory and affective) processing (Ehlers & Clark, 2000).

Clinicians have long documented the overlap between OCD and PTSD. Pitman (1993) presented the case of a veteran who developed OCD following traumatic experiences during his service, leading the author to coin the term Post-traumatic Obsessive Compulsive Disorder. Other case studies have since reported on similar presentations (e.g. De Moraes et al., 2008; Sasson et al., 2005). Empirical studies highlight the disorders can co-occur, with the National Comorbidity Survey Replication showing those with a current diagnosis of PTSD were 3.62 times more likely to have OCD than those without PTSD (Brown et al., 2001; albeit with a small sample  $n = 13$ ). Whilst Huppert et al., (2005) reviewed studies reporting rates of this comorbidity and found rates of PTSD between 6 and 22% in those with OCD.

With reference to Rachman (1991), it is clear these disorders share “psychological connectedness” (p.461) with several authors highlighting this link (Dinn et al., 1999; Gershuny et al., 2002; Huppert et al., 2005; Riggs, 2000). Examples of the logical and semantic relationship between certain traumatic experiences and subsequent OCD symptoms are evident in Pitman (1993) and the case series of Sasson et al., (2005), such as being covered by the blood and flesh of soldiers leading to excessive cleaning. Huppert et al., (2005) suggest rituals and avoidance behaviours develop to avoid intrusive memories associated with trauma. Through negative reinforcement via compulsions alleviating distress, these symptoms become independently maintained (Fostick et al., 2012). Once established, symptoms of PTSD and OCD have been shown to interact and exacerbate each other. In a case series, Gershuny et al., (2003) observed symptoms of OCD and PTSD were connected such that targeting OCD specific symptoms in treatment resulted in increases in PTSD symptoms, such as intrusive thoughts and flashbacks related to specific trauma memories.

Cognitive models of co-occurrence of these disorders (Riggs, 2000; Dinn et al., 1999) focus on the negative cognitive effects of exposure to trauma evoking anxiety. This leads to excessive labelling of stimuli as threatening which leads to compensatory mechanisms, such as compulsions or neutralising behaviours which reduce arousal and are therefore reinforced (Eysenck & Rachman, 1965). Gershuny et al., (2003) proposed OCD symptoms, such as excessive checking or hyper-attention, facilitate avoidance of emotional discomfort generated by trauma cues, and therefore the authors' conceptualised obsessive compulsive behaviours as serving a protective function against trauma-related memories and associated negative affect. Indeed, to paraphrase Riggs (2000), it's easier to worry about the dirt in your kitchen than the trauma of your combat experience.

Literature describing “mental contamination” which can be apparent in OCD also has relevance here given that this can relate to traumatic experiences. Mental contamination is a current internal sense of dirtiness that occurs in the absence of physical contact with a contaminant (Rachman, 2004) and is thought to develop following previous emotional or physical violations such as degradation, betrayal or abuse. Importantly, feelings of mental contamination can be evoked by memories and images, such as traumatic memories, and therefore such symptoms could be triggered by situations associated with such memories. As with cognitive theories of PTSD and OCD which contest that a misinterpretation of meaning can result in distressing symptoms, the cognitive theory of mental contamination postulates too that this arises due to a serious misinterpretation of the personal significance of a psychological or physical violation. This distress results in vigorous and repetitive washing to get rid of associated unpleasant feelings (Radomsky et al., 2017). Coughtry et al., (2012) found that 46% of individuals with OCD experienced mental contamination and that increased severity of mental contamination was associated with more severe OCD.

There is strong evidence for treatment of OCD from a cognitive behavioural framework. Cognitive behavioural Therapy (CBT) including Exposure and Response Prevention (ERP) has been shown to be effective for the treatment of OCD and is recommended as the first line treatment by the National Institute of Health and Care Excellence (NICE; 2005). CBT including ERP identifies and modifies dysfunctional appraisals of intrusions and symptom related beliefs. It also identifies avoided stimuli which evoke anxiety, and has individuals come into contact with these stimuli whilst refraining from engaging in compulsions or neutralising behaviours (ERP).

Trauma-focused therapies incorporating exposure such as cognitive behaviour therapy (Ehlers & Clark, 2008), Eye Movement Desensitisation and Reprocessing (EMDR; Shapiro, 2014) and Narrative Exposure Therapy (NET; Lely, et al., 2019; Robjant & Fazel, 2010) have been demonstrated to be effective in the treatment of PTSD (Bisson et al., 2013) and are recommended by NICE guidelines (although EMDR is only recommended if an individual has a preference for this and the trauma was not combat-related). The approaches being trauma-focused means that they directly focus on memories, thoughts and feelings related to the traumatic event (Watkins et al., 2018). NET was specifically developed for individuals who have experienced multiple traumatic events. It enables individuals to develop a coherent autobiographical narrative of their most significant experiences to help contextualise events that were highly arousing so that internal reminders and trauma related triggers have less dominance of the person's life (Schauer, Robjant, Elbert & Neuner, 2020).

Treatment of OCD has been found to be less efficacious when there is comorbid PTSD. Gershuny et al., (2002) found those with OCD alone improved with ERP while those with co-occurring PTSD did not, with decreases in symptoms of OCD during ERP leading to intensification of PTSD symptoms. Further, Gershuny et al., (2008) observed that for those with treatment resistant OCD, 82% had a history of trauma while 39% met criteria for PTSD,

suggesting its co-occurrence contributed to treatment resistance. Gershuny et al. (2003) suggested due to their dynamic connection, targeting either disorder in isolation may impede therapy effectiveness. Clinicians have therefore advocated for treatment strategies targeting symptoms of OCD and PTSD simultaneously (Gershuny et al., 2002), or consecutively (de Silva & Marks, 1999). A number of recent case studies have reported on the presentation and treatment of individuals with OCD and comorbid PTSD. Nijdam et al., (2013) reported the successful treatment of a male with PTSD and OCD with EMDR followed by ERP following sexual assault. It was documented that during treatment, remission of PTSD symptoms preceded that of OCD symptoms and the authors highlighted that applying EMDR first made it easier for the patient to reduce OCD symptoms because of decreased anxiety and trauma reminders. Stobie (2009) also reported on the case of women with PTSD and OCD which occurred following a physical assault. They used trauma-focused CBT initially to treat PTSD followed by CBT for OCD. They reported that they used the context of the trauma (e.g. where the event occurred) to provide rationale to help normalise the occurrence certain intrusions which were triggered by stimuli (a similar location to where the trauma occurred). This helped change symptoms meaning and in combination with CBT/ERP reduced the compulsions associated with this intrusion.

The current case study seeks to add to the literature on the treatment of those presenting with PTSD and OCD by presenting a case of a male in his late 20s with OCD and PTSD symptoms which emerged following a series of traumatic and highly aversive experiences. The case was initially conceptualised as OCD, however PTSD symptoms related to trauma became more prominent upon certain exposures and interfered with OCD treatment. The case study is novel in that it presents the use of concurrent NET and CBT with ERP as a treatment approach for comorbid OCD and PTSD symptoms.



## **Client Characteristics and Presenting Problems**

This study reports the case of Arwyn (anonymized), a caucasian male referred to a Psychological Therapies Team who had recently stopped working and become housebound due to exacerbation of OCD symptoms. Arwyn reported he had been struggling with OCD for around 10 years. His symptoms centred around rituals to ensure cleanliness and decontamination. He reported spending 6 hours a day on rituals. He used antibacterial spray on his clothes, bed, and furnishings and washed excessively. He had quarantined all items he perceived were contaminated to his utility room, which he described as representing all the unpleasant things that occurred during his life. He refrained from eating in his family home due to fears of contamination. This resulted in him losing 10kg of body weight.

### ***History of Traumatic and aversive events***

Arwyn described four significant events he perceived as contributing to his eventual presentation. Arwyn did not initially disclose all these events but reported that intrusive memories of these events were triggered in the course of ERP. When recalling these events to begin with he was highly avoidant of discussing them and his recall was fragmented. The details below became clearer following the completion of NET itself. Certain events meet criteria for a traumatic event according to the DSM-V definition (APA, 2013) in order to qualify for a diagnosis of PTSD (i.e. serious injury in this case). For others this may be less clear and might be more appropriately termed an aversive event (however, see Brewin et al., 2009 and Anders et al., 2011 for a discussion around the usefulness of Criterion A).

Arwyn was physically assaulted whilst visiting friends at a university he was planning to attend (1), this assault resulted in serious physical injury that required significant cosmetic treatment. As a result of this incident he stopped seeing this group of friends which including breaking up with his girlfriend at the time and generally became “isolative”. He also decided

not to go to university and instead started an apprenticeship in the local area which he saw as an opportunity to “toughen up”. During this he was subjected to significant verbal and physical bullying which occurred over a prolonged period of time and was perpetrated by individuals in a position of power over him (i.e. his supervisors). A number of bullying incidents resulted in Arwyn being put in a position of helplessness (2). He eventually dropped out of this apprenticeship and completed a university degree. By this time he reported displaying symptoms of OCD such as excessive cleaning. Following his degree he went travelling. Due to perceived differing hygiene norms in the country he visited and with OCD symptoms already present in the form of concerns about contamination with germs leading to illness or death, Arwyn was in a high state of stress throughout this time. It was in this context he believed he was cursed by a native of the country following a difficult interaction (3) and therefore engaged in neutralising rituals in line with the religions of that country to rid himself from the curse. Following his return, his OCD symptoms became more severe with large amounts of anxiety and SSBs. Arwyn experienced significant anxiety when confronted with persons he perceived as originating from a country he had visited during his travels, or people from his workplace, leading him to take measures to avoid contact with them. This culminated in the final difficult event in which Arwyn became overwhelmed by anxiety in work whilst trying to avoid contact with contaminants (proximity to individuals perceived to be from the country in which he went travelling or objects that could have come into contact with such individuals) and subsequently became trapped in his utility room, unable to leave due to the fear of spreading perceived contamination (4).

### ***Previous Treatment***

Arwyn received 12 private CBT sessions before entering the current service. Following these sessions he had reduced his spraying of disinfectant and delayed washing his hands after touching doors. However, he still presented with avoidant and compulsive behaviours such as

not eating which presented concerns about his health and therefore motivated a referral to secondary care. The exact nature of the CBT Arwyn received was unclear but his previous therapist (with Arwyn's consent) reported he had a good understanding of the CBT model of OCD and his formulation. They also highlighted that Arwyn was extremely avoidant of difficult emotions.

### ***Symptom Measures***

The Obsessive Compulsive Inventory revised (OCI-R; Foa et al., 2002) was used to measure OCD symptoms. It consists of 18 items assessing 6 symptoms of OCD (washing, checking, ordering, obsessing, hoarding and neutralising). The OCI-R has excellent internal consistency, test-retest reliability, convergent and discriminant validity (Foa et al., 2002). A score of 21 or above is used as a cut-off to indicate the likely presence of OCD. The mean score on the OCI-R in a large sample of individuals with OCD is 28 (SD = 13.53), while those without a diagnosis of OCD scored 18.82 (SD = 11.10; Foa et al., 2002).

The Impact of Events Scale-Revised (IES-R; Weiss, 2007), a 22-item self-report measure that assesses subjective distress caused by traumatic events, was used on two occasions during Arwyn's treatment, once before starting NET and once at his final session. A score of 33 and above represents best cut-off for probable PTSD (Creamer et al., 2003).

### ***Informed Consent***

The rationale and process for each aspect of treatment was provided to Arwyn, who consented to proceed with treatment and for his assessment and treatment to be written as a case study.

### ***Design***

Treatment initially used CBT (including ERP) for OCD (Salkovskis, 1985) without any focus on traumatic memories (Phase A) and later progressed to combine NET with CBT including ERP (Phase B). The choice to add NET to the treatment approach was a pragmatic one which was influenced by a plateau in symptom reduction and evidence of re-experiencing symptoms of PTSD (intrusive unwanted memories and arousal) upon exposure to certain OCD triggers. When questioned about these during exposure there was also a reluctance (avoidance) to discuss these events. Phase A was completed by a qualified Clinical Psychologist. For Phase B, NET was facilitated by the same qualified Clinical Psychologist, while CBT with ERP was carried out by a Trainee Clinical Psychologist.

## **Treatment**

### ***Phase A***

#### *Initial Formulation*

A formulation of Arwyn's difficulties was collaboratively developed (see Figure 1), based on a vicious flower maintenance cycle of OCD (Salkovskis, 1998). A longitudinal component was added to make sense of how the OCD developed, and a critical incident to account for the *why now?* question. It was hypothesised a combination of predisposing factors, such as perfectionist tendencies (which included high standards of cleanliness) and significant life events interactively combined to lead to an overwhelming experience of anxiety during the critical incident. During this incident Arwyn described that he became overwhelmed by anxiety at work due to beliefs that it was contaminated as a result of coming into contact with individuals who he perceived were from the country in which he was "cursed" during his travels. He believed that "germs" associated with people originating from the country in which he went travelling would result in illness and himself and his family dying. He perceived he lost control and "escaped" to his utility room at home and became "trapped" due to not

wanting to spread contamination. On Figure 1 the connection between the critical incident and maintenance factors is reciprocal due to the fact that this beliefs and anxiety around contamination were already present to a lesser extent and influenced his experience of this incident.

Arwyn developed SSBs in relation to both anticipatory and consequent anxiety (Salkovskis, 1991) in order to stop the feared outcome occurring. These avoidance and escape behaviours had the effect of preventing Arwyn discovering his beliefs were inaccurate (in line with Salkovskis, 1991), and prevented the extinction of the anxiety when exposed to the feared situations (de Silva & Rachman, 1984).

#### *Intervention directed by initial formulation*

Arwyn was initially treated using CBT (including ERP) which is the first line treatment recommended for OCD by NICE guidelines (2005). It was hypothesised Arwyn exposing himself to feared situations via ERP (following an exposure hierarchy) and challenging his beliefs about the consequence of this would lead him to learn his predictions were untrue and to a reduction in anxiety.

Sessions 1 to 20 involved the collaborative development of a formulation. Arwyn was introduced to the idea of Theory A (contact with contaminants (“germs” associated with country I went travelling in) in will lead to illness and possible death as well as intolerable levels of anxiety causing me to lose control) and Theory B (I am worried that contact with contaminants will lead to illness and possible death, anxiety is normal and will come down on its own accord) (Challacombe et al., 2011) and evidence for these were discussed using cognitive restructuring. Arwyn completed homework tasks of exposure work to explore which theory fitted best with his experience. While ERP was conducted on two occasions with the psychologist, these were on items lower down Arwyn’s hierarchy. Most of the exposure homework was done by Arwyn alone or with help of a support worker or family

members. The ERP tasks included not using his antibacterial spray and eating food prepared in the kitchen adjacent to the utility room.

Arwyn made progress on increasing his exposure to things lower down the hierarchy and reduced his ritualistic behaviour significantly. However, by session 20 Arwyn's OCI-R scores had plateaued, remaining within the clinical range and both he and his family were frustrated at his rate of progress. Arwyn reported that exposure to items more directly associated with traumatic events (e.g. items or persons associated with travelling, his old workplace or local areas where he was placed for his apprenticeship) he had experienced evoked intrusive memories and high levels of arousal he was unable to tolerate and therefore would quickly discontinue the exposure. When initially questioned about these events Arwyn was visibly distressed and reluctant to discuss them and reported that he had never spoken to anyone about them before. The account of what he was able to initially share about his memory was also very fragmented.

### ***Phase B***

#### *Revised formulation*

Arwyn's formulation was revised to highlight how intrusive memories and thoughts with negative interpretation about traumatic events brought about a sense of current threat and high levels of arousal. This further added to Arwyn's belief he would experience intolerable anxiety. The dotted arrows on Figure 1 represent aspects added to the revised formulation, whilst asterisks represent aspects directly associated with traumatic events Arwyn experienced. Arwyn reported voluntary suppression of the memories related to his past experiences and was not prepared to discuss these in depth at the start of therapy, a characteristic also shared with PTSD (Ehlers and Clark, 2000). According to the cognitive model of PTSD, traumatic memories fail to become encoded in autobiographical memory and

hence are experienced with a sense of current threat resulting in intrusions and physiological arousal. Due to the symptoms' aversive nature, strategies emerge to control them (e.g. avoidance of reminders of trauma) further preventing processing and encoding into autobiographical memory (Ehlers & Clark, 2000). Arwyn's symptoms could fit such a profile, as he experienced high levels of distress when placed in situations associated with trauma (i.e. sense of current threat) and developed strategies to control these symptoms such as avoiding his workplace and items and memories associated with travelling. Whilst these avoidance symptoms were related to his OCD, Arwyn also reported avoidance symptoms that were pre-morbid to his OCD in that he decided not to go university (where he was assaulted) and isolated himself from his friends and broke up with his girlfriend as a result of them reminding him of the first traumatic event (assault). Due to avoidance, Arwyn's sense of current threat (Ehlers & Clark, 2000) when exposed to situations associated with traumatic events remained high, making it difficult to complete ERPs.

*Figure 1.* Arwyn's formulation. The dashed lines were added at Phase B (revised formulation). - here

It was hypothesised applying NET to allow further processing of the traumatic memories and therefore elaboration and contextualisation (Ehlers & Clark, 2000) would lead to autobiographical storage of the memories, reducing sense of current threat and therefore arousal. This would increase the likelihood that Arwyn could tolerate exposures. NET was chosen primarily pragmatically above approaches such as TF-CBT or EMDR due to Arwyn having experienced a series of traumatic and aversive events throughout his early adulthood which meant that there were significant periods of Arwyn's history that he found it difficult to talk about. His traumatic experiences also shared some relationship with one another, such that his experience of a serious assault led to a decision not to go to university and to start an

apprenticeship. It was during this subsequent apprenticeship he experienced prolonged and significant physical and psychological bullying by individuals in a higher position of power than himself (supervisors). NET assists an individual to realign their traumatic experiences within the larger context of their life and their place in the world (Schauer, Schauer, Neuner & Elbert, 2011) and is specifically designed for individuals who have experienced multiple traumatic events (Robjant & Fazel, 2010) and therefore this was seen as a potential helpful approach for Arwyn. NET involves constructing a chronological account of someone's traumatic experiences allowing conversion of the disjointed recollection of traumatic experiences to a coherent narrative, providing contextualisation (Schauer et al., 2011). The aim of NET was to reduce Arwyn's current sense of threat and arousal levels experienced when exposed to avoided situations to help tolerate ERPs. The aim of CBT with ERP was to further challenge his beliefs and allow further anxiety response extinction (Rachman & Hodgson, 1980) and therefore reduce compulsive behaviour.

#### *NET for trauma memories*

Sessions 22, 25, 27, 28 and 31 used NET to allow increased processing of Arwyn's trauma memories of the assault at a university open day, his experience of bullying at his apprenticeship, his experience travelling, and his perception he was overwhelmed by anxiety whilst in work. Arwyn described his experiences to the Clinical Psychologist who wrote them out as a script. The psychologist asked questions of clarification around particular hotspots including around cognitions and affect that was experienced at the time with the aim of trying to re-connect "hot" (cognitive, emotional and physiological representations) and "cold" (context) memories (Neuner, Schauer, & Elbert, 2020). The writing of the script slowed down the pace to help processing of the sensory and emotional aspects of the memories. Arwyn was set between-session homework of reading through the script.



### *CBT with ERP*

Sessions 24, 26, 29, 30 and 32 used CBT with ERP on items of his hierarchy which shared a relationship with Arwyn's traumatic history (see Table 1). It was hypothesised that Arwyn exposing himself to avoided stimuli without engaging in neutralising behaviours would lead to a reduction in his obsessive beliefs, anxiety levels and compulsive behaviours when re-exposed to the same stimuli. During the first session, time was allocated to discuss the rationale for exposure with reference to his formulation (psychoeducation) and to update his exposure hierarchy (it should be emphasised all items were perceived as high up the hierarchy relative to previous ERPs he successfully managed during his previous 20 CBT sessions). The final hierarchy is depicted in Table 1.

#### *Table 1. Arwyn's exposure hierarchy - here*

An example session of exposure to the utility room is given below which was conducted according to the protocol outlined in Salkovskis (2007). Each ERP session followed a similar structure and due to the prolonged and field-based nature lasted 3 hours on average. An exposure practice form was used to identify Arwyn's predictions while also providing the opportunity to track his subjective units of distress (SUDs). Outlining predictions was important to address the cognitive aspect of OCD (e.g Meyer, 1966). Arwyn predicted he would become so anxious he would lose control (100% prediction). Prior to entering the utility room Arwyn was reminded to refrain from engaging in SSBs and the importance of not neutralising the exposure once we left the room was highlighted, as cognitively, going into a feared situation with the intention of undoing it can be considered as an avoidance strategy (Salkovskis, 1985). While exposure began with the Trainee Psychologist suggesting items to be touched, a shift was made to Arwyn independently selecting the items of exposure. Arwyn touched various items associated with his place of work and apprenticeship while his attempts to prevent spreading of the perceived

contamination (known as “tracking”) was also brought to his attention. It was important to ensure Arwyn was fully engaged with the stimuli (he had a tendency to use conversation as a distraction method), as Borkovec and Boulougouris (1982) suggests active engagement with the stimuli is crucial for extinction. Discussion followed the conclusion of the exposure in order to consolidate Arwyn’s belief the prediction will not come true on subsequent occasions (Salkovskis, 2007). Arwyn stated he learnt his anxiety reduced and he did not lose control. He re-rated the belief that he’d lose control if exposed to the same situation again at 20%. Arwyn’s SUDs also reduced from 95 to 40. A contract was collaboratively created listing the compulsions Arwyn would refrain from and the next natural time he would wash. Arwyn’s homework was to expose himself to the utility room every morning for half an hour without engaging in compulsions afterwards.

By the end of Phase B Arwyn had completed all exposures listed in his hierarchy.

### ***Outcome***

On his entry to the service, Arwyn’s OCI-R score was 57, 2.14 SDs above the average of an OCD sample (Foa et al., 2002). As shown in Figure 2, following CBT (including ERP) treatment for OCD (Phase A) Arwyn’s symptoms reduced to 28 indicating reliable improvement (Abramowitz et al., 2005) however was still above clinical cut-off. The addition of concurrent NET and CBT with ERP to the items at the top of Arwyn’s hierarchy (Phase B) led to a further reduction in OCI-R score to 9 (further reliable improvement) and well below the clinical cut-off and therefore indicating reliable recovery (Clark & Oates, 2014). Arwyn first completed the IES-R prior to beginning Phase B, scoring 24 specifically related to his experiences in his last job (this was chosen as this was the memory he reported as currently the most traumatic). This score represents “clinical concern” (Asukai et al., 2002) but is below the clinical cut-off of 33. By the end of Phase B Arwyn’s IES-R score reduced to 15, indicating reliable improvement (Clark & Oates, 2014).

From a qualitative perspective, Arwyn's functioning significantly improved by the end of therapy in that he was actively communicating with his workplace with a view to start work again, had holidayed with friends from work and had moved in with his girlfriend (something he had previously avoided due to fears of contamination). Arwyn had also managed to return to a healthy weight and was actively discussing his memories with others.

*Figure 2. Arwyn's OCI-R and IES Scores - Here*

### **Discussion**

We presented the use of CBT including ERP (Phase A) followed by concurrent NET and CBT with ERP (Phase B) to treat an individual with OCD whose PTSD symptoms were interfering with first line OCD treatment.

Despite Arwyn's 12 sessions of private CBT, his OCI-R remained within the clinical range and his compulsive behaviours were presenting a risk to his health. Phase A of CBT with ERP led to reliable improvement in Arwyn's OCD symptoms and in particular in relation to SSBs and compulsions related to items lower down his exposure hierarchy. Despite this, his reduction in symptoms plateaued and remained within the clinical range and frustration emerged in response to a lack of progress on exposures higher up the hierarchy. Exposure to items at the top of his hierarchy resulted in involuntary intrusive memories and significant levels of distress which prevented Arwyn from engaging in ERP. While Arwyn's scores on the IES-R did not meet the threshold indicating clinically significant PTSD symptoms, Arwyn presented with high levels of distress upon being asked about these memories, he was evidencing current avoidance symptoms in the form of an unwillingness to discuss the memories, and reported historical avoidance symptoms that were pre-morbid to his OCD. When he did begin to try to share his memories of these events with the therapist his recall was very fragmented. Together, this suggested processing of these memories could be important to further progress. When NET and CBT with ERP were applied concurrently,

Arwyn was able to complete exposure tasks to stimuli that shared a strong association with his trauma memories, something he was not able to do during Phase A. By the end of treatment Arwyn had achieved reliable recovery from OCD and reliable improvement in PTSD symptoms and he had returned to a level of functioning he was content with.

### *Clinical Implications*

Our approach involved adding NET to CBT with ERP which was already underway. This is as opposed to using a staggered approach of switching to NET alone first, before continuing with CBT with ERP. Therefore, it is difficult to separate out the individual contribution of NET and CBT with ERP to the reduction in symptoms evidenced above. However, we did show that following their concurrent application there was a reduction in Arwyn's OCD symptoms to non-clinical levels. If OCD thoughts and behaviours developed to help avoid the distress related to trauma related memories (Huppert et al., 2005; Gershuny et al., 2003), it is possible increased processing of the memories provided opportunity for Arwyn to consider their contribution to his life choices and this may have helped integrate it with his life story and lowered the level of arousal experienced when confronted with triggers to the memories. This may have contributed to his ability to participate in exposures to items at the top of his hierarchy whilst dropping previous compulsive behaviours. At the earlier stages of therapy, even following ERP on items lower down his hierarchy, Arwyn was not willing not do this. This appears consistent with research suggesting the experience of trauma makes the anxiety more severe and therefore more difficult to engage with ERP (Riggs, 2000).

As described earlier, the co-occurrence of PTSD in those with OCD has been thought to contribute to its treatment resistance (Gershuny et al., 2002). In the current case study we provide some evidence the combination of NET for trauma memories and CBT with ERP for OCD could be beneficial for the treatment of OCD with origins in traumatic events and first line treatment for OCD has not resulted in reliable improvement of symptoms.

### *Alternative approaches*

In the current study, the choice to use NET was partly pragmatic due to Arwyn experiencing multiple connected traumatic or aversive experiences over a number of years. In light of the aim of NET in the current study i.e. to reduce sense of current threat and arousal in response to trauma related triggers, it is possible (and likely, given the evidence base) that other trauma-focused therapies such as TF-CBT, EMDR, or Cognitive Processing Therapy could have resulted in a similar outcome to NET (particularly in cases where there is a single trauma). Indeed other case studies have described combining TF-CBT (Stobie, 2009) or EMDR (Nijdm et al., 2013) with CBT and ERP (albeit consecutively rather than concurrently) as an approach to treating individuals with comorbid OCD and PTSD.

It is also important to highlight that other studies have highlighted the role of “aversive” memories in the development and maintenance of OCD (e.g. Coughtrey et al., 2013) and have used imagery re-scripting as a treatment approach in such cases. Imagery re-scripting can be used to relive and restructure the meaning or course of events in a memory to something less catastrophic and this is thought to be particularly helpful when the predominant emotions associated with a memory relate to anger, shame or guilt as opposed to fear (Antz et al., 2012). In a case series, Veale et al., (2015) applied imagery re-scripting to a series of 12 cases with OCD with intrusive distressing images emotionally linked to a past aversive memory and found reliable improvement in 9 out of 12 cases and clinically significant change in 7 out of 12. Given the above, it is possible that using imagery re-scripting rather than NET in the current case could also have led to similar outcomes.

A further alternative treatment approach could have been to use a trauma-focused treatment alone, without concurrent CBT with ERP. If Arwyn’s OCD developed to avoid memories and associated distress, reducing the levels of arousal associated with exposure to certain triggers with NET alone could have been sufficient to absolve the need to engage in

compulsive behaviours. Therefore it is possible that Arwyn could have reduced compulsive behaviours spontaneously without the need for further CBT with ERP. Future studies could explore this hypothesis by delivering a phase of NET (or other trauma-focused therapy) first and measuring symptom change, before moving onto a phase of CBT with ERP.

### ***Limitations***

There are a number of limitations to this case study which makes it difficult to draw strong conclusions about the combined use of NET and CBT with ERP for OCD and PTSD symptom reduction.

First, throughout Arwyn's treatment there was a consistent downward trend in symptoms, despite a plateau around 20 weeks. Arwyn's OCI-R scores may have continued to reduce had Phase A continued, given enough time. Second whilst Arwyn had the support of his family members or support workers during exposures during Phase A, it is possible the change in Phase B to prolonged (3 hour) in vivo ERP which was facilitated by the clinician led to Arwyn's increased willingness to complete the ERPs and therefore the improvements observed. It is possible that the quality of ERP was higher when clinician-supported rather than self-directed (although an RCT has found no differences in outcome between self-directed and therapist-directed ERP; van Oppen et al., 2010), whilst the presence of the therapist may have improved trust in the approach and helped with emotional containment. In order to get a more accurate picture of the contribution of NET to Arwyn's willingness to complete ERPs during Phase B it may have been preferable to keep the support he had in completing the exposures the same as Phase A (i.e. self-directed with the support of family). Third, long-term follow-up measures were not collected due to this not being routine practice in the service, therefore it is not clear if the reliable recovery was maintained. It is possible the pattern of symptom reduction illustrated in Figure 2 was not a stable pattern. Nonetheless, it is important to emphasise the positive functional changes Arwyn had made to his life in this

time. Lastly, this study described the single case of an individual whose sub-threshold PTSD symptoms were thought to be impacting on OCD treatment and therefore the approach described may not be relevant to other cases of comorbid OCD and PTSD. Since the completion of this case study, Pinicotti et al., (In Press) have reviewed the comorbid presentation of OCD and PTSD and have made a number of recommendations for its conceptualisation, assessment and treatment. The interested reader is directed to this.

### **Ethical Statements**

All authors abided by the Ethical Principles of Psychologists and Code of Conduct as set out by the BABCP and BPS. No ethical approval was required for the study since only routine clinical data was used. The NHS trust in which Arwyn was under approved its publication conditional on Arwyn providing his written informed consent.

### **Key Practice Points**

- Clinicians treating OCD should consider how unprocessed trauma could impact on standard treatment for OCD
- Combining NET to process trauma memories interspersed with CBT including ERP to treat OCD symptomology may be helpful in reducing symptoms of both OCD and PTSD
- Further research in robust and controlled studies is required prior to conclusions about the effectiveness of this approach

### **Further Reading**

de Silva, P., & Marks, M. (1999). The role of traumatic experiences in the genesis of obsessive-compulsive disorder. *Behaviour Research and Therapy*, 37, 941–951.

Pinciotti, C. M., Fontenelle, L. F., Van Kirk, N., & Riemann, B. (In Press). Co-occurring

obsessive-compulsive and posttraumatic stress disorder: A review of conceptualisation, assessment, and cognitive behavioural treatment. *Journal of Cognitive Psychotherapy*.

Radomsky, A. S., Coughtrey, A Roz Shafran, R., & Rachman, S. (2017). Abnormal and Normal Mental Contamination, *Journal of Obsessive-Compulsive and Related Disorders*, <http://dx.doi.org/10.1016/j.jocrd.2017.08.011>



## References

- Abramowitz, J. S., Tolin, D. F., & Diefenbach, G. J. (2005).** Measuring change in OCD: Sensitivity of the Obsessive-Compulsive Inventory-Revised. *Journal of Psychopathology and Behavioral Assessment*, 27(4), 317-324. doi: 10.1007/s10862-005-2411-y
- American Psychiatric Association. (2013).** *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Anders, S., Frazier, P., & Frankfurt, S. (2011).** Variations in Criterion A and PTSD rates in a community sample of women. *Journal of Anxiety Disorders*, 25 (2), 175 - 184. doi:10.1016/j.janxdis.2010.08.018
- Arntz, A. (2012).** Imagery Restricting as a therapeutic technique: Review of clinical trials, basic studies, and research agenda. *Journal of Experimental Psychopathology*, 3(2), 189 - 208. DOI:10.5127/jep.024211
- Asukai, N. Kato, H. et al. (2002).** Reliability and validity of the Japanese-language version of the Impact of event scale-revised (IES-R-J). *Journal of Nervous and Mental Disease*. 190 (3): 175- 182.
- Bisson, J. I., Roberts, N. P., Andrew, M., Cooper, R., & Lewis, C. (2013).** Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *Cochrane Database of Systematic Reviews* (12). doi: 10.1002/14651858.CD003388.pub4
- Borkovec, T., & Boulougouris, J. (1982).** *Learning Theory Approaches to Psychiatry*.
- Brewin, C. R., Lanius, R. A., Novac, A., Schnyder, U., & Galea, S. (2009).** Reformulating for DSM-V: Life after criterion A. *Journal of Traumatic Stress*, 22, 366–373

- Challacombe, F., Oldfield, V. B., & Salkovskis, P. M.** (2011). *Overcoming obsessive compulsive disorder with CBT*. Random House.
- Clark, D. M.** (1986). A cognitive approach to panic. *Behaviour Research and Therapy*, 24(4), 461–470.
- Clark, D. M., & Oates, M.** (2014). Measuring Improvement and Recovery Adult Services. *Improving access to Psychological Therapies, Version 2*.
- Coughtrey, A. E., Shafran, R., Lee, M. and Rachman, S.** (2013) The treatment of mental contamination: a case series. *Cognitive and Behavioural Practice*, 20(2). 221-231. ISSN 1077-7229
- Creamer, M., Bell, R., & Failla, S.** (2003). Psychometric properties of the Impact of Event Scale - Revised. *Behaviour Research and Therapy*, 41(12), 1489–1496.  
<https://doi.org/10.1016/j.brat.2003.07.010>
- Cromer, K. R., Schmidt, N. B., & Murphy, D. L.** (2007). Do traumatic events influence the clinical expression of compulsive hoarding? *Behaviour Research and Therapy*, 45(11), 2581–2592. <https://doi.org/10.1016/j.brat.2007.06.005>
- De Moraes, E. C., Torresan, R. C., Trench, E. V., & Torres, A. R.** (2008). A possible case of posttraumatic obsessive-compulsive disorder. *Revista Brasileira de Psiquiatria*, 30(3), 290–301. <https://doi.org/10.1590/S1516-44462008000300018>
- de Silva, P., & Marks, M.** (1999). The role of traumatic experiences in the genesis of obsessive-compulsive disorder. *Behaviour Research and Therapy*, 37, 941–951.
- de Silva, P., & Rachman, S.** (1984). Does escape behaviour strengthen agoraphobic A preliminary study. *Behavior Research and Therapy*, 22(I), 87–91.
- Dinn, W. M., Harris, C. L., & Raynard, R. C.** (1999). Posttraumatic obsessive-compulsive

disorder: A three-factor model. *Psychiatry*, 62(4), 313–324.

<https://doi.org/10.1080/00332747.1999.11024877>

**Dykshoorn, K. L.** (2014). Trauma-related obsessive–compulsive disorder: a review. *Health Psychology and Behavioral Medicine*, 2(1), 517–528.

<https://doi.org/10.1080/21642850.2014.905207>

**Ehlers, A., & Clark, D.** (2000). A cognitive model of posttraumatic stress disorder. *Behaviour Research and Therapy*, 38, 319–345.

**Ehlers, A., & Clark, D. M.** (2008). Post-traumatic stress disorder: The development of effective psychological treatments. *Nordic Journal of Psychiatry*, 62(Suppl sup47), 11–18

**Eysenck, H. J., & Rachman, S.** (1965). *The causes and cases of neurosis: An introduction to modern behavior therapy based on learning theory and the principles of conditioning.* (Kegan Paul). London: Routledge.

**Foa, E. B., Huppert, J. D., Leiberg, S., Langner, R., Kichic, R., Hajcak, G., & Salkovskis, P. M.** (2002). The obsessive-compulsive inventory: Development and validation of a short version. *Psychological Assessment*, 14(4), 485–496.

<https://doi.org/10.1037//1040-3590.14.4.485>

**Fostick, L., Nacasch, N., & Zohar, J.** (2012). Acute obsessive compulsive disorder (OCD) in veterans with posttraumatic stress disorder (PTSD). *World Journal of Biological Psychiatry*, 13(4), 312–315. <https://doi.org/10.3109/15622975.2011.607848>

**Gershuny, B. S., Baer, L., Jenike, M. A., Minichiello, W. E., & Wilhelm, S.** (2002). Comorbid posttraumatic stress disorder: impact on treatment outcome for obsessive-compulsive disorder. *American Journal of Psychiatry*, 159, 852–854.

- Gershuny, B. S., Baer, L., Parker, H., Gentes, E. L., Infield, A. L., & Jenike, M. A.** (2008). Trauma and posttraumatic stress disorder in treatment-resistant obsessive-compulsive disorder. *Depression and Anxiety*, *25*, 69–71. <https://doi.org/10.1002/da>
- Gershuny, B. S., Baer, L., Radomsky, A. S., Wilson, K. A., & Jenike, M. A.** (2003). Connections among symptoms of obsessive-compulsive disorder and posttraumatic stress disorder: A case series. *Behaviour Research and Therapy*, *41*(9), 1029–1041. [https://doi.org/10.1016/S0005-7967\(02\)00178-X](https://doi.org/10.1016/S0005-7967(02)00178-X)
- Huppert, J. D., Moser, J. S., Gershuny, B. S., Riggs, D. S., Spokas, M., Filip, J., ... Foa, E. B.** (2005). The relationship between obsessive-compulsive and posttraumatic stress symptoms in clinical and non-clinical samples. *Journal of Anxiety Disorders*, *19*(1), 127–136. <https://doi.org/10.1016/j.janxdis.2004.01.001>
- Jordan, B. K., Schlenger, W. E., Hough, R., Fairbank, J. A., Marmar, C. R., Kulka, R. A., & Weiss, D.** (1991). Lifetime and current prevalence of specific psychiatric disorders in Vietnam veterans and controls. *Archives of General Psychiatry*, *48*, 207–215.
- Lely, J. C. G., Smid, G. E., Jongedijk, R. A., Knipscheer, J. W. & Kleber, R. J.** (2019). The effectiveness of narrative exposure therapy: a review, meta-analysis and meta-regression analysis. *European Journal of Psychotraumatology*, *10* (1), 1 - 13
- Lochner, C., Du Toit, P. L., Zungu-Dirwayi, N., Marais, A., Van Kradenburg, J., Seedat, S., ... Stein, D. J.** (2002). Childhood trauma in obsessive-compulsive disorder, trichotillomania, and controls. *Depression and Anxiety*, *15*(2), 66–68. <https://doi.org/10.1002/da.10028>
- Morina, N., Sulaj, V., Schnyder, U., Klaghofer, R., Müller, J., Martin-Sölch, C., & Rufer, M.** (2016). Obsessive-compulsive and posttraumatic stress symptoms among

civilian survivors of war. *BMC Psychiatry*, 16(115), 1–8.

<https://doi.org/10.1186/s12888-016-0822-9>

**Meyer V. (1966).** Modification of expectations in cases with obsessional rituals. *Behaviour research and therapy*, 4(4), 273–280. [https://doi.org/10.1016/0005-7967\(66\)90023-4](https://doi.org/10.1016/0005-7967(66)90023-4)

**Nacasch, N., Fostick, L., & Zohar, J. (2011).** High prevalence of obsessive-compulsive disorder among posttraumatic stress disorder patients. *European Neuropsychopharmacology*, 21(12), 876–879.  
<https://doi.org/10.1016/j.euroneuro.2011.03.007>

**National Institute of Health and Care Excellence. (2005).** *Obsessive-compulsive disorder and body dysmorphic disorder: treatment. Clinical guideline [CG31]*.

**Nijdam, M. J., van der Pol, M. M., Dekens, R. E, Olf, M., & Denys, D. (2013).** Treatment of sexual trauma dissolves contamination fear: a case report. *European Journal of Psychotraumatology*, 4. <http://dx.doi.org/10.3402/ejpt.v4i0.19157>

**Pitman, R. K. (1993).** Posttraumatic Obsessive-Compulsive A Case Study. *Comprehensive Psychiatry*, 34(2), 102–107.

**Rachman, S. (1991).** A psychological approach to the study of residential satisfaction. *Clinical Psychology Review*, 11, 461–464.

**Radomsky, A. S., Coughtry, A., Shafran, R., & Rachman, S. (2017).** Abnormal and Normal Mental Contamination. *Journal of Obsessive-Compulsive and Related Disorders*. 17, 46 - 51. <http://dx.doi.org/10.1016/j.jocrd.2017.08.011>

**Riggs, D. S. (2000).** Treatment of concurrent PTSD and OCD: A commentary on the case of Howard. *Cognitive and Behavioral Practice*, 7(1), 130–132.

[https://doi.org/10.1016/S1077-7229\(00\)80015-2](https://doi.org/10.1016/S1077-7229(00)80015-2)

**Robjant, K., & Fazel, M.** (2010). The emerging evidence for Narrative Exposure Therapy:

A review. *Clinical Psychology Review*, *30*(8), 1030–1039.

<https://doi.org/10.1016/j.cpr.2010.07.004>

**Pinciottim C. M., Fontenelle, L.F., Van Kirk, N., & Riemann, B. C.** (In press). Co-

occurring obsessive-compulsive and posttraumatic stress disorder: A review of conceptualization, assessment and cognitive-behavioural treatment. *Journal of Cognitive Psychotherapy*.

**Salkovskis, P. M.** (1985). Obsessional-Compulsive Problems: a Cognitive-Behavioural

Analysis. *Behaviour Research and Therapy*, *23*(5), 571–583.

[https://doi.org/10.1016/0005-7967\(85\)90105-6](https://doi.org/10.1016/0005-7967(85)90105-6)

**Salkovskis, P. M.** (1991). The Importance of Behaviour in the Maintenance of Anxiety and

Panic: A Cognitive Account. *Behavioural Psychotherapy*, *19*(1), 6–19.

<https://doi.org/10.1017/S0141347300011472>

**Salkovskis, P. M.** (1998). The devil is in the detail: conceptualising and treating obsessional

problems. In N. Tarrier, A. Wells, & G. Haddock (Eds.), *Treating Complex Cases: The Cognitive Behaviour Therapy Approach*. Chichester: Wiley.

**Salkovskis, P.M.** (2007). Psychological treatment of obsessive compulsive disorder.

*Psychiatry*, Vol. 6 No. 6, pp. 229–233.

**Sasson, Y., Dekel, S., Nacasch, N., Chopra, M., Zinger, Y., Amital, D., & Zohar, J.**

(2005). Posttraumatic obsessive-compulsive disorder: A case series. *Psychiatry*

*Research*, *135*(2), 145–152. <https://doi.org/10.1016/j.psychres.2004.05.026>

**Stobie, B.** (2009). Cognitive therapy for post-traumatic stress disorder and obsessive-

compulsive disorder. In Grey, N. (Ed). *A Casebook of Cognitive Therapy for Traumatic*

*Stress Reactions* (pp. 14-16). Routledge. <https://doi.org/10.4324/9780203874318>

**van Oppen, P., van Balkom, A., Smit, J.H., Schuurmans, J., van Dyck, R., & Emmelkamp, P.M.** (2010). Does the therapy manual or the therapist matter most in treatment of obsessive-compulsive disorder? A randomized controlled trial of exposure with response or ritual prevention in 118 patients. *The Journal of clinical psychiatry*, 71(9), 1158-67 .

**Veale, D., Page, N., Woodward, E., & Salkovskis, P.** (2015). Imagery Rescripting for Obsessive Compulsive Disorder: A single case experimental design in 12 cases. *Journal of Behavior Therapy and Experimental Psychiatry*, 49(Special Issue), 230-236. <https://doi.org/10.1016/j.jbtep.2015.03.003>

**Watkins, L. E., Sprang, K. R., & Rothbaum, B. O.** (2018). Treating PTSD: A Review of Evidence-Based Psychotherapy Interventions. *Frontiers in behavioral neuroscience*, 12, 258. <https://doi.org/10.3389/fnbeh.2018.00258>

**Weiss, D. S.** (2007). The Impact of Event Scale-Revised. In J.P. Wilson & T.M. Keane (Eds.) *Assessing psychological trauma and PTSD: a practitioner's handbook* (2<sup>nd</sup> ed., pp. 168 – 189). New York: Guilford Press.





Figure 2. OCI-R and IES over time. Orange transparent area represents scores above the clinical cut-off for OCI-R scores.



*Table 1. Arwyn's exposure hierarchy*

1. Visiting old apprenticeship sites
2. Visiting old workplace
3. Touching travelling souvenirs
4. Meal at Restaurant related to experience travelling
5. Touching passport and mobile phone which were taken travelling
6. Going in Utility room and touching work and travelling related items
7. Eating a particular food prepared at restaurant