

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

The Taxonomic Report of the International
Lepidoptera Survey

Lepidoptera Survey

3-22-2021

Pupal Polymorphism of the Bog Copper Butterfly *Lycaena* (*Epidemia*) *epixanthe* (Bsd. & Le C. [1835]) (Lycaenidae: Lycaeninae) in New Jersey

David M. Wright
Lansdale, Pennsylvania

Follow this and additional works at: <https://digitalcommons.unl.edu/taxrpt>




Part of the [Entomology Commons](#), [Population Biology Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Wright, David M., "Pupal Polymorphism of the Bog Copper Butterfly *Lycaena* (*Epidemia*) *epixanthe* (Bsd. & Le C. [1835]) (Lycaenidae: Lycaeninae) in New Jersey" (2021). *The Taxonomic Report of the International Lepidoptera Survey*. 19.

<https://digitalcommons.unl.edu/taxrpt/19>

This Article is brought to you for free and open access by the Lepidoptera Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in The Taxonomic Report of the International Lepidoptera Survey by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



The Taxonomic Report

OF THE INTERNATIONAL LEPIDOPTERA SURVEY



ISSN 2643-4776 (print) / ISSN 2643-4806 (online)

Pupal polymorphism of the Bog Copper butterfly *Lycaena (Epidemia) epixanthe* (Bsd. & Le C. [1835]) (Lycaenidae: Lycaeninae) in New Jersey.

David M. Wright
124 Heartwood Drive
Lansdale, PA 19446
wripenn@aol.com

ABSTRACT. The immature stages of the Bog Copper butterfly, *Lycaena epixanthe*, were described in detail by the author nearly forty years ago. Two different pupal colors were interpreted to represent distinct separate morphs with genetically-fixed incidences. Reassessment of the data discloses a pattern of progressive color change of each individual pupa throughout the pupal period. This transforming strategy is adaptive and may reduce predator recognition.

Key words: Bog Copper, cranberry bogs, pupal polymorphism

INTRODUCTION

The lycaenid Bog Copper butterfly, *Lycaena (Epidemia) epixanthe* (Boisduval & Le Conte [1835]) is restricted to acid bogs in northeastern North America, extending from the Canadian Maritime Provinces westward to Minnesota and Manitoba, and south to the Appalachian Mountains in West Virginia and the Coastal Plain of New Jersey. The butterfly in all stages is closely associated with its cranberry larval hosts (*Vaccinium macrocarpon* Ait. and *V. oxycoccos* L.). The morphology of the immature stages was previously described by the author (Wright, 1983). In that study the author stated the great majority of pupae were green with a few being blackish purple. These morphs were interpreted to represent two distinct varieties with fixed incidences. By comparison, Newcomb (1911) characterized the pupal colors of the Dorcas Copper, *Lycaena (Epidemia) dorcas* (W. Kirby, 1837), as “hardly any two are alike”. This notion prompted a reassessment of the pupal polymorphism of *L. epixanthe*.

MATERIALS & METHOD

The colony of *Lycaena epixanthe* in Forge Pond bog (Fig. 1) has been under investigation since 1976. A butterfly rearing study initiated in June, 1981, yielded 56 live larvae from overwintering ova. First instar emergences *ex ovum* commenced on April 16th in 1982. After four molts, pupation began on May 27th. The pupal period averaged 13 days and ranged from 11 to 15 days. Adult eclosions occurred from June 7th to 26th. The color patterns of twenty pupae are documented in the author’s notebook and photo files. Images were scanned into digital format for inclusion here.



Fig. 1. Forge Pond cranberry bog, Atlantic County, New Jersey. Large Cranberry in bloom in the foreground.
Fig. 2. Pupal color patterns of Bog Copper butterfly. **2a.** Green speckled. **2b.** Transitional. **2c.** Blackish purple.

RESULTS

Lycaena epixanthe pupae as a group display a substantial diversity of color. Yet, significantly, the colors of each individual pupa change as the pupal period progresses. Pupae are pale green at the start and finish as solid black before eclosion. Some usual progressive changes include black speckling after two days (Fig. 2a), partitioning of wing cases (cream or tan) and thorax (gray or black) after 5-6 days (Fig. 2b), and generalized darkening (black or blackish purple) after 11-13 days (Fig. 2c). Variation in the intensity of black maculations and streaking of veins on the wing cases are commonly observed. Occasionally, the blackish purple phase may appear earlier in the pupal period (e.g., day 3-6). Rarely, a pupa is devoid of all black speckling and streaking. The observed pupal patterns are cryptic within their natural habitat. When hidden among the cranberry leaves and sphagnum moss, these pupae are virtually inconspicuous to the human eye.

DISCUSSION

Pupal color plasticity generally occurs in species whose larvae have preferences for pupation sites which vary in color (West & Hazel, 1996). The adaptive significance of this plasticity is crypsis (Hazel *et al.*, 1998). The sharing of traits among closely-related species may shed light on when they evolved within the Lycaeninae (Coppers). To date, pupal plasticity is found chiefly in species of the subgenus *Epidemia*, which customarily overwinter in the egg stage with larvae emerging in spring. Pupal periods are short. Pupal polymorphism is reported in *L. helloides* by Coolidge (1911) and Scott (2008); in *L. dorcas* by Newcomb (1911) and Scott (1986, 2008); in *L. florus* by Scott (2014); in *L. nivalis* by Newcomer (1911, 1963); and in *L. epixanthe* by Wright (1983). No data is available for *L. mariposa*. The most pronounced assortment of color patterns occurs in wetland species like *L. dorcas* and *L. epixanthe*, which are confined to alkaline fens and acid bogs. In these species, pupation and adult emergence take place concurrently with the bloom period of their larval host plants (*Potentilla fruticosa* and *Vaccinium macrocarpon*). This crucial period attracts numerous insects into the open habitat seeking nectar and prey. A broad variety of pupal colors would reduce the likelihood that a predator would build a successful search pattern. The bogs of the New Jersey Pine Barrens came into existence at the end of the Pleistocene around 10,000 to 11,000 years ago (Buell, 1970). Genomic studies reveal the *Epidemia* clade experienced a burst of radiation around 5 Mya during the Pliocene (Zhang *et al.*, 2019). Rapid diversification regularly correlates with the sharing of beneficial genes among radiating species. Progressive pupal polymorphism is likely is one of these shared qualities.

ACKNOWLEDGMENTS

The author wishes to express his deep gratitude to the late Annie Carter (Naturalist, Batsto Historical Site, Wharton State Forest, Atlantic County, NJ), whose dedication to Pine Barrens preservation provided the impetus for this study. I thank Philip Marucci and Nick Vorsa (Rutgers University Blueberry-Cranberry Research Center, Jenkins, NJ) for insightful discussions on cranberry and peat bog biology.

LITERATURE CITED

- Buell, M.F. 1970. Time of origin of the New Jersey Pine Barrens bogs. *Bull. Torrey Bot. Club* 97:105-108.
- Coolidge, K.R. 1924. Life history of *Heodes helloides* Bdv. (Lepid.: Lycaenidae). *Ent. News* 35:306-312.
- Hazel, W.N., Ante, S. & B. Stringfellow. 1998. The evolution of environmentally-cued pupal colour in swallowtail butterflies: natural selection for pupation site and pupal colour. *Ecol. Ent.* 23:41-44.
- Newcomer, E.J. 1911. The life histories of two lycaenid butterflies. *Canad. Ent.* 43:83-88.
- _____. 1963. The synonymy, variability and biology of *Lycaena nivalis*. *J. Res. Lepid.* 2(4):271-280.
- Newcomb, W.W. 1911. The life-history of *Chrysophanus dorcas* Kirby. *Canad. Ent.* 43(5):160-168.
- Scott, J.A. 1986. *The Butterflies of North America*. CA: Stanford, Stanford University Press. 583 pp.
- _____. 2008. Hostplants & early stages of *Lycaena florus*. *Papilio (New Series)* #18, p. 41-43.
- _____. 2014. *Lycaena florus* (Lycaenidae): A blackish pupa. *Papilio (New Series)* #22, p. 62-63.
- West, D.A. & W.N. Hazel. 1996. Natural pupation sites of three North American swallowtail butterflies: *Eurytides marcellus* (Cramer), *Papilio cressphontes* Cramer, and *P. troilus* L. (Papilionidae). *J. Lepid. Soc.* 50:297-302.
- Wright, D.M. 1983. Life history and morphology of the immature stages of the Bog Copper butterfly *Lycaena epixanthe* (Bsd. & Le C.) (Lepidoptera: Lycaenidae). *J. Res. Lepid.* 22(1):47-100.
- Zhang, J., Cong, Q., Shen, J., Opler, P.A. & N.V. Grishin. 2019. Genomics of a complete butterfly continent. bioRxiv BIORXIV/2019/829887 Available online at: <https://doi.org/10.1101/829887>

The Taxonomic Report
is a publication of
The International Lepidoptera Survey (TILS)

The International Lepidoptera Survey is registered as a non-profit Limited Liability Company (LLC) in the state of Virginia, U.S.A. The Taxonomic Report (TTR) is published for the purpose of providing a public and permanent scientific record. Contents are peer-reviewed but not necessarily through the anonymous review and comment process preferred by some publishers of serial literature. It appears in digital, open-access form, is regularly disseminated in hardcopy form to select institutional repositories and is also available as printed copy upon request at the discretion of authors and/or the editor. Printing and postage costs may apply. An initial run of 25 copies is printed on paper to meet ICZN recommendation 8B. Copies of all TTR papers are available at the archival TTR website: (<http://lepsurvey.carolinanature.com/report.html>) and via the following digital repositories:

Internet Archive (<https://archive.org/>)
Biodiversity Heritage Library (<https://www.biodiversitylibrary.org>)
Zobodat (<https://www.zobodat.at/>)

TILS Purpose

TILS is devoted to the worldwide collection of Lepidoptera for the purpose of scientific discovery, determination, and documentation, without which there can be no preservation.

TILS Motto

“As a world community, we cannot protect that which we do not know”

Articles for publication are sought

They may deal with any area of research on Lepidoptera, including faunal surveys, conservation topics, methods, etc. Taxonomic papers are especially welcome. There are no page charges for authors. Before sending a manuscript, simply write to **TTR editor, Harry Pavulaan, 606 Hunton Place NE, Leesburg, VA, 20176, USA** to initiate discussion on how to best handle your material for publication, and to discuss peer review options; or email to intlepsurvey@gmail.com (cc: to harrypav@hotmail.com if you do not receive a reply within one week).

Visit *The International Lepidoptera Survey* on the World Wide Web at:

<http://lepsurvey.carolinanature.com>

&

Join the discussion at our list serve on Groups.io at:

<https://groups.io/g/TILS>

You can subscribe by sending an email to: TILS+subscribe@groups.io

&

Join The International Lepidoptera Survey on Facebook at:

<https://www.facebook.com/groups/1072292259768446>