

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

H. W. Manter Laboratory Library Materials

---

8-2020

## Taxonomy of *Echinostoma revolutum* and 37-Collar-Spined *Echinostoma* spp.: A Historical Review

Jong-Yil Chai

*Seoul National University College of Medicine*

Jaeun Cho

*Institute of Parasitic Diseases, Korean Association of Health Promotion*

Taehee Chang

*Institute of Parasitic Diseases, Korean Association of Health Promotion*

Bong-Kwang Jung

*Institute of Parasitic Diseases, Korean Association of Health Promotion*

Woon-Mok Sohn

*Gyeongsang National University College of Medicine*

Follow this and additional works at: <https://digitalcommons.unl.edu/manterlibrary>



Part of the [Parasitology Commons](#)

---

Chai, Jong-Yil; Cho, Jaeun; Chang, Taehee; Jung, Bong-Kwang; and Sohn, Woon-Mok, "Taxonomy of *Echinostoma revolutum* and 37-Collar-Spined *Echinostoma* spp.: A Historical Review" (2020). *H. W. Manter Laboratory Library Materials*. 86.

<https://digitalcommons.unl.edu/manterlibrary/86>

This Article is brought to you for free and open access by DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in H. W. Manter Laboratory Library Materials by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



# Taxonomy of *Echinostoma revolutum* and 37-Collar-Spined *Echinostoma* spp.: A Historical Review

Jong-Yil Chai<sup>1,2,\*</sup>  Jaeun Cho<sup>1</sup>, Taehee Chang<sup>1</sup>, Bong-Kwang Jung<sup>1</sup>, Woon-Mok Sohn<sup>3</sup>

<sup>1</sup>Institute of Parasitic Diseases, Korea Association of Health Promotion, Seoul 07649, Korea; <sup>2</sup>Department of Tropical Medicine and Parasitology, Seoul National University College of Medicine, Seoul 03080, Korea; <sup>3</sup>Department of Parasitology and Tropical Medicine, and Institute of Health Sciences, Gyeongsang National University College of Medicine, Jinju 52727, Korea

**Abstract:** *Echinostoma* flukes armed with 37 collar spines on their head collar are called as 37-collar-spined *Echinostoma* spp. (group) or ‘*Echinostoma revolutum* group’. At least 56 nominal species have been described in this group. However, many of them were morphologically close to and difficult to distinguish from the other, thus synonymized with the others. However, some of the synonymies were disagreed by other researchers, and taxonomic debates have been continued. Fortunately, recent development of molecular techniques, in particular, sequencing of the mitochondrial (*nad1* and *cox1*) and nuclear genes (ITS region; ITS1-5.8S-ITS2), has enabled us to obtain highly useful data on phylogenetic relationships of these 37-collar-spined *Echinostoma* spp. Thus, 16 different species are currently acknowledged to be valid worldwide, which include *E. revolutum*, *E. bolschewense*, *E. caproni*, *E. cinetorchis*, *E. deserticum*, *E. lindoense*, *E. luisreyi*, *E. mekongi*, *E. miyagawai*, *E. nasincovae*, *E. novaezealandense*, *E. paraensei*, *E. paraulum*, *E. robustum*, *E. trivolvis*, and *Echinostoma* sp. IG of Georgieva et al., 2013. The validity of the other 10 species is retained until further evaluation, including molecular analyses; *E. acuticauda*, *E. barbosai*, *E. chloephagae*, *E. echinatum*, *E. jurini*, *E. nudicaudatum*, *E. parvocirrus*, *E. pinnicaudatum*, *E. ralli*, and *E. rodriguesi*. In this review, the history of discovery and taxonomic debates on these 26 valid or validity-retained species are briefly reviewed.

**Key words:** *Echinostoma*, *Echinostoma revolutum*, ‘*revolutum*’ group, 37-collar-spined echinostome, historical review

## INTRODUCTION

Echinostomes, including families Echinostomatidae, Himasthlidae, and Echinochasmidae, are a large group of trematodes parasitizing the small intestines of fish, reptiles, birds, and mammals [1]. Among the Echinostomatidae, *Echinostoma* is the most important genus in public health as well as veterinary medical aspects. The type species of the genus *Echinostoma* is *E. revolutum* (Froelich, 1802) Dietz, 1909, and within this genus numerous species have been described from birds and mammals [2,3]. At least 120 species were listed by Yamaguti [1] until the 1960s which included 101 species infecting birds and 22 species infecting mammals; among them 3 species were reported from both mammals and birds [1]. Later, a lot of new species have been described from various parts of the

world. However, host specificity of many species was further studied and redefined, and many of them were synonymized with the others [4-6].

*E. revolutum* and other 37-collar-spined *Echinostoma* spp. are a large group of echinostomes representing the Echinostomatidae in various aspects. They are also called as 37-collar-spined *Echinostoma* spp. (group), ‘*E. revolutum* group’ [4,7], or simply ‘*revolutum*’ group [8-11]. Kanev [4] proposed that “worms with 37 collar spines belonging to the genus *Echinostoma* and occurring in naturally infected birds in Europe and Asia be referred to as ‘*E. revolutum* group’. However, in the present review, for comprehensiveness and convenience, all 37-collar-spined *Echinostoma* spp. ever reported from naturally infected birds and/or mammals around the world have been assigned as ‘37-collar-spined *Echinostoma* spp.’ or ‘*E. revolutum* group’.

More than 56 nominal species have been described in this group (Tables 1, 2). However, there have long been debates on the taxonomy and classification of these species. More than a half of them have been synonymized with the others (Table 2). At present, 16 species (14 of them have molecular data) (Figs. 1, 2) are regarded as valid species, and 10 should be further

•Received 6 July 2020, revised 29 July 2020, accepted 29 July 2020.

\*Corresponding author (cyj@snu.ac.kr)

© 2020, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1.** List of 16 valid and 10 validity-retained species of 37-collar-spined *Echinostoma* group

Species and nominator	Size of adults (mm)	Size of eggs ( $\mu$ m)	Country of first report	Reference no.	Validity
<i>E. revolutum</i> (Froelich, 1802) Dietz, 1909	9.5-11.8×1.5-2.1	108-125×57-75	Germany	[11]	valid (type species)
<i>E. acuticauda</i> Nicoll, 1914	9.0-12.0×0.9-1.0	112-126×63-75	Australia	[93]	retained
<i>E. barbosai</i> Lie & Basch, 1966	3.6-7.8×0.46-0.96	111-131×62-66	Brazil	[117]	retained
<i>E. bolschewense</i> (Kotova, 1939) Nasincova, 1991	7.4-12.5×0.99-1.6	138-162×75-85	Russia	[122]	valid
<i>E. caproni</i> Richard, 1964	5.7×1.5 (av.)	105-120×50-60	Madagascar	[80]	valid
<i>E. chloephagae</i> Sutton & Lunaschi, 1980	5.1-6.9×0.82-0.99	100-120×60-70	Argentina	[131]	retained
<i>E. cinetorchis</i> Ando & Ozaki, 1923	9.5-14.6×1.7-2.2	96-100×61-70	Japan	[59]	valid
<i>E. deserticum</i> Kechemir et al., 2002	5.6-15.2×0.75-3.1	58-74×36-46	Niger	[43]	valid
<i>E. echinatum</i> (Zeder, 1803) de Blairville, 1828	8.1-11.1×0.77-1.5	?	Germany	[99]	retained
<i>E. jurini</i> (Skvortsov, 1924) Kanev, 1985	6.6-14.0×0.58-1.3	96-132×72-88	Russia	[5]	retained
<i>E. lindoense</i> Sandground & Bonne, 1940	13.0-15.0×2.5-3.0	92-124×65-76	Indonesia	[63]	valid
<i>E. luisreyi</i> Maldonado et al., 2003	5.3-9.3×1.1-2.3	89-113×65-82	Brazil	[44]	valid
<i>E. mekongi</i> Cho et al., 2020	9.0-13.1×1.3-2.5	98-132×62-90	Cambodia	[49]	valid
<i>E. miyagawai</i> Ishii, 1932	9.2-11.0×1.2-1.5	94-96×59-60	Japan	[11]	valid
<i>E. nasincovae</i> Georgieva et al., 2014	4.3-4.8×0.99-1.1	100-105×61-70	Czech Republic	[11]	valid
<i>E. novaezealandense</i> Georgieva et al., 2017	9.6-10.5×0.7-1.1	81-87×42-53	New Zealand	[24]	valid
<i>E. nudicaudatum</i> Nasir, 1960	6.8-7.6×0.98-1.3	97-115×67-72	UK	[171]	retained
<i>E. paraensei</i> Lie & Basch, 1967	7.5-16.0×0.79-2.0	104-122×74-86	Brazil	[176]	valid
<i>E. paraulum</i> Dietz, 1909	5.6-6.9×1.2-1.7	104-122×53-70	Austria/Russia	[11]	valid
<i>E. parvocirrus</i> Nassi & Dupouy, 1988	6.7-8.4×1.3-1.5	105-120×64-71	Guadeloupe	[38]	retained
<i>E. pinnicaudatum</i> Nasir, 1961	5.5-7.1×0.8-1.1	97-115×67-72	UK	[174]	retained
<i>E. ralli</i> Yamaguti, 1934	8.6×1.1 (av.)	110-130×68-81	Japan	[68]	retained
<i>E. robustum</i> Yamaguti, 1935	7.8-9.8×1.3-2.2	111-129×60-69	Taiwan	[187]	valid
<i>E. rodriguesi</i> Hsu et al., 1968	3.9-6.8×0.5-1.3	96-128×56-68	Brazil	[37]	retained
<i>E. trivolvis</i> (Cort, 1914) Kanev, 1985	5.5-21.0×0.5-1.5	90-130×60-70	USA	[6]	valid
<i>Echinostoma</i> sp. IG Georgieva et al., 2013	-	-	Germany	[45]	valid <sup>a</sup>

<sup>a</sup>validity acknowledged based on molecular and morphological data of cercariae, but description of adult worms needed.

evaluated for their validity (Table 1). Among them, zoonotic species infecting humans are at least 8, including *E. revolutum*, *E. cinetorchis*, *E. echinatum* (needs confirmation), *E. lindoense*, *E. mekongi*, *E. miyagawai* (experimental), *E. paraensei* (from the coprolite of a human mummy), and *E. paraulum* [2,3,5,12-15]. In this review, the historical aspects and current status of 26 valid or validity-retained species of 37-collar-spined *Echinostoma* group are briefly reviewed.

## BRIEF HISTORY

After revision and set-up new systematics of echinostomes in the early 1900s by Dietz [16,17], hundreds of articles have dealt with taxonomy and biology of echinostomes [18]. Among the workers, Beaver [19] was the one who extensively studied and reviewed the taxonomy of 37-collar-spined echinostomes. He obtained adult flukes from experimentally infected birds and mammals originating from the freshwater snail *Helisoma trivolvis* and described the worm as *E. revolutum*

in USA (later turned out to be *Echinostoma trivolvis* by Kanev et al. [6]) and morphologically compared this species with previously reported species from the world. He synonymized 9 species with *E. revolutum*, including *Echinostoma armigerum* Barker & Irvine, 1915, *Echinostoma cinetorchis* Ando & Ozaki, 1923, *Echinostoma coalitum* Barker & Beaver, 1915, *Echinostoma columbae* Zunker, 1925, *Echinostoma echinatum* (Zeder, 1803) de Blairville, 1828, *Echinostoma limicoli* Johnson, 1920, *Echinostoma miyagawai* Ishii, 1932, *Echinostoma mendax* Dietz, 1909, and *Echinostoma paraulum* Dietz, 1909. In addition, he treated 11 species as synonyms inquirenda, which included *Echinostoma acuticauda* Nicoll, 1914, *Echinostomas armatum* (Molin, 1850) Yamaguti, 1971, *Echinostoma callawayensis* Barker & Noll, 1915, *Echinostoma dilatatum* (Miram, 1840) Cobbold, 1960, *Echinostoma echinocephalum* (Rudolphi, 1819) Cobbold, 1860, *Echinostoma erraticum* Lutz, 1924, *Echinostoma microrchis* Lutz, 1924, *Echinostoma neglectum* Lutz, 1924, *Echinostoma nephrocystis* Lutz, 1924, *Echinostoma oxycephalum* (Rudolphi, 1819) Raillet, 1896, and *Echinostoma sudanense* Odhner, 1911. Later,

**Table 2.** List of 37-collar-spined *Echinostoma* spp. (30 species) synonymized with other species


Species and nominator	Country of first report	Synonymized with	Synonymy proposed by
<i>E. armatum</i> (Mollin, 1858) Yamaguti, 1971	South America	<i>E. revolutum</i>	Beaver [19]
<i>E. armigerum</i> Barker and Irvine, 1915	North America	<i>E. trivolvis</i>	Kanev [4,20]
<i>E. audyi</i> Lie and Umathevy, 1965	Malaysia	<i>E. revolutum</i>	Kanev [20]
<i>E. callawayensis</i> Barker and Noll, 1915	North America	<i>E. trivolvis</i>	Kanev [4,20]
<i>E. coalitum</i> Barker and Beaver, 1915	North America	<i>E. trivolvis</i>	Kanev [4,20]
<i>E. columbae</i> Zunker, 1925	Germany	<i>E. revolutum</i>	Beaver [19]
<i>Echinoparyphium contiguum</i> Barker and Bastron, 1915	USA	<i>E. trivolvis</i>	Kanev et al. [6]
<i>E. dilatatum</i> (Miram, 1840) Cobbold, 1860	Russia	<i>E. revolutum</i>	Beaver [19]
<i>E. equinatus gigas</i> Marco del Pont, 1926	Argentina	<i>Echinoparyphium recurvatum</i>	Lunaschi et al. [29]
<i>E. echinocephalum</i> (Rudolphi, 1819) Cobbold, 1860	Egypt	<i>E. revolutum</i>	Beaver [19]
<i>E. erraticum</i> Lutz, 1924	Brazil	<i>E. revolutum</i>	Beaver [19]
<i>E. friedi</i> Toledo et al., 2000	Spain	<i>E. miyagawai</i>	Faltýnková et al. [11]
<i>E. ivaniosi</i> Mohandas, 1973	India	<i>E. revolutum</i>	Kanev [4]
<i>E. liei</i> Jeyarasasingam et al., 1972	Egypt	<i>E. caproni</i>	Huffman and Fried [7]
<i>E. limicoli</i> Johnson, 1920	Europe	<i>E. revolutum</i>	Beaver [19]
<i>E. londonensis</i> Khan, 1961	UK	<i>E. jurini</i>	Kanev [4]
<i>E. mendax</i> Dietz, 1909	Brazil	<i>E. revolutum</i>	Beaver [19]
<i>E. microrchis</i> Lutz, 1924	Brazil	<i>E. revolutum</i>	Beaver [19]
<i>E. multispinosum</i> Pérez Vigueras, 1944	Cuba	<i>E. trivolvis</i>	Kanev et al. [6]
<i>E. neglectum</i> Lutz, 1924	Brazil	<i>E. revolutum</i>	Beaver [19]
<i>E. nephrocystis</i> Lutz, 1924	Brazil	<i>E. revolutum</i>	Beaver [19]
<i>E. orlovi</i> Romashov, 1966	Russia	<i>E. jurini</i>	Kanev et al. [5]
<i>E. oxycephalum</i> (Rudolphi, 1819) Railliet, 1896	Europe	<i>E. revolutum</i>	Beaver [19]
<i>E. revolutum tenuicollis</i> Bashkirova, 1941	Azerbaijan	<i>E. revolutum</i>	This review
<i>E. revolutum</i> var. <i>japonicum</i> Kurisu, 1932	Japan	<i>E. revolutum</i>	Yamaguti [1]
<i>E. sisjakowi</i> (Skvortsov, 1935) Yamaguti, 1971 <sup>a</sup>	Russia	<i>E. jurini</i>	Kanev et al. [5]
<i>E. spiniferum</i> (La Valette, 1855) sensu Nasincova, 1992	Czechoslovakia	<i>E. nasincovae</i>	Faltýnková et al. [11]
<i>E. stromi</i> Bashkirova, 1946	Azerbaijan	<i>E. revolutum</i>	Yamaguti [1]
<i>E. sudanense</i> Odhner, 1910	Sudan	<i>E. revolutum</i>	Beaver [19]
<i>E. togoensis</i> Jourdan and Kulo, 1981	Togo	<i>E. caproni</i>	Huffman and Fried [7]

<sup>a</sup>formerly *Echinoparyphium sisjakowi* Skvortsov, 1935.


however, some of these species, including *E. revolutum* of Beaver [19], *E. armigerum*, *E. coalitum*, and *E. callawayensis*, were synonymized with *Echinostoma trivolvis* (Cort, 1914) Kanev, 1985 [6,20]. Kanev et al. [6] also synonymized *Echinostoma multispinosum* Pérez Vigueras, 1944, *E. paraulum* of Miller, 1937, and *Echinoparyphium contiguum* Barker & Bastron, 1915 with *E. trivolvis*. However, the validity of *E. miyagawai*, *E. cinetorchis*, and *E. paraulum* were re-evaluated and acknowledged by Kostadinova et al. [21], Chai [3,22], and Georgieva et al. [10], respectively. The morphologies of *E. miyagawai* and *E. paraulum* were redescribed [11,21]. The name *E. acuticauda* is still used by other researchers [23,24], although this species needs re-evaluation. The 4 species of 37-collar-spined *Echinostoma* from Brazil, including *E. erraticum*, *E. microrchis*, *E. neglectum*, and *E. nephrocystis*, and 2 non-37-collar-spined species, *Echinostoma exile* (43-45 collar spines) and *Echinostoma parses-*

*pinosum* (29-33 collar spines), were re-examined and redescribed by Kohn and Fernandes [25]. Kostadinova and Gibson [26] stated that the 4 species of 37-collar-spined species redescribed by them [25] exhibited significant morphological differences suggesting their taxonomic significance. In addition, a checklist of cercariae in molluscs from Brazil listed the names of *E. erraticum* (infecting *Spirulina* and *Drepanotrema* snails) and *E. nephrocystis* (infecting *Physa* sp. snails) [27]. However, in the present review, these 4 species were tentatively regarded as synonyms of *E. revolutum*, as suggested by Beaver [19].

*Echinostoma equinatus gigas* Marco del Pont, 1926 [28] was reported from Argentina, but the worms were later assigned to *Echinoparyphium recurvatum* by Lunaschi et al. [29]. *Echinostoma chloephaeae* Sutton and Lunaschi, 1980 also reported from Argentina attracted no much taxonomic attention but listed in a checklist of parasites of birds from Argentina [29]. *Echinostoma*



**Fig. 1.** A phylogenetic tree of *Echinostoma revolutum* (Southeast Asian and American lineages) and 6 other 37-collar-spined *Echinostoma* group constructed based on 184 bp of mitochondrial *cox1* sequences by maximum-likelihood method using the MEGA-X program employing Tamura-nei model of nucleotide substitution with 1,000 bootstrap replications. *Opisthorchis viverrini* was used as an out-group.



**Fig. 2.** A phylogenetic tree of *Echinostoma revolutum* (Eurasian and American lineages) and 13 other 37-collar-spined *Echinostoma* spp. constructed based on 472 bp of mitochondrial *nad1* sequences by maximum-likelihood method using the MEGA-X program employing Tamura-nei model of nucleotide substitution with 1,000 bootstrap replications. *O. viverrini* was used as an outgroup.

ralli Yamaguti, 1934 was reported in Japan and this name was used by Yoshino et al. [30]. The taxonomic validity of *Echinostoma robustum* Yamaguti, 1935 which was originally reported from Taiwan was supported by Detwiler et al. [31] through molecular analyses of specimens from USA and Brazil. However, the specific diagnosis of these worms as *E. robustum* needs reconfirmation. *Echinostoma goldi* Oschmarin, 1956 was reported from the intestine of the bird *Pernis apivorus* in Russian Far East [32]; this species was mentioned to have 37 collar spines by Yamaguti [1] but the figure presented by Skrjabin and Bashikirova [32] shows that it has 35 collar spines. Therefore, this species (species inquirenda) is excluded from the 37-collar-spined *Echinostoma* group until further studies are performed to confirm that it evidently has 37 collar spines.

*Echinostoma revolutum* var. *japonicum* Kurisu, 1932 and *Echinostoma stromi* Bashikirova, 1946 have been synonymized with *E. revolutum* by Yamaguti [1]. *Echinostoma revolutum tenuicolle* Bashikirova, 1941 seems to be a synonym of *E. revolutum*. Among the 3 species reported from UK, including *Echinostoma nudicaudatum* Nasir, 1960, *Echinostoma pinnicaudatum* Nasir, 1961, and *Echinostoma londonensis* Khan, 1961, the former 2 have seldom attracted taxonomic attention, whereas *E. londonensis* was suggested to be identical with *E. echinatum* by Kanev [4].

Kanev [4,20] studied on the life cycle of *E. revolutum* in Europe (from Germany) using the freshwater snail *Lymnaea stagnalis* as the starting point and redescribed the morphology of larval and adult *E. revolutum*. Kanev [4] and Kanev et al. [5,6] synonymized *Echinostoma audyi* Lie & Umathevy, 1965, *Echinostoma ivaniosi* Mahandas, 1973, and *E. paraulum* Dietz, 1909 with *E. revolutum*. Also, Kanev [4,20], Huffman and Fried [7], and Kanev et al. [5,6,33] synonymized *Echinostoma lindoense* Sandground & Bonne, 1940, *Echinostoma barbosa* Lie & Basch, 1966, *E. miyagawai* Ishii, 1932, and *E. revolutum* of Nasincova, 1986 with *E. echinatum*. Kanev et al. [5] also synonymized *Echinostoma sisjakowi* (Skvortsov, 1935) Yamaguti, 1971, *Echinostoma orlovi* Romashov, 1966, and *Echinostoma bolschewense* (Kotova, 1939) Nasincova, 1991 with *E. jurini* (Skvortsov, 1924) Kanev, 1985. Meanwhile, *Echinostoma liei* Jeyarasasingam et al., 1972, *Echinostoma togoensis* Jourdan & Kulo, 1981, and *Echinostoma paraensei* Lie & Basch, 1967 were synonymized with *Echinostoma caproni* Richard, 1964 by Huffman and Fried [7]. Thus, Kanev [4] listed only 5 species in the 37-collar-spined *Echinostoma* group, which included *E. revolutum* (sensu stricto), *E. trivolvis*, *E. caproni*, *E. jurini*, and *E. echinatum*. Later, however, *E. paraensei* was acknowledged as a dis-

tinct species because of its unique isoenzymatic patterns [34] and unique DNA sequences [35]. In addition, Kostadinova and Gibson [26] and Kostadinova et al. [21,36] reconsidered *E. miyagawai* as a distinct species, and rather questioned about the taxonomic status of *E. echinatum*.

In the meantime, *Echinostoma rodriguesi* Hsu et al., 1968 was reported as a new 37-collar-spined group from Brazil [37], and *Echinostoma parvocirrus* Nassi and Dupouy [38], 1988 from Guadeloupe, French West Indies. *Echinostoma friedi* Toledo et al. [39], 2000 was described as a new species in Spain. Thereby, Kostadinova et al. [26] listed 8 species in the 'revolutum' group, which included *E. revolutum* (sensu stricto), *E. jurini*, *E. trivolvis*, *E. paraensei*, *E. caproni*, *E. miyagawai*, *E. parvocirrus*, and *E. friedi* but did not include *E. rodriguesi*. According to Kostadinova et al. [40], a voucher specimen of *E. revolutum* designated by Morgan and Blair [41] from Australia was found to be affiliated to *E. robustum* (?) although they did not favor this specific diagnosis; later Georgieva et al. [10] assigned this isolate to *E. miyagawai*. However, the validity of *E. robustum* was supported by Detwiler et al. [31,42] based on materials from North America and Brazil. After then, *Echinostoma deserticum* Kechemir et al., 2002 was reported as a new 37-collar-spined species from Africa [43], and *Echinostoma luisreyi* Maldonado et al., 2003 as a new member from Brazil [44]. Fried and Graczyk [8] listed 10 species in the 37-collar-spined *Echinostoma* spp., including *E. revolutum*, *E. caproni*, *E. echinatum*, *E. friedi*, *E. jurini*, *E. luisreyi*, *E. miyagawai*, *E. paraensei*, *E. parvocirrus*, and *E. trivolvis*; however, they did not mention about *E. deserticum*, *E. rodriguesi*, and *E. robustum*. Toledo et al. [9] listed 10 species, including *E. revolutum*, *E. caproni*, *E. deserticum*, *E. friedi*, *E. jurini*, *E. luisreyi*, *E. miyagawai*, *E. paraensei*, *E. parvocirrus*, and *E. trivolvis*, but did not mention on *E. echinatum*, *E. rodriguesi*, and *E. robustum*.

Based on molecular, morphological, and ecological data, the classification of 37-collar-spined *Echinostoma* species became more diverse and have been continuously changing [10,11,24,31,42,45,46]. Faltýnková et al. [11] synonymized *E. friedi* with *E. miyagawai* based on morphological data. Mohanta et al. [46] suggested synonymy of *E. robustum*, *E. miyagawai*, and *E. friedi*, but the synonymy between *E. miyagawai* (*E. friedi*) and *E. robustum* is disagreed by other workers [47]. It is also noteworthy that several new cryptic 37-collar-spined *Echinostoma* spp. have been discovered in different parts of the world, which included *Echinostoma* sp. IG by Georgieva et al., 2013 [45], *Echinostoma* spp. clade 3 by *nad1* and ITS2 analyses by

Noikong et al. [48], *Echinostoma nasincovae* Georgieva et al., 2014 [10,11], *Echinostoma novaezealandense* Georgieva et al., 2017 [24], and *Echinostoma mekongi* Cho et al., 2020 [49]. Synonymies previously based only on morphological characters should be reconsidered until the results are firmly supported by molecular evidences.

## SPECIES REPORTED IN EACH CONTINENT

### Asia

At least 11 species, including *E. revolutum* [50-57], *E. audyi* [58], *E. cinetorchis* [59-61], *E. ivaniosi* [62], *E. lindoense* [63], *E. mekongi* [49], *E. miyagawai* [64,65], *E. paraulum* [14,66,67], *E. ralli* [68], *E. revolutum* var. *japonicum* [69], and *E. robustum* [70] were reported to have 37 collar spines. *E. audyi* and *E. ivaniosi* were synonymized with *E. revolutum* by Kanev [4], and *E. revolutum* var. *japonicum* with *E. revolutum* by Yamaguti [1]. Beaver [19] synonymized *E. cinetorchis* with *E. revolutum*; however, *E. cinetorchis* is characteristic in having testes which are mobile to other locations within the body; one or both testes even disappear from the body while they grow to be adults [2,3,13,22,71]. *E. lindoense* was synonymized with *E. echinatum* by Kanev et al. [33]; however, the status of *E. echinatum* is questioned by many workers [10,11,26]. In addition, *E. lindoense* has unique larval and adult morphology discriminating from *E. revolutum* (see *E. lindoense* section), and thus the name *E. lindoense* is retained in this review. *E. mekongi* was reported as a new species from human infections in 2 provinces of Cambodia along the Mekong River, which is morphologically close to *E. revolutum* and *E. miyagawai* but molecularly distinct from them and also 12 other 37-collar-spined *Echinostoma* spp. available in Gen-

Bank [49]. A recent mitochondrial DNA study reported that 2 distinct species, including *E. revolutum* and *E. miyagawai*, exist in Thailand and Lao PDR [72]. In China, complete mitochondrial genome of *E. miyagawai* has been obtained and characterized [73,74]. A synonymy between *E. miyagawai* and *E. robustum* was suggested based on molecular analysis in Bangladesh [46]; however, this synonymy is disagreed by Heneberg [47]. *Cytochrome B* gene was found to be useful to differentiate Asian *E. revolutum* from African *E. caproni* and South American *E. paraensei* [75]. Thus, at least 8 species, including *E. revolutum*, *E. cinetorchis*, *E. lindoense*, *E. mekongi*, *E. miyagawai*, *E. paraulum*, *E. ralli*, and *E. robustum*, are recognized to be existing in Asia (Table 3).

### Europe

More than 18 species of 37-collar-spined *Echinostoma* spp. were reported from European countries, including Germany, Bulgaria, UK, Austria, Czech Republic, Spain, and Russia; *E. revolutum*, *E. bolschewense*, *E. columbae*, *E. dilatatum*, *E. echinatum*, *E. friedi*, *E. goldi*, *E. jurini*, *E. londonensis*, *E. miyagawai*, *E. nasincovae*, *E. nudicaudatum*, *E. orlovi*, *E. oxycephalum*, *E. paraulum*, *E. pinnicaudatum*, *E. sijakowi*, and *Echinostoma* sp. IG of Georgieva et al., 2013 [45]. Among them, *E. columbae*, *E. dilatatum*, and *E. oxycephalum* were synonymized with *E. revolutum* or put as species inquirenda by Beaver [19]. *E. orlovi* and *E. sijakowi* were synonymized with *E. jurini* by Kanev et al. [5]. *E. goldi* had 35 collar spines in the figure of Skrijabin and Bashkirova [32] thus considered not a member of 37-collar-spined group. *E. bolschewense* and *E. londonensis* were synonymized with *E. jurini* and *E. echinatum*, respectively, and *E. paraulum* was synonymized with *E. revolutum* by Kanev [4] and Kanev et

**Table 3.** Continental distribution of 37-collar-spined *Echinostoma* spp.

Asia	Europe	Africa	North/Central America	South America	Oceania
<i>E. revolutum</i>	<i>E. revolutum</i>	<i>E. caproni</i>	<i>E. revolutum</i>	<i>E. revolutum</i>	<i>E. revolutum</i>
<i>E. cinetorchis</i>	<i>E. bolschewense</i>	<i>E. deserticum</i>	<i>E. parvocirrus</i>	<i>E. barbosai</i>	<i>E. acuticauda</i>
<i>E. lindoense</i>	<i>E. echinatum</i>		<i>E. robustum</i>	<i>E. chloephagae</i>	<i>E. miyagawai</i>
<i>E. mekongi</i>	<i>E. jurini</i>		<i>E. trivolvis</i>	<i>E. luisreyi</i>	<i>E. novaezealandense</i>
<i>E. miyagawai</i>	<i>E. miyagawai</i>			<i>E. paraensei</i>	<i>E. paraensei</i>
<i>E. paraulum</i>	<i>E. nasincovae</i>			<i>E. robustum</i>	
<i>E. ralli</i>	<i>E. nudicaudatum</i>			<i>E. rodriguesi</i>	
<i>E. robustum</i>	<i>E. paraulum</i>				
	<i>E. pinnicaudatum</i>				
	<i>E. robustum</i>				
	<i>Echinostoma</i> sp. IG				

This table has been revised and updated from Chai (2019) [3].



al. [5]. However, the synonymy of *E. bolschewense* with *E. jurini* was denied because of morphological differences in their cercariae [11]. Thus, *E. bolschewense* was regarded as a distinct species, and the status of *E. jurini* was retained [11]. In addition, *E. paraulum* was revalidated as distinct species [10,11]. On the other hand, *E. friedi* reported by Toledo et al. [39] was synonymized with *E. miyagawai* by Faltýnková et al. [11]. *E. robustum* Yamaguti, 1935 has been acknowledged as a distinct species based on specimens harvested from a duck in UK [76]; this species was also recorded from birds in Poland [77] and Russia [78,79]. *E. miyagawai* originally reported from Japan was later reported also in Bulgaria and Czech Republic [21,36]. Put together, in Europe, at least 11 species of 37-collar-spined echinostomes can be currently acknowledged to be valid or validity-retained; *E. revolutum* (sensu stricto), *E. bolschewense*, *E. echinatum*, *E. jurini*, *E. miyagawai*, *E. nasincovae*, *E. nudicaudatum*, *E. paraulum*, *E. pinnicaudatum*, *E. robustum*, and *Echinostoma* sp. IG [11,76] (Table 3).

#### Africa

More than 5 species of 37-collar-spined *Echinostoma* spp. were nominally described, including *E. caproni* [80,81], *E. deserticum* [43], *E. revolutum* [82], *E. liei* [83], and *E. togoensis* [84]. However, Kanev [4] suggested that *E. revolutum* described by Bisserru [82] was actually *E. caproni*. Fried and Huffman [85] also suggested that the studies performed under the name *E. revolutum*, *E. liei*, and *E. togoensis* using materials from Africa actually dealt with *E. caproni*. Thereby, only 2 species, *E. caproni* and *E. deserticum* can be currently recognizable as the 37-collar-spined *Echinostoma* spp. existing in Africa (Table 3).

#### North and Central America

The taxonomy of 37-collar-spined *Echinostoma* spp. in North and Central America has not yet been properly revised or settled [10]. In North America, 5 species were reported at an earlier time, including *E. trivolvis*, *E. armigerum*, *E. coalitum*, *E. collawayensis*, and *Echinoparyphium contiguum* [6,10]. Beaver [19] treated 4 of the above (*E. armigerum*, *E. coalitum*, *E. collawayensis*, and *E. contiguum*) as synonyms of *E. revolutum* or synonyms inquirenda. However, Kanev [20] pointed out that the *E. revolutum* of Beaver [19] and many articles published from North America under the name *E. revolutum* during 1968-1988 should be reconsidered to have been actually *E. trivolvis*. The true existence of *E. revolutum* in North America was later confirmed in 1997-1998 by Sorensen et al. [86,87] through morphological

as well as molecular studies. Later, based on *nad1* gene sequences, the American *E. revolutum* was shown to be genetically distinct from European populations, and further studies seemed necessary on the taxonomy of North American 37-collar-spined *Echinostoma* spp. [10,31]. In addition, Detwiler et al. [31] detected the presence of *E. robustum* from Indiana and Minnesota, USA, and also from Brazil. From Guadeloupe (French West Indies), *E. parvocirrus* was reported [38]. Thus, in North and Central America, at least 4 species can be listed as the 'revolutum' group; *E. revolutum*, *E. parvocirrus*, *E. robustum*, and *E. trivolvis* [38,42] (Table 3).

#### South America

More than 14 species, namely, *E. revolutum*, *E. armatum*, *E. barbosai*, *E. chloephagae*, *E. equinatus gigas*, *E. erraticum*, *E. luisreyi*, *E. mendax*, *E. microrchis*, *E. neglectum*, *E. nephrocystis*, *E. paraensei*, *E. robustum*, and *E. rodriguesi*, have been described in South America [19,31,42,44,88-90]. The presence of *E. revolutum* in Brazil has been reported by various workers, including Brasil and Amato [91], as reviewed by Maldonado and Lanfredi [92]. *E. barbosai* was synonymized with *E. echinatum* by Kanev et al. [33]; however, the status of *E. echinatum* is not recognized by other workers [10,11]. Therefore, the name *E. barbosai* is retained until further confirmatory studies are performed. Beaver [19] treated the 4 species reported from Brazil by Lutz [88], i.e., *E. erraticum*, *E. neglectum*, *E. microrchis*, and *E. nephrocystis*, as synonym inquirenda. These might be identical with *E. revolutum* or *E. trivolvis*. However, Kanev et al. [6] did not recognize the presence of *E. trivolvis* in South America; although, the possible existence of *E. trivolvis* in South America remains to be confirmed. In the present review, the 4 species reported by Lutz [88] were tentatively synonymized with *E. revolutum*. *E. equinatus gigas* reported by Marco del Pont [28] was later assigned to *Echinoparyphium recurvatum* [29]. Kanev [20] considered *E. paraensei* a synonym of *E. caproni*, and Huffman and Fried [7] accepted this synonymy. However, *E. paraensei* was acknowledged as a distinct species because of its unique isoenzymatic patterns [34] and DNA sequences [35]. The presence of *E. robustum* in South America was first suggested by molecular analysis of materials obtained from Brazil [31]. Therefore, the existence of at least 7 species of 37-collar-spined echinostomes, including *E. revolutum*, *E. barbosai*, *E. chloephagae*, *E. luisreyi*, *E. paraensei*, *E. robustum*, and *E. rodriguesi*, is currently recognized to be valid or validity-retained in South America (Table 3). It is of considerable interest that in a

mummified human body in Brazil, eggs presumed to be of *E. paraensei* (or *E. luisreyi*) were detected by a molecular technique [15].

### Oceania

A total of 5 species of 37-collar-spined echinostomes have been reported; *E. acuticauda*, *E. miyagawai*, *E. novaezealandense*, *E. paraensei*, and *E. revolutum* (Table 1). *E. acuticauda* was reported in 1914 in Australia [93], and soon synonymized (synonym inquirenda) with *E. revolutum* by Beaver [19]. However, this synonymy is not agreed by other workers, and the name *E. acuticauda* has been used by Jones and Anderson [23] and Georgieva et al. [24]. Morgan and Blair [41] reported that 3 metacercarial isolates (LMeta-1, PMeta-1, and PMeta-2) from Townsville showed greater than 98% sequence similarity with *E. revolutum* from Europe. Later, Kostadinova et al. [40] suggested all these isolates to be *E. robustum* (?). However, Georgieva et al. [10] placed all these isolates among the clusters of *E. miyagawai*. Thus, the presence of *E. robustum* in Australia needs reconfirmation. Morgan and Blair [41] also isolated *E. paraensei* (PCerc-1) cercariae from *Glyptophysa* sp. snails in Townsville, Australia by molecular analysis [41]. In New Zealand, a species closely allied to *E. revolutum* was found by molecular studies [41], and 2 more species, *E. novaezealandense* and *E. miyagawai*, were discovered [24]. Thus, at least 5 species are existing in Oceania, including *E. revolutum*, *E. acuticauda*, *E. miyagawai*, *E. novaezealandense*, and *E. paraensei* (Table 3).

## SPECIES OF 37-COLLAR-SPINED ECHINOSTOMA GROUP

### *Echinostoma revolutum* (Froelich, 1802) Dietz, 1909


[syn. *Echinostoma armatum* Barker & Irvine, 1915; *Echinostoma audyi* Lie & Umathevy, 1965; *Echinostoma columbae* Zunker, 1925; *Echinostoma dilatatum* (Miram, 1940) Cobbold, 1860; *Echinostoma echinocephalum* (Rudolphi, 1819) Cobbold, 1860; *Echinostoma erraticum* Lutz, 1924; *Echinostoma ivaniosi* Mahandas, 1973; *Echinostoma limicoli* Johnson, 1920; *Echinostoma mendax* Dietz, 1909; *Echinostoma microrchis* Lutz, 1924; *Echinostoma neglectum* Lutz, 1924; *Echinostoma nephrocystis* Lutz, 1924; *Echinostoma oxycephalum* (Rudolphi, 1819) Railliet, 1896; *Echinostoma revolutum tenuicollis* Bashikirova, 1941; *Echinostoma revolutum* var. *japonicum* Kurisu, 1932; *Echinostoma stromi* Bashikirova, 1946; *Echinostoma sudanense* Odhner, 1910]

This species (Fig. 3) was originally described under the

name *Fasciola revoluta* based on adult flukes found in the large intestine of wild ducks *Anas boschas fereae* dissected on July 20, 1798 in Germany [4]. It was about 11 mm long and had 37 collar spines, and these morphological features have served as the main basis for further definitions of *E. revolutum* [4]. A year later, *E. echinatum* (under the name *Distoma echinatum*) was described by Zeder in Germany, and this name (*D. echinatum* or *E. echinatum*) had been used for about a hundred years until Dietz [16,17] renamed *F. revoluta* as *Echinostoma revolutum* in his systematic reorganization of the Echinostomatidae and synonymized *E. echinatum* with *E. revolutum*. Dietz [16,17] also synonymized *Echinostoma dilatatum* (under the name *Distoma dilatatum* Miram, 1940) and *Echinostoma armatum* (under the name *Distoma armatum* Molin, 1858) with *E. revolutum*. This was a big milestone for the taxonomy of *E. revolutum* and 37-collar-spined *Echinostoma* group. After Dietz [16,17], adult worms and larval stages found in Europe, Asia, Africa, Australia, and North and South America have been described as *E. revolutum* for at least 70-80 years [1,4,19,20,32,88,94-98] (Table 4). On the other hand, Kanev [20] and Kanev et al. [33] suggested the taxonomic validity of *E. echinatum*, and Schuster [99], Christensen et al. [100], and Huffman and Fried [7] supported it (see *E. echinatum* section). However, the validity of *E. echinatum* was recently put to a question by Kostadinova and Gibson [26], Georgieva et al. [10], and Faltýnková et al. [11].

Human infection with *E. revolutum* was first described in Taiwan in 1929 [101]. Later, in 1982, the prevalence of *E. revolutum* among Taiwan population was reported to be 0.11-0.65% [102]. Further human infections were reported in mainland China [12], Indonesia [103], Thailand [104], Cambodia [51], and Lao PDR [56]. However, all of these reports are not based on molecular confirmation and need further verification regarding the species identification.

Kanev [4] studied on almost all previous records on *E. revolutum* and other 37-collar-spined group and delimited the definition of *E. revolutum*. He stated that 37-collar-spined echinostomes cannot be morphologically identified only by adult flukes but they can be more clearly discriminated by the morphology of larval stages, in particular, the cercariae, and host-parasite relationships. Thus, *E. revolutum* was delimited as those flukes having (1) lymnaeid snails as the 1st intermediate host, (2) various pulmonate and prosobranch snails, mussels, frogs, and freshwater turtles as the second intermediate host, (3) only birds as the definitive host, (4) cercariae and adults armed with 37 collar spines, (5) geographical distribution



**Fig. 3.** Adult specimens of *Echinostoma revolutum* (A) from Thailand (courtesy of Prof. Chalobol Wongsawad, Chiang Mai University), *E. cinetorchis* (B) from South Korea, *E. mekongi* (C) from Cambodia, and *E. robustum* (D) from India under the same magnification. Scale bar=2 mm.

only in Europe and Asia, (6) *Cercaria echinata* Siebold, 1937, *E. echinatum*, and *E. jurini* as closely related species in Europe, and (7) specific characteristics only in larvae and host-parasite relationships [4]. However, some of these delimitations had to be revised because of various new findings. Using molecular data, Detwiler et al. [31] confirmed that both lymnaeid and planorbis snails can serve as the 1st intermediate host of *E. revolutum*. Detwiler et al. [42] also confirmed that *E. revolutum* can infect a mammalian host, the muskrat *Ondatra zibethicus*, in USA. The geographical distribution of *E. revolutum* was extended to North and South America, and Oceania [31,41,42,91] (Tables 3, 4). Faltýnková et al. [11] redescribed the morphology of rediae, cercariae, and adults of *E. revolutum* based on European samples (from Germany, Finland, Czech Republic, Poland, Bulgaria, and Iceland). In comparison with the report of Kanev [4], they found in the cercariae 6 outlets of penetration gland-cells (4 outlets in Kanev [4]) and total 12 outlets of paraesophageal gland-cells (16-20 outlets in Kanev [4]) [11] (Ta-

ble 5).

Molecular analysis of *E. revolutum* was started by Morgan and Blair [35]. They compared the nucleotide sequences of ITS region (ITS1-5.8S-ITS2) of *E. revolutum* (originated from Germany) with those of *E. trivolvis*, *E. paraensei*, *E. caproni* (including *E. liei* and *Echinostoma* sp. II from Africa), and *Echinostoma* sp. I from Africa (*E. deserticum*). They concluded that *E. revolutum* was phylogenetically close to *E. trivolvis* and *E. paraensei* but distinct from one another, and far from *E. caproni* and *E. deserticum*. Morgan and Blair [41,105] further compared the usefulness of ITS region and 2 mitochondrial loci (*cox1* and *nad1*) to discriminate the species and found that *nad1* appeared to be the most informative locus for investigating phylogenetic relationships within the 37-collar-spined *Echinostoma* group. The molecularly distinct status of *E. revolutum* in relation to other species and other genera has been supported by Kostadinova et al. [40].

The existence of *E. revolutum* in North America (under the

**Table 4.** Geographical distribution of 26 valid or validity-retained 37-collar-spined *Echinostoma* species

Species and nominator	Continent and country where this species has been reported
<i>E. revolutum</i> <sup>a</sup>	Asia, Europe, North America, South America, Oceania
<i>E. acuticauda</i>	Australia
<i>E. barbosai</i>	Brazil
<i>E. bolschewense</i>	Czech Republic, Russia, Slovak Republic
<i>E. caproni</i>	Cameroon, Congo, Egypt, Madagascar, Togo
<i>E. chloephagae</i>	Argentina
<i>E. cinetorchis</i>	China, Indonesia, Japan, South Korea, Taiwan, Vietnam
<i>E. deserticum</i>	Algeria, Niger
<i>E. echinatum</i>	Europe (Germany and other countries)
<i>E. jurini</i>	Europe (Bulgaria, Russia, and other countries)
<i>E. lindoense</i>	Indonesia, Malaysia, The Philippines, Thailand
<i>E. luisreyi</i>	Brazil
<i>E. mekongi</i>	Cambodia
<i>E. miyagawai</i> <sup>b</sup>	Asia, Europe, Oceania
<i>E. nasincovae</i>	Czech Republic, Slovak Republic
<i>E. novaezealandense</i>	New Zealand
<i>E. nudicaudatum</i>	UK
<i>E. paraensei</i>	Australia, Brazil
<i>E. paraulum</i>	Austria, Bangladesh, China, Czech Republic, Germany, Pakistan, Russia
<i>E. parvocirrus</i>	Guadeloupe (West Indies)
<i>E. pinnicaudatum</i>	UK
<i>E. ralli</i>	Japan
<i>E. robustum</i> <sup>c</sup>	Asia, Europe, North America, South America
<i>E. rodriguesi</i>	Argentina, Brazil
<i>E. trivolvis</i>	North America (Canada, USA)
<i>Echinostoma</i> sp. IG Georgieva et al., 2013	Germany, Iceland, UK

This table has been updated from Chai (2019) [3].

<sup>a</sup>Asia (Bangladesh, China, India, Indonesia, Iran, Japan, Lao PDR, Malaysia, South Korea, Taiwan, Thailand, and Vietnam), Europe (Austria, Belarus, Bulgaria, Czech Republic, Finland, France, Germany, Greece, Hungary, Iceland, The Netherlands, Poland, Russia, Slovak Republic, UK, Yugoslavia), North America (USA), South America (Brazil), Oceania (New Zealand).

<sup>b</sup>Asia (China, Japan, Lao PDR, Thailand, South Korea, Vietnam), Europe (Austria, Bulgaria, Czech Republic, Hungary, Poland, Russia, Spain, Ukraine), Oceania (New Zealand).

<sup>c</sup>Asia (Bangladesh, China, Japan, Taiwan), Europe (Germany, Poland, Russia, UK, Uzbekistan), North America (USA), South America (Brazil).

name *E. echinatum*, actually *E. trivolvis*) was first documented in 1888 [106] followed by others in 1895 [107] and 1896 [108]. In the first document, the specimen was found in muskrats from Pennsylvania [19,42], and the next documents were about findings from rabbits and chickens, respectively [107,108]. In 1937, Beaver [19] reported *E. revolutum* from experimentally

infected birds and mammals, and Fried and coworkers performed extensive works thereafter (more than 70 studies during 1968-1988) under the name *E. revolutum* [7]. However, Kanev [4,20] mentioned that *E. revolutum* of Beaver [19] and all other North American literature dealing with *E. revolutum* until 1988 were supposed to have been *E. trivolvis*.

The true existence of *E. revolutum* in North America was first confirmed by Sorensen et al. [86] who found naturally infected freshwater snails *Lymnaea elodes* shedding 37-collar-spined cercariae in Indiana, USA. They experimentally infected various snail species to obtain the metacercariae, which were given to chicken and geese to obtain the adult flukes. Sorensen et al. [87] notified intraspecific variation in ITS loci sequences among isolates of *E. revolutum* (North American and European) and also *E. trivolvis* (different localities in Indiana). The existence of *E. revolutum* in North America was again confirmed by Detwiler et al. [31,42] through molecular analysis of ITS region, *cox1*, and *nad1* loci. In addition, Detwiler et al. [31] reported that both lymnaeid and planorbid snails can serve as the first intermediate host of *E. revolutum* and *E. trivolvis* in USA. Detwiler et al. [42] also confirmed that *E. revolutum* can infect a mammalian host, the muskrat in USA. Phylogenetic trees based on mitochondrial *cox1* (Fig. 1) and *nad1* (Fig. 2) sequences of 7 and 14 *Echinostoma* species, respectively, show their phylogenetic relationships.

The *nad1* sequence variation among the “*E. revolutum* group” indicated that *E. revolutum* haplotypes from Europe formed a monophyletic group which clustered closely with a monophyletic group of isolates from North America [45]. In Asia, molecular genetic studies were conducted on *E. revolutum* in comparison with *Hypoderaeum conoideum*, *Echinoparyphium recurvatum*, and *Artyfechinostomum malayanum* (under the name *Echinostoma malayanum*) using multilocus enzyme electrophoresis and sequencing of ITS1 and *cox1* [52,53,109,110]. However, isolates from Southeast Asia were not included in the study of Georgieva et al. [45] due to the lack of *nad1* data [72]. Subsequently, it was found that *E. revolutum* from Southeast Asia (Thailand and Lao PDR) clustered as a monophyletic clade with the European isolates, named as “Eurasian lineage”, and this was distinct from the American isolates, which was named as “American lineage” [72]. This *nad1*-based genetic variation of *E. revolutum* according to geographical locations has been supported by an analysis of *cytochrome B* (*CYTB*) gene [75]. Mohanta et al. [46] accepted the 2 genetic lineages of *E. revolutum* using ITS2 and *nad1* sequences. However, Buddhachat

**Table 5.** Characteristics of *Echinostoma revolutum* in comparison with closely related species<sup>a</sup>

	<i>E. revolutum</i>	<i>E. lindoense</i>	<i>E. miyagawai</i>	<i>E. robustum</i>	<i>E. trivolvis</i>
Definitive host	birds, mammals	mammals	birds, mammals	birds	birds, mammals
1st intermediate host	lymnaeid and planorbid snails	lymnaeid and planorbid snails	lymnaeid and planorbid snails	lymnaeid snails	lymnaeid and planorbid snails
No. of outlets of penetration gland-cells (cercaria)	6	6	6	?	6
No. of outlets of paraesophageal gland-cells (cercaria)	12	60-64	42-46	?	4-6
Size of metacercariae (µm)	132-152 (diameter)	120-130 (in diameter)	144-154 (in diameter)	?	135-170 (in diameter)
Collar spines (adult)	slender and sharply pointed	short and less sharply pointed	small with sharply pointed ends	innermost end group spines smaller than others	very sharply pointed
Body constriction near the ventral sucker level (adult)	no constriction	constricted at middle level of ventral sucker	constricted at posterior end level of ventral sucker	constricted at posterior end level of ventral sucker	no constriction
Shape of testes (adult)	smooth or slightly lobed	deeply lobed	subglobular and indented, 3 lobed	irregularly lobed and horizontally extended	smooth, oval or slightly irregular
Vitellaria confluence near the posterior end of body (adult)	not confluent	confluent	confluent	confluent	confluent

<sup>a</sup>Other closely related species include *E. cinetorchis* (reduced number and abnormal location of testes), *E. mekongi* (remarkably variable in body shape and morphology of testes, i.e., globular, slightly or deeply lobed; smaller head collar, collar spines, oral and ventral suckers, and cirrus sac compared to *E. revolutum* and *E. miyagawai*; vitellaria not confluent), *E. paraensei* (dorsalmost collar spines smaller than others), and *E. paraulium* (slight median indentation in 2 testes).

and Chontanarith [111] used DNA barcoding of *cox1* and *nad1* in combination with high-resolution melting analysis to identify *E. revolutum* and found that the *E. revolutum* clade of *nad1* phylogeny obtained from the Thai isolate formed a different lineage from the Eurasian lineage. Thus, the *nad1* phylogenetic tree revealed 3 lineages of *E. revolutum*; Asian, Eurasian, and American lineages [111]. The complete mitochondrial genome (17,030 bp) of *E. revolutum* was obtained and characterized by Le et al. [112] in comparison with 9 echinostome species, including *E. caproni*, *E. miyagawai*, and *E. paraensei*. They concluded that *E. revolutum* grouped in a monophyletic subclade as a sister taxon to *E. miyagawai* and paraphyletic to other echinostomatids in the Echinostomatidae [112]. A multiplex PCR technique targeting the *nad1* gene has been developed for differential detection of 4 medically important echinostome species, including *E. revolutum* [113].

After *E. miyagawai* was first described from domestic fowls and domestic and wild ducks in 1932 in Japan [114], Beaver [19] raised synonymy of *E. miyagawai* with *E. revolutum*. Yamaguti [1,97] accepted this synonymy but Russian researchers, including Bashkirova [96] and Skryabin and Bashkirova [32], denied the synonymy. Kosupko [115,116] was the first who ap-

plied complex morphological and ecological approaches to distinguish *E. revolutum* and *E. miyagawai* and validated the 2 species as separate taxa on the basis of their morphological differences in cercariae and adults, ontogenic development, first intermediate and final host preferences, and their distribution in the host intestine. Nevertheless, Kanev [4] denied the validity of *E. miyagawai* and synonymized it with *E. echinatum*. Kostadinova et al. [21] re-validated *E. miyagawai* to be distinct from *E. revolutum* and *E. echinatum* of Kanev [1,20] on the basis of experimental completion of its life cycle. According to them, the adult worms of *E. miyagawai* were characterized by having a very elongate body with a constriction at the posterior border of the ventral sucker (no constriction in *E. revolutum*), a large head collar with relatively small collar spines (larger collar spines in *E. revolutum*), a spherical ventral sucker which is only about half the maximum body width (larger ventral sucker in *E. revolutum*), a long cirrus-sac reaching posteriorly to the middle of the ventral sucker (to the anterior border of ventral sucker in *E. revolutum*), indented subglobular testes (globular or slightly lobed in *E. revolutum*), and vitellaria forming 2 lateral fields of follicles which are almost confluent in the post-testicular space (not confluent in *E. revolutum*) [21] (Table 5).

In addition, they [36] suggested 5 morphometric variables to distinguish *E. miyagawai* from *E. revolutum*, which included body width at posterior border of the ventral sucker, head collar width, length of the esophagus, width of the ventral sucker, and length of the pre-ovarian region.

#### *Echinostoma acuticauda* Nicoll, 1914

This species was originally described from a bird (straw-necked ibis *Carphibis spinicollis*) in North Queensland, Australia [93]. It was put to a synonymy with *E. revolutum* as a synonym inquirenda by Beaver [19]; however, this synonymy has not been followed by succeeding authors, for example, Mendheim [95], Jones and Anderson [23], Georgieva et al. [24], and Memon et al. [67]. This species is morphologically close to *E. novaezealandense* described from New Zealand but has a shorter forebody, more anteriorly located ovary and testes, and much larger eggs [24]. However, the taxonomic status of *E. acuticauda* should be re-evaluated by detailed morphological as well as molecular studies in the near future.

#### *Echinostoma barbosai* Lie & Basch, 1966

*E. barbosai* was originally described from pigeons, chicks, and ducklings experimentally infected with the metacercariae from *Biomphalaria glabrata* snails in Brazil [117]. This species resembled *E. audyi* (a synonym of *E. revolutum*) but was smaller in size of adults and different in cercarial characteristics and the first intermediate host [117]. *E. barbosai* was also found in Bulgaria by Kanev group [118]. However, Mutafova and Kanev [119] synonymized this species with *E. echinatum*, because the karyotype of *E. barbosai* was completely corresponded to that of *E. echinatum*. This synonymy was agreed by Huffman and Fried [7] and Fried and Graczyk [8]. However, Kostadinova and Gibson [26] doubted the status of both species, *E. echinatum* and *E. barbosai*. Meanwhile, Latin American authors continue to use the name *E. barbosai* [27,30,120,121]. Under this situation, the name *E. barbosai* should be retained until further confirmatory studies with molecular data are provided.

#### *Echinostoma bolschewense* (Kotova, 1939) Nasincova, 1991

This species originates from *Cercaria bolschewensis* Kotova, 1939 shed from prosobranch snails in the European part of Russia [122]. Later, the adult fluke, named as *E. bolschewense*, was collected from the small intestine of experimentally infected hamsters through a life cycle study in South Bohemia,

Czech Republic [122]. Kanev [4] and Kanev et al. [5] regarded this species a synonym of *E. jurini*. However, Georgieva et al. [10] studied on *nad1* and 28S rDNA sequences of *E. bolschewense* in the cercarial stage from Slovak Republic and found that *E. bolschewense* is genetically distinct from *E. revolutum*, *E. trivolvis*, *E. caproni*, *E. paraulum*, and *E. miyagawai* (but molecular data of *E. jurini* are yet unavailable). In addition, Faltýnková et al. [11] analyzed the differential morphological features of larval and adult stages of *E. bolschewense* in comparison with those of *E. juri* reported by Kanev et al. [5] and acknowledged the validity of *E. bolschewense*. However, they did not place *E. jurini* in synonymy with *E. bolschewense*. To re-validate the 2 closely related species, *E. bolschewense* and *E. jurini*, comparative molecular studies are required.

#### *Echinostoma caproni* Richard, 1964

[syn. *Echinostoma liei* Jeyarasasingam et al., 1972; *Echinostoma togoensis* Jourdan & Kulo, 1981]

This echinostome was originally reported from the small intestine of naturally infected birds (kestrels; *Falco newtoni*) and a domestic fowl experimentally infected with the metacercariae in a freshwater snail *Bulinus lirutus* in Madagascar [80]. The cercariae (under the name *E. liei*) were also found in *Biomphalaria alexandrina* snails collected in Egypt which were experimentally infected to *B. glabrata* and other snail species to obtain the metacercariae and then to chicks and hamsters to obtain the adult flukes [83]. This echinostome was also reported in Togo (under the name *E. togoensis*) which developed into rediae in *Biomphalaria pfeifferi*, metacercariae in aquatic pulmonate snails, and adults in laboratory mice [84]. The synonymy of *E. liei* and *E. togoensis* with *E. caproni* was proposed by Huffman and Fried [7] and Fried and Huffman [85] which was followed by other workers [3,26]. In addition, Kanev [20], Huffman and Fried [7], and Christensen et al. [100] synonymized *E. paraensei* with *E. caproni*. However, Sloss et al. [34] and Morgan and Blair [35] suggested that *E. paraensei* is distinct from *E. caproni* based on different isoenzyme electrophoretic patterns and DNA sequences of ITS region. *E. caproni* was studied also under the name of *E. revolutum* by many workers, including Barus et al. [123], Christensen [124], Christensen et al. [125-128], Bindseil and Christensen [129], and Simonsen and Andersen [130] as mentioned by Fried and Huffman [85].

The molecular data (*cox1* and *nad1*) of *E. caproni* have been available by Morgan and Blair [35,41,105] and Macilla [72]. *E. caproni* was phylogenetically close to *E. trivolvis* as shown by

*cox1* sequences [72]. *Cytochrome B* gene was also found to be useful to differentiate African *E. caproni* from South American *E. paraensei* and Asian *E. revolutum* [75].

#### *Echinostoma chloephae* Sutton & Lunaschi, 1980

*E. chloephae* was first discovered from the rectum of a bird *Chloepha picta melanopectera* in Argentina [131]. This species continued to be recorded from naturally infected birds, including *C. picta leucopectera*, in Argentina [29]. Metacercariae of seemingly *E. chloephae* were detected in a bivalve species *Diplodon chilensis* in Argentina, and adult flukes were obtained from experimentally infected chickens with the metacercariae [30]. The taxonomic validity of this species in comparison with other 37-collar-spined *Echinostoma* spp. should be evaluated in the near future, through morphological as well as molecular studies.

#### *Echinostoma cinetorchis* Ando & Ozaki, 1923

This echinostome (Fig. 3) was originally described based on specimens recovered from the small intestine of wild rats in Japan by Ando and Ozaki [59], and redescribed by Dollfus [132] in French. Kurisu [69] and Sugimoto [133] found this echinostome in the intestine of domestic fowls in Japan and a dog in Taiwan, respectively. Tanabe and Takeishi [134] reported a low prevalence (1.65%) of *E. cinetorchis* in rats *Rattus norvegicus* captured around Tokyo, Japan. In South Korea, this echinostome was reported from wild rats [135,136] and dogs [137]. Its life cycle was successfully completed in the laboratory using *Hippeutis cantori* snails as the source for cercariae and metacercariae [61]. Human infection with *E. cinetorchis* was first reported in Japan [138-140], and then in Korea [60,141-144]. Now, it is well known as a zoonotic echinostome infecting humans, rodents, and poultry in Asian countries, including Japan, South Korea, China, Taiwan, Indonesia, and Vietnam [3,92,145-148].

Beaver [19] synonymized *E. cinetorchis* with *E. revolutum* based on their morphologic similarities, including the number of collar spines ( $n=37$ ) and dimensions of internal organs and eggs. However, this synonymy was not justified by other workers due to various reasons, including, in particular, the unique feature of *E. cinetorchis* having testes that move to abnormal locations and even disappear from the body during the worms grow up to be adults (Table 5) as well as differences in selecting the snail host [1,3]. This species is thought to reproduce parthenogenetically without testes and sperms. Now

*E. cinetorchis* is acknowledged as a distinct species [2,3,92,149].

In 2016, *nad1* sequences of *E. cinetorchis* were deposited based on materials (adult flukes) obtained from South Korea by Lee et al. (to be published) in GenBank; the sequences are unique and clearly separated from those of *E. revolutum* Eurasian and American lineages.

#### *Echinostoma deserticum* Kechemir, Jourdane & Mas-Coma, 2002

*E. deserticum* was described as a new species from the small intestine of the African grass rat *Arvicanthis niloticus* from Niger [43]. The majority of adult flukes do not have testes; some have only 1 testis and very rarely 2 testes [43] which is highly similar to *E. cinetorchis*. This species is thought to reproduce parthenogenetically without testes and sperms [43]. Sporocysts and rediae were isolated from *Bulinus truncatus* and *B. globosus* snails from south of Algeria and Niger [43].

Molecular studies on this species were done under the name of *Echinostoma* sp. I from Niger by Morgan and Blair [35,41,105] in 1995 before it was formally described as a new species in 2002. The sequences of ITS region and mitochondrial *cox1* and *nad1* loci indicated that *E. deserticum* is the most distinct in comparison with other 37-collar-spined echinostomes, including *E. revolutum*, *E. trivolvis*, *E. paraensei*, and *E. caproni* (also *E. liei* and *Echinostoma* sp. II) [35,41,87,105]. In addition, the *nad1* sequence of *E. deserticum* appears to be markedly different from that of the morphologically similar species, *E. cinetorchis*, deposited in GenBank, with sequence homology of only 84.7%.

#### *Echinostoma echinatum* (Zeder, 1803) de Blainville, 1828 [syn. *Echinostoma londonensis* Khan, 1961]

*E. echinatum* was described in 1803 under the name *Distoma echinatum* from birds and mammals (?) in Germany, with detailed descriptions of collar spines [4,150]. A year before (1802), *Fasciola revoluta* (= *E. revolutum*) was described by Froelich from birds in Germany [4]. In 1809, Rudolphi erected a subgeneric group *Echinis* (*Echinostoma*) within the genus *Distoma* and placed *F. revoluta* as the type species [4]. After Rudolphi, however, *F. revoluta* was out of use, and its function as the type species was replaced by *Distoma echinatum* Zeder, 1803 [4]. Practically almost all 37-collar-spined echinostomes found in Europe from Rudolphi in 1809 until Dietz in 1909 were diagnosed as *D. echinatum* [4]. However, Dietz [16,17] restored the validity of *F. revoluta* renaming it as *E. revolutum* in his systemat-

ic reorganization of the echinostomatid flukes and synonymized *D. echinatum* with *E. revolutum*. A list of adult and larval worms described under the name *D. echinatum* was transferred to *E. revolutum* by Dietz [4]. Thereafter, over 500 articles dealing with this echinostome have described the species name as *E. revolutum* [4]. Beaver [19], Mendheim [95], Bashikirova [96], Skrjabin and Bashikirova [32], and Yamaguti [1,97] followed this synonymy, and the name *E. echinatum* disappeared for some time.

Later, however, *E. echinatum* was revived by Kanev [4,20] and Kanev et al. [5,33] based on various differential characters from *E. revolutum* and *E. jurini*; for example, *E. echinatum* took both lymnaeid and planorbid snails as the first intermediate host, whereas *E. revolutum* and *E. jurini* took only lymnaeid and viviparid snails, respectively. The definitive host of *E. echinatum* was birds and mammals, including humans, whereas *E. revolutum* infected only avian hosts, and *E. jurini* infected only mammalian animals [5]. The 3 species also differed in the cercarial chaetotaxy and developmental period of eggs and cercariae [5,33]. The validity of *E. echinatum* was supported by other workers, including Schuster [99], Christensen et al. [100], Huffman and Fried [7], and Fried and Graczyk [8].

However, some of the differential characters suggested above were denied by succeeding authors; for example, it turned out that *E. revolutum* can infect mammalian hosts too, i.e., muskrats in USA, based on molecular analysis of *nad1* sequences [42]. In addition, *E. revolutum* can take both lymnaeid and planorbid snails as the first intermediate host in USA [31,42]. Moreover, Kostadinova [151] criticized Kanev and coworkers' report of *E. echinatum* (sensu Kanev [4,20] and Kanev et al. [33]) stating that re-examination of the voucher specimens identified by Kanev et al. [33] as *E. echinatum* showed 2 other distinct species; *Echinostoma sarcinum* Dietz, 1909 possessing 47 collar spines and *Echinostoma* sp. (*E. echinatum*?) with 37 collar spines.

Regarding *E. lindoense*, its geographical distribution was reported to extend from the original Asian countries (Indonesia and Malaysia) to Brazil [152], Poland, Czechoslovakia (now Czech Republic and Slovak Republic) [153], and Spain [154]. Based on this, Kanev [4] described that *E. echinatum* is distributed in different geographical regions of Asia, South America, and Europe, including Germany, where *E. echinatum* was originally described. However, Kostadinova and Gibson [26] doubted the synonymy of *E. lindoense* with *E. echinatum* because of the remarkable differences in the cercarial morpholo-

gy. Moreover, the validity of *E. echinatum* remains to be re-validated because *E. echinatum* has not been formally described in a taxonomic publication [10,11,45]. In GenBank, molecular data under the name *E. echinatum* is so far unavailable and should be provided in order to re-validate its taxonomic status. Because of the unique cercarial and adult worm morphology (see *E. lindoense* section), the name *E. lindoense* should be retained until molecular data in comparison with *E. echinatum* (currently unavailable) become available.

#### *Echinostoma jurini* (Skvortsov, 1924) Kanev, 1985

[syn. *Echinostoma orlovi* Romashov, 1966; *Echinostoma sisjakowi* (Skvortsov, 1935) Yamaguti, 1971]

*E. jurini* was originally described under the name *Cercaria jurini* based on cercariae shed from viviparid snails collected along the Volga River in Russia [5]. Nasinkova [122] stated that the description of *C. jurini* Skvortsov, 1924 was so inaccurate and incomplete that it cannot be considered a description of a species. Later, this species was redescribed by Kanev [20] and Kanev et al. [5] based on larvae and adults obtained experimentally starting from naturally infected *Viviparus contectus* and *V. viviparus* snails collected along the Danube River in Bulgaria. The metacercariae were obtained from the renopericardial sac of laboratory-raised snails, *Physa acuta* and *P. fontinalis* after exposure to the cercariae, and adults were recovered from experimental hamsters, rats, and mice [5]. The morphology of adult flukes closely resembled that of *E. revolutum*; however, Kanev [4] and Kanev et al. [5] validated this species because of differences in selecting intermediate and definitive hosts and differences in the morphology of cercariae. They stated that *E. jurini* infects viviparid snails and mammals, whereas *E. revolutum* infects lymnaeid snails and avian hosts. They put a variety of echinostome species in possible synonymy with *E. jurini*, which included *E. orlovi* Romashov, 1966, *Echinostoma sisjakowi* (Skvortsov, 1934) Yamaguti, 1971, and *Echinoparyphium sisjakowi* (Skvortsov, 1934).

The validity of *E. jurini* was acknowledged by Kostadinova and Gibson [26], Fried and Graczyk [8], and Toledo et al. [9]. However, Kostadinova and Gibson [26] criticized that the conclusions made by Kanev et al. [5] that "Practically, all 37-collar-spined adults of *Echinostoma* spp. in mammals in Europe (including *E. bolschewense*) might be identical with *E. jurini*." appeared to be an overestimate of the degree of morphological variability in the absence of an experimental support. Moreover, Georgieva et al. [10] obtained *nad1* and 28S rDNA se-



quences of *E. bolschewense* adults originating from *Viviparus acerosus* snails in the river Danube, Slovak Republic and found that *E. bolschewense* is genetically distinct from *E. revolutum*, *E. trivolvis*, *E. caproni*, *E. paraulum*, and *E. miyagawai*. However, molecular data of *E. jurini* have not yet been reported [72]. Faltýnková et al. [11] analyzed differential morphological features of larval and adult stages of *E. bolschewense* in comparison with those of *E. juri* reported by Kanev et al. [5]. The cercariae of *E. jurini* had 6 outlets for the penetration gland-cells and 8-10 outlets for the paraesophageal gland-cells on the dorsal lip of the oral sucker [5]; however, the cercariae of *E. bolschewense* possessed 10 outlets for the penetration gland-cells (6 median and 4 medio-lateral) and lacks paraesophageal gland-cells [11]. Furthermore, the adults of *E. bolschewense* had much larger eggs (138-162 × 75-85 µm) than those of *E. jurini* (96-132 × 72-88 µm) [11]. Based on these characters, they denied the synonymy raised by Kanev et al. [5] giving remarks on the validity of *E. bolschewense*. They neither placed *E. jurini* in synonymy with *E. bolschewense*. To re-validate these 2 closely related species comparative molecular studies seem to be urgently needed.

#### *Echinostoma lindoense* Sandground & Bonne, 1940

This species was first described from human patients in the Lake Lindoe region of Central Celebes, Indonesia [63]. This echinostome was also discovered from humans in Jakarta, Indonesia and then animals, including rats in Malaysia, Thailand, and the Philippines [155-157]. An echinostome species similar in their morphology and biology to *E. lindoense* was also found in Brazil [152] and European countries, including Bulgaria, Poland, Czech Republic, Slovak Republic, Germany, Austria, UK, and Russia [153,158]. However, the *E. lindoense* reported from Europe and South America needs reconfirmation through molecular analysis with Asian (Indonesian) species.

The major differential points of *E. lindoense* from *E. revolutum* included morphology of larval stages (particularly the cercariae) and adults; in cercariae *E. lindoense* had 2 dorsal, 2 ventral, and 2 small ventrolateral tail fin-folds, whereas *E. revolutum* had only 1 small dorsal fin-fold near the tip of the tail [158]. In addition, the testes of *E. lindoense* in adults were deeply indented (Asian strain) or superficially lobed (Brazilian strain; this needs re-evaluation), but those of *E. revolutum* are smooth or only slightly lobed [4,152] (Table 5). *E. lindoense* has shorter and wider collar spines than *E. revolutum* which has longer and slender collar spines with more or less pointed

ends [103].

The larval and adult stages of *E. lindoense* were morphologically and biologically similar to those of *E. echinatum* originating from Germany, and thus *E. lindoense* was synonymized with *E. echinatum* by Kanev and coworkers [4-6,20,33]. This synonymy was accepted by Huffman and Fried [7], Fried and Graczyk [8], Toledo et al. [149], and Toledo and Esteban [148]. However, Kostadinova and Gibson [26] and Kostadinova et al. [21,36] questioned about the taxonomic validity of *E. echinatum*, because of its many uncertain points, for example, a wide range (more than 20) of the number of paraesophageal gland-cell outlets in cercariae, and *E. echinatum* has not been properly reported as a taxonomic publication. The questionable status of *E. echinatum* was agreed in subsequent publications, including Georgieva et al. [10,45] and Faltýnková et al. [11]. Unfortunately, *E. lindoense* has not yet been molecularly analyzed. Chai [3] tentatively acknowledged the validity of *E. lindoense*.

#### *Echinostoma luisreyi* Maldonado, Vieira & Lanfredi, 2003

*E. luisreyi* Maldonado et al., 2003 was originally described from mice and hamsters experimentally infected with metacercariae from *Physa marmorata* and *B. glabrata* snails in Brazil [44]. The most important morphological character of this species was the oral corner spines that increase in size from the latero-oral to the ventro-oral regions [44]. Later, a rodent species, *Akodon montensis*, was found to be a natural definitive host [159]. Fried and Graczyk [8], Toledo et al. [9], and Georgieva et al. [10] recognized this species to be distinct among the 37-collar-spined *E. revolutum* group, and Pinto and Melo [27] mentioned about the cercariae of this species that emerged from *P. marmorata* snails in Brazil. This echinostome might have been zoonotic 600-1,200 years ago in Brazil, since eggs morphologically suggestive of *E. luisreyi* were found from the coprolites of human mummies [160]. Another report suggesting *E. luisreyi* or *E. paraensei* infection in a mummified body (520-600 years before present) of a human in Brazil mentioned that the eggs in coprolites were morphologically similar to *E. luisreyi* but molecularly (*cox1* gene; 83 bp) closer to *E. paraensei* [15]. Molecular data of *E. luisreyi* are not yet available in GenBank.

#### *Echinostoma mekongi* Cho, Jung, Chang, Sohn, Sinuon & Chai, 2020

*E. mekongi* (Fig. 3) was reported based on adult flukes recovered from human infections after praziquantel treatment and

purging in Kratie and Takeo Province, Cambodia along the Mekong River, which is molecularly distinct from *E. revolutum* and 13 other 37-collar-spined *Echinostoma* spp. deposited in GenBank [49]. Adults of this species were 9.0-13.1 in length and 1.3-2.5 mm in maximum width and the eggs in feces and worm uterus were 98-132 µm long and 62-90 µm wide [49]. The adult worms closely resembled those of *E. revolutum*, *E. miyagawai*, and several other 37-collar-spined *Echinostoma* species [49]. However, this species revealed remarkable variation in body shape and morphology of the testes (globular, slightly or deeply lobed), and smaller head collar, collar spines, oral and ventral suckers, and cirrus sac compared to *E. revolutum* and *E. miyagawai* [49] (Table 5). Sequencing of 2 mitochondrial genes (*nad1* and *cox1*) and a nuclear ITS region (ITS1-5.8S rRNA-ITS2) revealed unique features distinct from *E. revolutum* (Southeast Asian, Eurasian, European, and American lineages) and other 37-collar-spined *Echinostoma* group available in GenBank (*E. bolschewense*, *E. caproni*, *E. cinetorchis*, *E. miyagawai*, *E. nasincovae*, *E. novaezealandense*, *E. paraensei*, *E. paraulum*, *E. robustum*, *E. trivolvis*, and *Echinostoma* sp. IG) [49]. Studies on biological and epidemiological characteristics, including the life cycle, geographical distribution, and source of human infections, are urgently needed [49].

#### *Echinostoma miyagawai* Ishii, 1932

[syn. *Echinostoma friedi* Toledo, Muñoz-Antolí & Esteben, 2000]

This species was first described from domestic fowls, domestic ducks, and wild ducks in Japan [114]. Later, an experimental human infection was found successful in China [12]. Beaver [19] synonymized *E. miyagawai* with *E. revolutum* because of their morphological similarities, including the size and arrangement of collar spines, morphology of 2 testes, and size of the oral and ventral suckers, pharynx, and other internal organs. Yamaguti [1,97] accepted this synonymy. However, Russian researchers, including Bashikirova [96] and Skryabin and Bashikirova [32], denied this synonymy. In particular, Kosupko [115-116] applied complex morphological and ecological approaches to distinguish *E. revolutum* and *E. miyagawai* and validated the 2 species on the basis of their morphological differences in the cercariae and adults, ontogenic development, first intermediate and final host preferences, and their distribution in the host intestine. Opposed to this, Cooper [161] and Kanev [4] denied the validity of *E. miyagawai* and synonymized it with *E. revolutum* or *E. echinatum*, respectively. Later, however,

Kostadinova [162] and Kostadinova et al. [21,36] re-validated *E. miyagawai* on the basis of unique cercarial chaetotaxy and morphometric data of *E. miyagawai* in comparison with *E. revolutum* through experimental completion of its life cycle. The adult worms of *E. miyagawai* were characterized by a considerably elongate body with a constriction at the posterior border of the ventral sucker, a large head collar with relatively small collar spines, a spherical ventral sucker which is only about half the maximum body width, a long cirrus-sac reaching posteriorly to the middle of the ventral sucker, indented subglobular testes, and vitellaria forming 2 lateral fields of follicles which are almost confluent in the post-testicular space [21] (Table 5). In addition, Kostadinova et al. [36] suggested 5 morphometric variables to distinguish *E. miyagawai* from *E. revolutum*, which included body width at the posterior border of ventral sucker, head collar width, length of esophagus, width of ventral sucker, and length of the pre-ovarian region. Fried and Graczyk [8] and Toledo et al. [9] acknowledged the taxonomic validity of this species. *E. friedi* reported as a new species by Toledo et al. [39] in 2000 was synonymized with *E. miyagawai* based on morphological data by Faltýnková et al. [11].

Molecular data (*nad1* and 28S rRNA) of *E. miyagawai* have been available since Georgieva et al. [10]. The *nad1* analysis showed that *E. miyagawai* is in a distinct clade clearly separated from *E. revolutum* (sensu stricto; European), *E. revolutum* (American lineages), *E. paraulum*, and *E. caproni* [10]. This *E. miyagawai* clade incorporates *E. friedi* reported in GenBank by Marcilla et al. (AJ564379, Valencia, Spain) [10] and *E. revolutum* German strain (AF065832) by Morgan and Blair [41,105]. Nagataki et al. [72] found that *E. miyagawai* from Southeast Asia was monophyletic with European isolates, named as “Eurasian lineage”, which was slightly different from “Australian lineage” with no significant genetic differentiation being observed between these lineages. Fu et al. [73] and Li et al. [74] obtained the complete mitochondrial genome of *E. miyagawai* (Hunan isolate or Heilongjiang isolate, China) which was 14,416 or 14,468 bp in size and consisted of 12 protein-coding genes, 22 transfer RNA genes, 2 ribosomal RNA genes, and 1 non-coding region. Mohanta et al. [46] suggested a synonymy of *E. robustum*, *E. miyagawai*, and *E. friedi* based on *nad1* sequence analysis. However, Heneberg [47] did not agree to this synonymy but suggested that G1-G11 isolates from Bangladesh designated as *E. robustum* were in fact *E. miyagawai*. The taxonomic validities of both *E. miyagawai* and *E. robustum* have

been acknowledged by various workers [21,31,36,42].

The geographical distribution of *E. miyagawai* has been extended from Japan [163] to South Korea [64,65], Russia [164], China [12,73,74], Vietnam [165], Thailand, Lao PDR [72], Bulgaria [21,36,162], Austria and Hungary [166], Poland [167], Spain (under the name of *E. friedi*) [39], Czech Republic [11], and New Zealand [24] (Table 4).

*Echinostoma nasincovae* Faltýnková, Georgieva, Soldánová & Kostadinova, 2015

[syn. *Echinostoma spiniferum* (La Valette, 1855) sensu Nasincova, 1992; *Echinostoma revolutum* of Nasincova, 1986]

Našincová [168] described the life cycle of an echinostome species she believed to be *E. revolutum* in Central Europe, which she had completed in the laboratory based on cercariae from *Planorbarius corneus* snails [11]. Later, she recognized that the cercariae described from *P. corneus* differs from the cercariae of *E. revolutum* in the pattern of paraesophageal gland-cells and used the name *Echinostoma spiniferum* in her Ph.D. thesis in 1992 [11]. Later, Georgieva et al. [10] described a new species of *Echinostoma* (named as *Echinostoma* n. sp.) based on molecular dataset of *nad1* and 28S rDNA sequences during the analyses of 37-collar-spined *Echinostoma* spp. in Europe. This *Echinostoma* n. sp. formed a unique genetic clade neighbored with *E. paraensei* and 3 lineages (A-C) of *E. trivolvis* [31,42] and remotely joined by 3 isolates of *E. caproni* [10]. Faltýnková et al. [11] described this as a new species, *E. nasincovae*, synonymizing *E. revolutum* of Našincová [168] and *E. spiniferum* (La Valette, 1855) sensu Našincová, 1992 with *E. nasincovae*. They [11] described that its snail intermediate host is *P. corneus*, the definitive host is experimental hamsters, and the geographical localities included Czech Republic and Slovak Republic. In 2008, Faltýnková et al. [166] found *Gyraulus albus*, *P. corneus*, *Planorbis planorbis*, and *Bathymphalus contortus* from Czech Republic, Slovak Republic, Germany, and Poland acting as the snail hosts for *E. spiniferum* (synonym of *E. nasincovae*). Schwelm et al. [169] also listed *G. albus* as a snail host for *E. nasincovae*.

The cercariae of *E. nasincovae* differed from the cercariae of other species of *Echinostoma* within the 'revolutum' group in small body size, 6 ducts of penetration gland-cells opening on the dorsal lip of the oral sucker, and 30-39 outlets of paraesophageal gland-cells, of which 8 are located in the region of the oral sucker and 22-31 are confined to the area between esophagus and the dilated portion of the main collecting ex-

cretory ducts [11]. The adult of *E. nasincovae* was most similar to *E. revolutum* and *E. bolschewense*. However, compared with *E. revolutum*, *E. nasincovae* had smaller body and organs, smaller angle, lateral, and dorsal collar spines, and smaller eggs [11]. Compared with *E. bolschewense*, *E. nasincovae* had smaller body length and width, smaller egg length, smaller lateral and dorsal spine length, and smaller ovary and testes [11].

*Echinostoma novaezealandense* Georgieva, Blasco-Costa & Kostadinova, 2017

In 1998, Morgan and Blair [41] discovered an unidentified 37-collar-spined *Echinostoma* sp. adults (NZ-Ad) from the gut of a wild Canada goose (*Branta canadensis*) in New Zealand. This isolate (AF026289) was molecularly distinct from *E. revolutum* and *E. paraensei* based on the sequence of *nad1* gene [41]. Later, the same flukes were recovered from the small intestine and rectum of 2 avian species, a wild duck (*Anas platyrhynchos*) and a black swan (*Cygnus atratus*), in Central Otago District, New Zealand, and morphological and molecular studies (*nad1* and 28S rDNA) were performed [24]. The *nad1* sequences of 2 isolates from the ducks (KY436398 and KY436399) and 1 isolate from the swan (KY436397) revealed strong association with that of NZ-Ad (AF026289) of Morgan and Blair showing a unique position in the phylogenetic tree neighboring with *E. miyagawai*, *E. paraulum*, and *E. robustum/friedi* lineages [24]. The 28S rDNA sequence of 1 isolate from the swan (KY436407) also revealed a unique position neighboring with *E. miyagawai*, *E. paraulum*, *E. revolutum*, *Echinostoma* sp. IG, *E. trivolvis*, *E. paraensei*, and *E. nasincovae* [24]. Thus, the adult specimens were described as a new species, *E. novaezealandense* [24]. This species was morphologically close to *E. acuticauda* described in Australia but differed in having a longer forebody, more posteriorly located ovary and testes, and much smaller eggs [24]. This species is now listed among the helminth fauna of birds in New Zealand [170].

*Echinostoma nudicaudatum* Nasir, 1960

This species was first described from the intestine of experimental pigeons infected with the metacercariae from *L. stagnalis* snails in Birmingham, UK [171]. The species name came from the fact that the cercariae had no fin-folds on the tail [171]. The cercariae of *E. revolutum* have a fin-fold on the tail, rod-like cystogenous gland-cells, and 36 (possibly more) flame cells, but those of *E. nudicaudatum* were in the complete lack of a fin-fold on the tail, granular gland-cells in the cercarial

body, and at least 94 flame cells [171]. The adults of *E. nudicaudatum* differed from those of *E. revolutum*, *E. robustum*, and *E. lindoense* in the absolute sizes of collar spines, and from *E. ralli* in the arrangement of collar spines [171]. Interestingly, intra-redial encystment of the cercariae (i.e., metacercariae) was observed within the rediae of *E. nudicaudatum*, which means precocious encystment in snails [172,173], as observed similarly in *E. trivolvis* [6].

During the experimental life cycle, production of at least 3 redial generations was observed [172]. A year after the discovery of *E. nudicaudatum*, *E. pinnicaudatum* (the name came from the cercariae having a well-developed dorso-ventral fin-fold) was reported as a new species, from the same place where *E. nudicaudatum* was discovered [174]. In adult flukes, only a minor difference was found in the arrangement of collar spines between *E. nudicaudatum* and *E. pinnicaudatum*; in the former there were 7 unalternating lateral spines but in the latter there were 5 unalternating collar spines [174]. The cercariae of these 2 species were recorded among the cercarial fauna from British freshwater molluscs [175]. However, taxonomic attention has seldom been paid to these 2 species of *Echinostoma* spp., and further studies to evaluate their taxonomic validity through molecular analysis are needed.

#### *Echinostoma paraensei* Lie & Basch, 1967

This species was originally described from Brazil based on experimentally obtained adults from mammalian hosts, including hamsters, mice, and rats infected with the metacercariae from *B. glabrata* snails [176]. Later, this fluke was detected from naturally infected water rats, *Nectomys squamipes*, in Brazil [89,90]. It is of note that the eggs of *E. paraensei* (less probably *E. luisreyi*) were detected in the coprolite of a mummified human body in Brazil using molecular techniques [15].

The adult *E. paraensei* resembled adults of *E. revolutum*, *E. lindoense*, and *E. barbosai*; however, the minute size of 5-11 dorsal-most collar spines was characteristic for *E. paraensei* [176] (Table 5). Kanev [20] considered *E. paraensei* a synonym of *E. caproni* based mainly on the fact that *E. paraensei* also uses species of *Biomphalaria* as the 1st intermediate host. Huffman and Fried [7] accepted this concept, and Kanev [4] listed only 5 species in the 37-collar-spined group, which included *E. revolutum* (sensu stricto), *E. trivolvis*, *E. caproni*, *E. jurini*, and *E. echinatum* excluding *E. paraensei*. However, *E. paraensei* was acknowledged as a distinct species because of its unique isoenzymatic patterns [34], DNA sequences [35], patterns of random

amplification of polymorphic DNA (RAPD) [177], host-parasite relationships [178], and unique metacercarial excystation patterns and morphological features [179]. The validity of *E. paraensei* was agreed by Fried and Huffman [85], Kostadinova and Gibson [26], Fried and Graczyk [8], and Toledo et al. [9].

Molecular data of *E. paraensei* have been available since Morgan and Blair [35,41,105] who analyzed the ITS region, *cox1*, and *nad1* of *E. paraensei* (from Australia) and other 37-collar-spined *Echinostoma* spp. The results evidenced that *E. paraensei* (PCer-1) is distinct from *E. caproni* and phylogenetically rather close to *E. trivolvis* [35,41,105]. In addition, Morgan and Blair [41] found an isolate (cercariae from *Glyptophysa* sp. snails; AF026282) of *E. paraensei* from Townsville, Australia, which diverged from the Brazilian species (AF025834) by only 2 nucleotide substitution out of 530 sequenced. Georgieva et al. [10] further reported that *E. paraensei* was grouped with *E. nasincovae* (under the name *Echinostoma* n. sp.) and *E. trivolvis* forming a second clade among the 37-collar-spined *Echinostoma* group. Recently, cytochrome *B* gene was found to be useful to differentiate *E. paraensei* from Asian *E. revolutum* and African *E. caproni* [75].

#### *Echinostoma paraulum* Dietz, 1909

This species was originally described from the bird *Colymbus cristatus* and several species of ducks from Central Europe, including Austria and Russia [4,16,17,19]. Miller [180] described this species in North America (Canada) and synonymized *Echinostoma columbae* Zunker, 1925 with *E. paraulum*. However, Kanev et al. [6] regarded Miller's specimens as *E. trivolvis*, whereas Beaver [19] synonymized *E. columbae* with *E. revolutum*. Human infection with *E. paraulum* was reported from Russia in 1938 [181] and also from Yunnan Province, China in 1979 [14]. The rediae and cercariae were found in *P. corneus*, and the metacercariae were found in *Planorbis carinatus* snails [182]. This host selection was different in *E. revolutum*; rediae and cercariae developed in *Lymnaea palustris* and metacercariae in *P. corneus* [182]. The metacercariae of *E. revolutum* were bigger having a thick cyst wall but those of *E. paraulum* were smaller having a thin cyst wall [182].

Skrjabin [183], Sprehn [184], Skrijabin and Bashikirova [32], and Iskova [185] supported the taxonomic validity of *E. paraulum*; however, Beaver [19], Yamaguti [1,97], and Kanev [4,20] denied the validity and synonymized it with *E. revolutum*. Beaver [19] mentioned that 37 collar spines of *E. paraulum* were arranged exactly as in *E. revolutum*, its relative size of various

organs was identical with that of *E. revolutum*, and there was not a single character by which *E. paraulum* can be separated from *E. revolutum*. Yamaguti [1,97] synonymized various species with *E. revolutum*, including *E. echinatum*, *E. armatum*, *E. revolutum* var. *japonicum*, *E. miyagawai*, and *E. paraulum*. Kanev [4,20] confirmed its complete life cycle from Europe, including Austria where *E. paraulum* was originally described and concluded that *E. paraulum* corresponded completely with *E. revolutum*. Thereafter, Huffman and Fried [7], Kostadinova and Gibson [26], Fried and Graczyk [8], and Toledo et al. [9] did not recognize *E. paraulum* as a distinct species.

Georgieva et al. [10] was the first who molecularly confirmed the unique status of *E. paraulum*. They obtained 5 isolates from cercariae (4; *L. stagnalis*) and an adult (1; *Aythya fuligula*) originating from Germany and Czech Republic, respectively, and analyzed *nad1* and 28S rDNA sequences together with other 37-collar-spined *Echinostoma* spp. The phylogenetic tree of *nad1* showed that *E. paraulum* was distinct from all other species analyzed and included within the 1st clade where *E. revolutum* (sensu lato), *E. miyagawai*, *E. novaeseelandense* (as *Echinostoma* sp. NZ-Ad), and *E. robustum/friedi* are located [10]. The phylogenetic tree of 28S rDNA supported this showing similar findings [10]. They also mentioned that although *E. paraulum* and *E. revolutum* use the same snail species (*L. stagnalis*) as their 1st intermediate host, the cercariae of *E. paraulum* were dissimilar to those of *E. revolutum* in the pattern of paraesophageal gland-cells [10]. Subsequently, the morphology and life cycle of *E. paraulum* were redescribed by Faltýnková et al. [11]. According to them, the cercariae of *E. paraulum* were much larger in size with a much longer tail than those of *E. revolutum*, possessing a larger number of paraesophageal gland-cell outlets than the cercariae of *E. revolutum*. The adults of *E. paraulum* were in comparison with the same age of *E. revolutum* much smaller in size, with a more robust body, longer forebody, and distinctly wider head collar, and notably their testes had a characteristic median constriction [11] (Table 5). Nagataki et al. [72] supported the validity of *E. paraulum* using *nad1* analysis of 37-collar-spined *Echinostoma* spp. and mentioned that *E. paraulum* formed a sister group with *E. robustum* and *E. miyagawai*. Mohanta et al. [46] also established a phylogenetic tree based on *nad1* which supported the taxonomic validity of *E. paraulum*. Adult *E. paraulum* was reported from the house crow in Pakistan [67] and ducks in Bangladesh [66]. However, the specimen from Pakistan showed testes with no distinct median constriction, and thus the specific diagnosis

needs reconfirmation.

#### *Echinostoma parvocirrus* Nassi & Dupouy, 1988

This species was described as a new species from the naturally infected *B. glabrata* snails in Guadeloupe, French West Indies [38]. This snail host served as both the 1st and 2nd intermediate host (metacercariae also in *P. marmorata* snails and tadpoles of *Bufo marinus*), and birds (ducklings of *Cairina moschata* and canaries) served as the experimental definitive host [38]. The adults morphologically resembled *E. trivolvis* (under the name *E. multispinosum* reported from birds in Cuba), *E. revolutum* (under the name *E. mendax* reported from birds in Brazil), and *E. barbosai* (from birds in Brazil) but differed in body length, size of oral and ventral suckers, pharynx, and other organs [38]. In the cercariae of *E. parvocirrus*, the number of penetration gland-cell pores was 6 (4 in *E. barbosai*), and the number of paraesophageal gland-cell pores at the cephalic portion was 8 ventral and 4 dorsal (11 ventral and 6 dorsal in *E. barbosai*) [38]. This species has been listed among the 37-collar spined '*E. revolutum*' spp. by Kostadinova and Gibson [26], Fried and Graczyk [8], Maldonado and Lanfredi [92], Toledo et al. [9], Nagataki et al. [72], and Le et al. [112]. However, taxonomic attention on this species has been scarce, and there has been no information on molecular characteristics.

#### *Echinostoma pinnicaudatum* Nasir, 1961

This species was described based on adult flukes obtained from a laboratory-raised pigeon experimentally fed the metacercariae in *L. stagnalis* snails from Birmingham, UK [174]. The snails had both cercarial and metacercarial stages, and collected in the same locality where *E. nudicaudatum* was discovered [171,174]. The adults of *E. pinnicaudatum* were morphologically close to *E. nudicaudatum* with only a recognizable difference in the arrangement of collar spines; in the former there were 5 unalternating lateral spines, whereas in the latter there were 7 of these [174]. However, the morphology of cercariae was distinctly different. In the cercaria of *E. pinnicaudatum* there was a well-developed, dorso-ventral fin-fold (as the name '*pinnicaudatum*' indicates) but in the cercaria of *E. nudicaudatum* the fin-fold was totally absent [174]. The cercariae of *E. pinnicaudatum* was recorded among the cercarial fauna from British freshwater molluscs [175]. However, taxonomic attention has never been paid to *E. pinnicaudatum*, and further studies to evaluate its taxonomic validity through molecular analysis should be performed.

*Echinostoma ralli* Yamaguti, 1934

This species was originally described from the small intestine of a waterfowl, *Rallus aquaticus indicus* in Aichi, Shiga, and Shizuoka Prefecture, Japan [68]. Later, this species was found again in the same species of birds in Fukuoka Prefecture, Japan [186] and in the bird *Fulica atra* in Azerbaidzhan [1]. The adult worms had total 37 collar spines with 4 end group spines on each side, cup-shaped ventral sucker, and vitellaria beginning at the level of the posterior end of the ventral sucker [68]. It resembled *E. revolutum* (including synonyms *E. mendax*, *E. erraticum*, *E. microrchis*, and *E. neglectum*), *E. paraulum*, and *E. acuticauda* but differed in the number of end group spines, size of eggs, and others [68]. No further information is available on this species, and its taxonomic validity should be re-evaluated providing molecular data.

*Echinostoma robustum* Yamaguti, 1935

*E. robustum* (Fig. 3) was first described from the small intestine of 2 naturally infected avian species, *Streptopelia chinensis formosa* and *Columba livia domestica*, in Taiwan [187]. Domestic pigeons, chickens, and ducks, wild ducks, and wild geese were also found to be definitive hosts [1,97]. This species is morphologically close to *E. miyagawai* (and *E. friedi*) but can be differentiated in several features, including the morphology of testes, irregularly and deeply lobed and horizontally extended in *E. robustum* whereas subglobular with irregular margins or 3 lobed in *E. miyagawai* [46,98,187] (Table 5). In addition, the size of the ventral sucker in comparison to the body width is smaller in *E. miyagawai* [21] than in *E. robustum* [187]. The existence of this species has been extended to Japan [18], China [188], Bangladesh [46,66,70], Russia [78,79], Uzbekistan, Azerbaidzhan, Georgia, Estonia, Bulgaria [1], Poland [77], UK [76], USA [31,42], and Brazil [31] (Table 4). Cercariae develop in freshwater snails (*Radix lagotis*, *R. auricularia*, and *Planorbis* sp.), and encyst in the same or different snails (*Radix* spp. or *Cristaria plicata*), fish (*Hemiculter leucisculus*, *Acheilognathus chankaensis*, or *Gobio gobio*), or frogs (*Rana temporaria*) [79].

This species was synonymized with *E. revolutum* by Bezubik in 1956, but soon acknowledged as a distinct species by Rayski and Fahmy [76] in 1962 and listed in Skrjabin and Bashikirova [32] and Yamaguti [1,97]. However, Huffman and Fried [7], Kanev [4], Kostadinova and Gibson [26], Fried and Graczyk [8], and Toledo et al. [9] did not recognize this as a valid species. Kanev [4] synonymized this species with *E. echinatum*; however, it appeared to be inappropriate since *E. echinatum* it-

self needs revalidation [10,11,45]. The validity of *E. robustum* was supported by Detwiler et al. [31,42] through molecular analyses (*cox1*, *nad1*, and ITS) of isolates from Indiana and Minnesota, USA and Minas Gerais, Brazil. Georgieva et al. [45] indicated that *E. friedi* sensu Marcilla et al. (unpublished, in GenBank only, AJ564379) and *E. revolutum* of Morgan and Blair [35,41,105] (AF025832) from Europe had strong association with *E. robustum* sensu Detwiler et al. [31] from USA (GQ463053, GQ463054) and Brazil (GQ463055). The clade comprising of these 5 isolates exhibited a complex structure suggesting the existence of at least 3 different species [45]. Further studies are needed to back up this suggestion.

In addition, the clones of *E. robustum* sensu Detwiler et al. [31] and *E. friedi* (Marcilla et al., unpublished) showed almost identical *nad1* sequences and formed together subclades 'a-d' (*E. robustum*/*E. friedi* lineage). However, *E. friedi* was synonymized with *E. miyagawai* based on morphological analyses [11]. Mohanta et al. [46] suggested a synonymy between *E. robustum*, *E. miyagawai*, and *E. friedi* based on analysis of *nad1* sequences; however, Heneberg [47] did not agree to this synonymy but suggested that the G1-G11 isolates designated as *E. robustum* by Mohanta et al. [46] were in fact *E. miyagawai*. In this respect, whether *E. miyagawai* and *E. robustum* are taxonomically distinct species, despite several morphological differences [46], needs additional studies. Moreover, whether the specimens from Indiana and Minnesota (USA) and Brazil [31] are conspecific with the original *E. robustum* from Taiwan requires reconfirmation.

*Echinostoma rodriguesi* Hsu, Lie & Basch, 1968

Adults of this species were originally reported from experimental chicks, pigeons, hamsters, and mice fed the metacercariae in laboratory-raised *B. glabrata* snails in California, USA; the cercariae were originally obtained from *Physa rivalis* snails in Minas Gerais, Brazil and the infected snails were shipped to California [37]. *P. rivalis* also served as the 2nd intermediate host [37]. In adults of *E. rodriguesi*, the arrangement of collar spines resembled those of *E. revolutum*, *E. lindoense*, *E. barbosai*, and *E. paraensei*, and the size and shape of collar spines and testes were indistinguishable from those of *E. audyi* (synonym of *E. revolutum*) and *E. barbosai* [37]. However, the number of flame cells on the cercariae was 21 pairs in *E. rodriguesi*, *E. lindoense*, and *E. paraensei* but *E. barbosai* had 24 pairs, and *E. revolutum* (under the name *E. audyi*) had 27 pairs [37]. This species was also reported from rodents in Argentina [189,190].

Regarding its taxonomic validity, Kanev [20] was of opinion that *E. rodriguessi* may be a synonym of *E. trivolvis*, and Kostadinova and Gibson [26] did not list this species as a distinct species. Fried and Graczyk [8] and Toledo et al. [9] also excluded *E. rodriguessi* from the list of 37-collar-spined *Echinostoma* group. However, Kanev et al. [6] suggested that *E. trivolvis* occurred only in North America and did not include *E. rodriguessi* among the synonyms of *E. trivolvis*. In addition, Kanev et al. [18] used the name *E. rodriguessi* and mentioned about the migration of sporocysts within the snail body as a characteristic feature. Thereafter, *E. rodriguessi* has been acknowledged as a distinct species and mentioned by various researchers, including Latin American workers [27,44,89,90,120,189-191]. However, molecular data have not been provided for *E. rodriguessi*, and thus further validation of this species is needed.

#### *Echinostoma trivolvis* (Cort, 1914) Kanev, 1985

[syn. *Echinostoma armigerum* Barker and Irvine, 1915; *Echinostoma callawayensis* Barker and Noll, 1915; *Echinostoma coalitum* Barker and Beaver, 1915; *Echinoparyphium contiguum* Barker and Bastron, 1915; *Echinostoma multispinosum* Pérez Viguera, 1944]

This species originated from *Cercaria trivolvis*, its rediae, and metacercariae in the North American planorbid snail, *Helisoma trivolvis* (under the name *Planorbis trivolvis*) from Urbana, Illinois, USA [192,193]. The adult flukes of this species were described earlier in North America under the name *E. echinatum* by Leidy from musk rats in 1888 [106] followed by Stiles and Hassall from cotton rabbits in 1895 [107] and Hassall from chickens in 1896 [108]. They were also described under different names *E. armigerum* and *E. coalitum* by Barker on the specimens from muskrats in Nebraska in 1915 [194] and from muskrats in Ontario, Canada by Law and Kennedy in 1932 [19,195].

Beaver [19] studied on *C. trivolvis* shed from *Helisoma trivolvis* snails, obtained adult flukes in the laboratory, and assigned them as *E. revolutum*. Later, however, Kanev [20] and Kanev et al. [6] pointed out that the *E. revolutum* of Beaver [19] was actually *E. trivolvis*, and many of the articles published from North America until 1988 under the name *E. revolutum* should be reconsidered as having been *E. trivolvis*. Later, the true existence of *E. revolutum* in North America was morphologically as well as molecularly confirmed by Sorensen et al. [86,87]; it was transmitted by *L. elodes* snails in Indiana, USA. In addition, based on *nad1* gene sequences, the American *E. revolutum* was shown to be genetically different from European popula-

tions [10,31].

The morphological details of *E. trivolvis* larvae and adults were described by Kanev et al. [6]. He used the *C. trivolvis* shed from *H. trivolvis* snails collected from Douglas Lake, Michigan, USA as the starting point and obtained metacercariae from the pericardial sac of laboratory-raised *P. acuta* and *P. fontinalis* snails and adults from golden hamsters, rats, mice, ducks, geese, chickens, pigeons, turkeys, partridges, and guinea fowls. The first intermediate host also included *Physa gyrina* snails at St. Joseph, Illinois, USA [196]. Metacercariae developed also in the pericardial sac and posterior kidney region of mussels (*Anodonta cygnea*) and various freshwater snails, including *Viviparus viviparus*, *V. contectus*, *L. stagnalis*, *L. tomentosa*, *L. truncatula*, *L. palustris*, *L. peregra*, *L. auricularia*, *P. corneus*, *B. glabrata*, *B. alexandriana*, *Bithynia tentaculata* and *B. leachi*, and kidney and eye socket of tadpoles and frogs and freshwater turtles [6]. Precocious encystment of metacercariae within rediae was observed [6]. *C. trivolvis* has 37 collar spines, 7 fin-folds on the tail, 2 pairs of 3 flame cells at the level of the pharynx and total 36 flame cells, 6 pores of penetration gland-cells along the esophagus, and 4-6 pores of paraesophageal gland-cells on the oral sucker [6]. The cercariae were also described under different names, *C. trisolenata* Faust, 1917, *C. acanthostoma* Faust, 1918, and *C. complexa* Faust, 1919 [6]. The metacercariae were 135-170 µm in diameter, having a thick outer cyst wall and a thin inner cyst wall [6]. The adult flukes had a variable body size, from 3-5 mm to 20-30 mm, variable body shape, from elongate, flattened dorso-ventrally, and tapering posteriorly, to broad, with a tapering anterior end and a bluntly rounded posterior end [6].

*E. trivolvis* is different from *E. revolutum* in various points, including the number and arrangement of protein fractions in adult worm homogenates, number and distribution of argen-tophilic structures [197], number and position of paraesophageal gland-cell ducts in cercariae (4-6 on the oral sucker only in the former and 12 on the oral sucker and body in the latter), and geographical distribution [4,6,11] (Table 5). Sloss et al. [34] showed different electrophoretic patterns between *E. trivolvis*, *E. caproni*, and *E. paraensei*.

Kanev et al. [6] synonymized *E. armigerum*, *E. callawayensis*, *E. coalitum*, *E. multispinosum*, and *Echinoparyphium contiguum* with *E. trivolvis*. Kanev [20] also considered *E. rodriguessi* a synonym of *E. trivolvis* but later Kanev et al. [6] denied the presence of *E. trivolvis* in South America and did not list *E. rodriguessi* among the synonyms of *E. trivolvis*. The distinct taxonomic

status of *E. trivolvis* has been acknowledged by Huffman and Fried [7], Kostadinova and Gibson [26], Fried and Graczyk [8], Maldonado and Lanfredi [92], Toledo et al. [9], Detwiler et al. [31,42], and Georgieva et al. [10,24,45].

The molecular data of *E. trivolvis* have been available after Morgan and Blair [35] who compared the nucleotide sequences of ITS region (ITS1-5.8S-ITS2) with those of *E. revolutum*, *E. paraensei*, *E. caproni* (including *E. liei* and *Echinostoma* sp. II), and *Echinostoma* sp. I (*E. deserticum*). They concluded that *E. trivolvis* was phylogenetically close to *E. revolutum* and *E. paraensei* but distinct from one another and far from *E. caproni* and *Echinostoma* sp. I (*E. deserticum*). Sorensen et al. [87] notified intraspecific variation in ITS loci among isolates of *E. trivolvis* (different localities in Indiana) and *E. revolutum* (North American and European). The *nad1* and *cox1* sequences became available after Morgan and Blair [41] and Detwiler et al. [31,42]. The clones used were 3 from the cercariae of *E. trivolvis* from *H. trivolvis* and *L. elodes* snails in Indiana and Minnesota and 1 from the adult worm recovered in muskrats in Wisconsin [31]. The *nad1* phylogeny demonstrated that the muskrats were infected with 5 echinostome lineages; 3 *E. trivolvis* lineages, 1 *E. revolutum*, and 1 *Echinoparyphium* lineage 1 [31]. High levels of intraspecific variation with 3 lineages of *E. trivolvis* suggested the existence of multiple cryptic species within *E. trivolvis* [31,42].

*Echinostoma* sp. IG of Georgieva, Selbach, Faltýnková, Soldánová, Sures, Skirnisson & Kostadinova, 2013

[syn. *Echinostoma* cf. *friedi* of Kostadinova et al. (2003)]

*Echinostoma* sp. IG is a tentative name of a unique species but not yet formally described [45]. It was based on a larval stage (cercariae) from *Radix peregra* in Iceland and *R. auricularia* in Germany [11,45]. Its molecular information has been available for *nad1* and 28S rDNA sequences [10,45]. Cercariae and rediae ex. *Planorbis* sp. snails found in Wales, UK showing unique molecular characteristics [40] were also assigned to this species [11]. The cercariae of this species were unique having distal dorsal fin-fold and 12 paraesophageal gland-cell outlets restricted to the region of the oral sucker [11]. The adult stage has not yet been found [11].

## CONCLUSION

A total of 26 *Echinostoma* species are recognized as valid (16 species) or validity-retained (10) among the 37-collar-spined

*Echinostoma* spp. around the world. The morphology of the cercarial stage as well as the size and shape of adult flukes, including the relative size of the oral and ventral suckers, size, shape, and arrangement of collar spines, and morphology of testes and cirrus sac as well as arrangement of vitelline follicles are important criteria for species differentiation. However, some of these species are difficult to distinguish one from the other only by morphology. Molecular techniques, in particular, gene sequencing of mitochondrial *cox1* and *nad1*, is a highly useful technique for discriminating the species and seems to be essential to determine the taxonomic validity of 37-collar-spined *Echinostoma* spp. group.

## ACKNOWLEDGMENT

We are grateful to the staff of Institute of Parasitic Diseases, Korea Association of Health Promotion, Seoul, Korea who helped echinostome studies.

## CONFLICT OF INTEREST

We have no conflict of interest related to this study.

## REFERENCES

1. Yamaguti S. Synopsis of digenetic trematodes of vertebrates. Vol. I. Tokyo, Japan. Keigaku Publishing Co. 1971, pp 1-1074.
2. Chai JY. Echinostomes in humans. In Fried B, Toledo R, eds, The Biology of Echinostomes. New York, USA. Springer. 2009, pp 147-183.
3. Chai JY. Human Intestinal Flukes. Chapter 2. Echinostomes. The Netherlands. Springer Nature B.V. 2019, pp 169-343.
4. Kanev I. Life-cycle, delimitation and redescription of *Echinostoma revolutum* (Froelich, 1802) (Trematoda: Echinostomatidae). Syst Parasitol 1994; 28: 125-144.
5. Kanev I, Dimitrov V, Radev V, Fried B. Redescription of *Echinostoma jurini* (Skvortzov, 1924) with a discussion of its identity and characteristics. Ann Naturhist Mus Wien 1995; 97B: 37-53.
6. Kanev I, Fried B, Dimitrov V, Radev V. Redescription of *Echinostoma trivolvis* (Cort, 1914) (Trematoda: Echinostomatidae) with a discussion on its identity. Syst Parasitol 1995; 32: 61-70.
7. Huffman JE, Fried B. *Echinostoma* and echinostomiasis. Adv Parasitol 1990; 29: 215-269.
8. Fried B, Graczyk TK. Recent advances in the biology of *Echinostoma* species in the 'revolutum' group. Adv Parasitol 2004; 58: 139-195.
9. Toledo R, Esteban JG, Fried B. Recent advances in the biology



- of echinostomes. *Adv Parasitol* 2009; 69: 147-204.
10. Georgieva S, Faltýnková A, Brown R, Blasco-Costa I, Soldánová M, Sitko J, Scholz S, Kostadinova A. *Echinostoma 'revoletum'* (Digenea: Echinostomatidae) species complex revisited: species delimitation based on novel molecular and morphological data gathered in Europe. *Parasit Vectors* 2014; 7: 520.
  11. Faltýnková A, Georgieva S, Soldánová M, Kostadinova A. A reassessment of species diversity within the '*revoletum*' group of *Echinostoma Rudolphi*, 1809 (Digenea: Echinostomatidae) in Europe. *Syst Parasitol* 2015; 90: 1-25.
  12. Mao SP. Protozoan and helminth parasites of humans in mainland China. *Int J Parasitol* 1991; 21:347-351.
  13. Chai JY. Chapter 19. Echinostomes. In Xiao L, Ryan U, Feng Y eds, *Biology of Foodborne Parasites*. Boca Raton, USA. CRC Press, Taylor & Francis Group. 2015, pp 427-443.
  14. Li T, He S, Zhao H, Zhao G, Zhu XQ. Major trends in human parasitic diseases in China. *Trends Parasitol* 2010; 26: 264-270.
  15. Leles D, Cascardo P, Freire A dos S, Maldonado Jr A, Sianto L, Araujo A. Insights about echinostomiasis by paleomolecular diagnosis. *Parasitol Int* 2014; 63: 646-649.
  16. Dietz E. Die Echinostomiden der Vögel. *Zool Anz* 1909; 34: 180-192 (in German).
  17. Dietz E. Die Echinostomiden der Vögel. *Zool Jarb* 1910; Suppl XII: 265-512 (in German).
  18. Kanev I, Sterner M, Radev V, Fried B. An overview of the biology of echinostomes. In Fried B, Graczyk TK eds, *Echinostomes as Experimental Models for Biological Research*. Dordrecht, Boston, London. Kluwer Academic Publishers, Amsterdam, The Netherlands. 2000, pp 1-29.
  19. Beaver PC. Experimental studies on *Echinostoma revolutum* (Froelich), a fluke from birds and mammals. *Illin Biol Monogr* 1937; 15: 1-96.
  20. Kanev I. On the morphology, biology, ecology and taxonomy of *E. revolutum* group (Trematoda: Echinostomatidae: *Echinostoma*). Ph.D. Thesis. University of Sofia, Bulgaria. 1985.
  21. Kostadinova A, Gibson DI, Biserkow V, Chipev N. Re-validation of *Echinostoma miyagawai* Ishii, 1932 (Digenea: Echinostomatidae) on the basis of the experimental completion of its life-cycle. *Syst Parasitol* 2000; 45: 81-108.
  22. Chai JY. Intestinal flukes. In Murrell KD, Fried B eds, *Food-Borne Parasitic Zoonoses. Fish and Plant-Borne Parasites*. New York, USA. Springer. 2007, pp 53-115.
  23. Jones A, Anderson TJC. Helminths of rodents and marsupials from Papua New Guinea, with the description of two new species, *Echinostoma echymiperæ* n. sp. (Digenea: Echinostomatidae) and *Vampirolepis peroryctis* n. sp. (Cestoda: Hymenolepididae). *Syst Parasitol* 1990; 15: 223-237.
  24. Georgieva S, Blasco-Costa I, Kostadinova A. Molecular characterization of four echinostomes (Digenea: Echinostomatidae) from birds in New Zealand, with descriptions of *Echinostome novaezealandense* n. sp. and *Echinoparyphium poulini* n. sp. *Syst Parasitol* 2017; 94: 477-497.
  25. Kohn A, Fernandes BMM. Sobre as espécies do género *Echinostoma Rudolphi*, 1909 descritas por Adolpho Lutz em 1924. *Mem Inst Oswaldo Cruz* 1975; 73: 77-89.
  26. Kostadinova A, Gibson DI. The systematic of the echinostomes. In Fried B, Graczyk TK eds, *Echinostomes as Experimental Models for Biological Research*. Dordrecht, Boston, London. Kluwer Academic Publishers, Amsterdam, The Netherlands. 2000, pp 31-57.
  27. Pinto HA, Melo AL. A checklist of cercariae (Trematoda: Digenea) in mulluscs from Brazil. *Zootaxa* 2013; 3666: 449-475.
  28. Marcó del Pont. Contribución al estudio de los zooparásitos de animales salvajes. *Semana Méd* 1926; 33: 16-22 (in Spanish).
  29. Lunaschi L, Cremonte F, Drago FB. Checklist of digenean parasites of birds from Argentina. *Zootaxa* 2007; 1403: 1-6.
  30. Yoshino T, Nakamura S, Endoh D, Onuma M, Osa Y, Teraoka H, Kuwana T, Asakawa M. A helminthological survey of four families of waterfowl (Ardeidae, Rallidae, Scolopacidae and Phalaropodidae) from Hokkaido, Japan. *J Yamashina Inst Ornithol* 2009; 41: 42-54.
  31. Detwiler JT, Bos DH, Minchella DJ. Revealing the secret lives of cryptic species: examining the phylogenetic relationships of echinostome parasites in North America. *Mol Phylogenet Evol* 2010; 55: 611-620.
  32. Skrjabin KI, Bashikirova EI. Family Echinostomatidae Dietz, 1909. In Skrjabin KI ed, *Trematodes and Animals and Man*. Moscow-Leningrad, Russia, 1956; Vol XII: 51-930 (in Russian).
  33. Kanev I, Vassilev I, Bayssade-Dufour C, Albaret JI, Gassone J. Cercarial chaetotaxy of *Echinostoma revolutum* (Froelich, 1802) and *E. echinatum* (Zeder, 1803) (Trematoda, Echinostomatidae). *Ann Parasitol Hum Comp* 1987; 62: 222-234 (in French).
  34. Sloss B, Meece J, Tomano M, Nollen P. The genetic relationships between *Echinostoma caproni*, *E. paraensei*, and *E. trivolvis* as determined by electrophoresis. *J Helminthol* 1995; 69: 243-246.
  35. Morgan JA, Blair D. Nuclear rDNA ITS sequence variation in the trematode genus *Echinostoma*: an aid to establishing relationships within the 37-collar-spine group. *Parasitology* 1995; 111: 609-615.
  36. Kostadinova A, Gibson DI, Biserkow V, Chipev N. A quantitative approach to the evaluation of the morphological variability of two echinostomes, *Echinostoma miyagawai* Ishii, 1932 and *E. revolutum* (Froelich, 1802), from Europe. *Syst Parasitol* 2000; 45: 1-15.
  37. Hsu KC, Lie KJ, Basch PF. The life history of *Echinostoma rodriguessi* sp. n. (Trematoda: Echinostomatidae). *J Parasitol* 1968; 54: 333-339.
  38. Nassi H, Dupouy J. Experimental study of the life history of *Echinostoma parvocirrus* n. sp. (Trematoda: Echinostomatidae), a larval parasite of *Biomphalaria glabrata* in Guadeloupe. *Ann Parasitol Hum Comp* 1988; 63: 103-108 (in French).
  39. Toledo R, Muñoz-Antoli C, Esteban JG. The life-cycle of *Echinostoma friedi* n. sp. (Trematoda: Echinostomatidae) in Spain and a discussion on the relationships within the '*revoletum*' group based on cercarial chaetotaxy. *Syst Parasitol* 2000; 45:

- 199-217.
40. Kostadinova A, Herniou EA, Barrett J, Littlewood DTJ. Phylogenetic relationships of *Echinostoma* Rudolphi, 1809 (Digenea: Echinostomatidae) and related genera re-assessed via DNA and morphological analyses. *Syst Parasitol* 2003; 54: 159-176.
  41. Morgan JA, Blair D. Mitochondrial ND1 gene sequences used to identify echinostome isolates from Australia and New Zealand. *Int J Parasitol* 1998; 28: 493-502.
  42. Detwiler JT, Zajac AM, Minchella DJ, Belden LK. Revealing cryptic parasite diversity in a definitive host: echinostomes in muskrats. *J Parasitol* 2012; 98: 1148-1155.
  43. Kechemir N, Jourdane J, Mas-Coma S. Life cycle of a new African echinostome species reproducing by parthenogenesis. *J Nat Hist* 2002; 36: 1777-1784.
  44. Maldonado Jr A, Vieira GO, Lanfredi RM. *Echinostoma luiseryi* n. sp. (Platyhelminthes: Digenea) by light and scanning electron microscopy. *J Parasitol* 2003; 89: 800-808.
  45. Georgieva S, Selbach C, Faltýnková A, Soldánová M, Sures B, Skirnisson K, Kostadinova A. New cryptic species of the 'revolutum' group of *Echinostoma* (Digenea: Echinostomatidae) revealed by molecular and morphological data. *Parasit Vectors* 2013; 6: 64.
  46. Mohanta UK, Watanabe T, Anisuzzaman, Ohari Y, Itagaki T. Characterization of *Echinostoma revolutum* and *Echinostoma robustum* from ducks in Bangladesh based on morphology, nuclear ribosomal ITS2 and mitochondrial *nad1* sequences. *Parasitol Int* 2019; 69: 1-7.
  47. Heneberg P. Taxonomic comments on the validity of *Echinostoma miyagawai* Ishii, 1932 (Trematoda: Echinostomatidae). *Parasitol. Int* 2020; 74: 101931.
  48. Noikong W, Wongsawad C, Chai JY, Saenphet S, Trudgett A. Molecular analysis of echinostome metacercariae from their second intermediate host found in a localised geographic region reveals genetic heterogeneity and possible cryptic speciation. *PLoS Negl Trop Dis* 2014; 8: e2778.
  49. Cho J, Jung BK, Chang T, Sohn WM, Sinuon M, Chai JY. *Echinostoma mekongi* n. sp. (Digenea: Echinostomatidae) from riparian people along the Mekong River in Cambodia. *Korean J Parasitol* 2020; 58: 431-443.
  50. Lee SH, Sohn WM, Chai JY. *Echinostoma revolutum* and *Echinoparyphium recurvatum* recovered from house rats in Yangyang-gun, Kangwon-do. *Korean J Parasitol* 1990; 28: 235-240.
  51. Sohn WM, Chai JY, Yong TS, Eom KS, Yoon CH, Sinuon M, Socheat D, Lee SH. *Echinostoma revolutum* infection in children, Pursat Province, Cambodia. *Emerg Infect Dis* 2011; 17: 117-119.
  52. Saijuntha W, Tantrawatpan C, Sithithaworn P, Andrews RH, Petney TN. Genetic characterization of *Echinostoma revolutum* and *Echinoparyphium recurvatum* (Trematoda: Echinostomatidae) in Thailand and phylogenetic relationships with other isolates inferred by ITS1 sequence. *Parasitol Res* 2011; 108: 751-755.
  53. Saijuntha W, Tantrawatpan C, Sithithaworn P, Andrews RH, Petney TN. Spatial and temporal genetic variation of *Echinostoma revolutum* (Trematoda: Echinostomatidae) from Thailand and the Lao PDR. *Acta Trop* 2011; 118: 105-109.
  54. Saijuntha W, Sithithaworn P, Duengngai K, Kiatsopit N, Andrews RH, Petney TN. Genetic variation and relationships of four species of medically important echinostomes (Trematoda: Echinostomatidae) in South-East Asia. *Infect Genet Evol* 2011; 11: 375-381.
  55. Chai JY Sohn WM, Na BK, De NV. *Echinostoma revolutum*: metacercariae in *Filopaludina* snails from Nam Dinh Province, Vietnam, and adults from experimental hamsters. *Korean J Parasitol* 2011; 49: 449-455.
  56. Chai JY, Sohn WM, Yong TS, Eom KS, Min DY, Hoang EH, Phammasack B, Insisiengmay B, Rim HJ. Echinostome flukes recovered from humans in Khammouane Province, Lao PDR. *Korean J Parasitol* 2012; 50: 269-272.
  57. Chantima K, Chai JY, Wongsawad C. *Echinostoma revolutum*: freshwater snails as the second intermediate hosts in Chiang Mai Province, Thailand. *Korean J Parasitol* 2013; 51: 183-189.
  58. Lie KJ, Umathevy T. Studies on Echinostomatidae (Trematoda) in Malaya. VIII. The life history of *Echinostoma audyi* sp. n. *J Parasitol* 1965; 51: 781-788.
  59. Ando R, Ozaki Y. On four new species of trematodes of the family Echinostomatidae. *Dobutsugaku Zasshi* 1923; 35: 108-119 (in Japanese).
  60. Seo BS, Cho SY, Chai JY. Studies on intestinal trematodes in Korea I. A human case of *Echinostoma cinetorchis* infection with an epidemiological investigation. *Seoul J Med* 1980; 21: 21-29.
  61. Lee SH, Chai JY, Hong ST, Sohn WM. Experimental life history of *Echinostoma cinetorchis*. *Korean J Parasitol* 1990; 28: 39-44.
  62. Mohandas A. Studies on the life history of *Echinostoma ivanosii* n. sp. *J Helminthol* 1973; 47: 421-438.
  63. Sandground JH, Bonne C. *Echinostoma lindoensis* n. sp., a new parasite of man in the Celebes with an account of its life history and epidemiology. *Am J Trop Med Hyg* 1940; 20: 511-536.
  64. Chu JK, Cho YJ, Chung SB, Won BO, Yoon MB. Study on the trematode parasites of the birds in Korea. *Korean J Parasitol* 1973; 11: 70-75 (in Korean).
  65. Eom KS, Rim HJ, Jang DH. A study on the parasitic helminths of domestic duck (*Anas platyrhynchos* var. *domestica* Linnaeus) in Korea. *Korean J Parasitol* 1984; 22: 215-221.
  66. Anisuzzaman, Alim MA, Rahman MH, Mondal MMH. Helminth parasites in indigenous ducks: seasonal dynamics and effects on production performance. *J Bangladesh Agric Univ* 2005; 3: 283-290.
  67. Memon M, Birmani NA, Shaikh AM, Memon KH. New host record *Echinostoma paraulum* (Digenea: Echinostomatidae) in house crow (*Corvus splendens* Vieillot, 1817) of district Khairpur, Sindh, Pakistan. *J Entomol Zool Stud* 2018; 6: 586-589.
  68. Yamaguti S. Studies on the helminth fauna of Japan. Part 3. Avian trematodes, II. *Jpn J Zool* 1934; 5: 542-583.
  69. Kurisu Y. Studies on trematodes which take Japanese house hen as the definitive host. *Kumamoto Igakkai Zasshi* 1932; 8:

- 283-298 (in Japanese).
70. Islam MR, Shaikh H, Baki MA. Prevalence and pathology of helminth parasites in domestic ducks of Bangladesh. *Vet Parasitol* 1988; 29: 73-77.
  71. Rim HJ. Echinostomiasis. CRC Handbook Series in Zoonoses, Section C: Parasitic Zoonoses, Vol. III. Trematode Zoonoses. Boca Raton, USA. CRC Press Inc., 1982, pp 53-69.
  72. Nagataki M, Tantrawatpan C, Agatsuma T, Sugiura T, Duengai K, Sithithaworn P, Andrews RH, Petney TN, Saijuntha W. Mitochondrial DNA sequences of 37 collar-spined echinostomes (Digenea: Echinostomatidae) in Thailand and Lao PDR reveals presence of two species: *Echinostoma revolutum* and *E. miyagawai*. *Infect Genet Evol* 2015; 35: 56-62.
  73. Fu YT, Jin JC, Li F, Liu GH. Characterization of the complete mitochondrial genome of the echinostome *Echinostoma miyagawai* and phylogenetic implications. *Parasitol Res* 2019; 118: 3091-3097.
  74. Li Y, Qiu YY, Zeng MH, Diao PW, Chang QC, Gao Y, Zhang Y, Wang CR. The complete mitochondrial genome of *Echinostoma miyagawai*: comparisons with closely related species and phylogenetic implications. *Infect Genet Evol* 2019; 75: 103961.
  75. Anucherngchai S, Chontanarith T, Tejangkura T. The study of *cytochrome B* (CYTB): species specific detection and phylogenetic relationship of *Echinostoma revolutum* (Froelich, 1802). *J Parasit Dis* 2019; 43: 66-74.
  76. Rayski C, Fahmy MAM. Investigation on some trematodes of birds from the East Scotland. *Z Parazitenkd* 1962; 22: 186-195.
  77. Fagasinski A. Helminth parasites of domestic galliform birds in Poland. *Acta Parasitol Pol* 1962; 10: 347-368 (in Polish).
  78. Koltelnikov GA. The biology of *Echinostoma robustum* Yamaguti, 1935 causing disease in ducks. *Sbornik Nauchno-Tekhnich Inform Vsess Inst Ghelmintol KI Skrjabina* 1961; 7/8: 29 (in Russian).
  79. Alekseev VM. The life cycles of *Echinostoma revolutum* (Froelich, 1802) and *E. robustum* Yamaguti, 1935 in the Primorsk Region. *Soobshch dal'nevost. Fil. V.L. Komarova sib. Otdel. Akad. Nauk SSSR* 1968; 26: 3-7 (in Russian).
  80. Richard J. Trematodes d'Oiseaux de Madagascar (Note III). Espèces de la famille Echinostomatidae Poche 1926. *Ann Parasitol Hum Comp* 1964; 34: 607-620.
  81. Richard J, Brygoo ER. Life cycle of the trematode *Echinostoma caproni* Richard, 1964 (Echinostomatidae). *Ann Parasitol Hum Comp* 1978; 53: 265-275.
  82. Bisseru B. Stages in the development of larval echinostomes recovered from schistosome transmitting molluscs in Central Africa. *J Helminthol* 1967; 2: 89-108.
  83. Jeyarasasingam U, Heyneman D, Lim HK, Mansour N. Life cycle of a new echinostome from Egypt, *Echinostome liei* sp. nov. (Trematoda: Echinostomatidae). *Parasitology* 1972; 65: 203-222.
  84. Jourdane J, Kulo SD. Etude expérimentale du cycle biologique de *Echinostoma togoensis* n. sp., parasite à l'état larvaire de *Biomphalaria pfeifferi* au Togo. *Ann Parasitol (Paris)* 1981; 56: 477-488.
  85. Fried B, Huffman JE. The biology of the intestinal trematode *Echinostoma caproni*. *Adv Parasitol* 1996; 38: 311-368.
  86. Sorensen RE, Kanev I, Fried B, Minchella DJ. The occurrence and identification of *Echinostoma revolutum* from North American *Lymnaea elodes* snails. *J Parasitol* 1997; 83: 169-170.
  87. Sorensen RE, Curtis J, Minchella DJ. Intraspecific variation in the rDNA ITS loci of 37-collar-spined echinostomes from North America: implications for sequence-based diagnoses and phylogenetics. *J Parasitol* 1998; 84: 992-997.
  88. Lutz A. Untersuchung über die Entwicklungsgeschichte brasilianischer Trematoden. Spezieller Teil. I. Echinostomatidae. *Mem Inst Oswaldo Cruz* 1924; 17: 75-93.
  89. Maldonado A Jr, Loker ES, Morgan JAT, Rey L, Lanfredi RM. Description of the adult worms of a new Brazilian isolate of *Echinostoma paraensei* (Platyhelminthes: Digenea) from its natural vertebrate host *Nectomys squamipes* by light and scanning electron microscopy and molecular analysis. *Parasitol Res* 2001; 87: 840-848.
  90. Maldonado A Jr, Vieira GO, Garcia JS, Rey L, Lanfredi RM. Biological aspects of a new isolate of *Echinostoma paraensei* (Trematoda: Echinostomatidae): susceptibility of sympatric snails and the natural vertebrate host. *Parasitol Res* 2001; 87: 853-859.
  91. Brasil MD, Amato SB. Faunistic analysis of the helminths of sparrows (*Passer domesticus* L, 1758) captured in Campo Grande, Rio de Janeiro, R. J. *Mem Inst Oswaldo Cruz* 1992; 87: 43-48.
  92. Maldonado A Jr, Lanfredi RM. Echinostomes in the wild. In Fried B, Toledo R eds, *The Biology of Echinostomes*. New York, USA. Springer Science+Business Media, LLC. 2009, pp 129-145.
  93. Nicoll W. The trematode parasites of North Queensland. II. Parasites of birds. *Parasitology* 1914; 7: 105-126.
  94. Johnson J. The life cycle of *Echinostoma revolutum* (Froelich). *Univ California Public Zool* 1920; 19: 335-388.
  95. Mendheim H. Beiträge zur Systematik und Biologie der Familie Echinostomatidae (Trematoda). *Archiv für Naturgeschichte* 1943; 12: 175-302.
  96. Bashikirova EI. Family Echinostomatidae Dietz, 1909. In Skryabin KI ed, *Trematodes and Animals and Man*. Moscow, Russia. 1947; 1: 310-391.
  97. Yamaguti S. *Systema Helminthum*. Vol. I. The digenetic trematodes of vertebrates (Part I). Interscience Publishers Inc., New York, USA. 1958, pp 1-979.
  98. McDonald ME. Key to trematodes reported in waterfowl. US Department of the Interior. Fish and Wildlife Service, Resource Publication 142. Washington DC. 1981, pp. 1-156.
  99. Schuster R. *Echinostoma echinatum*, *Notocotylus noyeri* and *Quinqueserialis quinqueserialis* als seltene Parasiten von *Rattus norvegicus*. *Angew Parasitol* 1986; 27: 221-225 (in German).
  100. Christensen NØ, Fried B, Kanev I. Taxonomy of 37-collar spines *Echinostoma* (Trematoda: Echinostomatidae) in studies on the population regulation in experimental rodent hosts. *Angew Parasitol* 1990; 31: 127-130.

101. Anazawa K. On a human case of *Echinostoma revolutum* and its infection route. Taiwan Igakkai Zasshi 1929; no. 288: 221-241 (in Japanese).
102. Lu SC. Echinostomiasis in Taiwan. Int J Zoon 1982; 9: 33-38.
103. Bonne C, Bras G, Lie KJ. Five echinostomes in man in the Malayan Archipelago. Am J Dig Dis 1953; 20: 12-16.
104. Radomyos P, Radomyos B, Tungtrongchitr A. Multi-infection with helminths in adults from northeast Thailand as determined by post-treatment fecal examination of adult worms. Trop Med Parasitol 1994; 45: 133-135.
105. Morgan JA, Blair D. Relative merits of nuclear ribosomal internal transcribed spacers and mitochondrial CO1 and ND1 genes for distinguishing among *Echinostoma* species (Trematoda). Parasitology 1998; 116: 289-297.
106. Leidy J. On the trematodes of the muskrat. Proc Acad Nat Sci Phila 1888; 40: 126.
107. Stiles CW, Hassall A. A new species of intestinal fluke (*Distoma tricolor*) in the cotton-tail rabbit (*Lepus sylvaticus* Bachman) and in the northern hare (*L. americanus* Erxl.); *Distoma (Polyorchis) molle* (Leidy, 1856) Stiles & Hassall, 1894. Notes on parasites 29-30. Vet Magaz 1895; 1: 73.
108. Hassall A. Check list of animal parasites of chicken (*Gallus domesticus*). US Department of Agriculture. Bureau of Animal Industry, Circular 9. 1896, pp 7.
109. Saijuntha W, Sithithaworn P, Andrews RH. Genetic differentiation of *Echinostoma revolutum* and *Hypoderaeum conoideum* from domestic ducks in Thailand by multilocus enzyme electrophoresis. J Helminthol 2010; 84: 143-148.
110. Saijuntha W, Tapdara S, Tantrawatpan C. Multilocus enzyme electrophoresis analysis of *Echinostoma revolutum* and *Echinostoma malayanum* (Trematoda: Echinostomatidae) isolated from Khon Kaen Province, Thailand. Asian Pac J Trop Med 2010; 3: 633-636.
111. Buddhachat K, Chontanarath T. Is species identification of *Echinostoma revolutum* using mitochondrial DNA barcoding feasible with high-resolution melting analysis? Parasitol Res 2019; 118: 1799-1810.
112. Le TH, Pham LTK, Doan HTT, Le XTK, Saijuntha W, Rajapakse RPJ, Lawton SP. Comparative mitogenomics of the zoonotic parasite *Echinostoma revolutum* resolves taxonomic relationships within the '*E. revolutum*' species group and the Echinostomata (Platyhelminthes: Digenea). Parasitology 2020; 147: 566-576.
113. Tantrawatpan C, Saijuntha W. Multiplex PCR development for the differential detection of four medically important echinostomes (Trematoda: Echinostomatidae) in Thailand. Acta Trop 2019; 204: 105304.
114. Ishii N. Studies on bird trematodes. I. Bird trematodes in Japan & II. Four new bird trematodes. Jpn J Exp Med 1932; 11: 91-100.
115. Kosupko GA. Morphological peculiarity of cercariae of *Echinostoma revolutum* and *Echinostoma miyagawai*. Trudy vsesoyuznogo Instituta Helminthologii 1969; 15: 159-165 (in Russian).
116. Kosupko GA. Studies on the morphological and biological peculiarity of *Echinostoma revolutum* (von. Froelich, 1802) and *Echinostoma miyagawai* Ishii, 1932 (Trematoda: Echinostomatidae) on experimental material. PhD Thesis, Moscow, Russia. 1972, pp 1-258 (in Russian).
117. Lie KJ, Basch PF. Life history of *Echinostoma barbosai* sp. n (Trematoda: Echinostomatidae). J Parasitol 1966; 52: 1052-1057.
118. Mutafova T, Kanev I. Karyotype of an echinostome with features of *Echinostoma barbosai* Lie and Basch, 1966 (Trematoda: Echinostomatidae) found in Bulgaria. Khelminthologiya 1983; 16: 42-45.
119. Mutafova T, Kanev I. Karyotype and chromosome analysis of collar spined parasitic worms (Trematoda: Echinostomatidae). Abstract in the 5th Nat Conf Parasitol, Varna, Bulgaria. 1987, pp 185.
120. Fernández MV, Hamann MI, Núñez MO. Echinostome cercariae from *Biomphalaria straminea* (Mollusca: Planorbidae) in a ricefield from northeastern Argentina. Rev Mex Biodiver 2014; 85: 1024-1031.
121. Sianto L, Duarte AN, Borba VH, Magalhães JG, de Souza SM, Chame M. Echinostomes in felid coprolites from Brazil. J Parasitol 2016; 102: 385-387.
122. Nasincova V. The life cycle of *Echinostoma bolschevense* (Kotova, 1939) (Trematoda: Echinostomatidae). Folia Parasitol 1991; 38: 143-164.
123. Barus V, Moravec F, Rysavy B. Antagonistic interaction between *Echinostoma revolutum* and *Echinoparyphium recurvatum* (Trematoda) in the definitive host. Folia Parasitol (Praha) 1974; 21: 155-159.
124. Christensen NØ. *Echinostoma revolutum*: labelling of miracidia with radioselenium in vivo and assay for host finding. Exp Parasitol 1980; 50: 67-73.
125. Christensen NØ, Fagbemi BO, Nansen P. Trypanosoma brucei-induced blockage of expulsion of *Echinostoma revolutum* and of homologous *E. revolutum* resistance in mice. J Parasitol 1984; 70: 558-561.
126. Christensen NØ, Knudsen J, Fagbemi BO, Nansen P. Impairment of primary expulsion of *Echinostoma revolutum* in mice concurrently infected with *Schistosoma mansoni*. J Helminthol 1985; 59: 333-335.
127. Christensen NØ, Knudsen J, Andreassen J. *Echinostoma revolutum*: resistance to secondary and superimposed infection in mice. Exp Parasitol 1986; 61: 311-318.
128. Christensen NØ, Odaibo AB, Simonsen PE. *Echinostoma* population regulation in experimental rodent definitive hosts. Parasitol Res 1988; 75: 83-87.
129. Bindseil E, Christensen NØ. Thymus-independent crypt hyperplasia and villous atrophy in the small intestine of mice infected with the trematode *Echinostoma revolutum*. Parasitology 1984; 88: 431-438.
130. Simonsen PE, Andersen BJ. *Echinostoma revolutum* in mice: Dynamics of the antibody attack to the surface of an intestinal trematode. Int J Parasitol 1986; 16: 475-482.
131. Sutton CA, Lunaschi L. Contribution to the knowledge of the

- parasite fauna of Argentina. VIII. A new digenean in *Chloephaga picta melanoptera* Gmelin. *Neotropica* 1980; 26: 13-17 (in Portuguese).
132. Dollfus PR. Distomens parasites de Muridae du genre *Mus*. *Ann Parasitol (France)* 1925; 3: 85-102 (in French).
  133. Sugimoto M. On a trematode parasite (*Echinostoma cinetorchis* Ando and Ozaki, 1923) from a Formosan dog. *Nippon Juigak-kai Zasshi* 1933; 12: 231-236 (in Japanese).
  134. Tanabe K, Takeishi H. A survey of helminths in the digestive system of the rat. *Keio Igakku* 1936; 16: 1767-1785 (in Japanese).
  135. Seo BS, Rim HJ, Lee CW. Studies on the parasitic helminths of Korea. I. Trematodes of rodents. *Korean J Parasitol* 1964; 2: 20-26.
  136. Seo BS, Cho SY, Hong ST, Hong SJ, Lee SH. Studies on parasitic helminths of Korea. V. Survey on intestinal trematodes of house rats. *Korean J Parasitol* 1981; 19: 131-136.
  137. Cho SY, Kang SY, Ryang YS. Helminthes infections in the small intestine of stray dogs in Ejungbu City, Kyunggi Do, Korea. *Korean J Parasitol* 1981; 19: 55-59 (in Korean).
  138. Takahashi S, Ishii T, Ueno N. A human case of *Echinostoma cinetorchis*. *Tokyo Iji Shinshi* 1930; 2757: 141-144 (in Japanese).
  139. Takahashi S, Ishii T, Ueno N. The second human case of *Echinostoma cinetorchis* and a case of tapeworm in a man. *Tokyo Iji Shinshi* 1930; 2678: 1326-1327 (in Japanese).
  140. Kawahara S, Yamamoto E. Human cases of *Echinostoma cinetorchis*. *Tokyo Iji Shinshi* 1933; 2840: 1794-1796 (in Japanese).
  141. Ryang YS, Ahn YK, Kim WT, Shin KC, Lee KW, Kim TS. Two cases of human infection by *Echinostoma cinetorchis*. *Korean J Parasitol* 1986; 24: 71-76 (in Korean).
  142. Lee SK, Chung NS, Ko IH, Ko HI, Sohn WM. A case of natural human infection by *Echinostoma cinetorchis*. *Korean J Parasitol* 1988; 26: 61-64 (in Korean).
  143. Son WY, Huh S, Lee SU, Woo HC, Hong SJ. Intestinal trematode infections in the villagers in Koje-myon, Kochang-gun, Kyongsangnam-do, Korea. *Korean J Parasitol* 1994; 32: 149-155.
  144. Jung WT, Lee KJ, Kim HJ, Kim TH, Na BK, Sohn WM. A case of *Echinostoma cinetorchis* (Trematoda: Echinostomatidae) infection diagnosed by colonoscopy. *Korean J Parasitol* 2014; 52: 287-290.
  145. Chai JY, Lee SH. Food-borne intestinal trematode infections in the Republic of Korea. *Parasitol Int* 2002; 51: 129-154.
  146. Chai JY, Shin EH, Lee SH, Rim HJ. Foodborne intestinal flukes in Southeast Asia. *Korean J Parasitol* 2009; 47 (suppl): 69-102.
  147. Anh NTL, Madsen H, Dalsgaard A, Phuong NT, Thanh DTH, Murrell KD. Poultry as reservoir hosts for fishborne zoonotic trematodes in Vietnamese fish farms. *Vet Parasitol* 2010; 169: 391-394.
  148. Toledo R, Esteban JG. An update on human echinostomiasis. *Trans R Soc Trop Med Hyg* 2016; 110: 37-45.
  149. Toledo R, Muñoz-Antoli C, Esteban JG. Intestinal trematode infections. *Digenetic Trematodes. Adv Exp Med Biol* 2014; 766: 201-240.
  150. Toledo R, Radev V, Kanev I, Gardner SL, Fried B. History of echinostomes (Trematoda). *Acta Parasitol* 2014; 59: 555-567.
  151. Kostadinova A. *Echinostoma echinatum* (Zeder, 1803) *sensu* Kanev (Digenea: Echinostomatidae): a note of caution. *Syst Parasitol* 1995; 32: 23-26.
  152. Lie KJ. Further studies on the life history of *Echinostoma lindoense* Sandground and Bonne, 1940 (Trematoda: Echinostomatidae) with a report on its occurrence in Brazil. *Proc Helminthol Soc Wash* 1968; 35: 74-77.
  153. Vasilev I, Kanev I, Shvetlikovski M, Bushta I. Establishment of an echinostome with the features of *Echinostoma lindoense* Sandground et Bonne (1940) (Echinostomatidae: Trematoda) in Poland and Czechoslovakia. *Khelminthologia* 1982; 13: 12-21 (in Bulgarian).
  154. Feliu C, Gracenea M, Montoliu I, Torres I. On the finding of *Echinostoma lindoense* Sandground and Bone, 1940 (Trematoda: Echinostomatidae) in *Mus musculus* Linnaeus, 1758 (Rodentia; Muridae) of the Ebro Delta (NE of the Iberian Peninsula). *Rev Iber Parasitol* 1987; 47: 125-126 (in Spanish).
  155. Lie KJ. Studies on Echinostomatidae (Trematoda) in Malaya. VII. The life history of *Echinostoma lindoense* Sandground and Bonne, 1940. *Trop Geogr Med* 1964; 16: 72-81.
  156. Lie KJ, Nasemary S, Impand P. Five echinostome species from Thailand. *Southeast Asian J Trop Med Public Health* 1973; 4: 96-101.
  157. Eduardo SL, Lee GQ. Some zoonotic trematodes from the Philippine field rat, *Rattus mindanensis mindanensis* (Mearns, 1905) (Mammalia: Rodentia) in Bay, Laguna, Philippines with description and new records of species. *Phil J Vet Med* 2006; 43: 33-45.
  158. Lie KJ, Kanev I. Identification and distribution of *Echinostoma lindoense*, *E. audyi* and *E. revolutum* (Trematoda: Echinostomatidae). *Z Parasitenkd* 1983; 69: 223-227.
  159. Simões RO, Souza JGR, Maldonado Jr A, Luque JL. Variation in the helminth community structure of three sympatric sigmodontine rodents from the coastal Atlantic Forest of Rio de Janeiro, Brazil. *J Helminthol* 2011; 85: 171-178.
  160. Sianto L, Reinhard KJ, Chame M, Chaves S, Mendonça S, Gonçalves MLC, Fernandes A, Ferreira LE, Araújo A. The finding of *Echinostoma* (Trematoda: Digenea) and hookworm eggs in coprolites collected from a Brazilian mummified body dated 600-1,200 years before present. *J Parasitol* 2005; 91: 972-975.
  161. Cooper CL. The helminth parasites of an insular avian passerine community: relationship to landscape epidemiology. Ph.D. Dissertation to Ohio State University, USA. 1975, pp 1-209.
  162. Kostadinova A. Cercarial chaetotaxy of *Echinostoma miyagawai* Ishii, 1932 (Digenea: Echinostomatidae), with a review of the sensory patterns in the 'revolutum' group. *Syst Parasitol* 1999; 44: 201-209.
  163. Uchida A, Uchida K, Kawakami Y, Nagatomo M, Shen M, Ooi HK. A helminthological survey of parasites in the waterfowl of Kanagawa prefecture. *J Jpn Vet Med Assoc* 2005; 58: 127-131 (in Japanese).

164. Haziev GZ, Khan SA. Helminths of guinea fowl (*Numida meleagris*) in Bashkir ASSR. *Vet Parasitol* 1991; 38: 349-353.
165. Schou TW, Permin A, Juul-Madsen HR, Sørensen P, Labouriau R, Nguyễn TLH, Fink M, Pham SL. Gastrointestinal helminths in indigenous and exotic chickens in Vietnam: association of the intensity of infection with the major histocompatibility complex. *Parasitology* 2007; 134: 561-573.
166. Faltýnková A, Našincová V, Kablásková L. Larval trematodes (Digenea) of planorbid snails (Gastropoda: Pulmonata) in Central Europe: a survey of species and key to their identification. *Syst Parasitol* 2008; 69: 155-178.
167. Kavetska KM, Rząd I, Kornyshev VV, Korol EN, Sitko J, Szatancka K. Enteric helminths of the mallard *Anas platyrhynchos* L., 1758 in the north-western part of Poland. *Wiadom Parazitol* 2008; 54: 23-29 (in Polish).
168. Nasincova V. Contribution to the distribution and the life history of *Echinostoma revolutum* (Trematoda) in Central Europe. *Vestnik Ceskoslovenske Spolecnosti Zool* 1986; 50: 70-80.
169. Schwelm J, Soldánová M, Vyhliadalová T, Sures B, Selbach C. Small but diverse: larval trematode communities in the small freshwater planorbids *Gyraulus albus* and *Segmentina nitida* (Gastropoda: Pulmonata) from the Ruhr River, Germany. *Parasitol Res* 2017; 117: 241-255.
170. McKenna PB. Additions to the checklists of helminth and protozoan parasites of terrestrial mammals and birds in New Zealand. *NZ J Zool* 2018; 45: 395-401.
171. Nasir P. Studies on the life history of *Echinostoma nudicaudatum* n. sp. (Echinostomatidae: Trematoda). *J Parasitol* 1960; 46: 833-847.
172. Nasir P. Further observations on the life cycle of *Echinostoma nudicaudatum* Nasir, 1960 (Echinostomatidae: Trematoda). *Proc Helminthol Soc Wash* 1962; 29: 115-127.
173. O'Dwyer K, Poulin R. Taken to the limit-Is desiccation stress causing precocious encystment of trematode parasites in snails? *Parasitol Int* 2015; 64: 632-637.
174. Nasir P. Observations on the life cycle of *Echinostoma pinnicaudatum* n. sp. (Echinostomatidae: Trematoda). *Proc Helminthol Soc Wash* 1961; 28: 207-212.
175. Nasir P, Erasmus DA. A key to the cercariae from British freshwater molluscs. *J Helminthol* 1964; 38: 245-268.
176. Lie KJ, Basch PF. The life history of *Echinostoma paraensei* sp. n (Trematoda: Echinostomatidae). *J Parasitol* 1967; 53: 1192-1199.
177. Petrie JL, Burg EF III, Cain GD. Molecular characterization of *Echinostoma caproni* and *E. paraensei* by random amplification of polymorphic DNA (RAPD) analysis. *J Parasitol* 1996; 82: 360-362.
178. Meece JK, Nollen PM. A comparison of the adult and miracidial stages of *Echinostoma paraensei* and *E. caproni*. *Int J Parasitol* 1996; 26: 37-43.
179. Fried B, Reddy A. Comparative studies on excystation and morphological features of the metacercariae of *Echinostoma paraensei* and *E. caproni*. *Int J Parasitol* 1997; 27: 899-901.
180. Miller MJ. The parasites of pigeons in Canada. *Can J Res* 1937; 15: 91-103.
181. Skrjabin KI. *Echinostoma paraulium*-nouveau parasite de l'homme. *Med Parazitol Parazitarn Bolezni* 1938; 7: 129-138 (in Russian).
182. Supperer R. Untersuchungen über Parasiten der Hausente, *Anas platyrhynchos* Dom. *Z Parasitenkd* 1959; 19: 259-277 (in German).
183. Skrjabin KI. Trematodes dans les oiseaux d'Oural. *Ann Mus Zool Acad Sci St.-Petersburg*, 1915; 20: 395-417 (in Russian).
184. Sprehn C. Echinostomiden bei Tauben. *Deutsche tierärztl. Wochenschr* 1927; 25: 451-455 (in German).
185. Iskova NI. Trematoda. Vol. 4. Echinostomes. Kiev: Naukova Dumka. Fauna Ukrainy 1985; 34: 1-198 (in Russian).
186. Uchida A, Uchida K, Itagaki H, Kamegai S. Checklist of helminth parasites of Japanese birds. *Jpn J Parasitol* 1991; 40: 7-85.
187. Yamaguti S. Studies on the helminth fauna of Japan. Part 5. Trematodes of birds, III. *Jpn J Zool* 1935; 6: 159-182.
188. Yang G, Di S. Reviews on the trematodes of poultry in China. *J Sichuan Agricul Univ.* 1999; 17: 85-89 (in Chinese).
189. Lunaschi L, Drago FB. Checklist of digenean parasites of wild mammals from Argentina. *Zootaxa* 2007; 1580: 35-50.
190. Lunaschi L, Drago FB, Núñez V. Two new species of *Echinostoma* (Digenea: Echinostomatidae) from Argentinean birds. *Rev Mex Biodivers* 2018; 89: 356-364 (in Spanish).
191. Pinto HA, Melo AL. *Physa marmorata* (Mollusca: Physidae) as intermediate host of *Echinostoma exile* (Trematoda: Echinostomatidae) in Brazil. *Neotrop Helminthol* 2012; 6: 291-299 (in Portuguese).
192. Cort WW. Larval trematodes from North American fresh-water snails. *J Parasitol* 1914; 1: 65-84.
193. Cort WW. Some North American larval trematodes. *Illin Biol Monogr* 1915; 1: 447-532.
194. Barker FD. Parasites of the American muskrat (*Fiber zibethicus*). *J Parasitol* 1915; 1: 184-197.
195. Law RG, Kennedy AH. Parasites of fur-bearing animals. *Bull Dept Game Fisher, Ontario, Canada* 1932; 4: 1-30.
196. Miller EL. Studies on North American cercariae. *Illin Biol Monogr* 1936; 14: 1-125.
197. Dimitrov V, Kanev I, Fried B, Radev V. Sensilla of the cercariae of *Echinostoma trivolvis* (Cort, 1914) (Trematoda: Echinostomatidae). *Parasite* 1997; 2: 153-158.

