



Article

Age-Dependent Differences in Frequent Mental Distress (FMD) of US Older Adults Living in Multigenerational Families versus Living Alone

Debasree Das Gupta ^{1,*} and David W. S. Wong ²

¹ Department of Kinesiology and Health Science, Emma Eccles Jones College of Education and Human Services, Utah State University, Logan, UT 84322, USA

² Department of Geography and Geoinformation Science, George Mason University, Fairfax, VA 22030, USA

* Correspondence: debasree.dasgupta@usu.edu

Abstract: Frequent mental distress (FMD) is prevalent among older Americans, but less is known about the disparities in the FMD of older adults living in multigenerational families versus living alone. We pooled cross-sectional data (unweighted, $n = 126,144$) from the Behavioral Risk Factor Surveillance System (BRFSS) between 2016 and 2020 and compared the FMD (≥ 14 poor mental health days in the past 30 days = 1; 0 otherwise) of older adults (≥ 65 years) living in multigenerational families versus living alone in 36 states. After controlling for covariates, the findings indicate 23% lower odds of FMD among older adults living in multigenerational households compared to counterparts living alone (adjusted odds ratio (AOR): 0.77; 95% confidence interval (CI): 0.60, 0.99). The findings also show that the reduction in the odds of FMD with each 5 year increase in age was larger among older adults living in multigenerational families by 18% (AOR: 0.56; 95% CI: 0.46, 0.70) compared to older adults living alone (AOR: 0.74; 95% CI: 0.71, 0.77), and this difference was significant at the 5% significance level. Multigenerational living may have a protective association with FMD among older adults. Further research is needed to identify multigenerational family and non-kin factors that translate into mental health advantages for older adults.



Citation: Das Gupta, D.; Wong, D.W.S. Age-Dependent Differences in Frequent Mental Distress (FMD) of US Older Adults Living in Multigenerational Families versus Living Alone. *Int. J. Environ. Res. Public Health* **2023**, *20*, 3747. <https://doi.org/10.3390/ijerph20043747>

Academic Editor: Paul B. Tchounwou

Received: 31 October 2022
Revised: 10 February 2023
Accepted: 15 February 2023
Published: 20 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: older adults; multigenerational; living alone; mental health; frequent mental distress (FMD)

1. Introduction

Poor mental health is widely prevalent among older adults and is linked to high rates of morbidity and mortality in this group [1–6]. Although empirical estimates vary, approximately a quarter to one-third of older Americans report some type of poor mental health condition, which is often closely associated with psychosocial factors, such as social support, loneliness, and isolation [1,7–10]. With the progressive aging of the US population, improving mental health among older adults is a critical public health priority [2,11,12].

Poor mental health comprises an array of behavioral conditions that include, for instance, anxiety, depression, stress, and/or distress [13]. Among these mental health conditions, frequent mental distress (FMD) is one specific measure that is tracked locally and nationally as part of the Centers for Disease Control and Prevention’s (CDC) surveillance of health-related quality of life (HRQOL) of US adults [14,15]. FMD—defined as “14 or more days of self-reported poor mental health in the past month”—strongly correlates with medical markers of depression and represents the burden of mental illness in population subgroups [14–16]. Among older adults, FMD is more prevalent than clinically diagnosed mental disorders, such as anxiety and depression, which largely remain under-diagnosed and under- or unreported in this group [1,2].

Thus, estimating FMD among older adults may be a particularly useful approach to capture the larger proportion of those 65 years and older suffering from mental health

disorders. However, among a diverse swath of studies examining the prevalence and risk factors of FMD among population subgroups [4,6,7,13,17–26], only a small fraction focuses on the FMD of older adults [6,7,22,25]. In this latter group, no previous study has examined the prevalence of FMD or age-associated differences of this condition among older adults living in multigenerational family households. This line of inquiry assumes significance for more than one reason. While the coresidence of older adults with adult children and/or grandchildren used to be the historical norm, a progressive decline in multigenerational living was set off and persisted after the first demographic transition of the US [27,28]. However, recent studies reveal an increasing trajectory in multigenerational living with one in five US older adults living in multigenerational family households [27,29–31]. The rising share of multigenerational living of the US older adult population places the spotlight on this demographic group [27,29].

The broader literature conceptualizing multigenerational living is currently divided between those that forward benefits versus those that highlight the disadvantages for the mental health of older adults. On the one hand, prior frameworks have emphasized and reported on the salience of multigenerational living for human development and mental well-being over the life span and in healthy aging [32–35]. Along the same line, age segregation is identified as a factor that limits and hinders mental health in older ages [36,37]. Collectively, these findings suggest that multigenerational living may have the potential to benefit mental health, particularly in older ages. However, on the other hand, challenges, including caregiving stress, intergenerational conflicts and strains, and detriments to the mental health of older adults from multigenerational living, are also highlighted in the literature [28,38–46]. Irrespective of this conceptual divide, the empirical evidence on the prevalence of FMD among older adults living in multigenerational families is nonexistent.

The literature also indicates that the subjective mental health of older adults may be different at different ages. Previous evidence identified resiliency and better mental health in later life compared to younger ages and underscored that the adaptive benefits from better perceived mental health in older ages have a role in healthy aging [2,47–49]. Indeed, perceived mental health may have a greater role to play as older adults adjust to and negotiate many later-life changes, including mounting physical health deficits and functional limitations, as well as shrinking occupational roles and geographic spheres of activities [6,50,51]. A prior investigation [6] examining FMD among older adults highlighted age-associated disparities and reported decreased odds of FMD with increasing age. However, the understanding of whether the prevalence and patterns of FMD among older adults in multigenerational families may (or may not) also exhibit advantageous age-associated differences is not known. Lastly, on the other end of the spectrum, the status of living alone has been associated with social isolation, as well as with various adverse health conditions, including poor mental health, in later life [1,11,36,37,51].

Therefore, our objective was to evaluate whether multigenerational living has a protective (or a risk) association with FMD among older adults. In so doing, we first estimated whether the likelihood of FMD was lower among older adults living in multigenerational families compared to older adults living alone. Additionally, we also compared and contrasted the age-associated disparities in FMD among older adults living in multigenerational family households against older adults living alone.

2. Materials and Methods

Data: We used data from BRFSS, which is a leading source of nationally representative health data in the US [52,53]. The CDC, together with state health departments, has conducted this annual cross-sectional survey since 1984 in all 50 states, the District of Columbia, Guam, and Puerto Rico [52]. Trained interviewers administer the BRFSS survey via telephone to participants randomly selected from noninstitutionalized adults 18 years and older. A standardized questionnaire is used to track chronic conditions and health risk behaviors, access to and use of health services, and socioeconomics and demographics of

noninstitutionalized adults. The BRFSS employs a probability-based multistage cluster sampling methodology to generate representative samples in all 50 US states and regions. Additional details on this complex survey design methodology are available from the CDC [54]. Each year, approximately 400,000 adults participate in the BRFSS, and between 2016 and 2020, the mean response rate was 47.8%. The institutional review board (IRB) at the CDC has approved the BRFSS for research [55]. The data made available to the public are not individually identifiable, and a secondary analysis of this data does not constitute human subject research.

Study sample: The study sample included BRFSS landline participants 65 years and older who were living alone or were living in multigenerational families (unweighted count, $n = 172,193$; weighted count, $N = 36,773,297$) in 36 states across the US. The respondents participating via landline were asked the question: “Excluding adults living away from home, such as students away at college, how many members of your household, including yourself, are 18 years of age or older?” A second BRFSS question asked participants: “How many children less than 18 years of age live in your household?” as part of the Random Child Selection module that was administered in 36 states between 2016 and 2020. We identified all landline participants 65 years and older who responded “adults = 1” and “children = 0” to these two questions as older adults “living alone” (93.9% of the total study sample, $n = 167,272$; $N = 34,546,148$) in our analysis. The status of “living alone” has been used in prior studies, including those that use BRFSS data, as an objective indicator to capture the risk of social isolation, lower social support, and poor mental health consequences [50,51,56–63].

Next, to identify multigenerational family households, we applied a framework employed in prior studies [30,43,64–66]. Accordingly, we designated “multigenerational” as grandparent–grandchild family structures with or without the middle generation parents present in the household. If a participant reported one or more children (≤ 17 years) living in the household, the BRFSS asked the respondents: “How are you related to the child?”, and “grandparent” was one of the response options. All BRFSS landline respondents 65 years and older who indicated “grandparent” to this question constituted older adults living in multigenerational family households (6.1% of the total study sample, $n = 4921$; $N = 2,227,149$) in our study. In prior studies, the BRFSS “relationship” question has been used to identify grandparent–grandchild multigenerational family structures and to investigate older adult health [62,67–69].

Of the 172,193 older adult respondents living alone or living in multigenerational families, 73.3% ($n = 126,144$; $N = 26,446,545$) had no missing information on the mental health outcome variable and the covariates and comprised the analytic sample in our study. The composition of this analytic sample was (1) multigenerational, 6.5% ($n = 3898$; $N = 1,722,633$), and (2) living alone, 93.5% ($n = 122,246$; $N = 24,723,912$). This composition was comparable to the proportion of older adults living alone versus living in multigenerational families among excluded participants.

Outcome and explanatory variables: The outcome variable in our analysis was FMD, which was constructed using the “poor mental health days” question included under the BRFSS 4-item Healthy Days Core Module (CDC HRQOL-4) developed by the CDC [70]. Since 1993, the BRFSS has administered this module to ask respondents questions representing the four domains of overall, physical, mental, and social health [70]. The CDC HRQOL-4 questions are established valid measures, with multiple studies attesting to their strong psychometric properties and high test–retest reliability [71–76]. The longstanding validity of these measures is also documented in older adult populations, and, in congruence, these questions have been incorporated in the Medicare Health Outcomes Survey since 1995 [77–80].

The “poor mental health days” question asks respondents: “Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?” The reference time period of “past 30 days” was included to capture the emerging chronic symptoms impacting HRQOL, and a 14 day cut-point was employed to dichotomize this numeric

measure (range: 0 to 30 days) to represent FMD [70,75,81,82]. The 14 day threshold closely mirrors the duration used in clinical research and practice to indicate and diagnose depressive and anxiety disorders [13,71,75,81–84]. Previous studies on poor mental health have adopted this binary conceptualization as a single-unit measure to capture frequent and persistent deficits in mental HRQOL across various demographics, and applications of FMD are also found in state- and county-level population health surveillance studies [4,6,7,13,17–23,25,26,85–89]. We adopted this 14 day dichotomous metric to construct the outcome variable representing FMD (≥ 14 poor mental health days = 1; 0 otherwise) in our analysis.

The status of living in multigenerational families (=1) versus living alone (=0) served as the primary explanatory variable in our study. We also adopted an alternative coding scheme motivated upon findings reported in prior studies [62,67–69,90,91]. Previous results have highlighted the unique health vulnerabilities of single or solo grandparents living with their grandchild only [62,67–69,90,91]. Thus, we created a secondary explanatory variable consisting of the three categories of older adults living alone (=0), living with their grandchildren only (=1), and living with their grandchildren, spouse, and/or other adults (=2). We used responses to the following BRFSS questions: (1) number of adults 18 years or older in the household = 1, and children less than 18 years of age in the household = 0, coded with a (0) = living alone; (2) number of adults 18 years or older in the household = 1, children less than 18 years of age in the household $\neq 0$, and the relationship to child = “grandparent” coded with a (1) = living with grandchild only; and (3) number of adults 18 years or older in the household > 1, children less than 18 years of age in the household $\neq 0$, and the relationship to child = “grandparent” coded with a (2) = living with grandchild, spouse, and/or other adults. The BRFSS questions we used to construct this outcome and the explanatory variables are indicated in Table 1 below.

Table 1. Details on the outcome and explanatory variables, and the BRFSS questions used to construct these variables.

Variable Name	Variable Description	Question(s) in BRFSS Survey
Outcome Frequent mental distress (FMD)	FMD: ≥ 14 poor mental health days = 1; 0 otherwise	Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?
Explanatory		
Primary: multi-generational (=1) living versus living alone (=0)	Multigenerational: (relationship to child = “grandparent”) = 1	How are you related to the child?
	Living alone: (adults = 1 and children = 0) = 0	Excluding adults living away from home, such as students away at college, how many members of your household, including yourself, are 18 years of age or older?
Secondary: living alone (=0) versus living with grandchildren only (=1) versus living with grandchildren, spouse, and/or other adults (=2)	Living alone: (adults = 1 and children = 0) = 0	
	Living with grandchildren only (adults = 1 and children $\neq 0$ and relationship to child = “grandparent”) = 1	How many children less than 18 years of age live in your household?
	Living with grandchildren, spouse, and/or other adults (adults > 1 and children $\neq 0$ and relationship to child = “grandparent”) = 2	

Covariates: Demographics (age, sex, marital status, race, and ethnicity) and socioeconomics (education and income) are significant correlates of individual health including perceived mental and physical health [92]. To adjust for demographic and socioeconomic variations in FMD, we included age (5 year age categories), sex (male or female), marital status (married, unmarried couple, versus divorce, widowed, separated, and never married), race/ethnicity (White non-Hispanic (NH), Black NH, other NH, and Hispanic), education

(<high school, high school, attended college or technical school, and \geq college or technical school), and household income (<\$15,000, \$15,000–24,999, \$25,000–34,999, \$35,000–49,999, and \geq \$50,000) as controls in our analysis.

In addition, chronic conditions, as well as functional, cognitive, and sensory limitations, are reported as significant independent predictors of self-reported mental health status [92]. We combined binary (0 = no/1 = yes) responses to all of the BRFSS questions on lifetime diagnosis of chronic conditions that included depressive disorders; diabetes; cardiovascular diseases: coronary heart disease, myocardial infarction, or stroke; arthritis; chronic kidney disease; asthma; chronic obstructive pulmonary disease; and skin or other cancers to obtain an aggregate number of the chronic health conditions (range: 0–7) of the participants. Based on the US Census's [93] disability module, the BRFSS includes a six-item questionnaire to ask about functional, cognitive, and sensory limitations. These six questions ask respondents about (1) deafness or having serious difficulty hearing; (2) blindness or having serious difficulty seeing; (3) having serious difficulty concentrating, remembering, or making decisions; (4) having serious difficulty walking or climbing stairs; (5) having difficulty bathing or dressing; and (6) having difficulty doing errands alone, such as visiting a doctor's office or shopping. We aggregated the binary (no = 0/yes = 1) responses to these six questions in the BRFSS to derive an aggregate number (range: 0–6) representing the "number" of the physical limitations of the participants. Lastly, we included a year dummy variable (2016–2019 = 0 versus 2020 = 1) to adjust for any potential differences in mental health perceptions during the pandemic year.

Statistical Analysis: Based on descriptive statistics (percentages and standard errors), we examined variations in the FMD reported by older adults. Consistent with the methodology used in prior studies, we present the self-reported prevalence of FMD, which is the weighted percentage of respondents indicating FMD [1,4,6]. To control for the influence of confounding variables, we conducted regression analysis in which we considered all of the covariates mentioned above. We estimated the logistic regression specification (Equations (1) and (2) below) to model the binary outcome variable—FMD—employing Stata (version 15, Statacorp LLC, College Station, TX, USA) and conducted all statistical tests using the appropriate methodology for survey data analysis [94]. The findings presented in Section 3 (Results) are the weighted estimates with the statistical significance set at $p \leq 0.05$ (unless otherwise specified).

$$\log\left(\frac{\Pi}{1-\Pi}\right) = \beta_0 + \beta_1 X_1 + \dots + \beta_k X_k \quad (1)$$

which states that the (natural) logarithm of the odds of an event, referred to as the log odds, is a linear function of the X variables, such that:

$$\theta = \frac{\left(\frac{\Pi}{(1-\Pi)}\right)\Big|_{X = X_1}}{\left(\frac{\Pi}{(1-\Pi)}\right)\Big|_{X = X_2}} \quad (2)$$

where θ is the odds ratio between the odds for two sets of predictors; $X_{(1)}$ and $X_{(2)}$, $\left(\frac{\Pi}{1-\Pi}\right)$ are the odds of an event such that Π is the probability of the event, and X_1, \dots, X_k is the set of predictor variables.

3. Results

Descriptive results: Table 2 shows the descriptive characteristics of the two groups of older adults—those living alone versus those living in multigenerational families. Compared to older adults living alone, a higher proportion of older adults in multigenerational families were males (34% versus 31%), younger in age (65–69: 38% versus 19%), and belonged to minority racial ethnic groups (37% versus 19%). Older adults in multigenerational families had lower levels of education (<high school: 19% versus 15%) and lower levels of

physical disability (≥ 1 disability: 47% versus 50%) but comparable levels of chronic health conditions (16% no chronic conditions) when compared to counterparts living alone.

Table 2. Descriptive characteristics of older adults living alone versus living in multigenerational families, BRFSS 2016–2020.

	Living Alone		Living in Multigenerational Families	
	n = 122,246; N = 24,723,912		n = 3898; N = 1,722,633	
	%	[95% CI]	%	[95% CI]
Sex				
Male	30.7	[30.0, 31.3]	33.7	[30.2, 37.2]
Female	69.3	[68.7, 70.0]	66.4	[62.8, 69.7]
Age groups				
65–69	19.4	[18.8, 19.9]	38.3	[35.0, 41.8]
70–74	21.8	[21.2, 22.3]	29.4	[26.4, 32.7]
75–79	21.0	[20.5, 21.6]	19.5	[16.0, 23.5]
80 and above	37.9	[37.2, 38.6]	12.8	[10.3, 15.8]
Marital Status				
Married/unmarried couple	6.2	[5.9, 6.5]	54.1	[50.3, 57.8]
Divorced/widowed/separated/ never married	93.8	[93.5, 94.1]	45.9	[42.2, 49.7]
Race/ethnicity				
Non-Hispanic (NH) White	81.2	[80.5, 81.8]	62.8	[58.8, 66.7]
NH Black	9.6	[9.1, 10.0]	17.6	[14.6, 21.1]
NH other	3.8	[3.3, 4.2]	5.8	[4.5, 7.6]
Hispanic	5.5	[5.1, 5.9]	13.8	[10.6, 17.6]
Education				
<High school	14.9	[14.4, 15.5]	19.4	[16.0, 23.2]
High school	33.9	[33.2, 34.5]	30.4	[27.3, 33.8]
Attended college or technical school	31.0	[30.3, 31.6]	34.2	[30.7, 37.9]
\geq College or technical school	20.2	[19.8, 20.7]	16.1	[14.1, 18.2]
Income				
<\$15,000	17.4	[16.8, 18.0]	11.9	[8.9, 15.8]
\$15,000–24,999	30.6	[30.0, 31.3]	23.4	[20.2, 26.8]
\$25,000–34,999	16.8	[16.3, 17.3]	15.4	[12.8, 18.3]
\$35,000–49,999	15.4	[14.8, 15.9]	15.8	[13.7, 18.1]
\geq \$50,000	19.9	[19.4, 20.4]	33.6	[30.5, 36.9]
Chronic conditions				
No	16.8	[16.3, 17.4]	16.2	[13.7, 19.0]
Yes	83.2	[82.6, 83.8]	83.8	[81.0, 86.3]
Physical limitations				
No	50.6	[49.9, 51.2]	53.1	[49.4, 56.8]
Yes	49.5	[48.8, 50.1]	46.9	[43.2, 50.6]

n, Unweighted observation count; N, weighted observation count; CI, confidence interval; %, weighted %.

In Table 3, we provide the self-reported prevalence (weighted %) of FMD among older adults living alone versus living in multigenerational families. The results in Table 3 indicate that the FMD reported by the two groups of older adults were comparable—approximately 9% in each group reported FMD. However, a comparison across 5 year age categories indicated that the FMD reported by older adults living in multigenerational families was lower than those living alone for all age groups, except the 70–74 year old group. This age-associated difference between the FMD of older adults living alone versus living in multigenerational families was statistically significant for the 75 years and older age groups ($p < 0.05$).

Table 3. Self-reported prevalence of frequent mental distress (FMD) among older adults living alone versus living in multigenerational families, BRFSS 2016–2020.

	Living Alone		Living in Multigenerational Families		
	% ¹	[95% CI]	% ¹	[95% CI]	
65 and older	9.1	[8.8, 9.5]	9.0	[7.6, 10.6]	
Age groups					
65–69	12.8	[11.8, 13.8]	10.5	[8.3, 13.1]	
70–74	10.0	[9.3, 10.8]	12.8	[9.6, 16.8]	
75–79	8.4	[7.6, 9.3]	3.7	[2.2, 6.2]	***
80 and above	7.2	[6.6, 7.8]	4.0	[2.2, 7.0]	**

¹ Weighted prevalence. CI, confidence interval. ** $p < 0.05$ and *** $p < 0.01$; probabilities compare the FMD among older adults living alone versus living in multigenerational families.

Multivariable regression results: In Table 4, we provide the unadjusted (Model 1) and adjusted (Model 2) odds of FMD among older adults. The adjusted model included all covariates presented earlier in Section 2 (Materials and Methods). The findings in Table 4—Model 2 indicate that the odds of FMD reported by older adults living in multigenerational families was lower than that reported by counterparts living alone (adjusted odds ratio (AOR): 0.81; 95% confidence interval (CI): 0.65, 1.02). However, this difference achieved statistical significance only at the 10% significance level.

Table 4. Logistic regression results—FMD among older adults, BRFSS 2016–2020.

Unadjusted Models	Model 1		Model 3	
	OR	[95% CI]	OR	[95% CI]
Multigenerational (Ref: living alone)	0.98	[0.81, 1.19]		
Multigenerational (Ref: living alone) With grandchild only			1.49	* [0.99, 2.22]
With grandchild, spouse, and/or other adults			0.93	[0.75, 1.15]
Adjusted Models	Model 2		Model 4	
	AOR	[95% CI]	AOR	[95% CI]
Multigenerational (Ref: living alone)	0.81	* [0.65, 1.02]		
Multigenerational (Ref: living alone) With grandchild only			1.19	[0.75, 1.87]
With grandchild, spouse, and/or other adults			0.77	** [0.60, 0.99]
Age (5 year age groups)	0.73	*** [0.70, 0.76]	0.73	*** [0.70, 0.76]
Female (Ref: male)	1.04	[0.93, 1.16]	1.03	[0.92, 1.16]
Divorced/widowed/separated/never married (Ref: married/unmarried couple)	0.88	[0.73, 1.05]	0.86	[0.72, 1.03]
Race/ethnicity (Ref: NH White)				
NH Black	0.87	[0.72, 1.06]	0.87	[0.71, 1.06]
NH Other	0.92	[0.71, 1.20]	0.92	[0.71, 1.20]
Hispanic	0.90	[0.71, 1.15]	0.91	[0.71, 1.15]
Education (Ref: <high school)				
High school	1.04	[0.88, 1.24]	1.04	[0.88, 1.24]
Attended college or technical school	1.11	[0.93, 1.32]	1.11	[0.93, 1.32]
≥College or technical school	0.99	[0.82, 1.20]	0.99	[0.82, 1.20]
Income (Ref: <\$15,000)				
\$15,000–24,999	0.93	[0.81, 1.07]	0.93	[0.81, 1.07]
\$25,000–34,999	0.80	*** [0.68, 0.94]	0.80	*** [0.68, 0.94]
\$35,000–44,999	0.85	* [0.71, 1.01]	0.85	* [0.71, 1.01]
≥\$50,000	0.72	*** [0.59, 0.87]	0.72	*** [0.59, 0.87]
Chronic conditions	1.34	*** [1.30, 1.39]	1.34	*** [1.30, 1.39]
Physical limitations	1.57	*** [1.51, 1.63]	1.57	*** [1.51, 1.63]
Year: 2020 (Ref: 2016–2019)	1.03	[0.89, 1.21]	1.03	[0.88, 1.20]

OR, odds ratio; AOR, adjusted OR; CI, confidence interval; Ref: reference category. *** $p < 0.01$; ** $p < 0.05$; and * $p < 0.1$.

To further assess the difference in FMD reported by older adults living alone versus living in multigenerational families, we utilized the secondary explanatory variable created using the alternative coding scheme presented in Section 2 (Materials and Methods). This variable consisted of the three categories of older adults living alone (=0), living with their grandchildren only (=1), and living with their grandchildren, spouse, and/or other adults (=2). We refer to this last category as “non-grandchild only multigenerational families” henceforth. The results in Table 4 indicate the unadjusted (Model 3) and adjusted (Model 4) odds of FMD among older adults for these three categories. The unadjusted models (Model 3) shows that older adults living with their grandchild only were more likely to experience FMD compared to older adults living alone. However, this difference was no longer significant after controlling for covariates in the adjusted model (Model 4). Table 4—Model 4 also shows that the odds of FMD among older adults in non-grandchild only multigenerational families was lower by approximately 23% (AOR: 0.77; 95% CI: 0.60, 0.99; $p < 0.05$) when compared to older adults living alone.

Additionally, the results in both adjusted models (Models 2 and 4) in Table 4 indicate statistically significantly lower odds of FMD with each 5 year increase in age (AOR: 0.73; 95% CI: 0.70, 0.76). To further examine this age-associated difference, we conducted stratified regression analyses and compared the odds of FMD with each 5 year increase in age among older adults in non-grandchild only multigenerational families versus those living alone. The results are presented in Table 5, with Model 1 showing the adjusted odds of FMD among older adults living alone and Model 2 estimating the adjusted odds of FMD among those 65 years and older in non-grandchild only multigenerational families. With each 5 year increase in age, the odds of FMD decreased by 26% (AOR: 0.74; 95% CI: 0.71, 0.77) among older adults living alone (Table 5—Model 1). In contrast, this statistic among older adults in non-grandchild only multigenerational families was a 44% reduction (AOR: 0.56; 95% CI: 0.46, 0.70) in the self-reported odds of FMD with each 5 year increase in age (Table 5—Model 2).

Table 5. Stratified logistic regression results—FMD among older adults living alone (Model 1) versus living in multigenerational families with grandchild, spouse, and/or other adults (Model 2), BRFSS 2016–2020.

	Model 1:			Model 2:		
	Living Alone			Living with Grandchild, Spouse, and/or Other Adults		
	n = 122,246; N = 24,723,912			n = 3046; N = 1,550,179		
	AOR		[95% CI]	AOR		[95% CI]
Age (5 year age groups)	0.74	***	[0.71, 0.77]	0.56	***	[0.46, 0.70]
Female (Ref: male)	0.99		[0.88, 1.11]	1.97	***	[1.22, 3.19]
Divorced/widowed/separated/never married (Ref: married/unmarried couple)	0.81	***	[0.67, 0.97]	1.26		[0.77, 2.09]
Race/ethnicity (Ref: NH White)						
NH Black	0.96		[0.78, 1.18]	0.20	***	[0.09, 0.46]
NH Other	0.97		[0.74, 1.28]	0.25	***	[0.12, 0.52]
Hispanic	0.99		[0.77, 1.27]	0.46		[0.18, 1.17]
Education (Ref: <high school)						
High school	1.05		[0.88, 1.26]	0.90		[0.41, 1.99]
Attended college or technical school	1.11		[0.93, 1.33]	1.04		[0.46, 2.38]
≥College or technical school	0.99		[0.81, 1.20]	1.29		[0.55, 3.05]
Income (Ref: <\$15,000)						
\$15,000–24,999	0.90		[0.78, 1.04]	2.37		[1.19, 4.74]
\$25,000–34,999	0.78	***	[0.67, 0.92]	1.60		[0.64, 4.02]
\$35,000–44,999	0.81	***	[0.67, 0.97]	2.30	***	[1.00, 5.30]
≥\$50,000	0.71	***	[0.58, 0.87]	1.14		[0.50, 2.61]

Table 5. Cont.

	Model 1:			Model 2:		
	Living Alone			Living with Grandchild, Spouse, and/or Other Adults		
	n = 122,246; N = 24,723,912			n = 3046; N = 1,550,179		
Chronic conditions	1.34	***	[1.30, 1.39]	1.44	***	[1.26, 1.65]
Physical limitations	1.58	***	[1.51, 1.64]	1.49	***	[1.26, 1.76]
Year: 2020 (Ref: 2016–2019)	1.01		[0.86, 1.19]	1.50		[0.77, 2.93]

AOR, adjusted OR; CI, confidence interval; Ref, reference category. *** $p < 0.01$

To evaluate the equality of the coefficients for the 5 year age group variable from the stratified regressions for the two older adult groups—those living alone versus those living in non-grandchild only multigenerational families—we applied the methodology of “seemingly unrelated estimation” for the testing of the equality of the coefficients using the generalized Hausman specification test (Mize, Doan, & Long, 2019; Stata, n.d.). The results of the Hausman test indicated that the coefficients for age (5 year age category) for the two older adult groups were statistically significantly different ($p < 0.05$).

4. Discussion

With the progressive decline in physical health and the increase in chronic conditions, perceived mental health has gained salience for the healthy aging of older adults [2,6,47–50]. Frequent mental distress (FMD) is one of the leading health indicators that represents the burden of poor mental health in population groups [5,14–16]. A growing body of work has investigated FMD across various demographics, but studies analyzing FMD among older adults are sparse, and no prior work has examined FMD among older adults living in multigenerational family households. The prevalence of multigenerational living and older grandparent–grandchild coresidence in the US has been rising over recent decades, especially during the COVID-19 pandemic years [27,29,31,95,96]. Prior research conceptualizes the multigenerational living of older adults, on the one hand, to be advantageous and, on the other, to have an adverse influence on later-life mental health and, therefore, is inconclusive [32–35,38–46].

We evaluated FMD among older adults living alone versus living in multigenerational families in 36 US states using a representative sample from the BRFSS pooled between 2016 and 2020. In this analysis, the specific research question we aimed to address was, “does multigenerational living have a protective (or a risk) association with FMD among older adults?” In relation, we report two main findings. The results indicate statistically significant ($p < 0.05$) lower likelihood of FMD among older adults living with grandchild, spouse, and/or other adults versus those living alone. The findings also show age-associated differences with the likelihood of FMD decreasing with each 5 year increase in the age of older adults. However, this age-associated decrease in FMD was larger among older adults living with grandchild, spouse, and/or other adults compared to older adults living alone. Together, these results may be indicative of a protective association between multigenerational living and FMD among older adults.

Multiple prior studies forward evidence on the adverse mental health consequences among older adults living alone and, therefore, at an elevated risk of loneliness, social isolation, and feelings of depression [1,11,36,37,51,97]. In conjunction, we extend the prior findings by showcasing the likely benefit of multigenerational living, in contrast to living alone, for FMD among older adults. Our results using FMD are meaningful, as this indicator captures the burden of poor mental health among older adults more accurately than clinical diagnosis of depression [1,2,98].

The findings from our analysis also have some overlap with prior evidence on FMD among older Americans, although no prior study in this area has examined FMD among

those 65 years and older living in multigenerational families. Segev, Arif, and Rohrer [6] and Strine, Hootman, Okoro, et al., [4] analyzed BRFSS data and report significantly lower odds of FMD with each 5 year increase in the age of adults 45 years and older. Our results add to this evidence base by revealing that the age-associated reduction in FMD was larger among older adults living with a grandchild, spouse, and/or other adults when compared to counterparts living alone. The literature identifies greater resiliency, lower depressive distress, and lower negative emotions, often despite life challenges and adversities in old age [1,2,47,48,99–102]. Studies have also shown positive associations between resilience, life satisfaction, and the self-reported mental health of older adults [102,103]. It is plausible that multigenerational living plays a role in the furtherance of later-life resiliency and the mental HRQOL life of older adults.

The limitations of our study need specific acknowledgement. Given that the BRFSS is a cross-sectional dataset, our analysis could only show an association between the outcome FMD and the explanatory variable—multigenerational living versus status of living alone of older Americans—at one point in time. However, determining the mental health of older adults over time would be critical to establish the role of multigenerational living over the long run using longitudinal data. In addition, based on the BRFSS data, we could not identify features of multigenerational living, for example, family resources and support, which may be particularly beneficial for the mental health and resiliency of older adults. Thus, additional research is needed to parse out the protective features of multigenerational living that translate into a mental health advantage among older adults. Lastly, our analysis constitutes findings on older adults that are generalizable across a subset (approximately two-thirds) of the states in the US. Notwithstanding these limitations, a primary contribution of our study has been to forward evidence on the subjective mental health, specifically frequent mental distress (FMD), of older adults living in multigenerational families using a national dataset.

5. Conclusions

The research suggests that better perceived mental health at older ages has a role in healthy aging. An important question is whether this later-life advantage is a characteristic that represents the subjective mental health of older adults living in multigenerational family households. Addressing this question becomes paramount as the share of older adults living in multigenerational family households continues to rise in the US, particularly since the pandemic years. Using a national dataset, we showed a lower likelihood of frequent mental distress among older adults living in multigenerational families compared to older adults living alone in 36 US states. Once formed, multigenerational families often tend to live together over time and, therefore, have the potential to facilitate aging in place, as well as lower the need for the institutional care of older adults. Future studies are needed to identify the specific features of multigenerational living that improve or limit mental health among older adults across life stages and over time.

Author Contributions: Conceptualization, D.D.G. and D.W.S.W.; methodology, D.D.G.; data curation and formal analysis, D.D.G.; resources, D.D.G.; writing—original draft preparation, D.D.G.; writing—review and editing, D.D.G. and D.W.S.W.; funding acquisition, D.D.G. and D.W.S.W. All authors have read and agreed to the published version of the manuscript.

Funding: This study was partly supported by a grant (award number: 0000001678) from the County Health Rankings and Roadmaps program, a program of the University of Wisconsin Population Health Institute, with support from the Robert Wood Johnson Foundation.

Data Availability Statement: Data are available from the authors upon request.

Acknowledgments: The authors would like to thank Madeline Abram-Moyle and Taya G. Jerman, graduate researchers at Utah State University, for their support with research tasks. The authors of this study are also grateful to the Kinesiology and Health Science Department, Utah State University, and George Mason University Libraries for their support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Das Gupta, D.; Kelekar, U.; Rice, D. Associations between Living Alone, Depression, and Falls among Community-Dwelling Older Adults in the US. *Prev. Med. Rep.* **2020**, *20*, 101273. [[CrossRef](#)] [[PubMed](#)]
2. Fiske, A.; Wetherell, J.L.; Gatz, M. Depression in Older Adults. *Annu. Rev. Clin. Psychol.* **2009**, *5*, 363–389. [[CrossRef](#)] [[PubMed](#)]
3. Reynolds, K.; Pietrzak, R.H.; El-Gabalawy, R.; Mackenzie, C.S.; Sareen, J. Prevalence of Psychiatric Disorders in U.S. Older Adults: Findings from a Nationally Representative Survey. *World Psychiatry* **2015**, *14*, 74–81. [[CrossRef](#)] [[PubMed](#)]
4. Strine, T.W.; Hootman, J.M.; Okoro, C.A.; Balluz, L.; Moriarty, D.G.; Owens, M.; Mokdad, A. Frequent Mental Distress Status among Adults with Arthritis Age 45 Years and Older, 2001. *Arthritis Rheum.* **2004**, *51*, 533–537. [[CrossRef](#)] [[PubMed](#)]
5. Strine, T.W.; Greenlund, K.J.; Brown, D.W.; Mokdad, A.; Balluz, L. Characteristics of People Aged 45 Years or Older with Heart Disease by Frequent Mental Distress Status, 2001. *Prev. Med.* **2004**, *39*, 191–196. [[CrossRef](#)]
6. Segev, Z.; Arif, A.A.; Rohrer, J.E. Activity Limitations and Healthcare Access as Correlates of Frequent Mental Distress in Adults 65 Years and Older: A Behavioral Risk Factor Surveillance Study—2008. *J. Prim. Care Community Health* **2012**, *3*, 17–22. [[CrossRef](#)]
7. McGuire, L.C.; Strine, T.W.; Vachirasudlekha, S.; Anderson, L.A.; Berry, J.T.; Mokdad, A.H. Modifiable Characteristics of a Healthy Lifestyle and Chronic Health Conditions in Older Adults with or without Serious Psychological Distress, 2007 Behavioral Risk Factor Surveillance System. *Int. J. Public Health* **2009**, *54* (Suppl. S1), 84–93. [[CrossRef](#)]
8. Schwarzbach, M.; Luppia, M.; Forstmeier, S.; König, H.-H.; Riedel-Heller, S.G. Social Relations and Depression in Late Life—a Systematic Review. *Int. J. Geriatr. Psychiatry* **2014**, *29*, 1–21. [[CrossRef](#)]
9. Santini, Z.I.; Jose, P.E.; York Cornwell, E.; Koyanagi, A.; Nielsen, L.; Hinrichsen, C.; Meilstrup, C.; Madsen, K.R.; Koushede, V. Social Disconnectedness, Perceived Isolation, and Symptoms of Depression and Anxiety among Older Americans (NSHAP): A Longitudinal Mediation Analysis. *Lancet Public Health* **2020**, *5*, e62–e70. [[CrossRef](#)]
10. Pronk, N.P.; Kleinman, D.V.; Richmond, T.S. Healthy People 2030: Moving toward Equitable Health and Well-Being in the United States. *EClinicalMedicine* **2021**, *33*, 100777. [[CrossRef](#)]
11. National Academies of Sciences, Engineering, and Medicine; Division of Behavioral and Social Sciences and Education; Health and Medicine Division; Board on Behavioral, Cognitive, and Sensory Sciences; Board on Health Sciences Policy; Committee on the Health and Medical Dimensions of Social Isolation and Loneliness in Older Adults. *Social Isolation and Loneliness in Older Adults: Opportunities for the Health Care System*; National Academies Press (US): Washington, DC, USA, 2020.
12. Druss, B.G.; Satcher, D. Bridging Mental Health and Public Health. *Prev. Chronic Dis.* **2009**, *7*, A03.
13. Blackwelder, A.; Hoskins, M.; Huber, L. Effect of Inadequate Sleep on Frequent Mental Distress. *Prev. Chronic Dis.* **2021**, *18*, E61. [[CrossRef](#)] [[PubMed](#)]
14. County Health Rankings and Roadmap. *Frequent Mental Distress*; County Health Rankings & Roadmaps: Madison, WI, USA. Available online: <https://www.countyhealthrankings.org/explore-health-rankings/county-health-rankings-model/health-outcomes/quality-of-life/frequent-mental-distress> (accessed on 7 February 2023).
15. *Leading Health Indicators 2030: Advancing Health, Equity, and Well-Being*; National Academies Press: Washington, DC, USA, 2020. [[CrossRef](#)]
16. Strine, T.W.; Balluz, L.; Chapman, D.P.; Moriarty, D.G.; Owens, M.; Mokdad, A.H. Risk Behaviors and Healthcare Coverage among Adults by Frequent Mental Distress Status, 2001. *Am. J. Prev. Med.* **2004**, *26*, 213–216. [[CrossRef](#)]
17. Bruning, J.; Arif, A.A.; Rohrer, J.E. Medical Cost and Frequent Mental Distress among the Non-Elderly US Adult Population. *J. Public Health* **2014**, *36*, 134–139. [[CrossRef](#)] [[PubMed](#)]
18. Chen, H.-Y.; Baumgardner, D.J.; Rice, J.P. Health-Related Quality of Life Among Adults With Multiple Chronic Conditions in the United States, Behavioral Risk Factor Surveillance System, 2007. *Prev. Chronic Dis.* **2010**, *8*, A09. [[PubMed](#)]
19. Chowdhury, P.P.; Balluz, L.; Strine, T.W. Health-Related Quality of Life among Minority Populations in the United States, BRFSS 2001–2002. *Ethn. Dis.* **2008**, *18*, 483–487.
20. Cree, R.A.; Okoro, C.A.; Zack, M.M.; Carbone, E. Frequent Mental Distress Among Adults, by Disability Status, Disability Type, and Selected Characteristics—United States, 2018. *MMWR Morb. Mortal. Wkly. Rep.* **2020**, *69*, 1238–1243. [[CrossRef](#)]
21. Liu, Y.; Croft, J.B.; Wheaton, A.G.; Perry, G.S.; Chapman, D.P.; Strine, T.W.; McKnight-Eily, L.R.; Presley-Cantrell, L. Association between Perceived Insufficient Sleep, Frequent Mental Distress, Obesity and Chronic Diseases among US Adults, 2009 Behavioral Risk Factor Surveillance System. *BMC Public Health* **2013**, *13*, 1–8. [[CrossRef](#)]
22. McGuire, L.C.; Strine, T.W.; Okoro, C.A.; Ahluwalia, I.B.; Ford, E.S. Modifiable Characteristics of a Healthy Lifestyle in U.S. Older Adults with or without Frequent Mental Distress: 2003 Behavioral Risk Factor Surveillance System. *Am. J. Geriatr. Psychiatry* **2007**, *15*, 754–761. [[CrossRef](#)]
23. Mukherjee, S. Comparing Adult Males and Females in the United States to Examine the Association between Body Mass Index and Frequent Mental Distress: An Analysis of Data from BRFSS 2011. *Psychiatry J.* **2013**, *2013*, 230928. [[CrossRef](#)]
24. Rohrer, J.E.; Borders, T.F.; Blanton, J. Rural Residence Is Not a Risk Factor for Frequent Mental Distress: A Behavioral Risk Factor Surveillance Survey. *BMC Public Health* **2005**, *5*, 46. [[CrossRef](#)]
25. Skarupski, K.A.; de Leon, C.F.M.; Bienias, J.L.; Scherr, P.A.; Zack, M.M.; Moriarty, D.G.; Evans, D.A. Black-White Differences in Health-Related Quality of Life among Older Adults. *Qual. Life Res.* **2007**, *16*, 287–296. [[CrossRef](#)] [[PubMed](#)]

26. Zhao, G.; Okoro, C.A.; Hsia, J.; Town, M. Self-Perceived Poor/Fair Health, Frequent Mental Distress, and Health Insurance Status Among Working-Aged US Adults. *Prev. Chronic Dis.* **2018**, *15*, E95. [CrossRef] [PubMed]
27. Cohn, D.V.; Horowitz, J.M.; Minkin, R.; Fry, R.; Hurst, K. *The Demographics of Multigenerational Households*; Pew Research Center's Social & Demographic Trends Project: Washington, DC, USA, 2022. Available online: <https://www.pewresearch.org/social-trends/2022/03/24/the-demographics-of-multigenerational-households/> (accessed on 7 February 2023).
28. Silverstein, M.; Giarrusso, R. Aging and Family Life: A Decade Review. *J. Marriage Fam.* **2010**, *72*, 1039–1058. [CrossRef]
29. Cohn, D.; Passel, J.S. *A Record 64 Million Americans Live in Multigenerational Households*; Pew Research Center: Washington, DC, USA, 2018. Available online: <https://www.pewresearch.org/fact-tank/2018/04/05/a-record-64-million-americans-live-in-multigenerational-households/> (accessed on 7 February 2023).
30. Do, D.; Rodgers, R.; Drew, J.A.R. Multigenerational Families and Food Insecurity in the United States, 1998–2013. *Ihis Data Brief* **2015**, *1*, 1–12.
31. Pilkauskas, N.V.; Amorim, M.; Dunifon, R.E. Historical Trends in Children Living in Multigenerational Households in the United States: 1870–2018. *Demography* **2020**, *57*, 2269–2296. [CrossRef]
32. Baltes, P.B.; Baltes, M.M. Psychological Perspectives on Successful Aging: The Model of Selective Optimization with Compensation. In *Successful Aging: Perspectives from the Behavioral Sciences*; Cambridge University Press: New York, NY, USA, 1990; pp. 1–34. [CrossRef]
33. Erikson, E.H. *Childhood and Society*; W. W. Norton & Co.: New York, NY, USA, 1950; p. 397.
34. Lee, H.; Ryan, L.H.; Ofstedal, M.B.; Smith, J. Multigenerational Households During Childhood and Trajectories of Cognitive Functioning Among U.S. Older Adults. *J. Gerontol. B Psychol. Sci. Soc. Sci.* **2021**, *76*, 1161–1172. [CrossRef]
35. Muennig, P.; Jiao, B.; Singer, E. Living with Parents or Grandparents Increases Social Capital and Survival: 2014 General Social Survey-National Death Index. *SSM Popul. Health* **2017**, *4*, 71–75. [CrossRef]
36. Erikson, E.H.; Erikson, J.M. *The Life Cycle Completed (Extended Version)*; W. W. Norton & Company: New York, NY, USA, 1998.
37. Hagestad, G.O.; Uhlenberg, P. Should We Be Concerned About Age Segregation?: Some Theoretical and Empirical Explorations. *Res. Aging* **2006**, *28*, 638–653. [CrossRef]
38. Baker, L.A.; Silverstein, M. Preventive Health Behaviors Among Grandmothers Raising Grandchildren. *J. Gerontol. B Psychol. Sci. Soc. Sci.* **2008**, *63*, S304–S311. [CrossRef]
39. Baker, L.A.; Silverstein, M. Depressive Symptoms Among Grandparents Raising Grandchildren: The Impact of Participation in Multiple Roles. *J. Intergener. Relatsh.* **2008**, *6*, 285–304. [CrossRef] [PubMed]
40. Bengtson, V.; Giarrusso, R.; Mabry, J.B.; Silverstein, M. Solidarity, Conflict, and Ambivalence: Complementary or Competing Perspectives on Intergenerational Relationships? *J. Marriage Fam.* **2002**, *64*, 568–576. [CrossRef]
41. Coall, D.A.; Hertwig, R. Grandparental Investment: Past, Present, and Future. *Behav. Brain Sci.* **2010**, *33*, 1–19, discussion 19–40. [CrossRef] [PubMed]
42. Coall, D.A.; Hertwig, R. Grandparental Investment: A Relic of the Past or a Resource for the Future? *Curr. Dir. Psychol. Sci.* **2011**, *20*, 93–98. [CrossRef]
43. Danielsbacka, M.; Křenková, L.; Tanskanen, A.O. Grandparenting, Health, and Well-Being: A Systematic Literature Review. *Eur. J. Ageing* **2022**, *19*, 341–368. [CrossRef]
44. Hughes, M.E.; Waite, L.J.; LaPierre, T.A.; Luo, Y. All in the Family: The Impact of Caring for Grandchildren on Grandparents' Health. *J. Gerontol. B Psychol. Sci. Soc. Sci.* **2007**, *62*, S108–S119. [CrossRef]
45. Lowenstein, A. Solidarity-Conflict and Ambivalence: Testing Two Conceptual Frameworks and Their Impact on Quality of Life for Older Family Members. *J. Gerontol. B Psychol. Sci. Soc. Sci.* **2007**, *62*, S100–S107. [CrossRef]
46. Ward, R.A. Multiple Parent-Adult Child Relations and Well-Being in Middle and Later Life. *J. Gerontol. B Psychol. Sci. Soc. Sci.* **2008**, *63*, S239–S247. [CrossRef]
47. Gooding, P.A.; Hurst, A.; Johnson, J.; TARRIER, N. Psychological Resilience in Young and Older Adults. *Int. J. Geriatr. Psychiatry* **2012**, *27*, 262–270. [CrossRef]
48. Hayslip, B., Jr.; Smith, G.C. *Resilient Grandparent Caregivers: A Strengths-Based Perspective*; Routledge & CRC Press: Boca Raton, FL, USA, 2014.
49. Stepko, B.; AARP. *Older Adults Reporting Better Mental Health Than Young*; AARP: Washington, DC, USA, 2021. Available online: <https://www.aarp.org/health/conditions-treatments/info-2021/older-mental-health.html> (accessed on 7 February 2023).
50. National Academies of Sciences, Engineering, and Medicine; Division of Behavioral and Social Sciences and Education; Committee on Population. *Future Directions for the Demography of Aging: Proceedings of a Workshop*; Majmundar, M.K., Hayward, M.D., Eds.; National Academies Press (US): Washington, DC, USA, 2018.
51. Stone, J.; Evandrou, M.; Falkingham, J. The Transition to Living Alone and Psychological Distress in Later Life. *Age Ageing* **2013**, *42*, 366–372. [CrossRef]
52. Centers for Disease Control and Prevention. *Behavioral Risk Factor Surveillance System*; Centers for Disease Control and Prevention: Atlanta, GA, USA, 2022. Available online: <https://www.cdc.gov/brfss/index.html> (accessed on 29 August 2022).
53. Silva, N.M. The Behavioral Risk Factor Surveillance System. *Int. J. Aging Hum. Dev.* **2014**, *79*, 336–338. [CrossRef]
54. Centers for Disease Control and Prevention. *CDC-BRFSS-Survey Data & Documentation*; Centers for Disease Control and Prevention: Atlanta, GA, USA, 2022. Available online: https://www.cdc.gov/brfss/data_documentation/index.htm (accessed on 29 August 2022).

55. Centers for Disease Control and Prevention. *CDC-BRFSS-BRFSS Frequently Asked Questions (FAQs)*; Centers for Disease Control and Prevention: Atlanta, GA, USA, 2023. Available online: https://www.cdc.gov/brfss/about/brfss_faqs.htm (accessed on 7 February 2023).
56. Holt-Lunstad, J.; Smith, T.B.; Baker, M.; Harris, T.; Stephenson, D. Loneliness and Social Isolation as Risk Factors for Mortality: A Meta-Analytic Review. *Perspect. Psychol. Sci.* **2015**, *10*, 227–237. [[CrossRef](#)]
57. Ortiz, H. Crossing New Frontiers: Benefits Access among Isolated Seniors | [advancingstates.org](http://www.advancingstates.org/hcbs/article/crossing-new-frontiers-benefits-access-among-isolated-seniors). Available online: <http://www.advancingstates.org/hcbs/article/crossing-new-frontiers-benefits-access-among-isolated-seniors> (accessed on 7 February 2023).
58. Havinghurst, R. Ageing in Western Society. *Soc. Chall. Ageing. Lond. Croom Helm* **1978**, 15–44.
59. Henning-Smith, C. Quality of Life and Psychological Distress among Older Adults: The Role of Living Arrangements. *J. Appl. Gerontol.* **2016**, *35*, 39–61. [[CrossRef](#)]
60. Pohl, J.S.; Cochrane, B.B.; Schepp, K.G.; Woods, N.F. Falls and Social Isolation of Older Adults in the National Health and Aging Trends Study. *Res. Gerontol. Nurs.* **2018**, *11*, 61–70. [[CrossRef](#)]
61. Shaw, J.G.; Farid, M.; Noel-Miller, C.; Joseph, N.; Houser, A.; Asch, S.M.; Bhattacharya, J.; Flowers, L. Social Isolation and Medicare Spending: Among Older Adults, Objective Social Isolation Increases Expenditures While Loneliness Does Not. *J. Aging Health* **2017**, *29*, 1119–1143. [[CrossRef](#)]
62. Whitley, D.M.; Fuller-Thomson, E.; Brennenstuhl, S. Health Characteristics of Solo Grandparent Caregivers and Single Parents: A Comparative Profile Using the Behavior Risk Factor Surveillance Survey. *Curr. Gerontol. Geriatr. Res.* **2015**, *2015*, 630717. [[CrossRef](#)] [[PubMed](#)]
63. Wenger, G.C.; Davies, R.; Shahtahmasebi, S.; Scott, A. Social Isolation and Loneliness in Old Age: Review and Model Refinement. *Ageing Soc.* **1996**, *16*, 333–358. [[CrossRef](#)]
64. Dunifon, R.E.; Ziolo-Guest, K.M.; Kopko, K. Grandparent Coresidence and Family Well-Being: Implications for Research and Policy. *ANNALS Am. Acad. Political Soc. Sci.* **2014**, *654*, 110–126. [[CrossRef](#)]
65. Keene, J.R.; Batson, C.D. Under One Roof: A Review of Research on Intergenerational Coresidence and Multigenerational Households in the United States. *Sociol. Compass* **2010**, *4*, 642–657. [[CrossRef](#)]
66. Lofquist, D.A. *Multigenerational Households: 2009–2011*; American Community Survey Briefs US Census Bureau: Washington, DC, USA, 2012. Available online: <https://www.census.gov/library/publications/2012/acs/acsbr11-03.html> (accessed on 7 February 2023).
67. Whitley, D.M.; Fuller-Thomson, E. Latino Solo Grandparents Raising Grandchildren: Health Risks and Behaviors. *Hisp. Health Care Int.* **2018**, *16*, 11–19. [[CrossRef](#)]
68. Whitley, D.M.; Fuller-Thomson, E. African-American Solo Grandparents Raising Grandchildren: A Representative Profile of Their Health Status. *J. Community Health* **2017**, *42*, 312–323. [[CrossRef](#)] [[PubMed](#)]
69. Whitley, D.M.; Fuller-Thomson, E. The Health of the Nation’s Custodial Grandfathers and Older Single Fathers: Findings From the Behavior Risk Factor Surveillance System. *Am. J. Men’s Health* **2017**, *11*, 1614–1626. [[CrossRef](#)] [[PubMed](#)]
70. Centers for Disease Control and Prevention, *Measuring Healthy Days: Population Assessment of Health-Related Quality of Life, 2000*; Centers for Disease Control and Prevention: Atlanta, GA, USA. Available online: <https://www.cdc.gov/hrqol/pdfs/mhd.pdf> (accessed on 7 February 2023).
71. Andresen, E.; Catlin, T.; Wyrwich, K.; Jackson-Thompson, J. Retest Reliability of Surveillance Questions on Health Related Quality of Life. *J. Epidemiol Community Health* **2003**, *57*, 339–343. [[CrossRef](#)] [[PubMed](#)]
72. Bann, C.M.; Kobau, R.; Lewis, M.A.; Zack, M.M.; Luncheon, C.; Thompson, W.W. Development and Psychometric Evaluation of the Public Health Surveillance Well-Being Scale. *Qual. Life Res.* **2012**, *21*, 1031–1043. [[CrossRef](#)] [[PubMed](#)]
73. Hsia, J.; Zhao, G.; Town, M.; Ren, J.; Okoro, C.A.; Pierannunzi, C.; Garvin, W. Comparisons of Estimates From the Behavioral Risk Factor Surveillance System and Other National Health Surveys, 2011–2016. *Am. J. Prev. Med.* **2020**, *58*, e181–e190. [[CrossRef](#)]
74. Mielenz, T.; Jackson, E.; Currey, S.; DeVellis, R.; Callahan, L.F. Psychometric Properties of the Centers for Disease Control and Prevention Health-Related Quality of Life (CDC HRQOL) Items in Adults with Arthritis. *Health Qual. Life Outcomes* **2006**, *4*, 66. [[CrossRef](#)]
75. Moriarty, D.G.; Zack, M.M.; Kobau, R. The Centers for Disease Control and Prevention’s Healthy Days Measures—Population Tracking of Perceived Physical and Mental Health over Time. *Health Qual. Life Outcomes* **2003**, *1*, 37. [[CrossRef](#)]
76. Pierannunzi, C.; Hu, S.S.; Balluz, L. A Systematic Review of Publications Assessing Reliability and Validity of the Behavioral Risk Factor Surveillance System (BRFSS), 2004–2011. *BMC Med. Res. Methodol.* **2013**, *13*, 49. [[CrossRef](#)] [[PubMed](#)]
77. Centers for Disease Control and Prevention. *Healthy Days Core Module: HRQOL-14 Measure* | HRQOL | CDC; Centers for Disease Control and Prevention: Atlanta, GA, USA. Available online: https://www.cdc.gov/hrqol/hrqol14_measure.htm (accessed on 7 February 2023).
78. Centers for Disease Control and Prevention. HRQOL Measures Can Help Track Health as Population Ages. Health-Related Quality of Life, Part I, 2003. *Chronic Dis. Notes Rep.* **2003**, *16*, 1–36.
79. Moriarty, D.G.; Kobau, R.; Zack, M.M.; Zahran, H.S. Tracking Healthy Days—A Window on the Health of Older Adults. *Prev. Chronic Dis.* **2005**, *2*, A16. [[PubMed](#)]
80. Ounpuu, S.; Chambers, L.W.; Chan, D.; Yusuf, S. Validity of the US Behavioral Risk Factor Surveillance System’s Health Related Quality of Life Survey Tool in a Group of Older Canadians. *Chronic Dis. Can.* **2001**, *22*, 93–101. [[PubMed](#)]

81. Fan, Z.J.; Bonauto, D.K.; Foley, M.P.; Anderson, N.J.; Yragui, N.L.; Silverstein, B.A. Occupation and the Prevalence of Current Depression and Frequent Mental Distress, WA BRFSS 2006 and 2008. *Am. J. Ind. Med.* **2012**, *55*, 893–903. [CrossRef]
82. Slabaugh, S.L.; Shah, M.; Zack, M.; Happe, L.; Cordier, T.; Havens, E.; Davidson, E.; Miao, M.; Prewitt, T.; Jia, H. Leveraging Health-Related Quality of Life in Population Health Management: The Case for Healthy Days. *Popul. Health Manag.* **2017**, *20*, 13–22. [CrossRef] [PubMed]
83. Edwards, V.J.; Anderson, L.A.; Thompson, W.W.; Deokar, A.J. Mental Health Differences between Men and Women Caregivers, BRFSS 2009. *J. Women Aging* **2017**, *29*, 385–391. [CrossRef]
84. Grenard, D.L.; Valencia, E.J.; Brown, J.A.; Winer, R.L.; Littman, A.J. Impact of Caregiving During Emerging Adulthood on Frequent Mental Distress, Smoking, and Drinking Behaviors: United States, 2015–2017. *Am. J. Public Health* **2020**, *110*, 1853–1860. [CrossRef]
85. Charara, R.; El Bcheraoui, C.; Kravitz, H.; Dhingra, S.S.; Mokdad, A.H. Mental Distress and Functional Health in the United States. *Prev. Med.* **2016**, *89*, 292–300. [CrossRef]
86. Dwyer-Lindgren, L.; Mackenbach, J.P.; van Lenthe, F.J.; Mokdad, A.H. Self-Reported General Health, Physical Distress, Mental Distress, and Activity Limitation by US County, 1995–2012. *Popul. Health Metr.* **2017**, *15*, 16. [CrossRef]
87. Hatzenbuehler, M.L.; Prins, S.J.; Flake, M.; Philbin, M.; Frazer, M.S.; Hagen, D.; Hirsch, J. Immigration Policies and Mental Health Morbidity among Latinos: A State-Level Analysis. *Soc. Sci. Med.* **2017**, *174*, 169–178. [CrossRef]
88. Moriarty, D.G.; Zack, M.M.; Holt, J.B.; Chapman, D.P.; Safran, M.A. Geographic Patterns of Frequent Mental Distress: U.S. Adults, 1993–2001 and 2003–2006. *Am. J. Prev. Med.* **2009**, *36*, 497–505. [CrossRef]
89. Price, J.D.; Barbour, K.E.; Liu, Y.; Lu, H.; Amerson, N.L.; Murphy, L.B.; Helmick, C.G.; Calanan, R.M.; Sandoval-Rosario, M.; Samanic, C.M.; et al. State-Specific Prevalence and Characteristics of Frequent Mental Distress and History of Depression Diagnosis Among Adults with Arthritis—United States, 2017. *MMWR Morb. Mortal. Wkly. Rep.* **2020**, *68*, 1173–1178. [CrossRef] [PubMed]
90. Taylor, M.F.; Marquis, R.; Coall, D.A.; Batten, R.; Werner, J. The physical health dilemmas facing custodial grandparent caregivers: Policy considerations. *Cogent Med.* **2017**, *4*, 1292594. [CrossRef]
91. Blustein, J.; Chan, S.; Guanais, F.C. Elevated depressive symptoms among caregiving grandparents. *Health Serv. Res.* **2004**, *39*, 1671–1690. [CrossRef]
92. Silva, M.; Loureiro, A.; Cardoso, G. Social Determinants of Mental Health: A Review of the Evidence. *Eur. J. Psychiatry* **2016**, *30*, 259–292.
93. US Census Bureau. *How Disability Data are Collected from The American Community Survey*; US Census Bureau: Washington, DC, USA, 2021. Available online: <https://www.census.gov/topics/health/disability/guidance/data-collection-acs.html> (accessed on 7 February 2023).
94. Hilbe, J.M. *Logistic Regression Models*, 1st ed.; Chapman and Hall/CRC: Boca Raton, FL, USA, 2009.
95. Firestone, S.K.; Butts, D. Changing the Narrative on Multigenerational Living. 2021. Available online: <https://generations.asaging.org/changing-narrative-multigenerational-living> (accessed on 7 February 2023).
96. Glaser, K.; Stuchbury, R.; Price, D.; Di Gessa, G.; Ribe, E.; Tinker, A. Trends in the Prevalence of Grandparents Living with Grandchild(Ren) in Selected European Countries and the United States. *Eur. J. Ageing* **2018**, *15*, 237–250. [CrossRef]
97. Xiu-Ying, H.; Qian, C.; Xiao-Dong, P.; Xue-Mei, Z.; Chang-Quan, H. Living Arrangements and Risk for Late Life Depression: A Meta-Analysis of Published Literature. *Int. J. Psychiatry Med.* **2012**, *43*, 19–34. [CrossRef]
98. Wetherell, J.L.; Petkus, A.J.; McChesney, K.; Stein, M.B.; Judd, P.H.; Rockwell, E.; Sewell, D.D.; Patterson, T.L. Older Adults Are Less Accurate than Younger Adults at Identifying Symptoms of Anxiety and Depression. *J. Nerv. Ment. Dis.* **2009**, *197*, 623–626. [CrossRef]
99. Coon, D.W. Resilience and Family Caregiving. *Annu. Rev. Gerontol. Geriatr.* **2012**, *32*, 231–249. [CrossRef]
100. Giarrusso, R.; Silverstein, M.; Feng, D. Psychological Costs and Benefits of Raising Grandchildren: Evidence from a National Survey of Grandparents. In *To Grandmother’s House We Go and Stay: Perspectives on Custodial Grandparents*; Springer: New York, NY, USA, 2000; pp. 71–90.
101. Hildon, Z.; Montgomery, S.M.; Blane, D.; Wiggins, R.D.; Netuveli, G. Examining Resilience of Quality of Life in the Face of Health-Related and Psychosocial Adversity at Older Ages: What Is “Right” about the Way We Age? *Gerontologist* **2010**, *50*, 36–47. [CrossRef]
102. MacLeod, S.; Musich, S.; Hawkins, K.; Alsgaard, K.; Wicker, E.R. The Impact of Resilience among Older Adults. *Geriatr. Nurs.* **2016**, *37*, 266–272. [CrossRef]
103. Lombardo, P.; Jones, W.; Wang, L.; Shen, X.; Goldner, E.M. The Fundamental Association between Mental Health and Life Satisfaction: Results from Successive Waves of a Canadian National Survey. *BMC Public Health* **2018**, *18*, 1–9. [CrossRef] [PubMed]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.