



2-2-2022

NF- κ B perturbation reveals unique immunomodulatory functions in Prx1 + Fibroblasts that Promote Development of Atopic Dermatitis

K. I. Ko

J. J. Merlet

B. P. DerGarabedian

H. Zhen

Y. Suzuki-Horiuchi

See next page for additional authors

Follow this and additional works at: https://repository.upenn.edu/dental_papers

Recommended Citation

Ko, K. I., Merlet, J. J., DerGarabedian, B. P., Zhen, H., Suzuki-Horiuchi, Y., Hedberg, M. L., Hu, E., Nguyen, A. T., Prouty, S., Alawi, F., & Walsh, M. C. (2022). NF- κ B perturbation reveals unique immunomodulatory functions in Prx1 + Fibroblasts that Promote Development of Atopic Dermatitis. *Science Translational Medicine*, 14 (630), <http://dx.doi.org/10.1126/scitranslmed.abj0324>

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/dental_papers/571
For more information, please contact repository@pobox.upenn.edu.

NF- κ B perturbation reveals unique immunomodulatory functions in Prx1⁺ Fibroblasts that Promote Development of Atopic Dermatitis

Abstract

Skin is composed of diverse cell populations that cooperatively maintain homeostasis. Up-regulation of the nuclear factor κ B (NF- κ B) pathway may lead to the development of chronic inflammatory disorders of the skin, but its role during the early events remains unclear. Through analysis of single-cell RNA sequencing data via iterative random forest leave one out prediction, an explainable artificial intelligence method, we identified an immunoregulatory role for a unique paired related homeobox-1 (Prx1)⁺ fibroblast subpopulation. Disruption of *Ikkb*-NF- κ B under homeostatic conditions in these fibroblasts paradoxically induced skin inflammation due to the overexpression of C-C motif chemokine ligand 11 (CCL11; or eotaxin-1) characterized by eosinophil infiltration and a subsequent T_H2 immune response. Because the inflammatory phenotype resembled that seen in human atopic dermatitis (AD), we examined human AD skin samples and found that human AD fibroblasts also overexpressed CCL11 and that perturbation of *Ikkb*-NF- κ B in primary human dermal fibroblasts up-regulated CCL11. Monoclonal antibody treatment against CCL11 was effective in reducing the eosinophilia and T_H2 inflammation in a mouse model.

Together, the murine model and human AD specimens point to dysregulated Prx1⁺ fibroblasts as a previously unrecognized etiologic factor that may contribute to the pathogenesis of AD and suggest that targeting CCL11 may be a way to treat AD-like skin lesions. © 2022 The Authors, some rights reserved

Keywords

Animals, Artificial Intelligence, Dermatitis, Atopic, Fibroblasts, Immunity, Mice, NF-kappa B, Skin

Author(s)

K. I. Ko, J. J. Merlet, B. P. DerGarabedian, H. Zhen, Y. Suzuki-Horiuchi, M. L. Hedberg, E. Hu, Ahn T. Nguyen, S. Prouty, F. Alawi, and M. C. Walsh

ATOPIC DERMATITIS

NF- κ B perturbation reveals unique immunomodulatory functions in Prx1⁺ fibroblasts that promote development of atopic dermatitis

Kang I. Ko^{1†}, Jean J. Merlet^{2†}, Brett P. DerGarabedian¹, Huang Zhen^{1,3}, Yoko Suzuki-Horiuchi⁴, Matthew L. Hedberg⁴, Eileen Hu¹, Anh T. Nguyen¹, Stephen Prouty⁴, Faizan Alawi⁵, Matthew C. Walsh⁶, Yongwon Choi⁶, Sarah E. Millar⁷, Ashley Cliff², Jonathon Romero², Michael R. Garvin⁸, John T. Seykora^{4*‡}, Daniel Jacobson^{8*‡}, Dana T. Graves^{1*‡}

Skin is composed of diverse cell populations that cooperatively maintain homeostasis. Up-regulation of the nuclear factor κ B (NF- κ B) pathway may lead to the development of chronic inflammatory disorders of the skin, but its role during the early events remains unclear. Through analysis of single-cell RNA sequencing data via iterative random forest leave one out prediction, an explainable artificial intelligence method, we identified an immunoregulatory role for a unique paired related homeobox-1 (Prx1)⁺ fibroblast subpopulation. Disruption of *Ikkb*-NF- κ B under homeostatic conditions in these fibroblasts paradoxically induced skin inflammation due to the overexpression of C-C motif chemokine ligand 11 (CCL11; or eotaxin-1) characterized by eosinophil infiltration and a subsequent T_H2 immune response. Because the inflammatory phenotype resembled that seen in human atopic dermatitis (AD), we examined human AD skin samples and found that human AD fibroblasts also overexpressed CCL11 and that perturbation of *Ikkb*-NF- κ B in primary human dermal fibroblasts up-regulated CCL11. Monoclonal antibody treatment against CCL11 was effective in reducing the eosinophilia and T_H2 inflammation in a mouse model. Together, the murine model and human AD specimens point to dysregulated Prx1⁺ fibroblasts as a previously unrecognized etiologic factor that may contribute to the pathogenesis of AD and suggest that targeting CCL11 may be a way to treat AD-like skin lesions.

INTRODUCTION

Skin disorders affect 85 million people in the United States and globally amount to 41.6 million years lost because of disease and living with disability (1), representing a critical public health burden. The factors that initiate pathogenesis of psoriasis and atopic dermatitis (AD) include disruption of the epithelial barrier and altered immune responses (2, 3). However, despite recent advances, clinical therapies often fail to achieve adequate resolution of symptoms in moderate to severe cases (4). Because the skin must respond rapidly to physical and microbial challenges, resident cells in the skin must respond quickly to perturbations to maintain a fine balance between pro- and anti-inflammatory processes (5). Thus, an improved understanding of cutaneous homeostasis is essential to better understand the early events in dermatologic diseases and to develop therapeutics targeting these early events.

¹Department of Periodontics, School of Dental Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA. ²Bredesen Center for Interdisciplinary Research and Graduate Education, University of Tennessee Knoxville, Knoxville, TN 37996, USA. ³Department of Periodontology, Peking University School and Hospital of Stomatology, Haidian District, Beijing 100081, China. ⁴Department of Dermatology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA. ⁵Department of Basic and Translational Sciences, School of Dental Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA. ⁶Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA. ⁷Black Family Stem Cell Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA. ⁸Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830, USA.

*Corresponding author. Email: seykora@penmedicine.upenn.edu (J.T.S.); jacobsonda@ornl.gov (D.J.); dtgraves@upenn.edu (D.T.G.).

†These authors contributed equally to this work.

‡These authors contributed equally to this work as co-senior authors.

Fibroblasts are the principal cell type in the dermis and exhibit considerable heterogeneity, including scar-inducing subsets marked by *En1* (6), *Prrx1* (7), or *Dpp4* (6, 8). Recent single-cell RNA sequencing (scRNA-seq) studies on human AD skin have revealed fibroblast subsets with unexpected immunomodulatory transcriptomes (9, 10), suggesting a potential role of these cells in the pathogenesis of inflammatory skin disorders. Synovial fibroblasts have been proposed to exhibit a proinflammatory phenotype to mediate joint destruction after induction of an autoimmune response in rheumatoid arthritis (11), whereas tumor-associated fibroblasts have been linked to immunosuppression and poor survival outcomes (12). In each of these cases, fibroblasts play a secondary role of responding to their microenvironment to modulate the immune response. Whether skin fibroblasts contribute to immune homeostasis in skin inflammatory disorders remains unknown.

Nuclear factor κ B (NF- κ B) is a master inflammatory transcription factor that regulates immune-responsive genes (13). Canonical NF- κ B proteins and their activator, the inhibitor of NF- κ B kinase complex, are ubiquitously expressed by cells including fibroblasts (14). In many dermatologic inflammatory conditions, up-regulation of NF- κ B activity is directly linked to proinflammatory events (15). Initiating events are not thought to be directly caused by NF- κ B but rather by the response of cells to proinflammatory signals that activate NF- κ B and up-regulate the pathogenic process (16).

Here, we used a genetically modified mouse model in which NF- κ B activation in fibroblastic cells was blocked by lineage-specific deletion of the NF- κ B regulator, inhibitor of NF- κ B kinase subunit β (*Ikkb*). Inhibition of NF- κ B in fibroblastic cells led to the formation of skin lesions, indicating that NF- κ B played a key role in maintaining homeostasis and preventing excessive inflammation.

Copyright © 2022
The Authors, some
rights reserved;
exclusive licensee
American Association
for the Advancement
of Science. No claim
to original U.S.
Government Works

Downloaded from https://www.science.org at University of Pennsylvania on June 14, 2022

With the use of scRNA-seq, we identified that paired related homeobox-1-positive ($Prx1^+$) cells containing *Ikkb* deletion played a key role in immune disruption by overexpressing the eosinophilic chemokine C-C motif chemokine ligand 11 (CCL11), which led to an inflammatory response. These results suggest that $Prx1^+$ cells are essential for suppressing myeloid inflammation initiated by *Ikkb* under basal conditions and that the lack of *Ikkb* leads to skin inflammatory disease characterized by early eosinophilia and a type 2 immune response frequently seen in AD-like lesions.

RESULTS

To investigate the role of NF- κ B activation in skin fibroblasts, we deleted *Ikkb*, a gene needed for activation of the canonical NF- κ B pathway (17), in $Prx1Cre^+Ikkb^{fl/fl}$ mice. $Prx1$ lineage cells are present exclusively in the ventral mesenchyme (fig. S1), consistent with

their embryological origin (18, 19). We first examined NF- κ B p65 nuclear localization, indicative of its activation, by immunofluorescence in vimentin⁺ fibroblasts and perilipin⁺ adipocytes in adult $Prx1Cre^+Ikkb^{fl/fl}$ and control littermates. Seventeen percent of fibroblasts and 20% of adipocytes from the wild-type mice had NF- κ B nuclear localization under standard physiologic conditions, whereas less than 4% had nuclear NF- κ B in the *Ikkb*-deleted mice (Fig. 1, A and B). We observed cutaneous lesions in the ventral but not dorsal skin of the *Ikkb*-deleted mice (Fig. 1C). The affected skin demonstrated hair loss, thickening, scaling, erythema, or focal crusting with scab formation ($n = 6$) that was not present in wild-type mice ($n = 6$). The histology of skin from four of six $Prx1Cre^+Ikkb^{fl/fl}$ mice demonstrated hyperkeratotic scale composed of ortho- or parakeratin and acanthosis associated with spongiosis, which was pronounced in three experimental mice and not present in wild-type mice (Fig. 1D). In the experimental mice, inflamed lesions




Fig. 1. Preventing basal NF- κ B activation in $Prx1^+$ mesenchymal cells leads to ventral skin anomalies. (A) Representative immunofluorescent images of ventral skin stained with NF- κ B (p65) antibody and vimentin (fibroblast marker) or perilipin (adipocyte marker) in 16-week-old control [wild type (WT)] or experimental ($Prx1Cre^+Ikkb^{fl/fl}$) mice. White arrows point to fibroblasts or adipocytes with nuclear NF- κ B expression. Scale bars, 50 μ m; insert scale bars, 5 μ m. DAPI, 4',6-diamidino-2-phenylindole. (B) Quantification of double-positive vimentin⁺/nuclear NF- κ B⁺ fibroblasts per total vimentin⁺ cells (left) and perilipin⁺/nuclear NF- κ B⁺ adipocytes per total perilipin⁺ cells (right), comparing WT and $Prx1Cre^+Ikkb^{fl/fl}$ [conditional knockout (cKO)] group. $n = 7$ to 8 mice in each group. (C) Representative photograph images of ventral and dorsal skin in 16- to 20-week-old WT and $Prx1Cre^+Ikkb^{fl/fl}$ mice. Scale bars, 1 cm. (D) Hematoxylin and eosin (H&E) and trichrome images of ventral skin from WT or cKO mice. d, dermis; ad, adipose layer. Scale bars, 0.5 mm. Insert, red arrows point to eosinophils. Scale bars, 50 μ m. (E and F) Epidermis, dermis, and adipose width in ventral (E) or dorsal (F) skin of WT and cKO mice. $n = 6$ to 10 mice in each group. (G) Mechanical testing of ventral skins from WT and cKO mice. Stiffness and maximum load until failure (tear) were quantified. $n = 4$ to 5 each. Data are represented as means \pm SEM of biological replicates. Animal experiments were repeated two to three times. * $P < 0.05$, Student's t test comparing WT to cKO groups. ns, not significant

had a prominent inflammatory infiltrate containing lymphocytes, eosinophils, and occasional neutrophils in all skin compartments (Fig. 1D), similar to those seen in human AD (20). Consistent with the distribution of skin lesions, *Prx1* lineage cells were found in ventral but not dorsal skin, and their cell percentage found in the ventral dermis of control and *Ikkb*-deleted mice was similar (fig. S1, A to C). Epidermal and dermal widths were significantly increased in ventral but not dorsal skin of *Prx1Cre⁺.Ikkb^{ff}* mice compared to control group ($P < 0.05$; Fig. 1, E and F, and fig. S1D). The skin with *Ikkb* deletion had decreased strength, as measured by reduced stiffness and mechanical load to failure (Fig. 1G), despite the histological evidence of fibroplasia and increased width of the dermis. The results suggest that the skin anomaly caused by the gene deletion is more complex than fibroplasia, possibly reflecting thickened dermis due to edema caused by inflammatory infiltrates.

To better understand the progression of the skin anomaly caused by *Ikkb* deletion, we tracked the lesion development throughout

postnatal growth from young (4 weeks old) to adult (20 weeks old) phase. In the *Ikkb*-deleted mice, alopecia developed as early as 10 weeks, always preceded ulceration (Fig. 2, A and B), and progressed without resolution (Fig. 2C). Because the initial clinical phenotype was hair loss, we tested whether the hair follicle niche was affected by the *Ikkb* deletion. The number of hair follicles and CD34⁺ hair bulges were not changed by *Ikkb* deletion (Fig. 2D), ruling out hair follicle niche disruption as an etiologic factor for the hair loss and alopecia.

To investigate early cellular events that may be responsible for the skin anomaly in adult mice, we examined neonatal (1 day after birth; P1) and young (4 weeks after birth) mice, time points at which no gross skin lesions were present. Histologically, the skin of P1 mice in control and *Ikkb* deletion groups showed no signs of acanthosis or fibrosis (Fig. 2E). Four-week-old mice with the *Ikkb* gene deletion exhibited an abundant subcutaneous and dermal leukocyte infiltration with multilobed or kidney-shaped nuclei, consistent




Fig. 2. Progressive dermatitis caused by *Ikkb* deletion worsens with aging. (A) Representative images of clinical lesion from *Prx1Cre⁺.Ikkb^{ff}* (cKO) mice over 15- to 19-week period in age. Scale bars, 1 cm. (B) Quantification of percent control (WT) or cKO animals affected by alopecia (solid line) and/or ulceration (dashed line) in function of age over 4- to 21-week time points. (C) Lesion area quantified in square millimeters over the observational period. (D) Representative immunofluorescent images of CD34⁺ hair follicle bulge from WT or cKO groups (left, dotted line) and quantification of hair follicle numbers from H&E images or by CD34⁺ staining (right) in 20-week-old mice. Scale bars, 50 µm. (E) H&E images of ventral skin in 1-day-old (P1) neonatal WT and cKO mice (left) and quantification of epidermis and dermis thickness in each group (right). Scale bars, 0.5 mm. (F) H&E images of ventral skin in 4-week-old young WT or cKO mice (left; scale bar, 0.5 mm), with insert images showing overt inflammatory infiltrate in the skin of cKO mice (right; scale bars, 50 µm). (G) Quantification of epidermis, dermis, and adipose layer thickness between WT and cKO mice at 4 weeks of age. Dashed lines demarcate dermal (d) and adipose (ad) borders in (E) and (F). Data are represented as means ± SEM of biological replicates. Animal experiments were repeated three times. $n = 6$ to 10 each. * $P < 0.05$, Student's *t* test comparing WT to cKO groups.

with myeloid cells (Fig. 2F), and an absence of dermal and epidermal thickening (Fig. 2G). We therefore sought to better characterize the type of inflammation and found significantly elevated numbers of myeloperoxidase (MPO)⁺ neutrophils and F4/80⁺ myeloid cells in both young and adult mice with *Ikkb* deletion, whereas the total number of CD4⁺ T cells was unchanged ($P < 0.05$; Fig. 3, A and B, and fig. S2A). *Ikkb* deletion did not alter the number of inflammatory cells in dorsal skin (fig. S2B), and there were no significant differences in MPO⁺ cells in the ventral skin of control and *Ikkb*-deleted P1 mice ($P > 0.05$; Fig. 3C). Similar results were obtained by flow cytometry analysis; Gr-1^{high} cells (equivalent to Ly6g⁺ neutrophils) and F4/80⁺ myeloid cells were significantly higher in the *Ikkb*-deleted mice, whereas CD3⁺ T cell numbers did not change ($P < 0.05$; Fig. 3, D and E). To rule out the possibility that the inflammation was caused by external irritation from the solid corn bedding, young mice were housed with soft paper bedding, which did not reduce skin inflammation (Fig. 3F). Together, *Ikkb* deletion in Prx1⁺ fibroblastic cells causes myeloid inflammation in the skin that precedes the clinical anomalies found in adulthood.

Because Prx1⁺ mesenchymal progenitors that line the bone marrow are critical for the maintenance of hematopoiesis (21), we tested whether *Ikkb* deletion in Prx1 lineage cells had an impact on systemic inflammation. We found a significant increase in circulating monocytes and eosinophils from the peripheral blood, whereas other cell metrics, including total red blood cell count, were largely unchanged ($P < 0.05$; Fig. 3G). Splenomegaly was always present in the *Ikkb* deletion group (Fig. 3H) with normal architecture of splenic germinal centers (Fig. 3I). There was a significant increase in myeloid splenocyte counts (neutrophils, monocytes, macrophages, and eosinophils) ($P < 0.05$; Fig. 3J), indicating systemic inflammation without abnormal erythropoiesis. A similar increase in myeloid cell numbers was found in axillary lymph nodes of *Ikkb*-deleted adult mice (fig. S3). However, the number of myeloid cells in the bone marrow was unchanged (Fig. 3K), suggesting that myelopoiesis was not affected by *Ikkb* deletion. Combined with the finding that dorsal skin was unaffected by *Ikkb* deletion, the lack of change in myelopoiesis suggests that changes in circulating myeloid cells were likely the result of severe skin inflammation and not vice versa.

Prx1 lineage cells in the adult skin include dermal fibroblasts and subcutaneous adipocytes (19, 22). We therefore generated mice that had *Ikkb*-specific deletion in adipocytes (*Adipoq-Cre⁺.Ikkb^{fl/fl}*) and pan-fibroblasts (*Col1a2CreER⁺.Ikkb^{fl/fl}*) and examined whether skin lesions developed. Adipocyte-specific deletion of *Ikkb* failed to elicit myeloid inflammation in the skin, spleen, or bone marrow of young mice (Fig. 4, A to C), indicating that skin adipocytes were not the source of immune dysregulation caused by the NF- κ B inhibition. In contrast, induced *Ikkb* deletion in Col1a2⁺ fibroblasts resulted in histological changes characterized by a mild increase in myeloid and eosinophilic cells, as well as enhanced F4/80⁺ cell numbers without inducing myeloid cell changes in the spleen or bone marrow (Fig. 4, D to F). These results rule out skin adipocytes as the source of immune dysregulation caused by the NF- κ B inhibition. We further tested whether postnatal Prx1⁺ (pnPrx1⁺) cells are responsible for maintaining immune homeostasis. In neonatal mice, pnPrx1⁺ cells represent early mesenchymal progenitors, and lineage tracing demonstrated that they became skin fibroblasts and subcutaneous adipocytes at 4 weeks (Fig. 4G). Induced deletion of *Ikkb* in P1 mice elicited mild skin inflammation at 4 weeks

(Fig. 4, H and I), whereas induced *Ikkb* deletion in 8-week-old mice failed to produce inflammation in mice aged 12 or 24 weeks despite the similar pnPrx1⁺ cell frequency for each induction time point (fig. S4). These results indicate that there is a critical perinatal growth period during which immune homeostasis is established in the skin and that Prx1⁺ skin fibroblasts, not adipocytes, are essential for this to occur through basal *Ikkb* activity.

To investigate the transcriptomic shift caused by the *Ikkb* deletion in Prx1 lineage cells, we performed scRNA-seq on skin cells from 4-week-old *Prx1Cre⁺.Ikkb^{fl/fl}* mice (4485 cells) and controls (6599 cells). We applied iterative random forest leave one out prediction (iRF-LOOP) (23), an explainable artificial intelligence method, to the combined samples by using each cell's gene expression as a feature vector to create a predictive cell expression network. Next, Markov clustering (MCL) (24) was applied at 1.30 inflation to the top 0.4% of edges in this network to create 316 separate clusters of cells (fig. S5), in which the gene expression profile of each cell within a cluster was highly predictive of the other cells in that cluster (data file S1). Cell types were probabilistically assigned to each cell via their marker genes by applying the CellAssign R package (25). This cell typing was projected onto the iRF-LOOP+MCL clusters and visualized using Cytoscape (Fig. 5A) (26). This high-granularity separation of the cells enabled us to identify group-based distributions between the control and *Ikkb*-deleted conditional knockout (cKO) groups (Fig. 5B).

To determine intercluster relatedness, we performed hierarchical clustering analysis with Ward's linkage method (27) on the mean expression vectors of each iRF-LOOP+MCL cluster and projected this separation onto a t-distributed stochastic neighbor embedding (t-SNE) plot of these vectors. Among the three major fibroblast groups identified by the hierarchical clustering (Fig. 5C), fibroblast group 2 was largely found in the *Ikkb* deletion group (Fig. 5D), indicating a substantial change in the transcriptome in fibroblasts caused by the *Ikkb* deletion. Fibroblasts from the control group had subtypes similar to those based on the work of Joost *et al.* (28), while the majority of fibroblasts from the *Ikkb*-deleted group exhibited a transcriptome profile similar to hypodermal fibroblasts (Fig. 5E). Furthermore, we used each iRF-LOOP+MCL cluster identity as a class label and used iRF to obtain an ordered list of genes predictive of each cluster. The largest fibroblast clusters were predicted by the expression of *Palld*, a microfilament gene ubiquitously expressed in mesenchymal cells (29) and by *Pi16*, a gene that regulates neuropathic pain (Fig. 5F) (30). In contrast, the top two fibroblast clusters from the *Ikkb*-deleted mice were predicted by *Col6a5* and *Coch*, which are highly up-regulated in human AD and in the inner ear, respectively (Fig. 5F) (31, 32). These results suggest that the fibroblast phenotypes in the *Ikkb* deletion and control groups are functionally distinct.

We next hypothesized that *Ikkb* deletion in Prx1 lineage fibroblasts caused a severe immune cell infiltration by aberrant chemokine expression. Pairwise differential expression comparisons between all fibroblast clusters in the control and *Ikkb* deletion groups were examined using the DESeq2 R package (33), resulting in 88 such comparisons (data file S2). Several chemokines were up-regulated in fibroblasts with *Ikkb* deletion, including *Ccl7*, *Cxcl12*, *Ccl8*, *Ccl19*, and *Ccl11* (Fig. 6A). Among these, three chemokines (*Cxcl12*, *Ccl11*, and *Ccl7*) were up-regulated in *Ikkb*-deleted fibroblasts with high *Prrx1* unique molecular identifier (UMI) count (constituting 42% of fibroblasts) compared to those in control group (48% of




Fig. 3. Skin inflammation in *Ikkb*-deleted mice is characterized by excessive myeloid cell infiltration. (A) Representative immunofluorescent images with myeloperoxidase antibody (MPO; a neutrophil marker) and F4/80 antibody (a pan-macrophage marker) in 4-week-old WT or *Prx1Cre⁺Ikkb^{fl/fl}* (cKO) mice. Scale bars, 50 μ m. (B) Quantification of immunopositive cells for MPO, F4/80, or CD4 expression in 4-week-old (left) or 20-week-old (right) mice between WT and cKO groups; IF, immunofluorescence experiment. $n = 6$ to 8 each. (C) Quantification of MPO⁺ cells in the ventral skin of 1-day-old (P1) mice between WT and cKO groups. $n = 4$ to 6 each. (D) Representative flow cytometry plot on myeloid cell gating (live, CD45⁺CD11b⁺ cells) to analyze Gr-1^{Hi} neutrophils and F4/80⁺ myeloid cells in 4-week-old mice. (E) Quantification of neutrophils (CD45⁺CD11b⁺F4/80⁻Gr-1^{Hi}), F4/80⁺ myeloid cells (CD45⁺CD11b⁺F4/80⁺Gr-1⁻), and CD3 T cells (SSC^{low}CD45⁺CD3) per total live cells in 4-week-old (left) or 20-week-old mice (right) by flow cytometry analysis (FC). (F) Quantification of neutrophils and F4/80⁺ myeloid cells from 4-week-old mice that were weaned in soft paper cage bedding. (G) Quantification of circulating hematopoietic cells from retro-orbital blood samples of the control (WT) or *Prx1Cre⁺Ikkb^{fl/fl}* (cKO) mice aged at 20 weeks old by complete blood count/differential analysis. WBC, white blood cell; RBC, red blood cell; $n = 6$ to 9 each. (H) Representative micrograph of spleen from 20-week-old WT and cKO mice (left; scale bars, 1 cm). Right: Quantification of spleen weight per weight. (I) Spleen cryosections stained for T cells (CD3 antibody) and B cells (B220 antibody) demonstrating intact germinal centers in both WT and cKO mice. Scale bars, 0.5 mm. (J and K) Quantification of neutrophils (Neut; CD45⁺CD11b⁺Gr-1^{Hi}), monocytes (MO; CD45⁺CD11b⁺CD115⁺), macrophages (M Φ ; CD45⁺CD11b⁺F4/80⁺), and eosinophils (Eos; CD45⁺CD11b⁺Siglec-F⁺) per total live cells from splenocyte preparations (J) and from bone marrow preparations (K) of WT and cKO mice; $n = 5$ to 6 each. Data are represented as means \pm SEM of biological replicates. Animal experiments were repeated three times. * $P < 0.05$, Student's *t* test comparing WT to cKO groups.

Downloaded from https://www.science.org at University of Pennsylvania on June 14, 2022




Fig. 4. *Ikkb* deletion in skin fibroblasts, not adipocytes, is responsible for myeloid inflammation during perinatal growth. (A) H&E images of ventral skin in 4-week-old control group (WT) or adipocyte-specific *Ikkb* deletion (*Adipoq-Cre⁺Ikkb^{ff/ff}*) mice. Scale bars, 50 μ m. (B) Flow cytometry analysis of neutrophils (CD45⁺CD11b⁺Gr-1^{hi}) and F4/80⁺ myeloid cells (CD45⁺CD11b⁺F4/80⁺) from enzymatically digested ventral skin of WT or *Adipoq-Cre⁺Ikkb^{ff/ff}* mice. (C) Quantification of neutrophils and F4/80⁺ myeloid cells per total live cell count in ventral skin (left), spleen (middle), and bone marrow (right) comparing WT to adipocyte-specific deletion of *Ikkb* group. $n = 7$ to 8 each. (D) H&E images of ventral skin in 4-week-old WT or experimental mice that had induced deletion of *Ikkb* in Col1a2-expressing fibroblasts (*Col1a2CreER⁺Ikkb^{ff/ff}*). Mice received tamoxifen twice at P1 and P3 (50 μ g per dose) and were euthanized at 4 weeks of age. Red arrows point at monocytic and/or eosinophilic cells. Scale bars, 50 μ m. (E) Flow cytometry plot for neutrophils and F4/80⁺ myeloid cells in WT or mice with *Ikkb* deletion in Col1a2⁺ fibroblasts. Tamoxifen was administered at P1 and P3 (intragastric, 50 μ g per dose). Representative plots were gated for CD11b⁺ cells. (F) Quantification of neutrophils and F4/80⁺ myeloid cells per total live cell count in ventral (left), spleen (middle), and bone marrow (right) comparing WT to *Ikkb* deletion in Col1a2-expressing pan-fibroblasts. $n = 5$ to 8 each. (G) Perinatal induction of *dTomato* reporter gene in postnatal Prx1⁺ cells (*Prx1CreER⁺R26R^{dTomato}*), labeling dermal mesenchymal cells and some subcutaneous adipocytes (insert, arrows; scale bar, 20 μ m) after 4-week tracing period. Scale bar, 0.5 mm. (H) Immunofluorescent images of F4/80⁺ cells in the ventral skin of 4-week-old WT mice and *Prx1CreER⁺Ikkb^{ff/ff}* mice that had perinatal deletion of *Ikkb* (left) and quantification of F4/80⁺ cells per square millimeter (right). $n = 8$ to 10 each. Both WT and *Prx1CreER⁺Ikkb^{ff/ff}* mice received tamoxifen. Scale bars, 50 μ m. (I) Flow cytometry analysis (FC) for F4/80⁺ myeloid cells from ventral skin of WT and *Prx1CreER⁺Ikkb^{ff/ff}* mice. $n = 5$ to 10 each. Data are represented as means \pm SEM of biological replicates. Animal experiments were repeated three times. * $P < 0.05$, Student's *t* test comparing WT to each respective experimental group.

fibroblasts) (Fig. 6A and fig. S6A; full data in data file S3). Gene ontology analysis revealed an up-regulation of biological processes related to cell migration, cytokine response, and inflammatory response in the *Prx1^{hi}* fibroblasts of the *Ikkb*-deleted group (Fig. 6B). Projecting gene expression heatmaps of *Cxcl12*, *Ccl11*, and *Ccl7* onto the t-SNE visualization of the iRF-LOOP+MCL clusters revealed that *Cxcl12* and *Ccl7* are highly expressed by endothelial cells and macrophages, respectively. In contrast, *Ccl11* expression was exclusive to

fibroblasts and enhanced in the *Ikkb* deletion clusters (Fig. 6C and fig. S6B). Up-regulation of *Ccl11* in fibroblasts from the *Ikkb*-deleted mice was verified by the Seurat pipeline (Fig. 6D and fig. S6, C to H) (34). Immunofluorescence experiments demonstrated that CCL11⁺ cells with fibroblastic morphology were enriched in the ventral skin of the *Ikkb*-deleted group and were localized to the border between the dermis and subcutaneous adipose layer (Fig. 6E). Because CCL11 is a potent chemoattractant for eosinophils, we quantified the





Fig. 5. scRNA-seq reveals marked changes in the fibroblast transcriptome caused by *Ikkb* gene deletion. (A) iRF-LOOP+MCL (iterative random forest–Markov clustering) of 11,084 cells from enzymatically digested ventral skin of WT and *Prx1^{Cre+}Ikkb^{fl/fl}* mice with cell type designation based on putative gene expression. Clusters with >50 cells are shown. NKT cells, natural killer T cells; NK cells, natural killer cells. (B) WT (blue) and cKO (*Prx1^{Cre+}Ikkb^{fl/fl}*, red) group designation demonstrating even distribution of sample (3566 cells from WT and 2510 cells from cKO are shown). (C) Hierarchical partitioning of MCLs and cell type designation using Ward's linkage method between Pearson correlations of mean expression vectors of the clusters. The groups generated by this partitioning have been t-SNE-projected in (D) and share the same color legend. (D) t-SNE plot with cell type designation (left) and WT/cKO designation (right) demonstrating clear separation of fibroblast clusters (encircled in dashed line) derived from *Ikkb*-deleted group. Mixed WT/cKO cluster threshold was determined at <0.90 cutoff. (E) Fibroblast subtyping based on (36) demonstrating a transcriptomic shift toward that of hypodermal fibroblasts when *Ikkb* is deleted (left of the dashed line). (F) Importance score for genes that determined the formation of top 2 clusters from WT and cKO groups as characterized in (A) and (B).

number of eosinophils in ventral skin and found that *Ikkb* deletion caused a marked increase in F4/80⁺Siglec-F⁺ eosinophil numbers (Fig. 6F). Because the F4/80⁺ myeloid cells that we detected in Fig. 3 could be either macrophages or eosinophils, we examined the

coexpression of markers associated with classically or alternatively activated macrophages (CD80 and CD206, respectively) and Siglec-F for eosinophilic identity and found that over 80% of F4/80⁺ cells were identified as eosinophils in the ventral skin of *Ikkb* deletion mice

Fig. 6. *lkkb* deletion causes excessive *Ccl11* expression by fibroblasts and leads to subsequent eosinophilia.

(A) Pairwise comparisons of chemokine genes from each fibroblast clusters (top) and from *Prx1*^{high} fibroblasts, comparing WT to *Prx1*^{Cre}*lkkb*^{fl/fl} (cKO) group. Notable up-regulated chemokine genes were *Cxcl12*, *Ccl11*, and *Cd7*. LFC2, log₂ fold change. (B) Gene ontology (GO) analysis for biological process related to chemotaxis and inflammation based on genes up-regulated in *Prx1*^{high} fibroblasts from cKO group compared to that of WT. (C) t-SNE plots of iRF-LOOP+MCL clusters with *Cxcl12*, *Cd7*, and *Ccl11* gene expression with notable up-regulation of *Ccl11* in fibroblast clusters derived from cKO group. (D) Volcano plot of genes up-regulated in WT and cKO groups using each fibroblast as a data point through single-cell analysis pipeline established by Seurat and DESeq2 packages. (E) In vivo validation of CCL11 protein up-regulation by immunofluorescence with CCL11 antibody in ventral skin of 4-week-old WT and cKO mice. Scale bars, 200 μm. Insert images (orange box) show CCL11⁺ cells at the border of dermis and subcutaneous layers, as indicated by the arrows. Scale bars, 50 μm. Representative images are from *n* = 7 each. Isotype IgG control shows autofluorescence in hair follicles. (F) Flow cytometry analysis of eosinophil count (CD45⁺CD11b⁺F4/80⁺SiglecF⁺) per total live cells, comparing WT to cKO groups. (G) Percentage of SiglecF⁺ eosinophils, classically activated macrophages (M1, CD80⁺), and alternatively activated macrophages (M2, CD206⁺) per total CD45⁺CD11b⁺F4/80⁺ cells. (H) t-SNE plot for 5136 fibroblasts (3609 WT and 1527 cKO) in the scRNA-seq dataset, comparing both WT versus cKO (left) and *Cebpb* UMI counts (normalized and log₁₀-transformed) (right). (I) Immunofluorescence with antibodies specific to CEBPB and PDGFRA (platelet-derived growth factor receptor alpha) in 4-week-old WT and experimental mice (right) and quantification of CEBPB⁺PDGFRA⁺ fibroblasts. Scale bars, 50 μm. (J) Quantitative real-time polymerase chain reaction (RT-qPCR) for *lkkb*, *Cebpb*, and *Ccl11* mRNA in skin fibroblasts with no treatment (sham), scrambled siRNA control (Scr), or *lkkb* siRNA (20 nM). Cells were isolated from ventral skin of 2-week-old *Prx1*^{Cre}*R26R*^{dTomato} mice and stimulated with low-dose IL4 (5 ng/ml) for 12 to 16 hours before analysis. (K) Flow cytometry analysis of *Prx1*⁺ fibroblasts for mean fluorescent intensity of CEBPB (left) or CCL11⁺ fibroblast numbers (right). Cells were gated for live/dTomato⁺ expression. (L) Quantification of CCL11⁺ fibroblasts after knockdown with *Cebpb* siRNA alone, *lkkb* siRNA alone, and *Cebpb*+*lkkb* siRNA. (M) Quantification of *CCL11* mRNA by RT-qPCR with *IKKB* knockdown alone or *IKKB*+*CEBPB* knockdown in human neonatal fibroblasts without IL4 treatment. (F and G) *n* = 6 each. (J to M) *n* = 3 each. Data are represented as means ± SEM of biological replicates. In vitro experiments were repeated twice, and animal experiments were repeated three times. **P* < 0.05, Student's *t* test comparing WT to cKO groups (F and G) or one-way ANOVA followed by post hoc test (J to M).



Downloaded from https://www.science.org at University of Pennsylvania on June 14, 2022

(Fig. 6G). Thus, *Ikkb* deletion in skin fibroblasts causes aberrant expression of CCL11, which is associated with overt eosinophilia in early skin inflammation.

We next explored possible mechanisms by which *Ikkb* deletion prevents NF- κ B activation to paradoxically induce inflammation. scRNA-seq data pointed to *Cebpb* (CCAAT/enhancer-binding protein β), a basic leucine zipper domain transcription factor implicated in inflammation (35), as highly up-regulated in fibroblasts with *Ikkb* deletion (Fig. 6H). Increased CEBPB protein expression by the fibroblasts with *Ikkb* deletion was validated by immunofluorescence studies (Fig. 6I). To test direct modulation of *Cebpb* and *Ccl11* expression, we examined Prx1⁺ fibroblasts from 2-week-old *Prx1^{Cre+}.Rosa^{dTomato}* mice and found that *Ikkb* knockdown with small interfering RNA (siRNA) enhanced CEBPB and CCL11 mRNA and protein expression in vitro (Fig. 6, J and K). We then used a publicly available chromatin immunoprecipitation sequencing database [Signaling Pathways Project (36)] and noted high binding scores of CEBPB to CCL11 promoter/enhancer region in mesenchymal stromal cells (table S1), suggesting direct transcriptional regulation. Concomitant knockdown of *Ikkb* and *Cebpb* by siRNA significantly reduced CCL11 up-regulation, as noted by the diminished numbers of CCL11⁺Prx1⁺ fibroblasts compared to *Ikkb* knockdown alone ($P < 0.05$; Fig. 6L), indicating that CEBPB is an important mechanistic link between *Ikkb* deletion and CCL11 up-regulation in these cells. To show relevance with human fibroblasts, we examined *CCL11* expression after *IKBKB* and *CEBPB* knockdown in two independent neonatal human fibroblast cell lines and found analogous results (Fig. 6M and fig. S6I).

We next examined the alteration in immune cell profiles by *Ikkb* deletion in fibroblasts. Within scRNA-seq data, type 2 lymphocytes [T helper 2 (T_H2)] constituted the largest proportion of lymphocytes in mice with *Ikkb* deletion, and their absolute numbers were also elevated (Fig. 7A). The largest leukocyte cluster (cluster 3) was mostly derived from the *Ikkb*-deleted group, and the formation of this cluster by iRF-LOOP+MCL was predicted by transcripts representing type 2 immunity (*Il5*, *Areg*, and *Il13*) (Fig. 7B and fig. S7A). Seurat analysis verified this result and identified type 2 cytokines (*Il5*, *Areg*, and *Il13*), *Il1rl1* (encoding for ST2 receptor) and transcription factor *Gata3* as up-regulated transcripts in *Ikkb*-deleted group (Fig. 7, C and D, and fig. S7, B to F). Other notable findings included myeloid cells from the cKO mice exhibiting increased proinflammatory *Il1b* gene and reduced proresolving *Tgfb1* gene signatures (Fig. 7E) and mast cells from the cKO group with increased expression of *Hdc* and *Fcer1a* (fig. S8, A to C), which are genes encoding for histamine decarboxylase and the Fc-IgE (immunoglobulin E) receptor, respectively, that are implicated in pruritus (37). In addition, serum IgE concentration was significantly higher in mice with *Ikkb* deletion ($P < 0.05$; fig. S8D), a characteristic finding in patients with AD (20). On the basis of the scRNA-seq results, we next hypothesized that early skin lesions are characterized by T_H2 in addition to eosinophils, which worsens as mice age from 4 weeks (young) to 20 weeks (adult). We found that high eosinophil counts from young *Ikkb*-deleted mice were reduced with time (Fig. 7F). In contrast, CD3⁺CD4⁺IL4⁺ T_H2 cells increased significantly as the mice aged ($P < 0.05$; Fig. 7F), although there was no overall increase in CD4⁺ lymphocytes (Fig. 3B). Because type 2 immunity and eosinophilia are classic signs of human AD (20), we investigated whether CCL11 is overexpressed by the fibroblasts in human AD lesions. Using RNAScope, we examined validated human AD skin with sex- and age-matched control and found that AD skin had a significant increase in the absolute and

relative numbers of CCL11-expressing fibroblasts (*CCL11⁺PDGFRA⁺*) per total *PDGFRA⁺* fibroblasts ($P < 0.05$; Fig. 7, G and H, and fig. S9A). Moreover, human AD skin had significantly higher numbers of *CCL11⁺CEBPB⁺* cells ($P < 0.05$; Fig. 7J and fig. S9B). This was consistent with our murine model of dysregulated fibroblasts as a source of CCL11 for early disease progression in AD-like lesions. We further examined published scRNA-seq data from human AD skin (10) for potential chemokine dysregulation by *PRRX1⁺* mesenchymal cells and found that *IKBKB* was down-regulated, whereas *CEBPB* and *CCL11* were up-regulated ($P < 0.05$; table S3 and data file S4), in agreement with our observations.

To test whether the aberrant up-regulation of CCL11 by *Ikkb*-deleted fibroblasts is responsible for downstream events of inflammation, we performed an in vivo neutralization experiment by injecting a monoclonal CCL11 antibody into 2-week-old mice (Fig. 8A) and examined the effect on preventing eosinophilia and T_H2 responses. Antibody treatment effectively reduced the number of eosinophil infiltrations by >50% in the skin of the *Ikkb*-deleted mice (Fig. 8, B to D). Subsequently, the number of T_H2 cells was significantly abrogated when CCL11 was inhibited in the skin of *Ikkb*-deleted mice ($P < 0.05$; Fig. 8, E to F), demonstrating a therapeutic effect of CCL11 neutralization on the degree of eosinophilia and T_H2 response in early AD-like skin lesions.

DISCUSSION

Epithelial barrier disruption and immune dysregulation are thought to be the primary etiologic factors in chronic inflammatory skin conditions such as AD (2, 20), based on the classical experimental models using tape-stripping, ectopic allergen, and/or gene deletion for immune function (38). Here, we show that skin fibroblasts, in addition to their traditional role in skin repair, are also critical regulators of immune homeostasis during early perinatal growth, which is regulated by *Ikkb* under basal physiologic conditions. Fibroblasts in which activation of the canonical NF- κ B pathway is blocked by *Ikkb* gene deletion resulted in a skin phenotype that mirrors human AD-like skin lesions that exhibit eosinophilia and large numbers of type 2 immune cells (2, 20, 39). The lesion was specific to Prx1⁺ fibroblast-enriched ventral skin, an area where mice have limited ability to directly scratch. Because regular corn bedding can be abrasive to murine skin (40), we housed mice in a soft bedding material to reduce exogenous irritation but found that it had no rescue effect. Thus, our mouse model likely reflects an “inside-out” mechanism of AD (20), where dysregulation of endogenous fibroblasts initiates a subsequent immune cascade, as opposed to external stimulation (“outside-in,” for example, by scratching). The AD-like lesions that we observed in Prx1-driven *Ikkb*-deleted mice is corroborated by a recent study with Nestin-driven *Ikkb* in facial mesenchymal cells, both of which target phenotypically similar fibroblastic cells (41). Although the latter study did not identify any mechanisms that led to the dermal lesions, Nestin⁺ and Prx1⁺ fibroblastic cells are thought to have several similarities (42). Moreover, we found that *Ikkb* deletion in adipocytes had minimal effect on subcutaneous inflammation in young mice as it did in another study that examined 16-week-old adult mice (43). We showed that *Ikkb*-deleted fibroblasts aberrantly expressed CCL11 to initiate an eosinophilic and T_H2 inflammation and that blocking CCL11 effectively reduced eosinophilia and type 2 immune response. Collectively, these data demonstrate that Prx1⁺ fibroblasts may have a unique role in maintaining homeostasis




Fig. 7. *Ikkb* deletion in fibroblasts causes a shift in inflammatory phenotype toward type 2 immune response, mimicking atopic skin in humans. (A) Leukocyte frequency distribution between control (WT) and *Prx1^{Cre+}Ikkb^{ff}* (cKO) groups demonstrating marked increase in Th2 proportion in cKO group. (B) Importance scores of genes that determined the largest immune cell cluster (cluster 3) that is predominated by cells of cKO group. *Il5*, *Areg* and *Il13* are classic type 2 immune cytokines. (C) Seurat clustering of immune cells from WT (blue) and cKO (red) mice; myeloid and lymphoid cell designation was determined by their putative gene expression as shown in fig. S7. (D) Volcano plot of genes up-regulated in lymphocyte population of WT and cKO groups using Seurat and DESeq2 packages. (E) Violin plot of *Il1b* and *Tgfb1* in myeloid cell population between WT and cKO mice. (F) Flow cytometry analysis of skin eosinophils and CD3⁺CD4⁺IL4⁺ Th2 cells from young (4-week-old) and adult (20-week-old) WT and cKO mice. *n* = 6 to 8 each. (G) Representative RNAScope images of human control and atopic skin specimens detecting CCL11 (red) and PDGFRA (green dots) signals in dermis. Dashed line demarcates epidermal-dermal junction. Black arrows point to some PDGFRA⁺ fibroblasts and red arrows to CCL11⁺ cells. Scale bars, 50 μm. (H) Quantification of double CCL11⁺PDGFRA⁺ cells per total spindle-shaped cells (left) and percentage of CCL11⁺ expressing fibroblasts by normalizing to total PDGFRA⁺ cells. (I) Quantification of CCL11⁺CEBPB⁺ spindle-shaped cells in matched control versus human AD skin. *n* = 8 each. Data are represented as means ± SEM of biological replicates. Animal experiments were repeated three times, and RNAScope was repeated twice. **P* < 0.05; (F) two-way ANOVA followed by Tukey's post hoc test; (H and I) Student's *t* test.

during development and that disruption of *Ikkb* in these cells may lead to the formation of dermal lesions that share many similarities to human AD. Moreover, the results suggest that blocking CCL11 is a potentially important therapeutic approach to treating early inflammatory skin conditions.

Pediatric patients have a propensity for AD and other inflammatory skin conditions (44), suggesting a disease mechanism linked to early development. We were able to detect histologic changes before the gross formation of skin lesions in mice, providing new insight

into the etiologic factors. Skin inflammation caused by *Ikkb* deletion occurred during perinatal development. The changes were not due to age-dependent reduction in pnPrx1⁺ cells because their frequency in the skin dermis was similar in both neonatal and adult mice. Rather, these findings suggest that early perinatal growth is a critical time period during which skin fibroblasts with basal NF-κB activity are essential for their immunoregulatory function, consistent with the AD susceptibility for children (20, 44, 45). Our findings highlight the previously unknown immunoregulatory function of young




Fig. 8. CCL11 neutralizing antibody treatment in young mice with *Ikkb* deletion reduces eosinophil infiltration and T_H2 cell numbers. (A) Experimental design for CCL11 neutralization in young (2- to 4-week-old) mice. (B) Flow cytometry plot of eosinophils ($F4/80^+Siglec-F^+$) from enzymatically digested ventral skin of control (WT) that received isotype IgG, *Prx1Cre⁺Ikkb^{-/-}* (cKO) mice that received either isotype IgG or CCL11 neutralizing antibody. (C) Quantification of eosinophils per total live cells comparing WT+IgG, cKO+IgG, and cKO+CCL11 neutralizing antibody group. (D) H&E images of ventral skin from WT mice that received control IgG and cKO mice that received IgG or CCL11 neutralizing antibody. Scale bars, 0.5 mm. Red arrows point to eosinophils. Insert scale bars, 50 μ m. (E) Flow cytometry plot of T_H2 cells ($CD4^+IL4^+$) on live $CD45^+CD3^+$ gated cells. (F) Quantification of T_H2 percent per total $CD3^+$ T cells (left) and that of T_H2 absolute count per 10^4 cells (right). Data are represented as means \pm SEM of biological replicates. Animal experiments were repeated three times. * $P < 0.05$, one-way ANOVA followed by Tukey's post hoc test. Ab, antibody; FMO, fluorescence-minus-one control.

fibroblasts without which skin immunity fails to stabilize and causes severe inflammatory skin condition that closely mimics human atopy.

NF- κ B activation is critical in modulating inflammatory immune responses (13), and its inhibition in chronic inflammatory conditions such as diabetes and autoimmunity alleviates the disease outcome (46, 47). However, we demonstrate that preventing NF- κ B activation in skin fibroblasts paradoxically induces severe skin inflammation, contrary to the expected anti-inflammatory outcome. Despite the well-established proinflammatory role of NF- κ B, there are recent examples where conditional deletion of *Ikkb* interferes with

homeostasis. This has been shown in hepatocytes for glucose homeostasis (48) and for steady-state function of migratory dendritic cells (49). Thus, our results further strengthen the concept that, in addition to mediating inflammatory responses, NF- κ B plays an essential role in maintaining homeostasis under physiologic conditions. In the dermis, NF- κ B specifically is needed in *Prx1⁺* fibroblasts to prevent the overexpression of factors that generate a dermal inflammatory condition. For example, CEBPB was highly up-regulated when *Ikkb* was deleted or knocked down in *Prx1⁺* fibroblasts in vivo and in vitro. CEBPB enhances transcription of inflammatory

cytokines (50) and is predicted to directly bind to *Ccl11* promoter region in stromal cells (table S1). Moreover, CEBPB may interact with STAT proteins to indirectly up-regulate CCL11 in fibroblasts (51). Double knockdown of *Ikkb* and *Cebpb* rescued *Ccl11* up-regulation induced by *Ikkb* knockdown alone, suggesting a modulatory role of *Ikkb*-NF- κ B axis on CEBPB and subsequent up-regulation of CCL11 expression by neonatal fibroblasts.

Here, we have focused on aberrant CCL11 expression as a potent etiologic factor for eosinophil recruitment. We did not capture eosinophils in the scRNA-seq experiments because of our initial exclusion of high side-scatter granulocytes through cell sorting. Nevertheless, using other approaches, we observed that the eosinophil population made up to 20 to 40% of cells obtained from the skin of young *Ikkb*-deleted mice and that eosinophil dominance was substantially reduced in 20-week-old mice. In contrast, TH2 population increased with age in adult *Ikkb*-deleted mice, consistent with the overt type 2 immune response seen in human AD (20). Up-regulation of CCL11 had been implicated in human atopic skin (52), although the cellular source is not well understood. In results similar to our findings, dermal fibroblasts from human atopic skin have been shown to exhibit higher CCL11 mRNA expression compared to control groups in response to IL4 (interleukin-4) stimulation in vitro (53). The etiology of human AD has been elusive, possibly because of the difficulty in detecting the subclinical phase of AD. The current therapeutics for moderate to severe AD use topical immunosuppressive drugs or monoclonal antibody against IL4R α (dupilumab) to suppress TH2 response (54). Our data suggest that alternative approaches to block eosinophilia through chemokine antagonists will reduce TH2 response. In this regard, clinical trials are ongoing using bertilimumab, a humanized antibody against CCL11, to treat bullous pemphigoid (NCT02226146), an autoimmune skin disorder characterized by eosinophilia (55). Neutralization of CCL11 in early AD may be a key to preventing emergence of clinical AD symptoms and possibly atopic march.

scRNA-seq studies on human AD samples have recently been reported (9, 10). When PRRX1⁺ mesenchymal cells were specifically examined, results consistent with our studies were observed, including a decrease in *IKBKB* and increases in *CEBPB* and *CCL11*. In contrast, another human study had *CCL11* values that were too low to analyze (9). We also found in human specimens that *CEBPB* was highly up-regulated in AD skin and in mice with *Ikkb* deletion. These studies support an immunomodulatory role of Prx1⁺ cells for eosinophil trafficking to the skin. Fibroblasts in developing human skin exhibit more pronounced inflammatory gene signatures than adult skin (10), suggesting an age-dependent mechanism. Our data indicate that the *Ikkb*-NF- κ B signaling axis plays a role in CEBPB-dependent CCL11 production in dermal fibroblasts. However, additional immune mechanisms may also contribute to lesion formation.

Our study has limitations. We found that reduced *Ikkb*-NF- κ B signaling through gene deletion triggered AD-like lesion formation in mice and that human fibroblasts from AD lesions had reduced *IKBKB* expression. However, the mechanisms that may be responsible for reduced *Ikkb*-NF- κ B signaling in human dermal fibroblastic cells are unknown. One possibility is AD susceptibility due to genetic mutation in *IKBKB*, similar to that of filaggrin loss-of-function mutation (20). We were able to trigger preclinical etiologic events in mice and temporally follow the steps that occur before the formation of established dermal lesions. Whether the same temporal sequence of events occurs in human AD lesions is not known and will be

difficult to establish because of their preclinical nature. However, the studies provide insight into the cellular dysregulation that can lead to a type 2 lesion in the skin.

In summary, there is similarity between our mouse model of AD-like lesions and human AD, demonstrating the participation of the immunoregulatory transcriptome of Prx1⁺ skin fibroblasts in lesion formation. Because AD-like lesions arise from *Ikkb*-NF- κ B perturbation in murine fibroblasts, it will be important to investigate similar mechanism in the early stages of human AD.

MATERIALS AND METHODS

Study design

The objective of this study was to investigate pathogenic mechanisms by which perturbation of *Ikkb*-NF- κ B signaling in skin fibroblasts causes an inflammatory skin disorder that closely mirrors human AD. Mice that lacked *Ikkb* gene in Prx1⁺ mesenchyme were examined from neonatal to adult stages to identify initial event that preceded skin lesion development. Fibrotic etiology was assessed by biomechanical testing. The extent and type of skin inflammation were examined by flow cytometry and immunofluorescence, and systemic inflammation was examined by complete blood count with differential analysis and by multicolor flow cytometry of splenic and bone marrow cells. To test whether the inflammatory effects were due to Prx1-derived skin fibroblasts or adipocytes, mice with lineage-specific deletion of *Ikkb* in pan-fibroblasts or adipocytes were examined. Prx1⁺ fibroblasts were further examined by scRNA-seq and iRF-LOOP analyses, which revealed a marked shift in chemotactic transcripts by the skin fibroblasts that lacked *Ikkb* gene. Further bioinformatics and flow cytometry analyses validated CCL11 as a primary component responsible for early eosinophilia that later transitioned to type 2 immunity, both of which are implicated in human AD. To establish clinical linkage, we used vetted human skin specimens to investigate whether human AD lesions overexpressed CCL11 in fibroblasts by using in situ hybridization. We further tested for translational value of targeting CCL11 by preclinical treatment with neutralizing antibody in young mice and examined its impact on early eosinophilia and type 2 immunity. The study was carried out following ARRIVE (Animal Research: Reporting of In Vivo Experiments) guidelines. Study size was calculated by power analysis using each animal as an experimental unit. Animals were randomly allocated to either control or experimental group. Results were analyzed in a blinded fashion such that the researcher was unaware of group designations. Validation of *Ikkb* deletion in animal models and flow cytometry gating strategies are provided in figs. S10 and S11, respectively.

Animal studies

All animal experiments were carried out conforming to a protocol approved by the University of Pennsylvania Institutional Animal Care and Use Committee (IACUC no. 804855). The following mice were purchased from the Jackson Laboratory: *B6.Cg-Tg(Prrx1-cre)1Cjt/J (Prx1Cre)*, *B6.Cg-Tg(Prrx1-cre/ERT2,-EGFP)1Smkm/J (Prx1CreER)*, *B6.Cg-Gt(ROSA)26Sor^{tm9(CAG-tdTomato)Hze/J (R26^{TdTomato})}*, *Tg(Adipoq-cre)1Evdv (Adipoq-Cre)*, and *Tg(Col1a2-cre/ERT,-ALPP)7Cpd/J (Col1a2-CreER)*. Floxed *Ikkb* mice (*Ikkb^{f/f}*) were obtained from M. Karin (University of California, San Diego). *Ikkb* deletion by Cre-mediated recombination was validated as in fig. S10. Mice were housed in regular corn bedding and were euthanized at 1-day-old, 4-week-old, and 16- to 20-week-old time points. In another set of experiments, animals

were housed in Cellu-nest (Shepherd Specialty Papers) soft bedding from birth to the fourth week euthanasia time point.

Statistical analysis

Statistical analysis was performed using Prism software (GraphPad v9). All data are expressed as the means \pm SEM. Power analysis with $\alpha = 0.05$ and 40% effect was used to determine the sample size. Unpaired Student's *t* test was performed in comparing two groups (wild-type versus *Ikkb*-deleted mice) under the assumption of normal Gaussian distribution. For experiments comparing among three or more groups, one-way analysis of variance (ANOVA) or two-way ANOVA followed by Tukey's post hoc analysis was used as indicated in the figure legends. $P < 0.05$ was considered statistically significant. The individual animal was used as a unit of measurement (*n*) and replicated twice or thrice as indicated in the figure legends, and sex was not considered as a factor in the analyses. Further details are provided in the Supplementary Materials.

SUPPLEMENTARY MATERIALS

www.science.org/doi/10.1126/scitranslmed.abj0324

Materials and Methods

Figs. S1 to S12

Tables S1 to S3

Data files S1 to S5

References (56–59)

[View/request a protocol for this paper from Bio-protocol.](#)

REFERENCE AND NOTES

- C. Karimkhani, R. P. Dellavalle, L. E. Coffeng, C. Flohr, R. J. Hay, S. M. Langan, E. O. Nsoesie, A. J. Ferrari, H. E. Erskine, J. I. Silverberg, T. Vos, M. Naghavi, Global skin disease morbidity and mortality: An update from the global burden of disease study 2013. *JAMA Dermatol.* **153**, 406–412 (2017).
- S. Weidinger, L. A. Beck, T. Bieber, K. Kabashima, A. D. Irvine, Atopic dermatitis. *Nat. Rev. Dis. Primers* **4**, 1 (2018).
- J. E. Greb, A. M. Goldminz, J. T. Elder, M. G. Lebwohl, D. D. Gladman, J. J. Wu, N. N. Mehta, A. Y. Finlay, A. B. Gottlieb, Psoriasis. *Nat. Rev. Dis. Primers* **2**, 16082 (2016).
- P. D. Arkwright, C. Motala, H. Subramanian, J. Spergel, L. C. Schneider, A. Wollenberg; Atopic Dermatitis Working Group of the Allergic Skin Diseases Committee of the AAAAI, Management of difficult-to-treat atopic dermatitis. *J Allergy Clin Immunol Pract* **1**, 142–151 (2013).
- M. Pasparakis, I. Haase, F. O. Nestle, Mechanisms regulating skin immunity and inflammation. *Nat. Rev. Immunol.* **14**, 289–301 (2014).
- Y. Rinkevich, G. G. Walmsley, M. S. Hu, Z. N. Maan, A. M. Newman, M. Drukker, M. Januszky, G. W. Krampitz, G. C. Gurtner, H. P. Lorenz, I. L. Weissman, M. T. Longaker, Skin fibrosis. Identification and isolation of a dermal lineage with intrinsic fibrogenic potential. *Science* **348**, aaa2151 (2015).
- T. Leavitt, M. S. Hu, M. R. Borrelli, M. Januszky, J. T. Garcia, R. C. Ransom, S. Mascharak, H. E. desJardins-Park, U. M. Litzenburger, G. G. Walmsley, C. D. Marshall, A. L. Moore, B. Duoto, S. Adem, D. S. Foster, A. Salhotra, A. H. Shen, M. Griffin, E. Z. Shen, L. A. Barnes, E. R. Zielins, Z. N. Maan, Y. Wei, C. K. F. Chan, D. C. Wan, H. P. Lorenz, H. Y. Chang, G. C. Gurtner, M. T. Longaker, Prrx1 fibroblasts represent a pro-fibrotic lineage in the mouse ventral dermis. *Cell Rep.* **33**, 108356 (2020).
- T. Tabib, C. Morse, T. Wang, W. Chen, R. Lafyatis, SFRP2/DPP4 and FMO1/LSP1 define major fibroblast populations in human skin. *J. Invest. Dermatol.* **138**, 802–810 (2018).
- H. He, H. Suryawanshi, P. Morozov, J. Gay-Mimbrera, E. Del Duca, H. J. Kim, N. Kameyama, Y. Estrada, E. Der, J. G. Krueger, J. Ruano, T. Tuschl, E. Guttman-Yassky, Single-cell transcriptome analysis of human skin identifies novel fibroblast subpopulation and enrichment of immune subsets in atopic dermatitis. *J. Allergy Clin. Immunol.* **145**, 1615–1628 (2020).
- G. Reynolds, P. Vegh, J. Fletcher, E. F. M. Poyner, E. Stephenson, I. Goh, R. A. Botting, N. Huang, B. Olabi, A. Dubois, D. Dixon, K. Green, D. Maunder, J. Engelbert, M. Efremova, K. Polanski, L. Jardine, C. Jones, T. Ness, D. Horsfall, J. McGrath, C. Carey, D. M. Popescu, S. Webb, X. N. Wang, B. Sayer, J. E. Park, V. A. Negri, D. Belokhvostova, M. D. Lynch, D. McDonald, A. Filby, T. Hagai, K. B. Meyer, A. Husain, J. Coxhead, R. Vento-Tormo, S. Behjati, S. Lisgo, A. C. Villani, J. Bacardit, P. H. Jones, E. A. O'Toole, G. S. Ogg, N. Rajan, N. J. Reynolds, S. A. Teichmann, F. M. Watt, M. Haniffa, Developmental cell programs are co-opted in inflammatory skin disease. *Science* **371**, eaba6500 (2021).
- A. P. Croft, J. Campos, K. Jansen, J. D. Turner, J. Marshall, M. Attar, L. Savary, C. Wehmeyer, A. J. Naylor, S. Kemble, J. Begum, K. Durholz, H. Perlman, F. Barone, H. M. McGettrick, D. T. Fearon, K. Wei, S. Raychaudhuri, I. Korsunsky, M. B. Brenner, M. Coles, S. N. Sansom, A. Filer, C. D. Buckley, Distinct fibroblast subsets drive inflammation and damage in arthritis. *Nature* **570**, 246–251 (2019).
- T. Liu, C. Han, S. Wang, P. Fang, Z. Ma, L. Xu, R. Yin, Cancer-associated fibroblasts: An emerging target of anti-cancer immunotherapy. *J. Hematol. Oncol.* **12**, 86 (2019).
- M. S. Hayden, S. Ghosh, NF- κ B in immunobiology. *Cell Res.* **21**, 223–244 (2011).
- Q. Zhang, M. J. Lenardo, D. Baltimore, 30 years of NF- κ B: A blossoming of relevance to human pathobiology. *Cell* **168**, 37–57 (2017).
- T. H. Leung, L. F. Zhang, J. Wang, S. Ning, S. J. Knox, S. K. Kim, Topical hypochlorite ameliorates NF- κ B-mediated skin diseases in mice. *J. Clin. Invest.* **123**, 5361–5370 (2013).
- T. Liu, L. Zhang, D. Joo, S. C. Sun, NF- κ B signaling in inflammation. *Signal Transduct. Target. Ther.* **2**, (2017).
- A. Oeckinghaus, M. S. Hayden, S. Ghosh, Crosstalk in NF- κ B signaling pathways. *Nat. Immunol.* **12**, 695–708 (2011).
- M. Logan, J. F. Martin, A. Nagy, C. Lobe, E. N. Olson, C. J. Tabin, Expression of Cre recombinase in the developing mouse limb bud driven by a Prx1 enhancer. *Genesis* **33**, 77–80 (2002).
- J. L. Durland, M. Sferlazzo, M. Logan, A. C. Burke, Visualizing the lateral somitic frontier in the Prx1Cre transgenic mouse. *J. Anat.* **212**, 590–602 (2008).
- S. M. Langan, A. D. Irvine, S. Weidinger, Atopic dermatitis. *Lancet* **396**, 345–360 (2020).
- A. Greenbaum, Y. M. Hsu, R. B. Day, L. G. Schuettelpelz, M. J. Christopher, J. N. Borgerding, T. Nagasawa, D. C. Link, CXCL12 in early mesenchymal progenitors is required for haematopoietic stem-cell maintenance. *Nature* **495**, 227–230 (2013).
- J. Sanchez-Gurmaches, W. Y. Hsiao, D. A. Guertin, Highly selective in vivo labeling of subcutaneous white adipocyte precursors with Prx1-Cre. *Stem Cell Rep.* **4**, 541–550 (2015).
- A. Cliff, J. Romero, D. Kainer, A. Walker, A. Furches, D. Jacobson, A high-performance computing implementation of iterative random forest for the creation of predictive expression networks. *Genes (Basel)* **10**, 996 (2019).
- S. van Dongen, A cluster algorithm for graphs. *Information Systems [INS]* (Centrum Wiskunde & Informatica, 2000); <https://ir.cwi.nl/pub/4463>.
- A. W. Zhang, C. O'Flanagan, E. A. Chavez, J. L. P. Lim, N. Ceglia, A. McPherson, M. Wiens, P. Walters, T. Chan, B. Hewitson, D. Lai, A. Mottok, C. Sarkozy, L. Chong, T. Aoki, X. Wang, A. P. Weng, J. N. McAlpine, S. Aparicio, C. Steidl, K. R. Campbell, S. P. Shah, Probabilistic cell-type assignment of single-cell RNA-seq for tumor microenvironment profiling. *Nat. Methods* **16**, 1007–1015 (2019).
- P. Shannon, A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B. Schwikowski, T. Ideker, Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).
- J. Ward, Hierarchical grouping to optimize an objective function. *J. Am. Stat. Assoc.* **58**, 236–244 (1963).
- S. Joost, K. Annusver, T. Jacob, X. Sun, T. Dalessandri, U. Sivan, I. Sequeira, R. Sandberg, M. Kasper, The molecular anatomy of mouse skin during hair growth and rest. *Cell Stem Cell* **26**, 441–457 e447 (2020).
- M. M. Parast, C. A. Otey, Characterization of palladin, a novel protein localized to stress fibers and cell adhesions. *J. Cell Biol.* **150**, 643–656 (2000).
- P. Singhar, R. T. P. Trinh, J. Ma, X. Huo, B. Peng, C. J. Heijnen, A. Kavelaars, The fibroblast-derived protein P116 controls neuropathic pain. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 5463–5471 (2020).
- C. Soderhall, I. Marenholz, T. Kerscher, F. Ruschendorf, J. Esparza-Gordillo, M. Worm, C. Gruber, G. Mayr, M. Albrecht, K. Rohde, H. Schulz, U. Wahn, N. Hubner, Y. A. Lee, Variants in a novel epidermal collagen gene (COL29A1) are associated with atopic dermatitis. *PLoS Biol.* **5**, e242 (2007).
- N. G. Robertson, C. W. Cremers, P. L. Huygen, T. Ikezono, B. Krastins, H. Kremer, S. F. Kuo, M. C. Liberman, S. N. Merchant, C. E. Miller, J. B. Nadol Jr., D. A. Sarracino, W. I. Verhagen, C. C. Morton, Cochlin immunostaining of inner ear pathologic deposits and proteomic analysis in DFNA9 deafness and vestibular dysfunction. *Hum. Mol. Genet.* **15**, 1071–1085 (2006).
- M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
- T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck III, Y. Hao, M. Stoekius, P. Smibert, R. Satija, Comprehensive integration of single-cell data. *Cell* **177**, 1888–1902.e21 (2019).
- M. B. McPeak, D. Youssef, D. A. Williams, C. L. Pritchett, Z. Q. Yao, C. E. McCall, M. El Gazzar, Frontline science: Myeloid cell-specific deletion of Cebpb decreases sepsis-induced immunosuppression in mice. *J. Leukoc. Biol.* **102**, 191–200 (2017).

36. S. A. Ochsner, D. Abraham, K. Martin, W. Ding, A. McOwiti, W. Kankanamge, Z. Wang, K. Andreano, R. A. Hamilton, Y. Chen, A. Hamilton, M. L. Gantner, M. Dehart, S. Qu, S. G. Hilsenbeck, L. B. Becnel, D. Bridges, A. Ma'ayan, J. M. Huss, F. Stossi, C. E. Foulds, A. Kralli, D. P. McDonnell, N. J. McKenna, The Signaling Pathways Project, an integrated 'omics knowledgebase for mammalian cellular signaling pathways. *Sci. Data* **6**, 252 (2019).
37. N. Hirasawa, Expression of histidine decarboxylase and its roles in inflammation. *Int. J. Mol. Sci.* **20**, 376 (2019).
38. H. Jin, R. He, M. Oyoshi, R. S. Geha, Animal models of atopic dermatitis. *J. Invest. Dermatol.* **129**, 31–40 (2009).
39. F. T. Liu, H. Goodarzi, H. Y. Chen, IgE, mast cells, and eosinophils in atopic dermatitis. *Clin. Rev. Allergy Immunol.* **41**, 298–310 (2011).
40. R. L. Carter, N. S. Lipman, Feed and bedding, in *Management of Animal Care and Use Programs in Research, Education, and Testing*. R. H. Weichbrod, G. A. H. Thompson, J. N. Norton, Ed. (CRC Press/Taylor & Francis, ed. 2, 2018), chap. 27.
41. S. Nunomura, N. Ejiri, M. Kitajima, Y. Nanri, K. Arima, Y. Mitamura, T. Yoshihara, K. Fujii, K. Takao, J. Imura, H. J. Fehling, K. Izuhara, I. Kitajima, Establishment of a mouse model of atopic dermatitis by deleting *Ik2* in dermal fibroblasts. *J. Invest. Dermatol.* **139**, 1274–1283 (2019).
42. Y. Kfoury, D. T. Scadden, Mesenchymal cell contributions to the stem cell niche. *Cell Stem Cell* **16**, 239–253 (2015).
43. S. H. Park, Z. Liu, Y. Sui, R. N. Helsley, B. Zhu, D. K. Powell, P. A. Kern, C. Zhou, *IKK β* is essential for adipocyte survival and adaptive adipose remodeling in obesity. *Diabetes* **65**, 1616–1629 (2016).
44. R. Chovatiya, J. I. Silverberg, Pathophysiology of atopic dermatitis and psoriasis: Implications for management in children. *Children* **6**, 108 (2019).
45. J. Kim, B. E. Kim, D. Y. M. Leung, Pathophysiology of atopic dermatitis: Clinical implications. *Allergy Asthma Proc.* **40**, 84–92 (2019).
46. K. I. Ko, A. L. Syverson, R. M. Kralik, J. Choi, B. P. DerGarabedian, C. Chen, D. T. Graves, Diabetes-induced NF- κ B dysregulation in skeletal stem cells prevents resolution of inflammation. *Diabetes* **68**, 2095–2106 (2019).
47. F. D. Herrington, R. J. Carmody, C. S. Goodyear, Modulation of NF- κ B signaling as a therapeutic target in autoimmunity. *J. Biomol. Screen.* **21**, 223–242 (2016).
48. J. Liu, D. Ibi, K. Taniguchi, J. Lee, H. Herrema, B. Akosman, P. Mucka, M. A. Salazar Hernandez, M. F. Uyar, S. W. Park, M. Karin, U. Ozcan, Inflammation improves glucose homeostasis through *IKK β -XBP1s* interaction. *Cell* **167**, 1052–1066.e18 (2016).
49. M. Baratin, C. Foray, O. Demaria, M. Habbadine, E. Pollet, J. Maurizio, C. Verthuy, S. Davanture, H. Azukizawa, A. Flores-Langarica, M. Dalod, T. Lawrence, Homeostatic NF- κ B signaling in steady-state migratory dendritic cells regulates immune homeostasis and tolerance. *Immunity* **42**, 627–639 (2015).
50. S. M. Rahman, R. C. Janssen, M. Choudhury, K. C. Baquero, R. M. Aikens, B. A. de la Houssaye, J. E. Friedman, CCAAT/enhancer-binding protein β (C/EBP β) expression regulates dietary-induced inflammation in macrophages and adipose tissue in mice. *J. Biol. Chem.* **287**, 34349–34360 (2012).
51. X. Zhou, H. Hu, S. Balzar, J. B. Trudeau, S. E. Wenzel, MAPK regulation of IL-4/IL-13 receptors contributes to the synergistic increase in CCL11/eotaxin-1 in response to TGF- β 1 and IL-13 in human airway fibroblasts. *J. Immunol.* **188**, 6046–6054 (2012).
52. W. Owczarek, M. Paplinska, T. Targowski, K. Jahnz-Rozyk, E. Paluchowska, A. Kucharczyk, B. Kasztalewicz, Analysis of eotaxin 1/CCL11, eotaxin 2/CCL24 and eotaxin 3/CCL26 expression in lesional and non-lesional skin of patients with atopic dermatitis. *Cytokine* **50**, 181–185 (2010).
53. N. Gahr, R. Folster-Holst, M. Weichenthal, E. Christophers, J. M. Schroder, J. Bartels, Dermal fibroblasts from acute inflamed atopic dermatitis lesions display increased eotaxin/CCL11 responsiveness to interleukin-4 stimulation. *Br. J. Dermatol.* **164**, 586–592 (2011).
54. M. Shirley, Dupilumab: First global approval. *Drugs* **77**, 1115–1121 (2017).
55. K. T. Amber, M. Valdebran, K. Kridin, S. A. Grando, The role of eosinophils in bullous pemphigoid: A developing model of eosinophil pathogenicity in mucocutaneous disease. *Front. Med.* **5**, 201 (2018).
56. L. Waltman, N. J. van Eck, A smart local moving algorithm for large-scale modularity-based community detection. *Eur. Phys. J. B* **86**, 471 (2013).
57. L. McInnes, J. Healy, J. Melville, UMAP: Uniform manifold approximation and projection for dimension reduction. arXiv:1802.03426 [stat.ML] (2018).
58. B. Vieth, S. Parekh, C. Ziegenhain, W. Enard, I. Hellmann, A systematic evaluation of single cell RNA-seq analysis pipelines. *Nat. Commun.* **10**, 4667 (2019).
59. T. Wang, B. Li, C. E. Nelson, S. Nabavi, Comparative analysis of differential gene expression analysis tools for single-cell RNA sequencing data. *BMC Bioinformatics* **20**, 40 (2019).

Acknowledgments: We thank M. Karin (UCSD) for providing transgenic mice. This research used resources of the Oak Ridge Leadership Computing Facility, which is a Department of Energy Office of Science User Facility supported under contract DE-AC05-00OR22725.

Funding: This study was supported by NIH grants R01-DE019108 (to D.T.G.), K08-DE027129 (to K.I.K.), R01-DE030415 (to K.I.K.), R01-DA051908 (to D.J., M.R.G., and J.J.M.), R01-ES028114 (to J.T.S.), and P30-AR069589 (to J.T.S.). M.L.H. and Y.H. are supported by T32-AR007465 (NIAMS), and M.L.H. is supported by the Wayne C. Johnson Endowed Research Fellowship.

Author contributions: Study concept and design: K.I.K., J.J.M., D.J., J.T.S., and D.T.G. In vivo mouse experiments: K.I.K., B.P.D., H.Z., E.H., and A.T.N. In vitro experiments with primary mouse and human fibroblasts: K.I.K. and M.L.H. RNAScope experiments: Y.H. and S.P. Data analysis: K.I.K., J.J.M., B.P.D., H.Z., Y.H., E.H., A.T.N., S.P., F.A., A.C., J.R., M.R.G., M.L.H., J.T.S., D.J., and D.T.G. scRNA-seq and bioinformatics analysis: K.I.K., J.J.M., A.C., J.R., M.R.G., D.J., and D.T.G. Supervision: D.J., J.T.S., and D.T.G. Funding acquisition: K.I.K., J.T.S., D.J., and D.T.G. Drafting of original manuscript: K.I.K., J.J.M., D.J., J.T.S., and D.T.G. Critical review and editing of draft: K.I.K., J.J.M., M.C.W., Y.C., S.E.M., A.C., J.R., J.T.S., D.J., and D.T.G.

Competing interests: The authors declare that they have no competing interests. **Data and materials availability.** All data associated with this study are found in the main text and the Supplementary Materials. scRNA-seq data are available through Gene Expression Omnibus (GEO: GSE172226). Computer codes for scRNA-seq analyses are available at <https://doi.org/10.5281/zenodo.5826191>.

Animals and human neonatal primary fibroblasts from this study can be made available via material transfer agreement upon request to the corresponding authors.

Submitted 16 April 2021

Resubmitted 16 September 2021

Accepted 12 January 2022

Published 2 February 2022

10.1126/scitranslmed.abj0324

NF- κ B perturbation reveals unique immunomodulatory functions in Prx1 fibroblasts that promote development of atopic dermatitis

Kang I. KoJean J. MerletBrett P. DerGarabedianHuang ZhenYoko Suzuki-HoriuchiMatthew L. HedbergEileen HuAnh T. NguyenStephen ProutyFaizan AlawiMatthew C. WalshYongwon ChoiSarah E. MillarAshley CliffJonathon RomeroMichael R. GarvinJohn T. SeykoraDaniel JacobsonDana T. Graves

Sci. Transl. Med., 14 (630), eabj0324. • DOI: 10.1126/scitranslmed.abj0324

Prx1 fibroblasts prime AD

The early events of atopic dermatitis (AD) are not fully elucidated, but immunomodulatory fibroblast subsets may contribute to disease. Here, Ko and colleagues deleted *Ikkb* in fibroblastic cells in mice to block NF- κ B activation, finding that these mice developed AD-like skin lesions. Single-cell RNA sequencing identified the presence of paired related homeobox-1–positive (Prx1) fibroblasts that overexpressed C-C motif chemokine ligand 11 (CCL11), leading to myeloid cell infiltration in the skin and a type 2 immune response that preceded development of the lesions. Treatment of young *Ikkb*-deleted mice with a monoclonal antibody blocking CCL11 resulted in decreased eosinophil skin infiltration and type 2 immune response, suggesting that CCL11 may be a target to treat early atopic dermatitis.

View the article online

<https://www.science.org/doi/10.1126/scitranslmed.abj0324>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)