

Case Report Rapport de cas

Mouse barley awn (*Hordeum murinum*) migration induced cystolithiasis in 2 male dogs

Javier Del Angel-Caraza, Carlos C. Pérez-García, Balázs Bende, Inmaculada Diez-Prieto, Belén García-Rodríguez

Abstract – Two male dogs were presented with cystic uroliths composed of magnesium ammonium phosphate (struvite). Each had an atypical nidus, a mouse barley awn (*Hordeum murinum*). To our knowledge, this is the first report of grass awns located in the bladder lumen of dogs. The composition of uroliths and the pathophysiology of grass awn migration to the urinary bladder are discussed.

Résumé – Cystolithiase induite par la migration de la barbe de l'orge des rats (*Hordeum murinum*) chez 2 chiens mâles. Deux chiens mâles ont été présentés avec des urolithes cystiques composés de phosphate double d'ammonium et de magnésium (struvite). Chacun avait un nidus atypique, une barbe d'orge des rats (*Hordeum murinum*). À notre connaissance, il s'agit du premier rapport de barbes d'herbe logées dans la lumière de la vessie de chiens. La composition des urolithes et la pathophysiologie de la migration de l'orge des rats vers la vessie sont discutées.

(Traduit par Isabelle Vallières)

Can Vet J 2011;52:67–69

Case description

A urinary stone (urolith # 1) in a male 3-year-old Maltese dog with a history of hematuria and dysuria, was removed by cystotomy. In another veterinary clinic, a 1.5-year-old male mixed-breed dog, with a similar clinical history was subjected to the same surgical procedure for removal of a urolith (urolith # 2). The uroliths were submitted separately by the veterinary practitioners. Neither dog was reported to have had a previous episode of urolithiasis. Unfortunately we were unable to obtain information on the imaging studies, urinalysis, and urine culture.

A urolith may contain a nidus (area of obvious initiation of urolith growth), stone (major body of the urolith), shell (layer of precipitated material that completely surrounds the stone)

Small Animal Teaching Hospital, Veterinary Faculty, Universidad Autónoma del Estado de México, Toluca-México (Del Angel-Caraza); Laboratory on Urolithiasis Research, Department of Medicine, Surgery and Anatomy, Universidad de León, León-Spain (Del Angel-Caraza, Pérez-García, Diez-Prieto, García-Rodríguez); Budapest Urolith Center, Budapest, Hungary (Bende).

Address all correspondence to Dr. Javier Del Angel-Caraza; e-mail: dlangel@uaemex.mx; delangelvet@hotmail.com

Dr. Del Angel-Caraza's current address is Hospital Veterinario para Pequeñas Especies-FMVZ, Universidad Autónoma del Estado de México. Jesús Carranza # 203 Col. Universidad, CP 50130. Toluca-México.

Use of this article is limited to a single copy for personal study. Anyone interested in obtaining reprints should contact the CVMA office (hbroughton@cvma-acmv.org) for additional copies or permission to use this material elsewhere.




Figure 1. Uroliths with a nidus of plant material. The grass awn is clearly visible in urolith # 1.

and surface crystals (incomplete covering of the outermost surface) (1).

After a careful visual examination, the uroliths were fractured in half for stereoscopic study (Zoom Stereomicroscope SWZ1500; Nikon Instruments, Tokyo, Japan). An ultramicrochemical examination (In vitro Diagnostic Reagent kit, Harzolith-I; Reanal Fine Chemical, Budapest, Hungary) and infrared spectroscopy (Infrared Spectrometer FT-IR 2000; Perkin Elmer, Buckinghamshire, United Kingdom) were used to




Figure 2. Scanning electron microscopy of uroliths. Asterisks – plant material.

determine the mineral composition. In both cases, a nidus with the typical morphology of plant material (in 1 case the image was clearly of a mouse barley awn) was observed (Figure 1).

Urolith # 1 measured $20 \times 8 \times 5$ mm, was whitish in color, and had an oval shape, a rough surface, and weighed 1.4 g. A grass awn was the nidus of the urolith. The layer surrounding the grass awn was composed of a mixture of struvite (80%) and calcium phosphate (20%). The outer layers were formed by struvite (100%).

Urolith # 2 also had a whitish color, an oval shape, and a rough surface; the dimensions were $25 \times 10 \times 6$ mm, and the weight 934 mg. All the layers of the urolith were composed of 100% struvite; a nidus of vegetable structure was identified. Bacteriological culture of the nidus of each urolith and surrounding mineral layers were negative.

Scanning electron microscopy (Scanning Electron Microscope Jeol JSM-6480LV, JEOL, Japan) showed the grass awns with microscopic spikes and hairs surrounded by calcium phosphate and struvite crystals (urolith # 1) or by struvite crystals alone (urolith # 2) (Figure 2).

The grass awns found inside both uroliths were identified as mouse barley (*Hordeum murinum*) (Figure 3) by the Department of Botany, Universidad de León.

Discussion

Foreign objects affecting the urinary tract have only rarely been documented in dogs and cats (2–7) and horses (8). In a recent report of the Minnesota Urolith Center, foreign objects were found inside the uroliths in 0.25% of all samples (3). Suture material was the most common foreign object found in uroliths; but urinary catheters, hair, bullets, needles and plant material have been reported (2–10). The foreign objects can arrive in the urinary bladder by iatrogenic mechanisms (during surgery or urinary tract instrumentation), by retrograde movement up the urethra (6), by a transabdominal route (by migration from the skin) (5), or by a transvesical route (by migration from other abdominal structures) (6,7).

Knowledge of the mineral composition of the urolith is important for the clinical management, treatment, and




Figure 3. The grass awns of mouse barley (*Hordeum murinum*) found inside the uroliths.

prevention of urolithiasis. It is critical to know the composition of the nidus because it is the place where the urolith begins to form and a foreign object can act as a heterogeneous nidus (11). In the 2 cases herein the heterogeneous nidus (grass awn) was surrounded by struvite. Although struvite uroliths can form in a sterile environment, they are more commonly found in dogs that have a urinary tract infection caused by urease-positive bacteria (such as *Staphylococcus* spp. or *Proteus* spp.) because the resultant alkaline urinary pH reduces the solubility of magnesium ammonium phosphate (2,3). The foreign objects may promote bacterial proliferation due to the alteration in the defense mechanisms of the urinary tract. The information available on both clinical cases does not include any reference to urinary culture and, although the culture of the uroliths was negative, we cannot reject the possibility of the presence of a urinary tract infection when the urolith was formed.

Grass awns are especially adapted to the anterograde migration (anterior barbet florets on the surface force the grass awn to move forward and prevent retrograde migration). Usually the front part of the spike penetrates through the skin or body orifices, after being retained in the coat of the animal (12,13). The most common orifice of penetration is the external ear canal; penetration at cutaneous locations such as the dorsal interdigital webs is also common. However, penetration can occur at any site such as the soft tissues of the conjunctiva, the nictitating membrane, inside the eye (14), the nasal sinuses and oral cavity, the bronchial (15,16) and esophageal mucosa, bones (17) or the central nervous system (12,13,18). In many cases, the lesions have the appearance of abscesses.

Although grass awn migration is a common cause of foreign object-related disease in dogs, grass awn migration in the urinary system has only been reported in the urethra and not in the urinary bladder (6,9,10). If migration is retrograde up the urethra, as it seems to be in this study, it is practically impossible for the grass awn to be expelled during micturition because of their “open umbrella” shape (6). Just how plant awns gain access to the lumen of the urinary bladder is the subject of conjecture (3). To our knowledge this is the first report of mouse barley awn (*Hordeum murinum*) in the urinary bladder in dogs. Retrograde

migration of grass awns up the urethra should be considered in the diagnosis of foreign objects in association with cystolithiasis.

Acknowledgments

The authors are grateful for the help of Dr. Antonio Javier Sánchez-Rodríguez (Service of Microscopy of Universidad de León) for the scanning electron microscopy study, Prof. Felix Llamas and Prof. Carmen Acedo (Department of Botany of Universidad de León) for the grass awn identification and Dr. Edgardo Soriano-Vargas (CIESA, Universidad Autónoma del Estado de México) for his editorial assistance. CVJ

References

- Osborne CA, Lulich JP, Polzin DJ, et al. Analysis of 77,000 canine uroliths. Perspectives from the Minnesota Urolith Center. *Vet Clin North Am Small Anim Pract* 1999;29:17–37.
- Osborne CA, Lulich JP, Bartges JW, et al. Canine and feline urolithiasis: Relationship of etiopathogenesis to treatment and prevention. In: Osborne CA, Finco DR eds. *Canine and Feline Nephrology and Urology*. Philadelphia: Williams and Wilkins, 1995:798–888.
- Ulrich LK, Osborne CA, Cokley A, et al. Changing paradigms in the frequency and management of canine compound uroliths. *Vet Clin North Am. Small Anim Pract* 2009;39:41–53.
- Appel SL, Lefebvre SL, Houston DM, et al. Evaluation of risk factors associated with suture-nidus cystoliths in dogs and cats: 176 cases (1999–2006). *J Am Vet Med Assoc* 2008;233:1889–1895.
- Houston DM, Eaglesome H. Unusual case of foreign body-induced struvite urolithiasis in a dog. *Can Vet J* 1999;40:125–126.
- Reimer SB, Kyles AE, Schulz KS, et al. Unusual urethral calculi in 2 male dogs. *J Am Anim Hosp Assoc* 2004;40:157–161.
- Wyatt KM, Marchevsky AM, Kelly A. An enterovesical foreign body in a dog. *Aust Vet J* 1999;77:27–29.
- Textor JA, Slone DE, Clark CK. Cystolithiasis secondary to intravesical foreign body in a horse. *Vet Rec* 2005;156:24–26.
- Morshead D. Submucosal urethral calculus secondary to foxtail awn migration in a dog. *J Am Vet Med Assoc* 1983;182:1247–1248.
- Spellman PG, Spencer JE, Knox AI. Penetrating grass seed. *Vet Rec* 1990;127:410–411.
- Khan SR, Hackett RL. Urolithogenesis of mixed foreign body stones. *J Urol Canadian* 1987;138:1321–1328.
- Brennan KE, Ihrke PJ. Grass awn migration in dogs and cats: A retrospective study of 182 cases. *J Am Vet Med Assoc* 1983;182:1201–1204.
- Gnudi G, Volta A, Bonazzi M, et al. Ultrasonographic features of grass awn migration in the dog. *Vet Radiol Ultrasound* 2005;46:423–426.
- Bussanich MN, Rootman J. Intraocular foreign body in a dog. *Can Vet J* 1981;22:207–210.
- Piek CJ, Robben JH. Pyothorax in nine dogs. *Tijdschr Diergeneesk* 1999;124:276–280.
- Schultz RM, Zwingenberger A. Radiographic, computed tomographic, and ultrasonographic findings with migrating intrathoracic grass awns in dogs and cats. *Vet Radiol Ultrasound* 2008;49:249–255.
- Johnston DE, Summers BA. Osteomyelitis of the lumbar vertebrae in dogs caused by grass-seed foreign bodies. *Aust Vet J* 1971;47:289–294.
- Dennis MM, Pearce LK, Norrdin RW, et al. Bacterial meningoenophalitis and ventriculitis due to migrating plant foreign bodies in three dogs. *Vet Pathol* 2005;42:840–844.