

Knowledge and the Use of Pain Assessment Among Critical Care and Surgical Nurses in the Management of Pain in Usmanu Danfodiyo University Teaching Hospital, Sokoto.

Akhimie Oyakilome ^{1,*}, Sani Dalhat Khalid ², Kombo Abdulrahman Salihu ², Auwalu Muhammed ³, Muhammad Awwal Ladan ²

¹ Usmanu Danfodiyo University Teaching Hospital, Sokoto Nigeria

² Ahmadu Bello University Zaria, Nigeria

³ Usmanu Danfodiyo University, Sokoto

* Corresponding Author: kourhage@gmail.com

Received August 13, 2022,

Revised October 9, 2022,

Accepted February 26, 2023,

Available online February 28, 2023

Abstract

Introduction: Effective pain assessment is attributed to be the basis upon which pain can be managed efficiently and should be routinely undertaken for all postoperative procedures. An individual's response to pain is peculiar and depends on the person and as such pain should be assessed on individual basis.

Aim: The purpose of the study was to examine the use of pain assessment scale in pain management among nurses in critical care and surgical areas of Usmanu Danfodiyo University Teaching Hospital, Sokoto.

Method: This study employed descriptive cross-sectional design. A stratified random sampling technique was used to select 148 nurses across 5 units within the area of study. Descriptive statistics like frequency, mean, standard deviation and percentage were used to describe distribution of data. Descriptive statistics were utilized in data assessing the distribution of data. Pearson's rank correlation was used in determining the relationship between knowledge and level of utilization of pain scales in pain management at the $p=0.05$ level of significance. A total of 148 study participants were involved in this study.

Result: More than half of the nurses, 79 (58.5%), were females; 53 (39.3%) of them were in the age category of 34-39 years. The magnitude of good knowledge towards pain management among nurses was 67.4% with a mean score of 16.41 ± 3.902 . The magnitude for the level of utilization of pain assessment scale in pain management was 80% with a mean score of 4.30 ± 0.381 . There is a positive relationship between knowledge and utilization of pain assessment scale in pain management with $r=0.215$.

Conclusion: Nurses working in critical care and surgical areas of Usmanu Danfodiyo University Teaching Hospital, Sokoto had good knowledge, and utilize pain assessment tool in pain management than those reported in previous studies in southern Nigeria.

Keywords: pain, knowledge, utilization

Introduction

An effective pain assessment is attributed to be the basis upon which pain can be managed effectively and should be routinely undertaken for all postoperative procedures. An Individual's response to pain is peculiar and depends on the person and as such it is recommended that pain should be assessed on individual basis. Screening used for pain should be a part of a regular assessment, and this has led the American Pain Society (APS) to proclaim pain as the "fifth vital sign" (Nuseir et al., 2016). However, there was a lack of knowledge about some key concepts such

as value for patients' volition in pain assessment and anticipatory analgesia concepts.

Internationally, it is argued that many nurses in common practical settings are deficient in the knowledge about basic pain evaluation and management principles, this also applies to the attitude nurse have towards pain and its appraisal (Ung et al., 2016). With this it might be assumed that inadequate assessment and treatment of pain will continue to be an issue in the care provided in the health care system. Also nurses are seen to be key persons who can improve the quality of pain management and who can provide nursing care to sufficiently meet the patient's needs (Liyew et al., 2020). In this regard (Chatchumni, M., Namvongprom, A., Eriksson, H., & Mazaheri, 2016) suggest that nurses with stronger knowledge and attitude tend to manage pain better with improved outcomes, and higher patient satisfaction scores. As such nurses play a vital role in pain assessment and management and must be knowledgeable regarding how best to assess and manage the pain (Chatchumni, M., Namvongprom, A., Eriksson, H., & Mazaheri, 2016),

Pain can only be managed by the use of pain assessment tools and these tools can be grouped into either single-dimensional or multidimensional scales. The measures involve patient self-report on a part or parts of the pain. Thus, the outcomes have to be viewed as guides and not absolutes. They ought to be seen as only aide to the history and physical examination of the patient. The frequently used tools to measure pain strength include verbal rating scales, numeric rating scales, and visual analogue scales (Honorio *et al* 2011).

The American Society of Pain in 2015 estimated that 25 million U.S. adults had on a daily basis unremitting pain, and 23 million more recount severe pain (Nahin, 2016). While (Uysal, 2018) reported that in Turkey, 20-50% of the patients with cancer feel the pain at first admission; 30-40% of them suffer pain all through the treatment; and 60-70 % of them experience the pain at terminal stage. In Saudi Arabia, (Issa & Awajeh, Adel Musbah and Khraisat, 2017) suggested that pain assessment and management are vital components of nursing care and two of the most essential patient rights. Undoubtedly, pain is the foremost challenge in critically ill patients admitted to intensive care units (ICU) and 40-77.4% of ICU patient whine about the experience of pain (Deldar et al., 2018). Most documented studies on pain in Sub-Sahara Africa has been studied mainly in relation to HIV/AIDS and cancer (Selman et al., 2013). For example in Uganda, (Kizza et al., 2016) observed that nurses generally had adequate knowledge about pain assessment principles. This has been argued to be unprincipled to allow a patient to endure without taking suitable actions to relieve his or her pain (Kizza et al., 2016). Elsewhere, nurses were reported to have poor knowledge levels and attitudes with respect to managing pain of adult medical patients in Zimbabwe (Manwere et al., 2015). In another study in Ghana, findings showed that more than 57% of nurses had deficient tools that can be used to assess and measure pain, 12% of the health care providers had never used any tool in the assessment of pain (Manwere et al., 2015).

Pain assessment requires health care providers to have a good understanding of it, because poor evaluation and documentation is ascribed to the high occurrence of acute pain experienced by patients (Menlah et al., 2018). As such, standard pain assessment methods, with emphasis on Numeric Rating scale, offer a reliable and valid appraisal of this individual symptom. While evaluating a patient's pain level, it is paramount for nurses to be unbiased; nurses should demonstrate that they

recognize the patient is in pain, and endeavour to start an empathetic relation (Vallerand et al., 2011). The purpose of the study was to examine the use of pain assessment scale in pain management among nurses in critical care and surgical areas of Usmanu Danfodiyo University Teaching Hospital, Sokoto

Methodology

A descriptive cross-sectional design was used to determine the level of knowledge, perceived attitude and reported utilization of pain assessment scale in the management of pain among nurses in critical care and surgical nurses of UDUTH. UDUTH Sokoto is a tertiary health institution serving as a referral centre to all primary and secondary health facilities in its catchment area which consist of Sokoto, Kebbi, Zamfara, Katsina and Niger states. It commenced operation in 1989 and is a 650 bed hospital with various specialties. The population for the study is a total of 148 nurses in critical and surgical areas of the hospital. Using existing strata, a stratified random sampling technique was used to select 148 nurses. The various departments in the hospital departments were classified into 5 main strata having nearly the same working conditions: (1) Accident and Emergency, (2) Trauma centre (3) Intensive care unit and anaesthesia, (4) Surgical wards, (5) Oncology ward.

The instrument for data collection was a questionnaire adapted from from Ferrell and McCaffery's (2014) Knowledge and Attitudes Survey Regarding Pain (KASRP) self-report questionnaire. The questionnaire comprised of 47 items in English, including 24 knowledge questions, and the interviewees will be asked to reply in two ways: 1 = True; 0 = False, with a total range of "0-24"; they were asked to answer 11 attitude questions according to five-point Likert scales: from "strongly agree = 5" to "strongly disagree = 1" with total score range of "11-55"; they were asked to answer 6 practice questions according to four-point Likert scales: from "always = 5" to "less often = 1" with total score range of "6-30" and they were asked to choose from 13 barriers questions that applies to them. Answers were evaluated considering the extent to which they are compatible with pain therapy standards commonly acknowledged by the international pain management guidelines. A knowledge score of 0-12 means poor knowledge and 13-24 means good knowledge.

The questionnaire was based on the contents of the objectives of the study. It was divided into five sections. Sections A, B, C, D and E contain 7, 24, 11, 6 and 12 questions respectively.

Section A	Elicited information on the respondents' demographic profile in terms of; age, gender, ethnic group, religion, marital status, rank and years of service.
Section B	Focused on the knowledge of pain assessment scale in pain management
Section C	Attitude of Nurses in the use of pain assessment scale in pain management
Section D	Practicability of the use of pain assessment scale in pain management
Section E	Factors affecting the use of pain assessment scale in pain management

Data collection and analysis

The copies of the questionnaire were then distributed to health care professionals in the various units and departments by the researcher and this lasted for four weeks. Therefore four weeks was used for data collection. Data obtained through the questionnaire was appropriately cleaned to ensure accuracy and consistency. The data collected was analyzed using the Statistical package for social science version 23. The respondents' demographics, pain assessment tool-related knowledge, attitude, and practice were analyzed using frequencies and percentages.

Ethical consideration

An ethical approval to conduct the research was obtained from the ethical and research committee of UDUTH, Sokoto with **UDUTH/HREC/2021.1108/V2**.

All information about the study was explained to the participants for them to understand the purpose of the study which is academic and therefore informed Participants are informed that participation is voluntary and they can withdraw at anytime.

Results

To determine the reliability of the instrument a pilot study of 10% of the sample size was carried out in Specialist Hospital Sokoto. The results was statistical analyzed and the results of the Cronbach's score presented below:

Table 1. Cronbach's Alpha Test for Reliability of Instrument

Section	Cronbach's Alpha	Cronbach's Alpha based on Standardized item	Number of Item
Knowledge of Pain Assessment Scale	0.889	0.896	24
Practice of Pain Assessment Scale in the Management of Pain	0.854	0.868	6
Factors Hindering the Practice of Pain Assessment Scale	0.746	0.770	12

Source: Own computation using SPSS 22 version, 2022

Table 2. Socio-demographic characteristics of respondents N=135

Variable	Categories	Frequency	Percentage (%)
Gender	Male	56	41.5
	Female	79	58.5
Age	22-27	12	8.9
	28-33	29	21.5
	34-39	53	39.3
	40-46	28	20.7
	47+	13	9.6
Educational Attainment	Registered Nurse Certificate	93	68.9
	BNSc	40	29.6
	MSc	2	1.5
Position/Rank	Nursing Officer	40	29.6
	Senior Nursing Officer	26	19.3
	Principal Nursing Officer	24	17.8
	Assistant Chief Nursing Officer	21	15.6
	Chief Nursing Officer	18	13.3
	Others	6	4.4
Duration in position	Less than 1	20	14.8
	2-3 years	70	51.9
	4-5 years	24	17.8
	6 years and above	21	15.6
Number of years as Registered Nurse	0-5 years	22	16.3
	6-10 years	32	23.7
	11-15 years	54	40.0
	16-20 years	12	8.9
	21 years and above	15	11.1
Departments	A and E	20	14.8
	Trauma Center	14	10.4
	Oncology	8	6.0
	Anesthesia and Intensive Care Unit	28	20.7
	Surgical Wards	65	48.1

Table 3. Level of Nurses’ Knowledge, utilization and relationship between knowledge and utilization of pain assessment scale in Pain Management (N=135)

Level of Nurses’ Knowledge in Pain Management			
Knowledge status	N	Percentage	Mean ± SD
Good Knowledge	91	67.4%	16.41 ± 3.902
Poor Knowledge	44	32.6%	10.68 ± 1.653
Total	135	100	
Level of reported utilization of pain scales towards pain management			
Variable	N	Percentage	
Good utilization	108	80.0%	
Poor utilization	27	20.0%	
Total	135	100	
Relationship between knowledge and Utilization Pain Assessment Scale in Pain Management			
Knowledge versus Utilization	Mean ± SD	r-value	P-value
Knowledge	14.47±4.430	0.215	0.011
Utilization	4.03±0.623		

Table 4. Factors Affecting the Use of Pain Assessment Scale in Pain Management (N=135)

S/N	Factor	True (%)	False (%)
1	Unavailability of pain assessment tools	104 (77)	31 (23)
2	Too much patients and nursing workload	115 (85.2)	20 (14.8)
3	Patients inability to communicate	81 (60)	54 (40)
4	Poor documentation of pain assessment and management	103 (76.3)	32 (23.7)
5	Lack of pain assessment protocols	106 (78.5)	29 (21.5)
6	No designated area for charting pain	104 (77)	31 (23)
7	Lack of familiarity with assessment tools	98 (72.6)	37 (27.4)
8	Pain assessment tools is difficult to use and complex to interpret	36 (26.7)	99 (73.3)
9	The use of pain assessment tools is time consuming and not practicable	50 (37)	85 (63)
10	Sedation interfering with pain assessment	108 (80)	27 (20)
11	Unconducive working environment	82 (60.7)	52 (39.3)
12	Lack of knowledge about pain tools.	96 (71.1)	38 (28.9)

A total of 148 questionnaires were self administered to the respondents and 142 were retrieved out of which 135 (95.1%) were completely filled and 7 incompletely filled. The analysis was based on the 135 completely filled questionnaires retrieved.

Most of the nurses were within the age bracket of 34 – 39 and 28 – 33 representing 39.5% and 21.5% respectively. Female nurses were the majority representing 58.5% while males were 41.5%. Most of the respondents 68.9% had registered nurse certificate while 29.6% of them had Bachelor of Nursing Science. Masters of Science in Nursing was the highest educational attainment of the nurses representing 1.5% of the respondents. Most of the respondents were nursing officer’s representing 29.6% while others (Assistant Director Nursing) were 4.4% of the respondents. The duration in the position (years) by nurses was mostly 2 – 3 years representing 51.9% of the respondents while those who had spent 6years and above were 15.6%. The respondents who have over 20 years experience are 11.1% while those with less than 5 years are 16.3%, while majority of the respondents have been registered for 11 – 15 years representing 40%.The majority of the nurses were in Surgical Wards representing 48.1% respondents while Anesthesia and Intensive Care Unit representing 33.3% of the respondents.

From table 3, it can be seen that 67.4% of the respondents have good knowledge about pain management scale while 32.6% of them have poor knowledge about pain management scale. Also, it can be seen that 80.0% of the respondents use pain scales in assessing pain prior to pain management while 20.0% of them do not use

pain assessment scales in assessing pain prior to pain management. It is evident that there is a positive relationship between knowledge and utilization of pain assessment scale in pain management with $r=0.215$. This suggested that the more the knowledge, the more the utilization of assessment scale in pain management among Nurses.

Table 4 shows factors affecting the use of pain assessment scale in pain management. The three most leading factors were; too much patients and nursing workload, with 85.2%, Sedation interfering with pain assessment with 80% and Lack of pain assessment protocols with 78.5% while pain assess tools been difficult to use and complex to interpret and the use of pain assessment tool been time consuming and not practicable were the lease factors with 26.7% and 37% respectively.

Discussion

Findings in the study shows 67.4% of Nurses in critical care and surgical area had good knowledge about pain management. This finding was consistent with the studies done in United Kingdom (UK) 73.8%, and Chicago, United States of America, 74%, Saudi Arabia 87.5%, Italy (55%), Iran (66.6%), Gondar Ethiopia (66.9%) where these percentages of the study participants had good knowledge towards pain management as reported by (Liyew et al., 2020) In addition, the findings were consistent that of Karamjeet (2017) where 66% had good knowledge about pain management also (Umuhoza et al., 2019) reported 74% of nurses had good knowledge. Furthermore, the study is in tandem with findings of (Oyetunji & Popoola, 2020) were 59.7% of Nurses in south west Nigeria had good knowlegde about pain management. This suggests that the respondents show knowledge consistent with their contemporaries in different parts of the world with respect to knowledge about pain management.

Furthermore, findings suggest too much patients and nursing workload 85.2%, lack of pain assessment protocols 78.5% and sedation interfering with pain assessment 80% was the three leading factors militating against the use of pain assessment scales in pain management. This was similar to findings by (Mędrzycka-Dąbrowska W, Dąbrowski S, Basiński A., 2015) where heavy workload and lack of time were factors that militate against the use of pain scales, (Kizza & Muliira, 2015) where poor documentation and communication of pain assessment priorities, workload and familiarities with assessment tool limit the use of pain assessment tools. Also (Toba et al., 2019) reported insufficient information about pain management, harsh regulation on the use of opioids and poorly assessed pain.

Also, findings shows there was positive relationship between knowledge and utilization of pain assessment scale in pain management with $r=0.215$. This is consistent with findings of (Alzghoul et al., 2016), in which nurses knowledge of pain management had a strong correlation with the use of pain assessment tool in pain management practice. The implication of this is that: the more the knowledge, the more the utilization of assessment scale in pain management among Nurses this in turn will alleviate patients/client suffering as regards pain management. It is therefore recommended that nurses and other health care workers ensure the use of a pain assessment tool prior to pain management as this will ensure effective and efficient therapy.

Limitation

Firstly, the study was carried out in UDUTH it is pertinent to note that the results cannot be generalized for the entire nurses in the states. However, the study can be extended to cover the entire state in Northern Nigeria. Secondly, the study assessed reported utilization of pain scales in pain management among nurses. Observation on the use of pain scales among nurses would have been more appropriate. Finally, factors identified to have impinged on the use of pain assessment scale in pain management are skewed negatively as it was only on the staff without noting what the management has to say.

Contribution to global practice

The study will improve global nursing practice as it will add to the existing body of literature and also nurses will use a pain assessment tool towards evaluating pain prior to its management.

Conclusion

On the basis of the findings of this study, the following conclusions were drawn: It was seen that of Nurses in critical care and surgical area had good knowledge about pain management and their knowledge is consistent with nurses in southern Nigeria and other parts of the world. Furthermore, the study revealed that patient's workload and lack of assessment protocols were the two leading factors militating against the use of pain scale in pain management. Also, nurses exhibited a positive relationship between knowledge and the use of pain assessment scale in pain management among critical care and surgical areas of UDUTH, Sokoto

Author contribution

The contributions of the various authors are as follows; Akhimie Oyakilome research concept and design, Sani Dalhat Khalid discussant of the findings, Kombo Abdulrahman Salihu pilot testing of tool, Auwal Mohammad critical revision of the article and Muhammad Awwal Ladan data analysis and interpretation.

Conflict of interest

There is no conflict of interest in this paper.

Acknowledgement

My profound gratitude goes to the entire team for their technical support towards the success of the study. The entire critical care nurses of the hospital for taking time to respond to the questionnaire.

Reference

- Alzghoul, B. I., Azimah, N., & Abdullah, C. (2016). Pain Management Practices by Nurses: An Application of the Knowledge, Attitude and Practices (KAP) Model. *Global Journal of Health Science*, 8(6), 1-7. <https://doi.org/10.5539/gjhs.v8n6p>
- Chatchumni, M., Namvongprom, A., Eriksson, H., & Mazaheri, M. (2016). Thai Nurses' experiences of post-operative pain assessment and its' influence on pain management decisions. *BMC Nursing*, 15(12). <https://doi.org/10.1186/s12912-016-0136-8>
- Deldar, K., Froutan, R., & Ebadi, A. (2018). Challenges faced by nurses in using pain assessment scale in patients unable to communicate: A qualitative study. *BMC Nursing*, 17(11), 1-8. <https://doi.org/10.1186/s12912-018-0281-3>
- Ferrell, B. and McCaffery, M., (2014). Knowledge and attitudes survey regarding pain, viewed 31 March 2021, [http://prc.coh.org/Knowledge & Attitude Survey 7-14 \(1\).pdf](http://prc.coh.org/Knowledge & Attitude Survey 7-14 (1).pdf).

- Honorio T. B., Srinivasa N. R., Robert E. M., Spencer S. L., Scott M.F., (2011). Pain assessment in Essentials of Pain Medicine. USA: Elsevier Saunders; 3rd edition, 28-33.
- Issa, M. R., & Awajeh, Adel Musbah and Khraisat, F. S. (2017). Knowledge and Attitude about Pain and Pain Management among Critical Care Nurses in a Tertiary Hospital Abstract. *Journal of Intensive and Critical Care*, 3(1:12). <https://doi.org/10.21767/2471-8505.100071>
- Karamjeet, K. (2017). Knowledge and attitude regarding pain management among staff nurses. *Nurse Care Open Acces J*, 2(1), 5-7. <https://doi.org/10.15406/ncoaj.2017.02.00024>
- Kizza, B., & Muliira, K. (2015). Nurses ' pain assessment practices with critically ill adult patients. *International Nursing Review*, 62, 573-582.
- Kizza, I. B., Muliira, J. K., Kohi, T. W., & Nabirye, R. C. (2016). Nurses' knowledge of the principles of acute pain assessment in critically ill adult patients who are able to self-report. *International Journal of Africa Nursing Sciences*, 4. <https://doi.org/10.1016/j.ijans.2016.02.001>
- Liyew, B., Dejen Tilahun, A., Habtie Bayu, N., & kassew, T. (2020). Knowledge and Attitude towards Pain Management among Nurses Working at University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia. *Pain Research and Management*, 2020, 1-9. <https://doi.org/10.1155/2020/6036575>
- Mahama, F., & Ninnoni, J. P. K. (2019). Assessment and Management of Postoperative Pain among Nurses at a Resource-Constraint Teaching Hospital in Ghana. *Hindawi Nursing Research and Practice*. <https://doi.org/10.1155/2019/9091467>
- Manwere, A., Chipfuwa, T., & Mukwamba, M. M. (2015). Knowledge and Attitudes of Registered Nurses towards Pain Management of Adult Medical Patients: A Case of Bindura Hospital. *iMedPubJournals*1-6.
- Mędrzycka-Dąbrowska W, Dąbrowski S, Basiński A. (2015). Problems and Barriers in Ensuring Effective Acute and Post-Operative Pain Management – an International Perspective. *Adv Clin Exp Med*, 25(4), 905-910. <https://doi.org/10.17219/acem/26394>
- Menlah, A., Garti, I., Amoo, S. A., Atakro, C. A., Amponsah, C., & Agyare, D. F. (2018). Knowledge, Attitudes, and Practices of Postoperative Pain Management by Nurses in Selected District Hospitals in Ghana. *SAGE Open Nursing*, 4. <https://doi.org/10.1177/2377960818790383>
- Nahin, R. L. (2016). Estimates of pain prevalence and severity in adults: United states 2012. *Journal of Pain Instructions*, 16(8), 769-780. <https://doi.org/10.1016/j.jpain.2015.05.002>
- Nuseir, K., Kassab, M., & Almomani, B. (2016). Healthcare providers' knowledge and current practice of pain assessment and management: How much progress have we made? *Pain Research and Management*, 2016, 18-23. <https://doi.org/10.1155/2016/8432973>
- Oyetunji, F. O., & Popoola, R. O. (2020). Knowledge As Predictors Of Nurses Attitude Towards. *African Journal of Health, Nursing and Midwifery*, 3(2), 53-66.
- Selman, L., Simms, V., Penfold, S., Powell, R. A., Mwangi-powell, F., Downing, J., Gikaara, N., Munene, G., Higginson, I. J., & Harding, R. (2013). ' My dreams are shuttered down and it hurts lots '– a qualitative study of palliative care needs and their management by HIV outpatient services in Kenya and Uganda. *BMC Palliative Care*, 12(35).
- Toba, H. A., Samara, A. M., & Zyoud, S. H. (2019). Nurses ' knowledge , perceived barriers , and practices regarding cancer pain management : a cross-sectional study from Palestine. *BMC Medical Education*, 19(167), 1-7.
- Umuhoza, O., Chironda, G., Katende, G., & Mukeshimana, M. (2019). International Journal of Africa Nursing Sciences Perceived knowledge and practices of nurses regarding immediate post-operative pain management in surgical wards in Rwanda . A descriptive cross-sectional study. *International Journal of Africa Nursing Sciences*, 10(April), 145-151.
- Ung, A., Salamonson, Y., Hu, W., & Gallego, G. (2016). Assessing knowledge , perceptions and attitudes to pain management among medical and nursing students : a review of the literature. *British Journal of Pain*, 10(1), 8-21. <https://doi.org/10.1177/2049463715583142>

- Uysal, N. (2018). Clearing Barriers in Cancer Pain Management : Roles of Nurses. *International Journal of Caring Sciences*, 11(2), 1323–1327.
- Vallerand, A. H., Musto, S., & Polomano, R. C. (2011). Nursing's role in cancer pain management. *Current Pain and Headache Reports*, 15(4), 250–262. <https://doi.org/10.1007/s11916-011-0203-5>