

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

---

Biology Faculty Publications and Presentations

College of Sciences

---

5-2021

## Understanding water and energy fluxes in the Amazonia: Lessons from an observation-model intercomparison

Natalia Restrepo-Coupe

Loren P. Albert

Marcos Longo

Ian Baker

Naomi M. Levine

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.utrgv.edu/bio\\_fac](https://scholarworks.utrgv.edu/bio_fac)



Part of the [Biology Commons](#)

---

### Recommended Citation

Restrepo-Coupe, N., Albert, L.P., Longo, M., Baker, I., Levine, N.M., Mercado, L.M., da Araujo, A.C., Christoffersen, B.O., Costa, M.H., Fitzjarrald, D.R., Galbraith, D., Imbuzeiro, H., Malhi, Y., von Randow, C., Zeng, X., Moorcroft, P. and Saleska, S.R. (2021), Understanding water and energy fluxes in the Amazonia: Lessons from an observation-model intercomparison. *Glob. Change Biol.*, 27: 1802-1819. <https://doi.org/10.1111/gcb.15555>

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Biology Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact [justin.white@utrgv.edu](mailto:justin.white@utrgv.edu), [william.flores01@utrgv.edu](mailto:william.flores01@utrgv.edu).

---

**Authors**

Natalia Restrepo-Coupe, Loren P. Albert, Marcos Longo, Ian Baker, Naomi M. Levine, Lina M. Mercado, Alessandro C. da Araujo, Bradley O. Christoffersen, Marcos H. Costa, and David R. Fitzjarrald

DR. NATALIA RESTREPO-COUBE (Orcid ID : 0000-0003-3921-1772)

DR. LOREN ALBERT (Orcid ID : 0000-0002-9674-6071)

DR. MARCOS LONGO (Orcid ID : 0000-0001-5062-6245)

DR. NAOMI LEVINE (Orcid ID : 0000-0002-4963-0535)

DR. DAVID R GALBRAITH (Orcid ID : 0000-0002-5555-4823)

PROF. PAUL R MOORCROFT (Orcid ID : 0000-0002-2876-4673)

DR. SCOTT R SALESKA (Orcid ID : 0000-0002-4974-3628)

Article type : Primary Research Articles

## **Understanding water and energy fluxes in the Amazonia: Lessons from an observation-model intercomparison**

### **Running head: Seasonal water-energy flux in Amazon forests**

Natalia Restrepo-Coupe<sup>1,2</sup>, Loren P. Albert<sup>1,3</sup>, Marcos Longo<sup>5,6</sup>, Ian Baker<sup>4</sup>, Naomi M. Levine<sup>5,7</sup>, Lina M. Mercado<sup>8,9</sup>, Alessandro C. da Araujo<sup>10,11</sup>, Bradley O'Donnell Christoffersen<sup>12,19</sup>, Marcos H. Costa<sup>13</sup>, David R. Fitzjarrald<sup>14</sup>, David Galbraith<sup>15</sup>, Hewley Imbuzeiro<sup>13</sup>, Yadvinder Malhi<sup>16</sup>, Celso von Randow<sup>17</sup>, Xubin Zeng<sup>18</sup>, Paul Moorcroft<sup>5</sup>, Scott R. Saleska<sup>1</sup>

(1) Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, USA.

(2) School of Life Sciences, University of Technology Sydney, Ultimo, NSW, Australia.

(3) Biology Department, West Virginia University, Morgantown, WV, USA.

(4) Colorado State University, Atmospheric Science, Fort Collins, CO, USA.

(5) Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA.

(6) Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA.

(7) College of Letters, Arts, and Science, University of Southern California, Los Angeles, CA, USA.

**This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/GCB.15555](https://doi.org/10.1111/GCB.15555)**

This article is protected by copyright. All rights reserved

- (8) University of Exeter, College of Life and Environmental Sciences, Exeter, Devon, EX4 4RJ UK.
- (9) Centre for Ecology and Hydrology, Wallingford, Oxfordshire, OX10 8BBUK, UK.
- (10) Embrapa Amazônia Oriental, Belém, Pará, Brazil.
- (11) Programa LBA, Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil.
- (12) Department of Biology, University of Texas Rio Grande Valley, Edinburg, TX USA.
- (13) Department of Agricultural Engineering, Federal University of Vicosa, Vicosa, MG, Brazil.
- (14) University at Albany SUNY, Albany, NY, USA.
- (15) School of Geography, University of Leeds, Leeds, UK.
- (16) Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, UK.
- (17) National Institute for Space Research, Brazil (INPE), Center for Earth Systems Science, São José dos Campos, SP, Brazil.
- (18) Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ, USA.
- (19) Earth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, NM, USA.

**Correspondence:** Natalia Restrepo-Coupe

email: nataliacoupe@gmail.com, phone: +1 647 328 1494

**Key words:** Evapotranspiration, energy balance, eddy covariance flux seasonality, Amazonia, tropical forests, land surface models, ecosystem-climate interactions.

**Type of Paper:** Primary Research Article.

### Abstract

Tropical forests are an important part of global water and energy cycles, but the mechanisms that drive seasonality of their land-atmosphere exchanges have proven challenging to capture in models. Here, we (1) report the seasonality of fluxes of latent heat ( $LE$ ), sensible heat ( $H$ ), and outgoing short and longwave radiation at four diverse tropical forest sites across Amazonia -- along the equator from the Caxiuana and Tapajós National Forests in the eastern Amazon to a forest near Manaus, and from the equatorial zone to the southern forest in Reserva Jaru; (2) investigate how vegetation and climate influence these fluxes; and (3) evaluate land surface model (LSM) performance by comparing simulations to observations. We found that previously identified failure of models to capture observed dry-season increases in evapotranspiration was associated with model over-estimations of (1) magnitude and seasonality of Bowen ratios (relative to aseasonal observations in which sensible was only 20-30% of the latent heat flux)

This article is protected by copyright. All rights reserved

indicating model exaggerated water limitation, (2) canopy emissivity and reflectance (albedo was only 10 to 15% of incoming solar radiation, compared to 0.15-0.22% simulated), and (3) vegetation temperatures (due to underestimation of dry-season evapotranspiration and associated cooling). These partially compensating model-observation discrepancies (e.g. higher temperatures expected from excess Bowen ratios were partially ameliorated by brighter leaves and more interception/evaporation) significantly biased seasonal model estimates of net radiation ( $R_n$ ), the key driver of water and energy fluxes ( $LE \sim 0.6R_n$  and  $H \sim 0.15R_n$ ). Though these biases varied among sites and models. A better representation of energy-related parameters associated with dynamic phenology (e.g. leaf optical properties, canopy interception, and skin temperature) could improve simulations and benchmarking of current vegetation-atmosphere exchange and reduce uncertainty of regional and global biogeochemical models.

## 1. Introduction

Tropical forests play a major role in the global water and energy cycles, and modulate tropical atmospheric circulation processes, cloud formation and precipitation (Hagos & Leung, 2011; Held & Soden, 2006; Jasechko et al., 2013; Silva Dias et al., 2002; Wei et al., 2017; Worden et al., 2007). Water and energy fluxes are intrinsically linked, as energy is required for the phase transition from liquid to vapor. Tropical forests evapotranspire the energy equivalent of more than half of the total solar energy absorbed by earth's land surfaces (Trenberth et al., 2009), helping to maintain high atmospheric water content, increase moisture recycling, and mediate cloud development (Peters, 2016; Tan et al., 2019). Evapotranspiration ( $ET$ ) mitigates heating as part of the incoming radiation is primarily "consumed" as latent heat ( $LE$ ) rather than as sensible heat flux ( $H$ ). High  $ET$  rates can offset the warming effect associated with tropical forest low albedo (the ratio of reflected to incoming shortwave radiation,  $SW_{out}/SW_{down}$ ) driven by its relatively dark surface (Bonan, 2008; Yanagi & Costa, 2011). Therefore, land use change, fire, climate and extreme weather events (Aragão et al., 2007, 2008; Chagnon & Bras, 2005; Davidson et al., 2012) are listed as key factors determining subsequent changes in tropical forest albedo's (negative climate forcing) and alterations of the evaporative cooling flux (positive feedbacks – reducing warming) (Bonan, 2008; Li et al., 2015; Liu et al., 2019). Measuring and understanding water, radiation, and energy seasonal fluxes under present climatological

conditions is thus needed to forecast the future of tropical forests and global atmospheric cycles (Fu et al., 2013; Sena et al., 2018; Spracklen et al., 2018).

Land-surface models (LSMs) represent our mechanistic understanding of cause-effect relationships between the surface and the atmosphere and constitute ideal tools to forecast water, energy and other biogeochemical fluxes (Pitman, 2003). However, given that ecosystem characteristics are diverse and that land-climate interactions are heterogeneous and complex, it is not surprising that LSMs have difficulty in reproducing the seasonality of rainforest *ET* (Baker et al., 2008; Christoffersen et al., 2014; Costa et al., 2010; Fisher et al., 2014; Restrepo-Coupe et al., 2017). A consistent problem is that models simulate reductions in *ET* during the dry season (when precipitation is less than  $\sim 100$  mm month<sup>-1</sup>), when most observations from eddy covariance towers in Amazonia show no reductions or even increases in *LE*, consistent with control by the availability of energy (net radiation), and inconsistent with limitation by available water (Baker et al., 2008; Christoffersen et al., 2014; Costa et al., 2010; R. A. Fisher et al., 2007; Restrepo-Coupe et al., 2017; Shuttleworth, 1988).

Previous attempts to improve the dry-season *LE* discrepancies between LSM simulations and observations of tropical forests, have been focused on the parameterization of higher soil water holding capacity, hydraulic redistribution (vegetation control mechanisms), deeper roots that can access the lower soil layers and/or increase root mass (enhanced pathways) and dynamics of stem-water storage (plant hydraulics) (Baker et al., 2008; Christoffersen et al., 2014; Harper et al., 2010; Lee et al., 2005; Yan et al., 2020). Unfortunately, some of these model modifications appear to drive LSMs to (1) overestimate annual and/or dry-season *ET* and/or (2) model simulations could become insensitive to drought conditions.

Christoffersen et al. (2014) previously analyzed simulations from the same model-data intercomparison investigated here, focusing on modeled mechanisms of water supply (rooting depth, access to groundwater sources, and soil water availability) and vegetation demand (intrinsic water use efficiency (*iWUE*) and stomatal conductance) that drive the simulated dry-season reductions in *ET*. Christoffersen et al. (2014) identified model underrepresentation of phenological processes (including leaf development and associated changes in *iWUE*) as a cause

of the bias. When these same LSM simulations were evaluated for their ability to represent the seasonal dynamics of carbon fluxes in these same tropical forests (Restrepo-Coupe et al., 2017), the analysis found that although water limitation was represented in models as the primary driver of the seasonality of photosynthesis across Amazonia, the LSMs did not accurately represent that seasonality. Observations showed incoming radiation and phenological cycles that included allocation lags between wood, leaf and non-structural carbon, and light harvesting adaptations (e.g., leaf demography) dominated carbon exchange and in some instances, were not well represented in LSMs. Both carbon and water fluxes are significantly influenced by tropical forest phenology (Chen et al., 2020; Restrepo-Coupe et al., 2017). However, the relationship between vegetation seasonal cycles and the radiation and energy exchange is not well documented.

Here, we extend the prior work of Christoffersen et al (2014) and Restrepo-Coupe et al., (2017), building on the consistent finding that  $LE$  appears to be controlled by net radiation ( $R_n$ ). If this finding is correct, then inherent to the challenge of accurate modeling of  $ET$  (equivalent  $LE$ ) is the accurate simulation of the other radiation components ( $LW_{out}$  and  $SW_{out}$ ), as well as the accurate partitioning of the relevant energy fluxes (e.g. energy allocated to  $LE$  and  $H$ ) (Bony et al., 2013; Getirana et al., 2014; Longo et al., 2019a), in addition to the accurate representation of phenological attributes (e.g. leaf-age driving seasonal canopy conductance values) (Lin et al., 2015; Medlyn et al., 2011) (see Figure 1). Yet, in tropical forests and across Amazonia there is scarce information on the seasonal cycle of energy-relevant components  $H$ , albedo ( $\alpha$ ), emissivity ( $\epsilon_s$ ), the Bowen ratio ( $Bowen=H/LE$ ), and the outgoing and incoming longwave radiation ( $LW_{out}$  and  $LW_{down}$ ).

Focusing on energy dynamics, we compare forest characteristics and water and energy fluxes from eddy covariance (EC) and meteorological observations at four tropical forest sites from the Brasil flux network, three Amazonian forests close to the Amazon river (Manaus-K34, Tapajós-K67, and Caxiuaná-CAX) and one southern location (Reserva Jaru-RJA) to four state-of-the-art land surface models (IBIS, ED2, JULES, and CLM3.5) (Restrepo-Coupe et al., 2017). The aim of this work is threefold: (1) to quantify and characterize the seasonal fluxes (timing and amplitude) and surface properties of the different water, energy and radiation cycle components;

(2) to determine the relationships between these energy-related fluxes and vegetation and climate drivers, as we investigate the ability of other simple models and relations to predict ecosystem-level fluxes (e.g. linear regressions between  $Rn$  and  $LE$ ); and (3) to identify areas to refine current LSM model formulations and to enhance seasonal  $LE$ ,  $H$  and  $Rn$  simulations by including vegetation characteristics (e.g. albedo) in the analysis and improving the derivation of radiative fluxes (e.g. outgoing  $SW$  and  $LW$ ), with special attention to the inherent coupling of carbon, energy and water cycles (Figure 1).

## 2. Methods

### 2.1. Site descriptions

Data were obtained at four EC flux tower tropical forest locations (Figure 2). All sites were established by the Brazilian-led Large-Scale Biosphere-Atmosphere Experiment in Amazonia (LBA-ECO) (Keller et al., 2004) and members of the Brasil flux network (da Rocha et al., 2004; Restrepo-Coupe et al., 2013). Three EC stations comprise a longitudinal transect close to the equator ( $\sim 3^\circ S$ ) along the Amazon river from east to west, from high to low mean annual net radiation (Figure 2) and different seasonal patterns of monthly precipitation: Caxiuanã (CAX), the Tapajós National forest near Santarém (K67) and the Reserva Cuieiras near Manaus (K34). The fourth site, the Ji-Paraná Reserva Jaru (RJA) forest, is located at the southern margins of the basin, at latitude  $10^\circ S$ . For a detailed site description refer to previous works by da Rocha et al. (2009), Restrepo-Coupe et al. (2013, 2017) and Table S1.

### 2.2 Eddy flux (EC), meteorological and biometric data

Sensible heat ( $H$ ), water ( $ET$ ) and carbon fluxes ( $F_c$ ) were measured using the EC method (Baldocchi et al., 1988; Wofsy et al., 1993). Hourly average covariances were obtained from high frequency observations (20 Hz) of vertical wind velocity, virtual temperature ( $T_{son}$ ;  $^\circ C$ ), and water ( $H_2O_{mix}$ ;  $mmol\ mol^{-1}$ ), and carbon dioxide ( $CO_2$ ; ppm) mixing ratios measured with a 3D sonic anemometer (CSAT) and an infrared gas analyzer (LI6262) (Burba, 2010; Foken et al., 2012). The  $LE$  was calculated as the product of water mass transfer ( $ET$ ;  $mm\ day^{-1}$ ) and latent heat of vaporization ( $\lambda$ ;  $MJ\ kg^{-1}$ ), where  $LE = ET\ \lambda$ . The  $\lambda$  calculated as a function of air temperature (Brutsaert, 1982).



Meteorological observations included: air temperature ( $T_{air}$ ; °C), relative humidity ( $RH$ ; %), precipitation ( $Precip$ ; mm), wind speed ( $ws$ ;  $m\ s^{-1}$ ), turbulence measured as friction velocity ( $u_*$ ;  $m\ s^{-1}$ ), and the following radiation fluxes in  $W\ m^{-2}$ : incoming ( $SW_{down}$ ) and outgoing shortwave ( $SW_{out}$ ), and incoming ( $LW_{down}$ ) and outgoing longwave ( $LW_{out}$ ). Net radiation ( $Rn$ ;  $W\ m^{-2}$ ) was defined as the balance between incoming and outgoing fluxes ( $Rn = SW_{down} - SW_{out} + LW_{down} - LW_{out}$ ). A four-dome net radiometer, CNR1 (Kipp & Zonen CM3 ISO-class, thermopile pyranometer, CG3 pyrgeometer, PT100 RTD) was used for the measurement of  $SW_{down}$ ,  $SW_{out}$ ,  $LW_{down}$  and  $LW_{out}$ , at all sites. The shortwave ( $SW$ ) or solar radiation was defined as broadband radiation between 0.3 to 3  $\mu m$  and the longwave ( $LW$ ) as radiation with a spectral range from 3 and 300  $\mu m$ . An independent  $Rn$  measurement from a single-component radiometer was available at K34 and K67.

Hourly data were subject to various quality control procedures: Values found to be outside  $\pm 3$ -standard deviations from the mean were removed for  $ws$ ,  $RH$ , and  $T_{air}$ . Analogous and concurrent measurements were used to identify periods of instrument malfunction (e.g.  $T_{son}$  and  $T_{air}$ ) recognized by observations outside 2-times the standard deviations from the linear relationship between the variables. Similarly to processing carbon flux data, we removed  $LE$  fluxes measured during low turbulence conditions (given a site-specific  $u_*$  threshold,  $u_{*threshold}$ ), thus the EC method's no-advection assumption does not apply (see Restrepo-Coupe et al. 2013) (Table S1).

The energy balance was defined as  $Rn - \Delta = LE + H + \Delta Sh + \Delta Sc + \Delta Sb$ , where  $\Delta Sh$  is the sensible heat storage on the canopy layer storage,  $\Delta Sc$  is the energy change due to photosynthetic activity,  $\Delta Sb$  is the biomass heat storage, and  $\Delta$  is the imbalance (Figure S1 and S2). The  $\Delta$  term includes measurement errors (e.g. differences between the footprint of the radiation sensor and the EC and loss of low frequency large-scale eddies) and unaccounted fluxes: ground heat flux ( $G$ ) and changes in the latent heat flux stored on the air column below the EC system ( $\Delta Sle$ ). At K34 where profile temperature observations were not available, the  $\Delta$  included  $\Delta Sh$  and  $\Delta Sb$ , as well. The  $\Delta Sh$  was calculated as the hourly change in temperature across the air column (eight, five and four height levels at K67, RJA, and CAX, respectively) multiplied by air density and specific heat at constant pressure (Figure S3). The  $\Delta Sc$  was defined as the product of gross ecosystem

productivity (see Sec. 2.4.) and the specific energy of conversion due to photosynthesis ( $1.088 \times 10^4 \text{ J gCO}_2^{-1}$ ) (Moderow et al., 2009). We calculated  $\Delta Sb$  as the product of canopy-specific heat capacity ( $C_{veg} = 2958 \text{ J kg}^{-1} \text{ K}^{-1}$ ), live wet biomass ( $m_{veg}$ ;  $\text{kg m}^{-2}$ ) and the change in temperature at canopy level ( $T_{cpy}$ ; K). See SI for  $m_{veg}$  values and  $T_{cpy}$  heights. To flag possible outliers, as part of our QA procedures, we used the slope of the regression ( $Rn$  vs.  $LE+H+\Delta Sh+\Delta Sc+\Delta Sb$ ) assuming the observations outside 2-times the standard deviations from the linear relationship (see Figure S6).

We reviewed the seasonality of the energy balance residual as to improve the confidence in our analysis rather than determine  $LE$ -corrected values (i.e., we did not force energy balance closure). Note that we observed no statistically significant differences in the seasonal (monthly) energy balance closure (Figure S1 and S5). For an extensive review of the energy balance problem, the reader is invited to refer to the work of Foken (2008), subsequent studies (Mauder et al., 2018; Reed et al., 2018) and our supporting information (SI).

At each EC site, meteorological drivers for the LSMs were generated from the standard suite of climatic variables available for periods between 1999 and 2006. We analyzed data for 2000-2005 for K34, 2002-2004 for K67, 2000-2002 for RJA and 1999-2003 for CAX. Drivers included:  $LW_{down}$ ,  $SW_{down}$ ,  $T_{air}$ ,  $ws$ , near surface specific humidity ( $Q_{air}$ ;  $\text{g kg}^{-1}$ ), rainfall ( $Precip$ ;  $\text{mm month}^{-1}$ ), and surface atmospheric pressure ( $P_a$ ; hPa) (Figure 3). The  $\text{CO}_2$  concentration ( $\text{CO}_{2air}$ ; ppm) was fixed at 375 ppm, the average value during the period of measurements (de Goncalves et al., 2009). Observational data were filled using other nearby meteorological sites and/or the mean monthly diurnal cycle; however, only successive years with gaps no larger than two consecutive months were accepted. Although model drivers were gap-filled, regressions, and other calculations presented in this manuscript were implemented using only non-filled flux observations and meteorological values. We sampled the EC data to match the timing of the model drivers and output.

Biogeochemical fluxes can be sensitive to canopy structure and function. For our analysis we used 16-day values of leaf area index ( $LAI$ ), net primary productivity ( $NPP$ ) allocated to leaves ( $NPP_{leaf}$ ;  $\text{gC m}^{-2} \text{ d}^{-1}$ ), wood ( $NPP_{wood}$ ;  $\text{gC m}^{-2} \text{ d}^{-1}$ ) and litterfall ( $NPP_{litter}$ ;  $\text{gC m}^{-2} \text{ d}^{-1}$ ). Litterfall

data were available for all forests and included recently published values by Freire et al. (2020) for RJA. We used previously published  $LAI$  values -- see Table S1 for references, values and methods. For a description of biometric sampling methods see the original works of Metcalfe et al. (2007), Brando et al. (2010), Rice et al. (2004), and Fisher et al. (2007) and for calculations and a description of the  $NPP$  seasonal values see Restrepo-Coupe et al. (2017).

### 2.3. Surface emissivity ( $\epsilon_s$ ), Bowen ratio, outgoing longwave radiation ( $LW_{out}$ ), and other calculations

We used observations of the longwave radiation balance ( $LW_{down}$  and  $LW_{out}$ ) as per the integral of the Planck radiation function, the Stefan-Boltzmann equation, to obtain the measure of the surface's ability to emit energy by radiation, the Earth's surface spectral emissivity ( $\epsilon_s$ ):

$$LW_{out} = \epsilon_s \sigma_{SB} T_{skin}^4 + (1 - \epsilon_s) LW_{down} \quad \text{Equation 1}$$

where  $\sigma_{SB}$  is  $5.6704 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$  the Stefan-Boltzmann constant,  $T_{skin}$  is the skin temperature (K) and  $\epsilon_a$  is the effective emissivity of the atmosphere (Jin & Liang, 2006). The equation included the reflected fraction of  $LW_{down}$  the second term ( $(1 - \epsilon_s) LW_{down}$ ), following Kirchhoff's law, which assumes that absorptivity and emissivity are the same for each spectral band (Liou, 2002). We used canopy level temperature measurements (lagged as to reach a maximum four hours after peak  $T_{air}$ ) as a proxy for  $T_{skin}$  (Moderow et al., 2009) (see SI section 4). No contact thermometry was installed at any of the study sites. We solved for  $\epsilon_s$ :

$$\epsilon_s = \frac{LW_{out} - LW_{down}}{\sigma_{SB} T_{skin}^4 - LW_{down}} \quad \text{Equation 2}$$

The derivation of  $\epsilon_s$  is a simplification of a complex process: We did not account for the vertical variations of  $T_{air}$ , and we neglected the re-emission of  $LW$  radiation by water vapor.

Nonetheless, we are measuring  $LW_{down}$  and  $LW_{out}$  at the four forests and we see this calculation as an improvement over the assumed emissivity values used by some LSMs. Similarly, to identify possible bias on model  $LW_{out}$  calculations, we solved Equation 2 for  $T_{skin}$  assuming  $\epsilon_s$  values of 0.99 (see SI section 4).

Here we include 1-km grid MOD11A2.v6 (Wan et al., 2015) the land surface temperature (*LST*) product to scale and compare  $T_{air}$  measurements to satellite-derived land-surface temperature used by some models on their emissivity calculations (Figure S8).

To describe the forest optical brightness, we calculated the daytime albedo (top of the atmosphere radiation,  $TOA > 200 \text{ W m}^{-2}$ ) as the unitless ratio of outgoing to incoming solar radiation ( $\alpha = SW_{out}/SW_{down}$ ). We computed the TOA following Goudriaan (1986) and set a threshold of  $TOA$  and  $SW_{down} > 200 \text{ W m}^{-2}$  to constrain daytime observations. To characterize the heat transfer and the partition between water and sensible heat fluxes, we used the Bowen ratio calculated as the fraction of  $H$  to  $LE$  ( $Bowen = H/LE$ ). The Bowen ratio is used by some models as a driver in stomatal conductance and photosynthesis calculations (Berry et al., 2013; Sellers, 1985).

#### 2.4. Vegetation contributions to *ET*

To quantify the vegetation response to meteorology, we evaluated the seasonal differences between observed *ET* and the reference *ET* ( $ET_{ref}$ ) (also known as potential *ET*). The  $ET_{ref}$  is solely driven by atmospheric demand and climatic parameters and independent of the vegetation water use and soil factors. The  $ET_{ref}$  was calculated following the FAO Penman-Monteith method as:

$$ET_{ref} = \frac{\delta}{\delta + \gamma} Rn \quad \text{Equation 3}$$

where  $\gamma$  is the psychrometric coefficient ( $C_p P_a 10^3 / 0.622 \lambda$ ;  $\text{kPa K}^{-1}$ ), and  $\delta$  is the slope of vapor pressure curve ( $\delta = 4098 e_{sat} / T_{air}^{-2}$ ;  $\text{kPa K}^{-1}$ ), and  $C_p$  is the specific heat of air at constant pressure ( $\text{J kg}^{-1} \text{K}^{-1}$ ).

We calculated the ecosystem water use efficiency (*WUE*) as the ratio between daytime photosynthetic activity ( $TOA > 200 \text{ W m}^{-2}$ ) measured as the gross primary productivity ( $GPP_{day\&dry}$ ;  $\text{gC m}^{-2} \text{d}^{-1}$ ) to  $ET_{day\&dry}$  over a 16-day period ( $WUE = GPP_{day\&dry}/ET_{day\&dry}$ ;  $\text{gC mm}^{-1}$ ). The  $ET_{day\&dry}$  ( $\text{mm d}^{-1}$ ) was measured excluding observations during and 12-hours after precipitation, and using only daytime data, and was assumed to be the *ET* dominated by

transpiration ( $T$ ) fluxes rather than by direct evaporation ( $E$ ) from interception (e.g. after rain) and from condensation (e.g. dawn measurements). Similarly, the TOA threshold removed all early morning - late afternoon values from the  $WUE$  calculations, thus small  $ET$  values translated into abnormally high efficiencies without physical merit. Here, we use the term gross primary productivity ( $GPP$ ) interchangeably with gross ecosystem productivity ( $GEP$ ;  $\text{gC m}^{-2} \text{d}^{-1}$ ) and negative gross ecosystem exchange ( $GEE$ ;  $\text{gC m}^{-2} \text{d}^{-1}$ ), where  $GPP \sim GEP = -GEE$  (Stoy et al., 2006). The  $GEE$  was estimated from the measured daytime net ecosystem exchange ( $NEE$ ;  $\text{gC m}^{-2} \text{d}^{-1}$ ) by subtracting estimates of ecosystem respiration ( $R_{eco}$ ;  $\text{gC m}^{-2} \text{d}^{-1}$ ), which in turn were derived from nighttime  $NEE$  ( $GEE = -NEE + R_{eco}$ ). The  $NEE$  was calculated as the sum of the fluxes measured at the top of the tower and the  $\text{CO}_2$  storage flux ( $NEE = F_c + S_{\text{CO}_2}$ ) and filtered for low turbulence periods (site-specific  $u_{*thresh}$ ).  $R_{eco}$  was calculated as the average within a centered 5-day wide moving window, assuming at least 8 valid hours of nighttime  $NEE$  (we expanded the window up to 30 days until sufficient valid data were included). The selected  $R_{eco}$  moving window accounts for sensitivity to seasonally varying soil moisture. Daytime  $R_{eco}$  was assumed to be equal to nighttime  $R_{eco}$ . See SI and Restrepo-Coupe et al. (2013, 2017) for uncertainty analysis and additional methods.

To better understand the contribution of vegetation to  $LE$  and consequently to the partition of turbulent heat fluxes (Figure 1), we calculated the canopy stomatal resistance to water vapor ( $rsV$ ;  $\text{s m}^{-1}$ ) and the corresponding canopy conductance ( $G_S$ ;  $\text{mmol m}^{-2} \text{s}^{-1}$ ) following the flux-gradient method as described by Wehr and Saleska (2015; 2020, 2021) (see SI section 6 for calculations and sensitivity analysis).

## 2.5. Land surface models (LSMs)

We present output from four process-based land surface models that were part of the ‘Interactions between Climate, Forests, and Land Use in the Amazon Basin: Modeling and Mitigating Large Scale Savannization’ project (Powell et al., 2013; Restrepo-Coupe et al., 2017). We used the Community Land Model-Dynamic Global Vegetation Model version 3.5 (CLM3.5) (Gotangco Castillo et al., 2012; Oleson et al., 2008; Stockli et al., 2008), the Ecosystem Demography model version 2 (ED2) (Longo et al., 2018; Longo et al., 2019b; Medvigy et al., 2009), the Integrated Biosphere Simulator (IBIS) (Foley et al., 1996; Kucharik et al., 2000) and

the Joint UK Land Environment Simulator (JULES v.2.1) (Best et al., 2011; Clark et al., 2011). The LSMs energy and water cycle dynamics, including how radiation and conductances were calculated by models are presented in Table S2.

Models compute  $Rn$  as the sum of  $LW_{down}$  and  $SW_{down}$  (forcing drivers) minus the outgoing energy flux, the  $LW_{out}$  and  $SW_{out}$  calculated using parameters assigned to a plant functional type (PFT) and/or via different canopy radiation transfer models and equations (e.g. the two-stream model and the Beer-Lambert law) (Fisher et al., 2018). Later,  $Rn$  is partitioned into  $LE$  and  $H$ . This partition is determined by atmospheric demand and the amount of water available for evaporation and transpiration (if the water supply is exhausted, energy will ultimately be spent exclusively on  $H$ ). If water is available,  $LE$  will be driven by temperature, wind velocity, available radiant energy and will be modulated by  $G_s$  and aerodynamic conductance ( $Gi$ ) (Figure 1). The  $G_s$ , representing the exchange of  $CO_2$  and  $H_2O$  between multiple canopy leaves and the atmosphere, is controlled by meteorological and edaphic conditions given the ecosystem's structure, and by plant trait expressions that determine the photosynthetic capacity (e.g. quality and quantity of leaves and stomatal behavior). Therefore,  $G_s$  links the energy, carbon and water cycles and constitutes a key vegetation status descriptor for LSMs.

LSMs calculated the down-regulation factor for stomatal conductance due to soil water stress (FSW) (also known as the  $\beta$  term) following Oleson et al. (2008) (CLM3.5) and Castanho et al. (2016) (ED2, IBIS, and JULES). The FSW factor ranges from 0 (maximum stress) to 1 (no stress).

Model diagnostic variables complied with radiation energy and water conservation equations (Equation 6 and 7). The energy balance residual was always smaller than  $1 \text{ W m}^{-2}$ :

$$SW_{down} - SW_{out} + LW_{down} - LW_{out} - H - LE - G = \Delta S_b + \Delta S_h \quad \text{Equation 9}$$

And the water balance residual was less than  $1 \times 10^{-6} \text{ kg m}^{-2} \text{ s}^{-1}$ , defined by:

$$Prec - ET - R - GW + F = (\Delta_{intercept} + \Delta_{srfstor} + \Delta_{soilmoist})/dt \quad \text{Equation 10}$$

where  $R$  is surface runoff,  $GW$  is subsurface runoff,  $F$  is recharge from rivers, and the  $\Delta_{intercept}$ ,  $\Delta_{srfstor}$  and  $\Delta_{soilmoist}$  are changes in interception, surface storage, and soil moisture, respectively (all values in units of  $\text{kg m}^{-2} \text{s}^{-1}$ ).

## 2.6. Calculating seasonality and comparing models to observations

For each hour on the 16-day period we used all available measurements (minimum four observations per hour) (Figure S7). We calculated the mean of the average daily cycle (minimum 22/24 hours of the cycle were required for calculation of seasonal mean). This method avoids assigning less weight to those periods where we have fewer measurements. For example, at K34 precipitation was common in the late afternoon; therefore,  $LE$ ,  $H$ , and other measurements that depend on the sonic anemometer were unavailable during rainfall events (Figure S9). Seasonal  $WUE$  ( $GEP_{day\&dry}/ET_{day\&dry}$ ) and  $ET/ET_{ref}$  were calculated using 16-day ratios. The average annual cycle was calculated from all available 16-day periods when at least two measurements were available (2-years of data for each period).

Models were compared to observations based on the timing and amplitude metrics of their annual cycle. Correlation coefficient ( $r$ ), root-mean-square difference of model-observations (RMSE), and the ratio of their variances were determined for the 16-day multiple years' time series and the difference in amplitude and timing of the seasonal cycle were summarized using the unitless normalized standard deviation calculated as the ratio between model ( $\sigma_m$ ) and observations ( $\sigma$ ) standard deviation via Taylor diagrams (Taylor, 2001) (see Figure 3e for its interpretation). Sites missing from figures indicate that the model overestimated the seasonality of observations and  $\sigma$  was greater than two.

We used Type II linear regressions between fluxes, parameters and variables to understand and quantify the relationships between flux drivers and meteorological variables (e.g.  $H$  vs.  $Rn$ ) and between ecosystem characteristics and processes (e.g.  $LAI$  vs. albedo), thus acknowledging both variables carried some degree of uncertainty. To describe the statistical significance of regressions, we calculated p-values and the coefficient of determination ( $r^2$ ), and the Akaike's Information Criterion (AIC), among other descriptors. We compared the resulting linear models

to simulations (benchmark) to identify key flux drivers and determine when and how LSMs can be under-utilizing the available variable information (Abramowitz, 2005; Best et al., 2015).

### 3. Results

#### 3.1. Seasonal meteorology and evapotranspiration (*ET*)

All sites showed contrasting degrees of seasonality in terms of rain, temperature, insolation, and/or day-length; including differences in the amplitude of the radiation and precipitation annual cycles and the timing metrics that define the start, end, peak and dry season length (Figure 3). Mean annual precipitation at RJA and K67 was close to 2000 mm compared to 2500 mm at CAX and K34. The dry season varied in length and strength from the 1-month long at K34 to the 5-month at K67 and RJA (Figure 3). Although the dry season at K34 only lasted for one month (August), there was a period from July to October when the precipitation was lower than the annual mean and when we observed above average incoming radiation values (similar seasonality to K67 and CAX). The number and intensity of precipitation events was different: (1) CAX with frequent-low intensity rainfall ( $\geq 250$  events month<sup>-1</sup> of  $< 0.5$  mm hr<sup>-1</sup>), (2) strong seasonal changes at RJA (dry-season with few lower than  $0.5$  mm hr<sup>-1</sup> intensity events and wet-season with  $\sim 50$  events higher than  $2.5$  mm hr<sup>-1</sup>), and (3) K67 and K34 close to aseasonal intensities ( $2.5$  mm hr<sup>-1</sup>); however, there were fewer events at K67 ( $\leq 50$  events month<sup>-1</sup>) compared to K34 ( $\leq 100$  events month<sup>-1</sup>) (Figure S9).

The observed annual cycle of *ET* showed three different patterns across forests: (1) maximum water vapor flux at the beginning of the dry season declining as the season progressed at the two wettest locations (K34 and CAX); (2) a well-defined *ET* cycle, with a middle of the dry-season peak at K67; and (3) an aseasonal *LE* flux at the southern forest of RJA (Figure 3c and 4a). Modeled *ET* showed seasonal synchronicity with observations at the two wettest sites (K34 and CAX); however, LSMs overestimated the dry-season flux by 150-20 mm month<sup>-1</sup> (Figure 3c). At K67 and RJA, models exaggerated the amplitude of the water flux seasonal cycle by 180-20 mm month<sup>-1</sup>. At these drier locations, LSM's predicted reductions in dry season *ET* that were generally driven by the available soil moisture, as demonstrated by the statistically significant relationship between flux and the plant available water model diagnostic FSW (p-value  $< 0.01$   $r^2$  from 0.1 (IBIS) to 0.7 (ED2) at K67 and 0.3 (ED2) to 0.7 (CLM3.5) at RJA) (Figure 3d and



S10). By contrast, observations showed available energy driving  $ET$  at all sites (Table S3). The slope of the regression between seasonal  $LE$  vs.  $Rn$  (type II, zero intercept) was  $\sim 0.6$  (Figure S11) ( $r^2 = 0.7$  at CAX,  $0.8$  at K34,  $0.5$  at K67 and  $0.1$  at RJA). Seasonal  $T_{air}$  and  $LE$  showed a significant positive correlation ( $r^2 = 0.42$ ,  $p$ -value $<0.01$ ) at only one site, K67 (Table S3). The  $ET_{day}$  was close to constant ( $7.7 \text{ mm day}^{-1}$ ) at the southern forest of RJA. RJA was the only forest where we observed no significant correlation between  $Rn$  and  $ET$  ( $r^2 < 0.1$ ,  $p$ -value $=0.9$ ) however, the linear model had a low RMSE value ( $7.78 \text{ W m}^{-2}$ ). Moreover, all site regressions between  $Rn_{day}$  and  $LE_{day}$  showed RJA observations following the general trend (Figure 8).

### 3.2. Partition of net radiation into turbulent fluxes

At the equatorial Amazon forests (K34, CAX, and K67), the 16-day cycle of  $H$  showed a maximum at the beginning and a minimum at the end of the dry season (Figure 4b). By contrast,  $H$  was close to aseasonal at RJA (a slight increase by the middle of the dry period). Models were able to capture the seasonal cycle of  $H$  at CAX; however, the dry-season  $H$  was underestimated by most of the LSMs at K34. LSMs overestimated  $LE$  and were out of phase with observations at K67 and RJA (Figure 4b). At K34 and RJA the relationship between observed  $H$  and  $LE$  was weak ( $r^2 < 0.2$ ,  $p$ -value $<0.01$ ) and significant at CAX and K67 ( $r^2 = 0.6$ ,  $p$ -value $<0.01$ ) (Figure S11). At RJA and CAX measurements of  $Rn$  explained 50% of the  $H$  seasonal variability. Moreover,  $H$  was significantly correlated with  $Rn$ , the slope (zero intercept) varying from  $0.12$  at K67,  $0.15$  at CAX and RJA, to  $0.22$  at K34 ( $r^2 \sim 0.4$ ,  $p$ -value $<0.01$ ) (Figure S11).

Observations showed that Bowen ratio values were nearly constant at  $\sim 0.32$  for K34 (highest) and at  $\sim 0.21$  for RJA and K67 (lowest among forests). We found that the Bowen ratio for the four LSMs was lower than the observed value at the two wettest locations (K34 and CAX) and above measurements at the two driest forests (K67 and RJA). Simulations showed a strong increase in Bowen ratio during the dry season at K67 (IBIS and ED2) and at RJA (all models) (Figure 4c).

Hourly and seasonal observations showed a good seasonal energy balance closure (slope  $LE + H$  vs.  $Rn$ ) ranging from 90% (CAX), 88% (K67 and K34) to 83% (RJA) (Figure S1 and S2). By comparison, FLUXNET sites have an average imbalance of  $\sim 20\%$  (Wilson et al., 2002). Where

profile temperature data were available, the introduction of canopy and biomass heat storage improved the overall hourly balance, especially the energy closure at dawn and dusk (see supplemental material, Figure S3). The  $\Delta$  showed a statistically significant correlation to  $Rn$  ( $\Delta \sim 0.1Rn$ ,  $r^2 > 0.8$ ,  $p$ -value  $< 0.01$ ) and no correlation to turbulence,  $T_{air}$  or rainfall (Figure S3 and S4). Therefore, we had no indication of lost fluxes due to advection (low  $u^*$ ) or errors associated to turbulence bursts (high  $u^*$ ). At CAX, frequent rainfall events made EC measurements challenging, and extensive periods of data needed to be removed (causing gaps in many regressions and figures). Rainfall events at CAX were less intense, however more frequent than at any other site (see Figure S9).

### 3.3. Radiation balance: Outgoing longwave ( $LW_{out}$ ) and reflected shortwave ( $SW_{out}$ ) radiation

The  $SW_{out}$  is determined by the surface reflectance (e.g. we see low  $SW_{out}$  values in dark bodies, and high values in bright bodies) and its relation to  $SW_{down}$  is measured as albedo ( $\alpha$ ) (Figure 5). Seasonality of  $\alpha$  showed modest increases as the dry-season progressed at all sites and was in-phase with the radiation seasonal cycle (Figure S15). Peak  $\alpha$  values (when forest was at its brightest) were observed by the middle of the dry season at the equatorial Amazon sites (CAX, K34, and K67) and at the end of the dry period at RJA (Figure 6a). The average  $\alpha$  was 0.12 at RJA, K34 and K67 and 0.09 at CAX. Negative regressions between precipitation and  $\alpha$  (the forest was darkest at the peak of the wet season) were statistically significant at all forests ( $p$ -values  $< 0.01$  with  $r^2$  values up to 0.4 at K67 and K34) (Figure S13). The forest characteristics showed some degree of correlation: (1) low  $LAI$  to high  $\alpha$  (negative slope) at CAX, and (2) high  $NPP_{leaf}$  to high canopy brightness (positive slope) at K67, RJA, and K34 (Figure S14). However, at all sites, the timing of maximum  $\alpha$  did correlate with peak leaf-flush greenness index phenocam observations (e.g. Lopes et al., 2016). Models overestimated  $\alpha$  annual mean across sites and underestimated the amplitude of the  $\alpha$  seasonal cycle.

Observations showed mean monthly values of  $SW_{out}$  close to  $20 \text{ W m}^{-2}$  at most forests (Figure 5a). The models captured the seasonal cycle of  $SW_{out}$  at all sites except RJA. The  $SW_{out}$  was significantly correlated with  $SW_{down}$  ( $r^2 = 0.9$  at K34 and RJA,  $0.7$  at K67 and  $r^2 = 0.5$  at CAX;  $p$ -values  $< 0.01$ ), with the slope of their linear relationship increasing from wet to dry forests, such

as 0.12 at K34 and CAX, 0.13 at K67 and 0.14 at RJA (Figure S15). Seasonal  $LW_{out}$  was significantly correlated with  $LW_{down}$ , however  $R^2$  values were low ( $r^2 = 0.34$  at K34, 0.5 at K67 and  $r^2 = 0.2$  at CAX and RJA,  $p$ -values  $< 0.01$ ) with a positive slope at K34 and RJA and a negative regression ( $LW_{down}$  increased faster than  $LW_{out}$  and surface-canopy temperature warming at a lower rate than the air) at CAX and K67 (Figure S17). At K67, CAX, and RJA, models captured the amplitude of the seasonal  $LW_{out}$  cycle, however at K34 the  $LW_{out}$  all models' simulations were out of phase with observations (Figure 5b).

The amplitude of the annual surface emissivity ( $\epsilon_s$ ) cycle representing the ability of the surface to emit longwave radiation, showed high dry-season values at RJA and CAX (Figure 6b). By contrast at CAX, observations showed low wet season  $\epsilon_s$  values. At K34 and K67 observed  $\epsilon_s$  were higher than 0.98 and close to 0.95 at RJA. We found statistically significant correlations ( $p < 0.01$ ,  $r^2$  range 0.3 to 0.8) between  $\epsilon_s$  and rainfall (positive) and  $T_{air}$  (negative) at K34 and vice versa at CAX -- no significant correlation was observed at K67 and RJA (Figure S13). LSMs generally did not capture the magnitude or seasonality of  $\epsilon_s$ , and no LSM aligned with observations across all sites (Figure 6b). Assuming constant  $\epsilon_s$  values of  $\sim 0.99$  in agreement with satellite measurements (Figure S8), showed models either overestimated  $T_{skin}$  ( $\sim 1$  to  $5^\circ\text{C}$ ) or underestimated  $\epsilon_s$  (Figure S12).

### 3.4. Ecosystem characteristics and contributions to the water and energy flux seasonality

The ratio between observed  $ET$  and  $ET_{ref}$  can be used to identify the periods when  $ET$  does not show any signs of water-supply limitation and the flux is mostly driven by atmospheric demand and solar radiation (Figure 3c and S19). Only during the wettest months at K34 we observed  $ET$  equivalent to  $ET_{ref}$  ( $ET/ET_{ref} \sim 100\%$ ) and  $\sim 70\%$  during the driest period (Figure 7a). In general, the slope of the regression between  $ET$  and  $ET_{ref}$  varied from 0.66 (RJA) to 0.74 (K67 and K34), with statistically significant differences between wet and dry season values only seen at RJA and K34 (Figure S19).

The vegetation control over  $ET$ , here represented by  $G_s$ , showed different degrees of seasonality and trends across forests (Figure 7b); nevertheless minimum values were observed at various

times during the dry-season at all sites: (1) At CAX the dry-season  $G_s$  was close to  $0.4 \text{ mmol m}^{-2} \text{ s}^{-1}$  and up to  $1.4 \text{ mmol m}^{-2} \text{ s}^{-1}$  -- the highest  $G_s$  values were observed at this site; (2) at K34 and K67, the  $G_s$  gradually decreased from the transition wet-to-dry period to reach minimum values at the onset of the rainy season. (3) RJA experienced a reduction in  $G_s$  mid wet-season to mid dry-season (an all site minima of  $3 \text{ mmol m}^{-2} \text{ s}^{-1}$ ). Models were able to capture  $G_s$  at most forests, however they underestimated the amplitude of the annual cycle at K34 and CAX (Figure 7b). The tradeoff between losing water through transpiration and gaining carbon showed different patterns across sites, suggesting leaf-level adaptations and ecosystem-level variation. For example, seasonal  $G_s$  showed a negative relationship to incoming radiation at K34, RJA, and during the dry season of K67 ( $r^2 < 0.3$ ,  $p\text{-value} < 0.01$ ). By contrast, higher  $SW_{down}$  correlated to high  $G_s$  at the very seasonal forest of CAX (where we observed the highest wet-period rainfall values among the four forests) and during the wet-season at K67 (Figure 8b and S20). In general,  $G_s$  was positively related to precipitation (Figure S21).

The ratio between ecosystem carbon-uptake and transpiration-dominated  $ET$ , here presented as  $WUE$  was correlated to  $G_s$  at CAX (negative,  $r^2=0.25$ ,  $p\text{-value} < 0.01$ ) and RJA (positive,  $r^2=0.48$ ,  $p\text{-value} < 0.01$ ) (Figure S22). A significant regression was observed at K67 only if  $WUE$  was lagged 2-months (minimum  $WUE$  preceded minimum  $G_s$ ) (Figure 7b). The  $WUE$  changes were non-significantly correlated to  $G_s$  at K34. Minimum  $WUE$  values were observed at the beginning of the dry season at equatorial sites (CAX, K34 and K67) and at the end of the dry period at RJA. The largest values of  $WUE$ , indicative of the highest photosynthetic rate per water use, were observed at different times for different sites when precipitation was  $> 100 \text{ mm month}^{-1}$  (start of at K34 and K67 and end of the wet season at RJA all at  $\sim 2.6 \text{ gC mm}^{-1}$ ) (Figure 7c). Most models were able to correctly estimate seasonal values of  $WUE$  and  $G_s$ , some overestimating  $G_s$  values at K34 and  $WUE$  at K67.

We used the Bowen ratio to describe the dominant type of heat transfer across the forests -- where  $LE$  clearly dominated the turbulent flux ( $H < 0.2 LE$ ). The relationship between Bowen ratio and  $G_s$  showed that at relatively high Bowen values  $> 0.3$ , the  $G_s$  reached a minimum of  $\sim 0.35 \text{ mmol m}^{-2} \text{ s}^{-1}$  (no further reductions were observed) (Figure 8a).

## 4. Discussion

This study identified three main tropical forest properties (relationships among fluxes and between fluxes and vegetation characteristics) that if understood and implemented in LSMs equations and/or benchmarking exercises could reduce the differences between observations and model estimates of seasonal  $ET$ ,  $Rn$  and  $H$  exchange: (1) Turbulent flux partitioning (e.g. high correlation between  $Rn$  and both turbulent fluxes, and nearly aseasonal Bowen ratio values), (2) representation of canopy reflectance and emissivity (e.g. albedo's annual cycle showed significantly lower absolute values and greater than expected amplitudes) and (3) endogenous ecosystem or physiology-related seasonality (e.g., leaf-level stomatal and  $WUE$  dynamics driven by leaf ontogeny and demography). These processes are related to surface energy properties, canopy-atmosphere water dynamics, their interactions, and more importantly the coupling between energy-carbon and water exchange. Here, we discuss some of our findings and suggest future observational and modeling work to improve simulations of tropical water and energy fluxes.

### 4.1. Determinants and distribution of net radiation into turbulent fluxes

Observations showed  $ET$  to be driven by radiation rather than by moisture availability as predicted by models. The  $R_n$  was able to explain more than 60% of the 16-day  $LE$  values and although we report a low  $r^2$  for the  $LE$  vs.  $Rn$  regression at the southern forest of RJA, the coefficient of determination was driven by the low amplitude of the seasonal  $LE$  and  $Rn$  flux rather than the linear regressions not being able to predict  $LE$ .

Analysis of variability of the observed Bowen number annual cycle showed a nearly aseasonal ratio ( $\sim 0.3$  at the wet sites of K34 and CAX, and 0.21 at the dry sites K67 and RJA, Figure 4c). This suggests a proportional scaling of the forest's energy balance at each location ( $H$  was a constant fraction of  $LE$ ). There was a relationship between the direction of bias in Bowen ratio estimates and site annual precipitation. LSMs overestimated dry-season Bowen values at the driest locations of K67 and RJA and underestimated the ratio at the wettest forests of K34 and CAX (models overestimated  $LE$  and underestimated  $H$ ) (similar to Best et al., 2015; Haughton et al., 2016; Morales et al., 2005). The expectation of a higher Bowen ratio (increase importance of

$H$  over  $LE$ ) at the drier sites did not apply at these tropical forests and could be explained by: (1) LSMs had a negative bias in dry-season  $R_n$ . (2) Models underestimated dry season  $LE$ , probably based on the incorrect assumption that water limitation (supply) rather than radiation (demand) drove the water flux (Federer, 1982). (3) LSMs may have difficulties simulating access to soil water at clay soils (e.g. K67) and although some recent model improvements have addressed this issue (e.g. ED2 see Longo et al., 2019a), measurements of field capacity and hydraulic conductivity were unavailable at our and other similar study sites. (4) Transpiration estimates may require to include processes related with plant hydraulics, like the addition of stem-water and other additional storage terms (e.g., CLM5 see Yan et al., 2020). (5) The time of rainfall, precipitation intensity and number of events (here we report significant differences among forest sites), rather than absolute precipitation values; may significantly influence the  $H/LE$  partition. Thus as rainfall characteristics and forest canopy structure (see item 6) can be key in defining how much water would be intercepted (directly evaporated), drained, and/or infiltrated (stored and later supplied). (6) Models may be assuming excess  $E$  from leaves surfaces (e.g. because of the high  $LAI$  forest values) and not enough water would be reaching the soil for infiltration during the wet season. This “water deficit” would be carried out into the dry season, limiting the moisture available for transpiration and artificially increasing  $H$ .

#### 4.2. Representation of canopy reflectance (albedo) and thermal properties

Although significant, the differences between modeled-observed  $ET$  cannot be explained solely by the way models partition  $H$  and  $LE$  fluxes (Haughton et al., 2016). This study shows that correct turbulent flux estimations require reliable  $R_n$  estimates. Most LSMs were able to capture the seasonal cycle of  $R_n$ . Thus,  $SW_{down}$  was provided to all models as a meteorological driver and dominated  $R_n$ . However, at CAX and RJA, both model  $LW_{out}$  and  $SW_{out}$  were higher than observations and consequently, seasonal values of  $R_n$  were underestimated. In some instances, the model-observation alignment was the result of obtaining the right answer for the incorrect reasons (e.g. LSMs overestimated  $SW_{out}$  and underestimated  $LW_{out}$  at K34). Models that consistently estimated higher than observed  $LW_{out}$  values may have to address the following issues: (1) the vegetation storage pool/heat capacity may be too low and/or (2) underestimated transpiration values, both causing  $T_{skin}$  to be too high. Additional measurements (e.g. thermal

cameras, sapflow sensors, soil moisture profiles, and H<sub>2</sub>O isotopes) would be necessary to measure  $T_{skin}$ , to infer the relationships between  $LE$ ,  $H$  and vegetation temperature and as to understand the mechanisms driving the relations between  $LW_{down}$  and  $LW_{out}$ .

Biases in LSMs  $Rn$  can also be attributed to  $SW_{out}$  calculations. Observed low albedos did contrast with model simulations resulting in more reflective (brighter) forest surfaces. Models underestimated the amount of canopy absorbed energy and may be imposing an “artificial” cooling effect. Surface albedo will be highly dependent on the leaf spectral properties and in general, canopy reflectance models relate higher  $LAI$  values to low albedo values (e.g. PROSAIL (Féret et al., 2017) assumes albedos  $\sim 0.2$  for a  $LAI > 4$ ) or albedos are parameterized as a constant (Hollinger et al., 2010). Nevertheless, we observed opposite sign regressions between  $LAI$  and albedo at CAX. Thus, indicating that  $\alpha$  was not only driven by the quantity of leaves, but by leaf quality and vegetation reflective surfaces (e.g. wood and epiphylls) (Chavana-Bryant et al., 2016; Wu et al., 2017). Across the Amazon, leaf phenology has shown to be a key driver of  $ET$  and carbon uptake (Albert et al., 2018; Chen et al., 2020; Manoli et al., 2018; Restrepo-Coupe et al., 2013; Wu et al., 2017) and should be incorporated/improved on the derivation of energy, radiation and water fluxes, as well.

### 4.3. Ecosystem characteristics and their contributions to the water and energy flux seasonality

Our results showed that when the  $H$  was higher than 20%  $LE$ , the  $G_s$  reached a minimum of  $\sim 0.35 \text{ mmol m}^{-2} \text{ s}^{-1}$ , with no further reductions. Indicating that the vegetation continued to transpire at the same or higher rate under relatively high Bowen ratio conditions. This finding may be not surprising as Stahl et al. (2013) found that during low precipitation periods 50% of a sample of 65 large tropical trees relied on soil water below 1-m depth, and others have reported hydraulic redistribution, stem-water storage and additional processes that may explain forests access to water during the dry-season (Christoffersen et al., 2014; Oliveira et al., 2005; Yan et al., 2020). Moreover, the gradual dry season decrease in  $G_s$  (as similarly reported in Christoffersen et al. (2014) and Costa et al. (2010)) and increase in  $LE$  observed at the equatorial forests, highlights the very significant role of evaporation during this period. However, only seasonal inventories of leaf age and traits, and evaporation vs. transpiration measurements (e.g.

H and O isotopes) will offer models validation data to avoid misrepresentation of the plant water exchange (e.g. under/over estimating photosynthesis and water use efficiency) (Lawrence et al., 2007).

Leaf-level stomatal conductance ( $g_s$ ) is expected to maximize carbon uptake while also reducing water loss from leaves (or reducing the carbon cost of hydraulic failure) when water is limiting (Anderegg et al., 2018; Medlyn et al., 2011; Sperry et al., 2017), and generally is site-specific and driven by adaptation to the different atmospheric seasonal drivers (Brum et al., 2018). Ecosystem level vegetation controls (e.g. *LAI* and leaf age and position across the canopy profile) determine the water flux, rate of photosynthesis and the “acceptable” degree of water stress the forest can tolerate during the dry season (Albert et al., 2018; Restrepo-Coupe et al., 2013; Wu et al., 2017, 2017). Similar to  $G_s$ , at all four forests we observed contrasting degrees of seasonality in terms of *WUE* (with a range of  $\pm 25\%$  of all year mean) and its timing metrics. Like *GEP*, across equatorial forests *WUE* increased as the dry season progressed and vice versa at RJA. At ecosystem scale we found that the regression between *WUE* and  $G_s$  was not statistically significant at K34 and K67, negatively correlated at CAX and positively at RJA (Figure S22). The lack of correlation between  $G_s$  and *WUE* would be driven by seasonal differences in intercellular  $\text{CO}_2$  concentrations, atmospheric pressure and humidity, vegetation growth temperature and other canopy characteristics (Lin et al., 2015; Medlyn et al., 2011, 2012). For example, higher *VPD* can increase transpiration and reduce *WUE* without any change in  $G_s$  and vice versa.

#### 4.4. Considerations for model improvement

This paper describes the seasonal patterns of different energy and water flux constituents and examines the relationships between them and different forest characteristics and climate variables at four tropical forests. We compared eddy covariance and biometric measurements to LSM simulations, as models represent our current understanding of the different atmosphere-biosphere processes at global and continental scales and are the ideal tool to predict vegetation responses to changes in climate. Our analysis highlights *forest phenology* as a significant driver of vegetation-atmosphere exchange and in particular, our data showed LSMs: (1) underestimated the amount of solar radiation the forests absorb and dry-season increases because we lack



information regarding the relationship between leaf density and reflectance properties at high  $LAI$  values; (2) similarly, interception and direct evaporation may be overestimated at high  $LAI$  forests, and consequently LSMs may be underestimating infiltration and transpiration fluxes, overestimating canopy temperature, and consequently driving LSMs output (3) to inaccurate estimations of  $LW_{out}$  (e.g., reducing the soil moisture content and increased canopy temperature would lead to unrealistically high  $T_{skin}$  and hence incorrect estimates of  $LW_{out}$ ) and  $SW_{out}$  (e.g. if we incorrectly characterize forest structure albedo will be too high). This seasonal bias on the outgoing flux (emissivity and albedo) dominated the model-observation  $Rn$  differences and will have an effect in the estimation of  $H$ ,  $LE$  fluxes and the Bowen ratio. Our findings can be used to benchmark LSMs and develop more robust plant functional type parametrization. Improvements in model development will translate into better predictions of future surface-atmosphere exchange.

### Acknowledgments

This research was funded by the Gordon and Betty Moore Foundation "Simulations from the Interactions between Climate, Forests, and Land Use in the Amazon Basin: Modeling and Mitigating Large Scale Savannization" project, the National Aeronautics and Space Administration (NASA) LBA CD-32, NASA LBA-DMIP project (NNX09AL52G), and NASA-ROSES (NNX17AF65G). N.R.C. acknowledges the support from the City of Toronto Children and Human Services, Dr. Debra Wunch, Dr. Dylan Jones, and the Department of Physics at the University of Toronto. B.O.C. was funded in part by the U.S. DOE (BER) NGEE-Tropics project to LANL. L.M.M acknowledges the UK Natural Environment Research Council funding (UK Earth System Modelling Project, UKESM, Grant NE/N017951/1). L.P.A. was supported by a Voss postdoctoral fellowship at the Institute at Brown for Environment and Society. M.L. was supported by the NASA postdoctoral program, administered by the Universities Space Research Association under contract with NASA. Research carried out at the JPL-California Institute of Technology, was under a contract with NASA. The authors would like to thank the staff of each tower site for their technical support and extensive fieldwork.

### References

Abramowitz, G. (2005). Towards a benchmark for land surface models. *Geophysical Research*

- Letters*, 32(22). <https://doi.org/10.1029/2005GL024419>
- Albert, L. P., Wu, J., Prohaska, N., Camargo, P. B. de, Huxman, T. E., Tribuzy, E. S., Ivanov, V. Y., Oliveira, R. S., Garcia, S., Smith, M. N., Junior, R. C. O., Restrepo-Coupe, N., Silva, R. da, Stark, S. C., Martins, G. A., Penha, D. V., & Saleska, S. R. (2018). Age-dependent leaf physiology and consequences for crown-scale carbon uptake during the dry season in an Amazon evergreen forest. *New Phytologist*, 219(3), 870–884. <https://doi.org/10.1111/nph.15056>
- Anderegg, W. R. L., Wolf, A., Arango-Velez, A., Choat, B., Chmura, D. J., Jansen, S., Kolb, T., Li, S., Meinzer, F. C., Pita, P., Dios, V. R. de, Sperry, J. S., Wolfe, B. T., & Pacala, S. (2018). Woody plants optimise stomatal behaviour relative to hydraulic risk. *Ecology Letters*, 21(7), 968–977. <https://doi.org/10.1111/ele.12962>
- Aragão, L. E. O. C., Malhi, Y., Barbier, N., Lima, A., Shimabukuro, Y., Anderson, L., & Saatchi, S. (2008). Interactions between rainfall, deforestation and fires during recent years in the Brazilian Amazonia. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 363(1498), 1779–1785. <https://doi.org/10.1098/rstb.2007.0026>
- Aragão, L. E. O. C., Malhi, Y., Roman-Cuesta, R. M., Saatchi, S., Anderson, L. O., & Shimabukuro, Y. E. (2007). Spatial patterns and fire response of recent Amazonian droughts. *Geophysical Research Letters*, 34, 5 PP. <https://doi.org/200710.1029/2006GL028946>
- Baker, I. T., Prihodko, L., Denning, A. S., Goulden, M., Miller, S., & Rocha, H. R. da. (2008). Seasonal drought stress in the Amazon: Reconciling models and observations. *Journal of Geophysical Research*, 113(G1), 1–10. <https://doi.org/10.1029/2007JG000644>
- Baldocchi, D. D., Hincks, B. B., & Meyers, T. P. (1988). Measuring Biosphere-Atmosphere Exchanges of Biologically Related Gases with Micrometeorological Methods. *Ecology*, 69(5), 1331–1340. <https://doi.org/10.2307/1941631>
- Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., Seibt, U., Stimler, K., Yakir, D., & Zhu, Z. (2013). A coupled model of the global cycles of carbonyl sulfide and CO<sub>2</sub>: A possible new window on the carbon cycle. *Journal of Geophysical Research: Biogeosciences*, 118(2), 842–852. <https://doi.org/10.1002/jgrg.20068>

- Best, M. J., Abramowitz, G., Johnson, H. R., Pitman, A. J., Balsamo, G., Boone, A., Cuntz, M., Decharme, B., Dirmeyer, P. A., Dong, J., Ek, M., Guo, Z., Haverd, V., van den Hurk, B. J. J., Nearing, G. S., Pak, B., Peters-Lidard, C., Santanello, J. A., Stevens, L., & Vuichard, N. (2015). The Plumbing of Land Surface Models: Benchmarking Model Performance. *Journal of Hydrometeorology*, *16*(3), 1425–1442. <https://doi.org/10.1175/JHM-D-14-0158.1>
- Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. . L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., & Harding, R. J. (2011). The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes. *Geosci. Model Dev.*, *4*(3), 677–699. <https://doi.org/10.5194/gmd-4-677-2011>
- Bonan, G. B. (2008). Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests. *Science*, *320*(5882), 1444–1449. <https://doi.org/10.1126/science.1155121>
- Bony, S., Bellon, G., Klocke, D., Sherwood, S., Fermepin, S., & Denvil, S. (2013). Robust direct effect of carbon dioxide on tropical circulation and regional precipitation. *Nature Geoscience*, *6*(6), 447–451. <https://doi.org/10.1038/ngeo1799>
- Brando, P. M., Goetz, S. J., Baccini, A., Nepstad, D. C., Beck, P. S. A., & Christman, M. C. (2010). Seasonal and interannual variability of climate and vegetation indices across the Amazon. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(33), 14685–14690. <https://doi.org/10.1073/pnas.0908741107>
- Brum, M., Vadeboncoeur, M. A., Ivanov, V., Asbjornsen, H., Saleska, S., Alves, L. F., Penha, D., Dias, J. D., Aragão, L. E. O. C., Barros, F., Bittencourt, P., Pereira, L., & Oliveira, R. S. (2018). Hydrological niche segregation defines forest structure and drought tolerance strategies in a seasonal Amazon forest. *Journal of Ecology*. <https://doi.org/10.1111/1365-2745.13022>
- Brutsaert, W. (1982). *Evaporation Into the Atmosphere: Theory, History and Applications*. <http://www.powells.com/biblio?isbn=9789027712479>
- Burba, G. (2010). *A brief practical guide to eddy covariance flux measurements: Principles and workflow examples for scientific and industrial applications* /. LI-COR,.
- Castanho, A. D. de A., Galbraith, D., Zhang, K., Coe, M. T., Costa, M. H., & Moorcroft, P.

- (2016). Changing Amazon biomass and the role of atmospheric CO<sub>2</sub> concentration, climate, and land use. *Global Biogeochemical Cycles*, 30(1), 18–39.  
<https://doi.org/10.1002/2015GB005135>
- Chagnon, F. J. F., & Bras, R. L. (2005). Contemporary climate change in the Amazon. *Geophysical Research Letters*, 32(13), L13703. <https://doi.org/10.1029/2005GL022722>
- Chavana-Bryant, C., Malhi, Y., Wu, J., Asner, G. P., Anastasiou, A., Enquist, B. J., Cosio Caravasi, E. G., Doughty, C. E., Saleska, S. R., Martin, R. E., & Gerard, F. F. (2016). Leaf aging of Amazonian canopy trees as revealed by spectral and physiochemical measurements. *The New Phytologist*. <https://doi.org/10.1111/nph.13853>
- Chen, X., Maignan, F., Viovy, N., Bastos, A., Goll, D., Wu, J., Liu, L., Yue, C., Peng, S., Yuan, W., Conceição, A. C. da, O'Sullivan, M., & Ciais, P. (2020). Novel Representation of Leaf Phenology Improves Simulation of Amazonian Evergreen Forest Photosynthesis in a Land Surface Model. *Journal of Advances in Modeling Earth Systems*, 12(1), e2018MS001565. <https://doi.org/10.1029/2018MS001565>
- Christoffersen, B. O., Restrepo-Coupe, N., Arain, M. A., Baker, I. T., Cestaro, B. P., Ciais, P., Fisher, J. B., Galbraith, D., Guan, X., Gulden, L., van den Hurk, B., Ichii, K., Imbuzeiro, H., Jain, A., Levine, N., Miguez-Macho, G., Poulter, B., Roberti, D. R., Sakaguchi, K., ... Saleska, S. R. (2014). Mechanisms of water supply and vegetation demand govern the seasonality and magnitude of evapotranspiration in Amazonia and Cerrado. *Agricultural and Forest Meteorology*, 191(0), 33–50. <https://doi.org/10.1016/j.agrformet.2014.02.008>
- Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., & Cox, P. M. (2011). The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics. *Geosci. Model Dev.*, 4(3), 701–722. <https://doi.org/10.5194/gmd-4-701-2011>
- Costa, M. H., Biajoli, M. C., Sanches, L., Malhado, A. C. M., Hutyra, L. R., Rocha, H. R. da, Aguiar, R. G., & Araújo, A. C. de. (2010). Atmospheric versus vegetation controls of Amazonian tropical rain forest evapotranspiration: Are the wet and seasonally dry rain forests any different? *Journal of Geophysical Research: Biogeosciences*, 115(G4). <https://doi.org/10.1029/2009JG001179>
- da Rocha, H. R., Goulden, M. L., Miller, S. D., Menton, M. C., Pinto, L. D. V. O., de Freitas, H.

- C., & e Silva Figueira, A. M. (2004). Seasonality of Water and Heat Fluxes over a Tropical Forest in Eastern Amazonia. *Ecological Applications*, *14*(sp4), 22–32. <https://doi.org/10.1890/02-6001>
- da Rocha, H. R., Manzi, A. O., Cabral, O. M., Miller, S. D., Goulden, M. L., Saleska, S. R., R.-Coupe, N., Wofsy, S. C., Borma, L. S., Artaxo, P., Vourlitis, G., Nogueira, J. S., Cardoso, F. L., Nobre, A. D., Kruijt, B., Freitas, H. C., von Randow, C., Aguiar, R. G., & Maia, J. F. (2009). Patterns of water and heat flux across a biome gradient from tropical forest to savanna in Brazil. *Journal of Geophysical Research: Biogeosciences*, *114*(G1), G00B12. <https://doi.org/10.1029/2007JG000640>
- Davidson, E. A., de Araújo, A. C., Artaxo, P., Balch, J. K., Brown, I. F., C. Bustamante, M. M., Coe, M. T., DeFries, R. S., Keller, M., Longo, M., Munger, J. W., Schroeder, W., Soares-Filho, B. S., Souza, C. M., & Wofsy, S. C. (2012). The Amazon basin in transition. *Nature*, *481*(7381), 321–328. <https://doi.org/10.1038/nature10717>
- de Goncalves, L. G. de, Baker, I., Christoffersen, B., Costa, M., Restrepo-Coupe, N., Rocha, H. da, Saleska, S., & Nobre Muza, M. (2009). *The Large Scale Biosphere-Atmosphere Experiment in Amazônia, Model Intercomparison Project (LBA-MIP) protocol*.
- Eva, H. D., & Huber (eds), O. (2005). *A Proposal for Defining the Geographical Boundaries of Amazonia: Synthesis of the results from an Expert Consultation Workshop organized by the European Commission in collaboration with the Amazon Cooperation Treaty Organization—JRC Ispra, 7-8 June 2005*. Office for Official Publications of the European Communities, Luxemburg.
- Federer, C. A. (1982). Transpirational supply and demand: Plant, soil, and atmospheric effects evaluated by simulation. *Water Resources Research*, *18*(2), 355–362. <https://doi.org/10.1029/WR018i002p00355>
- Féret, J.-B., Gitelson, A. A., Noble, S. D., & Jacquemoud, S. (2017). PROSPECT-D: Towards modeling leaf optical properties through a complete lifecycle. *Remote Sensing of Environment*, *193*, 204–215. <https://doi.org/10.1016/j.rse.2017.03.004>
- Fisher, J. B., Huntzinger, D. N., Schwalm, C. R., & Sitch, S. (2014). Modeling the Terrestrial Biosphere. *Annual Review of Environment and Resources*, *39*(1), 91–123. <https://doi.org/10.1146/annurev-environ-012913-093456>
- Fisher, R. A., Koven, C. D., Anderegg, W. R. L., Christoffersen, B. O., Dietze, M. C., Farnior, C.

- E., Holm, J. A., Hurtt, G. C., Knox, R. G., Lawrence, P. J., Lichstein, J. W., Longo, M., Matheny, A. M., Medvigy, D., Muller-Landau, H. C., Powell, T. L., Serbin, S. P., Sato, H., Shuman, J. K., ... Moorcroft, P. R. (2018). Vegetation demographics in Earth System Models: A review of progress and priorities. *Global Change Biology*, *24*(1), 35–54. <https://doi.org/10.1111/gcb.13910>
- Fisher, R. A., Williams, M., Costa, D., Lola, A., Malhi, Y., Costa, D., F, R., Almeida, S., & Meir, P. (2007). The response of an Eastern Amazonian rain forest to drought stress: Results and modelling analyses from a throughfall exclusion experiment. *Global Change Biology*, *13*(11), 2361–2378. <https://doi.org/10.1111/j.1365-2486.2007.01417.x>
- Foken, T. (2008). The energy balance closure problem: An overview. *Ecological Applications: A Publication of the Ecological Society of America*, *18*(6), 1351–1367.
- Foken, T., Aubinet, M., & Leuning, R. (2012). The Eddy Covariance Method. In M. Aubinet, T. Vesala, & D. Papale (Eds.), *Eddy Covariance* (pp. 1–19). Springer Netherlands. [https://doi.org/10.1007/978-94-007-2351-1\\_1](https://doi.org/10.1007/978-94-007-2351-1_1)
- Foley, J. A., Prentice, I. C., Ramankutty, N., Levis, S., Pollard, D., Sitch, S., & Haxeltine, A. (1996). An integrated biosphere model of land surface processes, terrestrial carbon balance, and vegetation dynamics. *Global Biogeochemical Cycles*, *10*(4), 603–628. <https://doi.org/10.1029/96GB02692>
- Freire, G. A. P., Ventura, D. J., Fotopoulos, I. G., Rosa, D. M., Aguiar, R. G., & Araújo, A. C. de. (2020). DINÂMICA DE SERAPILHEIRA EM UMA ÁREA DE FLORESTA DE TERRA FIRME, AMAZÔNIA OCIDENTAL. *Nativa*, *8*(3), 323–328. <https://doi.org/10.31413/nativa.v8i3.9155>
- Fu, R., Yin, L., Li, W., Arias, P. A., Dickinson, R. E., Huang, L., Chakraborty, S., Fernandes, K., Liebmann, B., Fisher, R., & Myneni, R. B. (2013). Increased dry-season length over southern Amazonia in recent decades and its implication for future climate projection. *Proceedings of the National Academy of Sciences*, *110*(45), 18110–18115. <https://doi.org/10.1073/pnas.1302584110>
- Getirana, A. C. V., Dutra, E., Guimberteau, M., Kam, J., Li, H.-Y., Decharme, B., Zhang, Z., Ducharme, A., Boone, A., Balsamo, G., Rodell, M., Toure, A. M., Xue, Y., Peters-Lidard, C. D., Kumar, S. V., Arsenault, K., Drapeau, G., Ruby Leung, L., Ronchail, J., & Sheffield, J. (2014). Water Balance in the Amazon Basin from a Land Surface Model

- Ensemble. *Journal of Hydrometeorology*, 15(6), 2586–2614.  
<https://doi.org/10.1175/JHM-D-14-0068.1>
- Gotangco Castillo, C. K., Levis, S., & Thornton, P. (2012). Evaluation of the New CNDV Option of the Community Land Model: Effects of Dynamic Vegetation and Interactive Nitrogen on CLM4 Means and Variability\*. *Journal of Climate*, 25(11), 3702–3714.  
<https://doi.org/10.1175/JCLI-D-11-00372.1>
- Goudriaan, J. (1986). A simple and fast numerical method for the computation of daily totals of crop photosynthesis. *Agricultural and Forest Meteorology*, 38(1–3), 249–254.  
[https://doi.org/doi:DOI: 10.1016/0168-1923\(86\)90063-8](https://doi.org/doi:DOI:10.1016/0168-1923(86)90063-8)
- Hagos, S., & Leung, L. R. (2011). On the Relationship between Uncertainties in Tropical Divergence and the Hydrological Cycle in Global Models. *Journal of Climate*, 25(1), 381–391. <https://doi.org/10.1175/JCLI-D-11-00058.1>
- Harper, A. B., Denning, A. S., Baker, I. T., Branson, M. D., Prihodko, L., & Randall, D. A. (2010). Role of deep soil moisture in modulating climate in the Amazon rainforest. *Geophysical Research Letters*, 37(5). <https://doi.org/10.1029/2009GL042302>
- Haughton, N., Abramowitz, G., Pitman, A. J., Or, D., Best, M. J., Johnson, H. R., Balsamo, G., Boone, A., Cuntz, M., Decharme, B., Dirmeyer, P. A., Dong, J., Ek, M., Guo, Z., Haverd, V., van den Hurk, B. J. J., Nearing, G. S., Pak, B., Santanello, J. A., ... Vuichard, N. (2016). The Plumbing of Land Surface Models: Is Poor Performance a Result of Methodology or Data Quality? *Journal of Hydrometeorology*, 17(6), 1705–1723.  
<https://doi.org/10.1175/JHM-D-15-0171.1>
- Held, I. M., & Soden, B. J. (2006). Robust Responses of the Hydrological Cycle to Global Warming. *Journal of Climate*, 19(21), 5686–5699. <https://doi.org/10.1175/JCLI3990.1>
- Hollinger, D. Y., Ollinger, S. V., Richardson, A. D., Meyers, T. P., Dail, D. B., Martin, M. E., Scott, N. A., Arkebauer, T. J., Baldocchi, D. D., Clark, K. L., Curtis, P. S., Davis, K. J., Desai, A. R., Dragoni, D., Goulden, M. L., Gu, L., Katul, G. G., Pallardy, S. G., U, K. T. P., ... Verma, S. B. (2010). Albedo estimates for land surface models and support for a new paradigm based on foliage nitrogen concentration. *Global Change Biology*, 16(2), 696–710. <https://doi.org/10.1111/j.1365-2486.2009.02028.x>
- Jasechko, S., Sharp, Z. D., Gibson, J. J., Birks, S. J., Yi, Y., & Fawcett, P. J. (2013). Terrestrial water fluxes dominated by transpiration. *Nature*, 496(7445), 347–350.

- <https://doi.org/10.1038/nature11983>
- Jin, M., & Liang, S. (2006). An Improved Land Surface Emissivity Parameter for Land Surface Models Using Global Remote Sensing Observations. *Journal of Climate*, *19*(12), 2867–2881. <https://doi.org/10.1175/JCLI3720.1>
- Kato, S., Loeb, N. G., Rose, F. G., Doelling, D. R., Rutan, D. A., Caldwell, T. E., Yu, L., & Weller, R. A. (2012). Surface Irradiances Consistent with CERES-Derived Top-of-Atmosphere Shortwave and Longwave Irradiances. *Journal of Climate*, *26*(9), 2719–2740. <https://doi.org/10.1175/JCLI-D-12-00436.1>
- Keller, M., Alencar, A., Asner, G. P., Bobby Braswell, Bustamante, M., Davidson, E., Feldpausch, T., Fernandes, E., Goulden, M., Kabat, P., Kruijt, B., Luizão, F., Miller, S., Markewitz, D., Nobre, A. D., Nobre, C. A., Filho, N. P., Rocha, H. da, Dias, P. S., ... Vourlitis, G. L. (2004). Ecological Research in the Large-Scale Biosphere-Atmosphere Experiment in Amazonia: Early Results. *Ecological Applications*, *14*(4), S3–S16.
- Kucharik, C. J., Foley, J. A., Delire, C., Fisher, V. A., Coe, M. T., Lenters, J. D., Young-Molling, C., Ramankutty, N., Norman, J. M., & Gower, S. T. (2000). Testing the performance of a dynamic global ecosystem model: Water balance, carbon balance, and vegetation structure. *Global Biogeochemical Cycles*, *14*(3), 795–825. <https://doi.org/10.1029/1999GB001138>
- Lawrence, D. M., Thornton, P. E., Oleson, K. W., & Bonan, G. B. (2007). The Partitioning of Evapotranspiration into Transpiration, Soil Evaporation, and Canopy Evaporation in a GCM: Impacts on Land–Atmosphere Interaction. *Journal of Hydrometeorology*, *8*(4), 862–880. <https://doi.org/10.1175/JHM596.1>
- Lee, J.-E., Oliveira, R. S., Dawson, T. E., & Fung, I. (2005). Root functioning modifies seasonal climate. *Proceedings of the National Academy of Sciences*, *102*(49), 17576–17581. <https://doi.org/10.1073/pnas.0508785102>
- Li, Y., Zhao, M., Motesharrei, S., Mu, Q., Kalnay, E., & Li, S. (2015). Local cooling and warming effects of forests based on satellite observations. *Nature Communications*, *6*(1), 6603. <https://doi.org/10.1038/ncomms7603>
- Lin, Y.-S., Medlyn, B. E., Duursma, R. A., Prentice, I. C., Wang, H., Baig, S., Eamus, D., Dias, V. R. de, Mitchell, P., Ellsworth, D. S., Beeck, M. O. de, Wallin, G., Uddling, J., Tarvainen, L., Linderson, M.-L., Cernusak, L. A., Nippert, J. B., Ocheltree, T. W.,



- Tissue, D. T., ... Wingate, L. (2015). Optimal stomatal behaviour around the world. *Nature Climate Change*, 5(5), 459–464. <https://doi.org/10.1038/nclimate2550>
- Liou, K. N. (2002). *An Introduction to Atmospheric Radiation* (2nd ed., Vol. 84). Academic Press. <https://www.elsevier.com/books/an-introduction-to-atmospheric-radiation/liou/978-0-12-451451-5>
- Liu, Z., Ballantyne, A. P., & Cooper, L. A. (2019). Biophysical feedback of global forest fires on surface temperature. *Nature Communications*, 10(1), 214. <https://doi.org/10.1038/s41467-018-08237-z>
- Longo, M., Knox, R. G., Levine, N. M., Alves, L. F., Bonal, D., Camargo, P. B., Fitzjarrald, D. R., Hayek, M. N., Restrepo-Coupe, N., Saleska, S. R., Silva, R. da, Stark, S. C., Tapajós, R. P., Wiedemann, K. T., Zhang, K., Wofsy, S. C., & Moorcroft, P. R. (2018). Ecosystem heterogeneity and diversity mitigate Amazon forest resilience to frequent extreme droughts. *New Phytologist*, 219(3), 914–931. <https://doi.org/10.1111/nph.15185>
- Longo, M., Knox, R. G., Levine, N. M., Swann, A. L. S., Medvigy, D. M., Dietze, M. C., Kim, Y., Zhang, K., Bonal, D., Burban, B., Camargo, P. B., Hayek, M. N., Saleska, S. R., Silva, R. da, Bras, R. L., Wofsy, S. C., & Moorcroft, P. R. (2019). The biophysics, ecology, and biogeochemistry of functionally diverse, vertically and horizontally heterogeneous ecosystems: The Ecosystem Demography model, version 2.2 – Part 2: Model evaluation for tropical South America. *Geoscientific Model Development*, 12(10), 4347–4374. <https://doi.org/10.5194/gmd-12-4347-2019>
- Longo, M., Knox, R. G., Medvigy, D. M., Levine, N. M., Dietze, M. C., Kim, Y., Swann, A. L. S., Zhang, K., Rollinson, C. R., Bras, R. L., Wofsy, S. C., & Moorcroft, P. R. (2019). The biophysics, ecology, and biogeochemistry of functionally diverse, vertically and horizontally heterogeneous ecosystems: The Ecosystem Demography model, version 2.2 – Part 1: Model description. *Geoscientific Model Development*, 12(10), 4309–4346. <https://doi.org/10.5194/gmd-12-4309-2019>
- Lopes, A. P., Nelson, B. W., Wu, J., Graça, P. M. L. de A., Tavares, J. V., Prohaska, N., Martins, G. A., & Saleska, S. R. (2016). Leaf flush drives dry season green-up of the Central Amazon. *Remote Sensing of Environment*, 182, 90–98. <https://doi.org/10.1016/j.rse.2016.05.009>
- Manoli, G., Ivanov, V. Y., & Fatichi, S. (2018). Dry-Season Greening and Water Stress in

- Amazonia: The Role of Modeling Leaf Phenology. *Journal of Geophysical Research: Biogeosciences*, 123(6), 1909–1926. <https://doi.org/10.1029/2017JG004282>
- Mauder, M., Genzel, S., Fu, J., Kiese, R., Soltani, M., Steinbrecher, R., Zeeman, M., Banerjee, T., Roo, F. D., & Kunstmann, H. (2018). Evaluation of energy balance closure adjustment methods by independent evapotranspiration estimates from lysimeters and hydrological simulations. *Hydrological Processes*, 32(1), 39–50. <https://doi.org/10.1002/hyp.11397>
- Medlyn, B. E., Duursma, R. A., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V. M., Crous, K. Y., Angelis, P. de, Freeman, M., & Wingate, L. (2012). Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*, 18(11), 3476–3476. <https://doi.org/10.1111/j.1365-2486.2012.02790.x>
- Medlyn, B. E., Duursma, R. A., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V. M., Crous, K. Y., Angelis, P. D., Freeman, M., & Wingate, L. (2011). Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*, 17(6), 2134–2144. <https://doi.org/10.1111/j.1365-2486.2010.02375.x>
- Medvigy, D., Wofsy, S. C., Munger, J. W., Hollinger, D. Y., & Moorcroft, P. R. (2009). Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2. *Journal of Geophysical Research: Biogeosciences*, 114(G1), G01002. <https://doi.org/10.1029/2008JG000812>
- Metcalf, D. B., Meir, P., Aragão, L. E. O. C., Malhi, Y., Costa, A. C. L. da, Braga, A., Gonçalves, P. H. L., Athaydes, J. de, Almeida, S. S. de, & Williams, M. (2007). Factors controlling spatio-temporal variation in carbon dioxide efflux from surface litter, roots, and soil organic matter at four rain forest sites in the eastern Amazon. *Journal of Geophysical Research*, 112(G4), 1–9. <https://doi.org/10.1029/2007JG000443>
- Moderow, U., Aubinet, M., Feigenwinter, C., Kolle, O., Lindroth, A., Mölder, M., Montagnani, L., Rebmann, C., & Bernhofer, C. (2009). Available energy and energy balance closure at four coniferous forest sites across Europe. *Theoretical and Applied Climatology*, 98(3), 397–412. <https://doi.org/10.1007/s00704-009-0175-0>
- Morales, P., Sykes, M. T., Prentice, I. C., Smith, P., Smith, B., Bugmann, H., Zierl, B., Friedlingstein, P., Viovy, N., Sabaté, S., Sánchez, A., Pla, E., Gracia, C. A., Sitch, S., Arneth, A., & Ogee, J. (2005). Comparing and evaluating process-based ecosystem

- model predictions of carbon and water fluxes in major European forest biomes. *Global Change Biology*, 11(12), 2211–2233. <https://doi.org/10.1111/j.1365-2486.2005.01036.x>
- NASA. (2019). *Clouds and the Earth's Radiant Energy System Information and Data (CERES)*. <http://ceres.larc.nasa.gov/>
- Oleson, K. W., Niu, G.-Y., Yang, Z.-L., Lawrence, D. M., Thornton, P. E., Lawrence, P. J., Stöckli, R., Dickinson, R. E., Bonan, G. B., Levis, S., Dai, A., & Qian, T. (2008). Improvements to the Community Land Model and their impact on the hydrological cycle. *Journal of Geophysical Research*, 113(G1), 1–26. <https://doi.org/10.1029/2007JG000563>
- Oliveira, R. S., Dawson, T. E., Burgess, S. S. O., & Nepstad, D. C. (2005). Hydraulic redistribution in three Amazonian trees. *Oecologia*, 145(3), 354–363. <https://doi.org/10.1007/s00442-005-0108-2>
- Peters, T. (2016). Water Balance in Tropical Regions. In L. Pancel & M. Köhl (Eds.), *Tropical Forestry Handbook* (pp. 391–403). Springer. [https://doi.org/10.1007/978-3-642-54601-3\\_40](https://doi.org/10.1007/978-3-642-54601-3_40)
- Pitman, A. J. (2003). The evolution of, and revolution in, land surface schemes designed for climate models. *International Journal of Climatology*, 23(5), 479–510. <https://doi.org/10.1002/joc.893>
- Powell, T. L., Galbraith, D. R., Christoffersen, B. O., Harper, A., Imbuzeiro, H. M. A., Rowland, L., Almeida, S., Brando, P. M., da Costa, A. C. L., Costa, M. H., Levine, N. M., Malhi, Y., Saleska, S. R., Sotta, E., Williams, M., Meir, P., & Moorcroft, P. R. (2013). Confronting model predictions of carbon fluxes with measurements of Amazon forests subjected to experimental drought. *New Phytologist*, 200(2), 350–365. <https://doi.org/10.1111/nph.12390>
- Reed, D. E., Frank, J. M., Ewers, B. E., & Desai, A. R. (2018). Time dependency of eddy covariance site energy balance. *Agricultural and Forest Meteorology*, 249, 467–478. <https://doi.org/10.1016/j.agrformet.2017.08.008>
- Restrepo-Coupe, N., da Rocha, H. R., da Araujo, A. C., Borma, L. S., Christoffersen, B., Cabral, O. M. R., de Camargo, P. B., Cardoso, F. L., da Costa, A. C. L., Fitzjarrald, D. R., Goulden, M. L., Kruijt, B., Maia, J. M. F., Malhi, Y. S., Manzi, A. O., Miller, S. D., Nobre, A. D., von Randow, C., Sá, L. D. A., ... Saleska, S. R. (2013). What drives the seasonality of photosynthesis across the Amazon basin? A cross-site analysis of eddy flux

- tower measurements from the Brasil flux network. *Agricultural and Forest Meteorology*, 182–183, 128–144.
- Restrepo-Coupe, N., Levine, N. M., Christoffersen, B. O., Albert, L. P., Wu, J., Costa, M. H., Galbraith, D., Imbuzeiro, H., Martins, G., da Araujo, A. C., Malhi, Y. S., Zeng, X., Moorcroft, P., & Saleska, S. R. (2017). Do dynamic global vegetation models capture the seasonality of carbon fluxes in the Amazon basin? A data-model intercomparison. *Global Change Biology*, 23(1), 191–208. <https://doi.org/10.1111/gcb.13442>
- Rice, A. H., Pyle, E. H., Saleska, S. R., Hutyyra, L. R., Palace, M., Keller, M., Camargo, P. B. de, Portilho, K., Marques, D. F., & Wofsy, S. C. (2004). Carbon balance and vegetation dynamics in an old-growth Amazonian forest. *Ecological Applications*, 14(4), S55–S71.
- Sellers, P. J. (1985). Canopy reflectance, photosynthesis and transpiration. *International Journal of Remote Sensing*, 6(8), 1335–1372. <https://doi.org/10.1080/01431168508948283>
- Sena, E. T., Dias, M. A. F. S., Carvalho, L. M. V., & Dias, P. L. S. (2018). Reduced Wet-Season Length Detected by Satellite Retrievals of Cloudiness over Brazilian Amazonia: A New Methodology. *Journal of Climate*, 31(24), 9941–9964. <https://doi.org/10.1175/JCLI-D-17-0702.1>
- Shuttleworth, W. J. (1988). Evaporation from Amazonian Rainforest. *Proceedings of the Royal Society of London. Series B, Biological Sciences*, 233(1272), 321–346.
- Silva Dias, M. a. F., Rutledge, S., Kabat, P., Silva Dias, P. L., Nobre, C., Fisch, G., Dolman, A. J., Zipser, E., Garstang, M., Manzi, A. O., Fuentes, J. D., Rocha, H. R., Marengo, J., Plana-Fattori, A., Sá, L. D. A., Alvalá, R. . c. S., Andreae, M. O., Artaxo, P., Gielow, R., & Gatti, L. (2002). Cloud and rain processes in a biosphere-atmosphere interaction context in the Amazon Region. *Journal of Geophysical Research: Atmospheres*, 107(D20), 8072. <https://doi.org/10.1029/2001JD000335>
- Sperry, J. S., Venturas, M. D., Anderegg, W. R. L., Mencuccini, M., Mackay, D. S., Wang, Y., & Love, D. M. (2017). Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. *Plant, Cell & Environment*, 40(6), 816–830. <https://doi.org/10.1111/pce.12852>
- Spracklen, D. V., Baker, J. C. A., Garcia-Carreras, L., & Marsham, J. H. (2018). The Effects of Tropical Vegetation on Rainfall. *Annual Review of Environment and Resources*, 43(1), 193–218. <https://doi.org/10.1146/annurev-environ-102017-030136>

- Stahl, C., Hérault, B., Rossi, V., Burban, B., Bréchet, C., & Bonal, D. (2013). Depth of soil water uptake by tropical rainforest trees during dry periods: Does tree dimension matter? *Oecologia*, *173*(4), 1191–1201. <https://doi.org/10.1007/s00442-013-2724-6>
- Stockli, R., Rutishauser, T., Dragoni, D., O’Keefe, J., Thornton, P., Jolly, M., Lu, L., & Denning, A. (2008). Remote sensing data assimilation for a prognostic phenology model. *Journal of Geophysical Research-Biogeosciences*, *113*(G4). <https://doi.org/10.1029/2008JG000781>
- Tan, Z.-H., Zhao, J.-F., Wang, G.-Z., Chen, M.-P., Yang, L.-Y., He, C.-S., Restrepo-Coupe, N., Peng, S.-S., Liu, X.-Y., da Rocha, H. R., Kosugi, Y., Hirano, T., Saleska, S. R., Goulden, M. L., Zeng, J., Ding, F.-J., Gao, F., & Song, L. (2019). Surface conductance for evapotranspiration of tropical forests: Calculations, variations, and controls. *Agricultural and Forest Meteorology*, *275*, 317–328. <https://doi.org/10.1016/j.agrformet.2019.06.006>
- Taylor, K. E. (2001). Summarizing multiple aspects of model performance in a single diagram. *Journal of Geophysical Research*, *106*(D7), PP. 7183-7192. <https://doi.org/200110.1029/2000JD900719>
- Trenberth, K. E., Fasullo, J. T., & Kiehl, J. (2009). Earth’s Global Energy Budget. *Bulletin of the American Meteorological Society*, *90*(3), 311–324. <https://doi.org/10.1175/2008BAMS2634.1>
- Wan, Z., Hook, S., & Hulley, G. (2015). MOD11A2 MODIS/Terra Land Surface Temperature/Emissivity 8-Day L3 Global 1km SIN Grid V006. *Distributed by NASA EOSDIS Land Processes DAAC*. <https://doi.org/10.5067/MODIS/MOD11A2.006>
- Wehr, R., & Saleska, S. R. (2015). An improved isotopic method for partitioning net ecosystem–atmosphere CO<sub>2</sub> exchange. *Agricultural and Forest Meteorology*, *214–215*, 515–531. <https://doi.org/10.1016/j.agrformet.2015.09.009>
- Wehr, Richard, & Saleska, S. R. (2020). Calculating Canopy Stomatal Conductance from Eddy Covariance Measurements, in Light of the Energy Budget Closure Problem. *Biogeosciences Discussions*, 1–16. <https://doi.org/10.5194/bg-2020-154>
- Wehr, Richard, & Saleska, S. R. (2021). Calculating canopy stomatal conductance from eddy covariance measurements, in light of the energy budget closure problem. *Biogeosciences*, *18*(1), 13–24. <https://doi.org/10.5194/bg-18-13-2021>
- Wei, Z., Yoshimura, K., Wang, L., Miralles, D. G., Jasechko, S., & Lee, X. (2017). Revisiting

- the contribution of transpiration to global terrestrial evapotranspiration. *Geophysical Research Letters*, 44(6), 2792–2801. <https://doi.org/10.1002/2016GL072235>
- Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., Bernhofer, C., Ceulemans, R., Dolman, H., Field, C., Grelle, A., Ibrom, A., Law, B. E., Kowalski, A., Meyers, T., Moncrieff, J., Monson, R., Oechel, W., Tenhunen, J., ... Verma, S. (2002). Energy balance closure at FLUXNET sites. *Agricultural and Forest Meteorology*, 113(1–4), 223–243. [https://doi.org/10.1016/S0168-1923\(02\)00109-0](https://doi.org/10.1016/S0168-1923(02)00109-0)
- Wofsy, S., Goulden, M., & Munger, J. W. (1993). Net Exchange of CO<sub>2</sub> in a Mid-Latitude Forest. *Science*, 260, 1314–1317.
- Worden, J., Noone, D., Bowman, K., Beer, R., Eldering, A., Fisher, B., Gunson, M., Goldman, A., Herman, R., Kulawik, S. S., Lampel, M., Osterman, G., Rinsland, C., Rodgers, C., Sander, S., Shephard, M., Webster, C. R., & Worden, H. (2007). Importance of rain evaporation and continental convection in the tropical water cycle. *Nature*, 445(7127), 528–532. <https://doi.org/10.1038/nature05508>
- Wu, J., Guan, K., Hayek, M., Restrepo-Coupe, N., Wiedemann, K. T., Xu, X., Wehr, R., Christoffersen, B. O., Miao, G., da Silva, R., de Araujo, A. C., Oliviera, R. C., Camargo, P. B., Monson, R. K., Huete, A. R., & Saleska, S. R. (2017). Partitioning controls on Amazon forest photosynthesis between environmental and biotic factors at hourly to interannual timescales. *Global Change Biology*, 23(3), 1240–1257. <https://doi.org/10.1111/gcb.13509>
- Yan, B., Mao, J., Dickinson, R. E., Thornton, P. E., Shi, X., Ricciuto, D. M., Warren, J. M., & Hoffman, F. M. (2020). Modelling tree stem-water dynamics over an Amazonian rainforest. *Ecohydrology*, 13(1), e2180. <https://doi.org/10.1002/eco.2180>
- Yanagi, S. N. M., & Costa, M. H. (2011). Simulations of tropical rainforest albedo: Is canopy wetness important? *Anais Da Academia Brasileira de Ciências*, 83(4), 1171–1180. <https://doi.org/10.1590/S0001-37652011005000047>

## List of Figures

Figure 1. Parameters and mechanisms that govern ecosystem water fluxes (evapotranspiration) and their seasonality included in this study. Here classified as: water availability, vegetation

response, radiation balance, and distribution of turbulent fluxes. Colored lines show which drivers interact with which parameters/mechanisms.

Figure 2. Locations of eddy covariance tower study sites in the Amazon Basin *sensu-stricto* (Eva & Huber (eds), 2005). Mean monthly net radiation ( $W\ m^{-2}$ ) from the Clouds and the Earth's Radiant Energy System v4.0 (CERES) (Kato et al., 2012) for the years 2003 to 2018 (NASA, 2019).

Figure 3. Annual cycle 16-day average observed climatic drivers (a) precipitation (*Precip*;  $mm\ month^{-1}$ ) (gray bars), air temperature ( $T_{air}$ ;  $degC$ ) (blue line left y-axis) and (b) incoming shortwave ( $SW_{down}$ ;  $W\ m^{-2}$ ) (black line right y-axis) and longwave radiation ( $LW_{down}$ ;  $W\ m^{-2}$ ) (blue line left y-axis). (c) Modeled and observed daytime evapotranspiration ( $ET_{day}$ ,  $mm\ month^{-1}$ ), dashed line corresponds to the reference evapotranspiration (model  $ET$  driven only by meteorology); and (d) model ecosystem-scale of model soil moisture 'stress' ( $FSW$ , where 1 = no stress). From left to right study sites (from wettest to driest) near Manaus (K34), Caxiuanã (CAX), Santarém (K67), and Reserva Jaru southern (RJA) forests. Gray-shaded area is dry season as defined using satellite-derived measures of precipitation (TRMM: 1998-2018). (e) Right hand plots correspond to Taylor diagrams for a statistical summary of model (color coded) fluxes compared to observations of seasonal fluxes (16-day). Missing sites indicate that the model overestimates the seasonality of observations; the ratio between model ( $\sigma_m$ ) to observation standard deviations ( $\sigma$ ) is  $>2$ . Simulations from ED2 (blue), IBIS (red), CLM3.5 (green), and JULES (purple).

Figure 4. Annual cycle 16-day average observations (black line) and modeled (color lines) of (a) latent heat flux ( $LE$ ;  $W\ m^{-2}$ ) -- energy equivalent of evapotranspiration ( $ET$ ;  $mm\ d^{-1}$ ), and model results from a linear regression between  $LE$  and  $Rn$  (dashed black line). (b) Sensible heat flux ( $H$ ;  $W\ m^{-2}$ ) and (c) unitless Bowen ratio ( $Bowen = H/LE$ ). From left to right study sites (from wettest to driest) near Manaus (K34), Caxiuanã (CAX), Santarém (K67), and Reserva Jaru southern (RJA) forests. Gray-shaded area is dry season as defined using satellite-derived measures of precipitation (TRMM: 1998-2018). Right hand plots correspond to Taylor diagrams for a statistical summary of model (color coded) fluxes compared to observations of seasonal

fluxes (16-day). Missing sites indicate that the model overestimates the seasonality of observations; the ratio between model ( $\sigma_m$ ) to observation standard deviations ( $\sigma$ ) is  $>2$ . Simulations from ED2 (blue), IBIS (red), CLM3.5 (green), and JULES (purple).

Figure 5. Annual cycle 16-day average observations (black line) and modeled (color lines) of (a) outgoing shortwave radiation ( $SW_{out}$ ;  $W m^{-2}$ ); (b) outgoing longwave radiation ( $LW_{out}$ ;  $W m^{-2}$ ) and (c) net radiation ( $R_n$ ;  $W m^{-2}$ ) (black continuous line) and the sum of turbulent fluxes, sensible plus latent heat flux ( $H+LE$ ;  $W m^{-2}$ ) (black dotted line). From left to right study sites (from wettest to driest) near Manaus (K34), Caxiuanã (CAX), Santarém (K67), and Reserva Jaru southern (RJA) forests. Gray-shaded area is dry season as defined using satellite-derived measures of precipitation (TRMM: 1998-2018). Right hand plots correspond to Taylor diagrams for a statistical summary of model (color coded) fluxes compared to observations of seasonal fluxes (16-day). Missing sites indicate that the model overestimates the seasonality of observations; the ratio between model ( $\sigma_m$ ) to observation standard deviations ( $\sigma$ ) is  $>2$ . Simulations from ED2 (blue), IBIS (red), CLM3.5 (green), and JULES (purple).

Figure 6. Annual cycle 16-day average observations (black line) and modeled (color lines) of (a) albedo -- the ratio of outgoing to incoming shortwave radiation ( $\alpha=SW_{out}/SW_{down}$ ), and (b) surface emissivity ( $\epsilon_s$ ). From left to right study sites (from wettest to driest) near Manaus (K34), Caxiuanã (CAX), Santarém (K67), and Reserva Jaru southern (RJA) forests. Gray-shaded area is dry season as defined using satellite-derived measures of precipitation (TRMM: 1998-2018). Right hand plots correspond to Taylor diagrams for a statistical summary of model (color coded) fluxes compared to observations of seasonal fluxes (16-day). Missing sites indicate that the model overestimates the seasonality of observations; the ratio between model ( $\sigma_m$ ) to observation standard deviations ( $\sigma$ ) is  $>2$ . Simulations from ED2 (blue), IBIS (red), CLM3.5 (green), and JULES (purple).


Figure 7. Annual cycle 16-day average observations (black line) and modeled (color lines) of (a) ratio between the observed and reference evapotranspiration ( $ET/ET_{ref}$ ), (b) canopy stomatal conductance ( $G_s$ ;  $mmol m^{-2} s^{-1}$ ) and (c) daytime water use efficiency where  $ET$  and  $GEP$  were sampled during dry conditions (no rain in prior 12 hours) assuming transpiration drives water




fluxes ( $WUE = GEP_{dry\&dry}/ET_{dry\&dry}$ ;  $gC\ mm^{-1}$ ). From left to right study sites (from wettest to driest) near Manaus (K34), Caxiuanã (CAX), Santarém (K67), and Reserva Jaru southern (RJA) forests. Gray-shaded area is dry season as defined using satellite-derived measures of precipitation (TRMM: 1998-2018). Right hand plots correspond to Taylor diagrams for a statistical summary of model (color coded) fluxes compared to observations of seasonal fluxes (16-day). Missing sites indicate that the model overestimates the seasonality of observations; the ratio between model ( $\sigma_m$ ) to observation standard deviations ( $\sigma_x$ ) is  $>2$ . Simulations from ED2 (blue), IBIS (red), CLM3.5 (green), and JULES (purple).


Figure 8. Relationships between seasonal 16-day average values of (a) canopy stomatal conductance ( $G_s$ ;  $mm\ s^{-1}$ ) and the ratio between sensible ( $H$ ;  $W\ m^{-2}$ ) and latent heat flux ( $LE$ ;  $W\ m^{-2}$ ) the unitless Bowen ratio ( $Bowen=H/LE$ ), (b)  $G_s$  and daytime net radiation ( $Rn_{day}$ ;  $W\ m^{-2}$ ); (c) daytime  $LE$  ( $LE_{day}$ ;  $W\ m^{-2}$ ) and  $Rn_{day}$ , and (d) daytime  $H$  ( $H_{day}$ ;  $W\ m^{-2}$ ) and  $Rn_{day}$ . Panels may include a linear regression for all available data (black line) and single regressions fitted for each site and seasons (color lines): Manaus (K34), Caxiuanã (CAX), Santarém (K67), and Reserva Jaru southern (RJA) forests if statistically significant. Seasons classified using satellite precipitation TRMM values (1996-2018), dry (rainfall  $<100\ mm\ month^{-1}$ ) and wet-period ( $> 100\ mm\ month^{-1}$ ).

Improving model *versus* observation  
water flux discrepancies




gcb\_15555\_f1.png






gcb\_15555\_f3.png




gcb\_15555\_f4.png




gcb\_15555\_f5.png

Author Manuscript



gcb\_15555\_f6.png


Author Manuscript



gcb\_15555\_f7.png

Author Manuscript





gcb\_15555\_f8.png