

# Evaluation of an Extended-duration Chemoprophylaxis Regimen for Venous Thromboembolism after Microsurgical Breast Reconstruction

Eric M. Pittelkow, MD\*

Will C. DeBrock, MD\*

Brian Mailey, MD†

Tarah J. Ballinger, MD‡

Juan Socas, MD\*

Mary E. Lester, MD\*

Aladdin H. Hassanein, MD, MMSc\*

**Background :** Patients undergoing free flap breast reconstruction are at a high risk for venous thromboembolism based upon Caprini scores. Guidelines for venous thromboembolism prophylaxis recommend high-risk groups receive extended chemoprophylaxis for several weeks after gynecological, orthopedic, and surgical oncology cases. Extended prophylaxis has not been studied in free flap breast reconstruction. The purpose of this study was to compare outcomes of free flap breast reconstruction patients who received extended venous thromboembolism (VTE) prophylaxis with those who received standard inpatient-only prophylaxis.

**Methods :** Patients undergoing microsurgical breast reconstruction were divided into two groups: standard VTE prophylaxis (Group I) and extended prophylaxis (Group II). Both groups received prophylactic subcutaneous heparin or enoxaparin preoperatively and enoxaparin 40mg daily postoperatively while inpatient. Group II was discharged with a home regimen of enoxaparin 40mg daily for an additional 14 days.

**Results :** In total, 103 patients met inclusion criteria (36 patients in Group I, 67 patients in Group II). The incidence of VTE was 1.5% in Group II compared with 2.8% in Group I ( $P = 0.6$ ). There was no difference in reoperative hematoma between Group I ( $n = 0$ ) and Group II ( $n = 1$ ) ( $P = 0.7$ ). Total flap loss was 2.2%.

**Conclusions :** Although this retrospective pilot study did not show statistical significance in VTE between those receiving extended home chemoprophylaxis (1.5% incidence) compared with inpatient-only chemoprophylaxis (2.8%), the risk of bleeding complications was similar. These results indicate that a larger, higher powered study is justified to assess if an extended home chemoprophylaxis protocol should be standard of care post free flap breast reconstruction. (*Plast Reconstr Surg Glob Open* 2021;9:e3741; doi: [10.1097/GOX.0000000000003741](https://doi.org/10.1097/GOX.0000000000003741); Published online 6 August 2021.)

## INTRODUCTION

Venous thromboembolism (VTE) causes mortality in 100,000 patients annually in the United States.<sup>1</sup> VTE is among the few life-threatening complications that can affect plastic surgery patients. Individualized risk stratification has been recommended for VTE prevention by the American College of Chest Physicians (ACCP) guidelines

and is widely accepted among surgical disciplines.<sup>2</sup> The Caprini risk assessment model is a validated tool for calculating VTE risk and guiding prophylaxis strategy.<sup>3-6</sup> The ACCP initially published VTE prevention guidelines in 2005 and updated recommendations in 2012 for orthopedic, nonorthopedic, and nonsurgical patients.<sup>2,7,8</sup> The 2012 ACCP guidelines addressed plastic surgery patients.<sup>2</sup> Recommendations were limited by the paucity of level I evidence in plastic surgery. Although the 2012 ACCP guidelines recommended that individuals with a high VTE risk (Caprini score  $\geq 5$ ) undergoing abdominal or pelvic surgery for cancer should receive extended-duration pharmacologic chemoprophylaxis, postmastectomy microsurgical breast free reconstruction was excluded from this recommendation.<sup>2</sup> Data from the Venous Thromboembolism Prevention Study, a consortium of five centers, were used to extrapolate VTE risk for plastic surgery patients based upon Caprini scores.<sup>4</sup> However, the Venous Thromboembolism Prevention Study did not report on

From the \*Division of Plastic Surgery, Indiana University School of Medicine, Indianapolis, Ia.; †Institute for Plastic Surgery, Southern Illinois University, Springfield, Ill.; and ‡Division of Hematology/Oncology, Indiana University School of Medicine, Indianapolis, Ia.

Received for publication April 3, 2021; accepted June 9, 2021.

Presented (as an oral presentation) at the American College of Surgeons Clinical Congress 2019, October 30, 2019, San Francisco, Calif.

Copyright © 2021 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 \(CCBY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: [10.1097/GOX.0000000000003741](https://doi.org/10.1097/GOX.0000000000003741)

**Disclosure:** The authors have no sources of funding to disclose. The authors have no financial interest in relation to the content of this article. The study did not receive any funding.

chemoprophylaxis beyond inpatient hospitalization. For Caprini scores higher than 8, the American Association of Plastic Surgeons recommends chemoprophylaxis.<sup>9</sup>

Current recommendations for VTE chemoprophylaxis in the plastic surgery literature following microsurgical breast reconstruction are not uniform. Prophylactic anticoagulation typically started 6–8 hours postprocedure and continued through the inpatient hospital duration.<sup>5,6,10</sup> The Caprini model stratifies the vast majority of patients undergoing microsurgical breast reconstruction as “high-risk.”<sup>5,11,12</sup> Patients who undergo microsurgical breast reconstruction have at least a 3.4% incidence of a deep vein thrombosis (DVT).<sup>13</sup> Factors that contribute to high VTE incidence for microsurgical breast reconstruction include cancer status, elevated body mass index (BMI), prolonged operative times, decreased mobility, extended length of inpatient stay, and increased intra-abdominal compartment pressure.<sup>13–15</sup> The purpose of this study was to assess extended home regimen VTE chemoprophylaxis to standard, inpatient-only chemoprophylaxis after microsurgical breast reconstruction.

## METHODS

### Study Population

Following approval from the institutional review board at Indiana University, patients undergoing microsurgical breast reconstruction from 2013 to 2016 at our institution were retrospectively evaluated. Individuals who had breast reconstruction with an abdominally-based free flap (deep inferior epigastric artery perforator (DIEP), superficial inferior epigastric artery, and muscle-sparing transverse rectus abdominis myocutaneous) were included. Patients with breast reconstruction from alternative donor sites (eg, profunda artery perforator, transverse upper gracili, and superior gluteal artery perforator) were excluded. All patients received mechanical compression devices placed preoperatively and during the inpatient stay.<sup>6,16</sup> Group I was categorized as “standard regimen” and received 5000 units of subcutaneous heparin or 40mg of subcutaneous enoxaparin preoperatively followed by 40mg of enoxaparin (first dose given 12 hours postoperatively) daily while inpatient until discharge.<sup>17</sup> Group II was designated “extended prophylaxis” and was administered preoperative chemoprophylaxis (5000 units of subcutaneous heparin or 40mg of subcutaneous enoxaparin) and daily enoxaparin 40mg (first dose given 12 hours postoperatively) while inpatient, followed by 2 weeks of enoxaparin 40mg daily after discharge. Patients in both groups received 325mg daily of aspirin postoperatively continuing until 30 days after discharge. During the study period, a postoperative protocol management change was adopted to use extended chemoprophylaxis for all patients undergoing microsurgical breast reconstruction. Group I was comprised of consecutive patients from the beginning of the study, and group II patients were subsequently after the practice change.

Outcome variables were VTE, reoperative hematoma, surgical site infection, seroma, and flap loss in

the postoperative period. Records were monitored for VTE occurrence for 4 months postoperatively or until the patient’s next surgery after free flap reconstruction. Predictive variables were age, sex, BMI, hospital length of stay, Caprini score, and comorbidities (diabetes, smoking). VTE risk was assessed using the 2005 Caprini model for VTE which stratified patients into “very low” (0 points), “low” (1–2 points), “moderate” (3–4 points), and “high” ( $\geq 5$  points) risk categories based upon ACCP guidelines.<sup>2,11</sup>

### Statistical Analysis

Data are presented as means and percentages. Chi-squared and Fischer exact test were used to analyze dichotomous dependent variables (VTE, reoperative hematoma, surgical site infection, seroma, flap loss) and comorbidities as appropriate given the low number of events. Independent samples *t*-tests were used to analyze continuous variables (hospital length of stay, BMI, age). Data were managed using the IU REDCAP data capture tool and analyzed with SPSS version 24.0 (IBM Corp., Armonk, N.Y.).

## RESULTS

There were 103 patients (180 flaps) who met inclusion criteria: 36 patients in Group I (standard inpatient prophylaxis) and 67 patients in Group II (extended chemoprophylaxis). One hundred patients underwent DIEP flap reconstruction, one patient underwent muscle-sparing transverse rectus abdominis myocutaneous flap reconstruction, and two patients underwent superficial inferior epigastric artery flap reconstruction. The average age was 49.4 ( $\pm 9.8$ ) years for Group I and 48.7 ( $\pm 9.2$ ) years for Group II ( $P = 0.7$ ). Mean BMI in Group I was 31.7 ( $\pm 6.0$ ) kg/m<sup>2</sup> compared with 30.7 ( $\pm 6.4$ ) kg/m<sup>2</sup> in Group II ( $P = 0.4$ ). There were eight smokers in Group I and 15 smokers in Group II ( $P = 0.99$ ) (Table 1). Group I had six diabetics, and Group II had eight diabetics ( $P = 0.6$ ). Seventy-seven patients (75%) underwent bilateral reconstruction, with the majority of those patients belonging to the extended regimen group ( $P = 0.02$ ): 22 bilateral (61%), 14 unilateral (39%) in Group I and 55 bilateral (82%), and 12 unilateral (18%) in Group II. Average length of stay was 5.6 ( $\pm 1.3$ ) days in Group I and 5.6 ( $\pm 2.4$ ) days in Group II ( $P = 0.8$ ). Mean Caprini score was 6.6 ( $\pm 7$ ) in Group I and 6.6 ( $\pm 9$ ) in Group II ( $P = 0.9$ ). One patient in each group developed VTE (2.8% in Group I, 1.5% in Group II) ( $P = 0.6$ ). One patient (Group II) experienced postoperative surgical hematoma, requiring takeback to the operating room ( $P = 0.7$ ). Four patients (11.1%) in Group I developed surgical site infections, compared with 15 (22.4%) patients in Group II ( $P = 0.2$ ). There were 14 seromas among the 103 patients (16.2% Group I, 11.9% Group II,  $P = 0.6$ ). Group I flap loss was 3.4% (two flaps) compared with Group II (1.6%, two flaps) for a total flap loss rate of 2.2%. The average operative time was 528  $\pm$  117 minutes.

The two patients who experienced VTE both had a prior history of VTE. The patient from Group I presented

**Table 1. Risk Factors and Outcomes for Patients Undergoing Abdominal Free Flap Breast Reconstruction in the Standard Chemoprophylaxis and Extended Chemoprophylaxis Groups**

Predictive Variables	Standard Chemoprophylaxis (Group I)	Extended Chemoprophylaxis (Group II)	P
Smoking, n (%)	8 (22.2%)	15 (22.4%)	0.99
Age, y (mean)	49.4 ± 9.8	48.7 ± 9.2	0.73
BMI, kg/m <sup>2</sup> (mean)	31.7 ± 6.0	30.7 ± 6.4	0.40
Caprini score (mean)	6.6 ± 0.7	6.6 ± 0.9	0.88
Length of stay, d (mean)	5.6 ± 1.3	5.6 ± 2.4	0.84
VTE, n (%)	1 (2.8%)	1 (1.5%)	0.58
Hematoma	0	1 (1.5%)	0.65
Flap loss, n (%)	2 (5.6%)	2 (2.3%)	0.61

with distal external iliac DVT on postoperative day 19 confirmed by venous duplex ultrasound. Computed tomography scan of her chest showed right upper and right lower lobe pulmonary emboli (PE). After evaluation by the hematology-oncology service, she was treated with bivalirudin, a direct thrombin inhibitor, received an inferior vena cava filter, and underwent a workup for heparin-induced thrombocytopenia, which was found to be positive. The patient was a 58-year-old woman with a BMI of 24.7. Her comorbidities included a recent history of smoking. She had undergone a bilateral mastectomy with immediate DIEP flap reconstruction with a total operating time of 704 minutes. Group II had one patient who experienced VTE. This patient presented with calf pain on postoperative day 22. A venous duplex ultrasound was ordered, which showed distal popliteal vein DVT. There was no PE. The patient was treated with 80 mg of enoxaparin twice daily for 2 weeks followed by 20 mg of rivaroxaban daily and eventually placed on 5 mg of apixaban twice daily. The apixaban was stopped 17 months after her presentation for DVT, and aspirin was started indefinitely. The patient was a 54-year-old woman with a BMI of 27.6. She did not have any significant medical problems. She had undergone a bilateral mastectomy with immediate DIEP flap breast reconstruction with a total operating time of 620 minutes.

## DISCUSSION

DVT and PE are complications that can cause severe morbidity and mortality. Other surgical disciplines have evolved to institute more aggressive, extended prophylactic measures to prevent VTE for high-risk patients based on Caprini assessment. There are limited data on extended chemoprophylaxis in high-risk plastic surgery patients. We present a pilot series on an extended home regimen of chemoprophylaxis (enoxaparin 40 mg subcutaneously daily) for 2 weeks post discharge for patients undergoing microsurgical breast reconstruction. These patients otherwise received a similar treatment to the standard group (preoperative chemoprophylaxis before induction, enoxaparin 40 mg daily while inpatient, and 325 mg daily oral aspirin for 30 days). We did not find a higher rate of complications such as hematoma. Patients who received enoxaparin for an extended duration in our study had a

1.5% risk of VTE compared with 2.8% for patients who received standard, inpatient-only enoxaparin. Although this difference did not reach statistical significance, our study's 1.5% risk is lower than the historical more than 3% rate that has been found in other studies for breast reconstruction patients only receiving prophylaxis while in-house.<sup>11,13</sup> A recent retrospective review reported a VTE rate among microsurgical breast reconstruction patients as 1.3%, similar to our findings. However, specific chemoprophylaxis regimens were not assessed.<sup>18</sup>

One concern cited for the use of chemoprophylaxis in microsurgical reconstruction is increased postoperative hematoma risk, given the large volume of dissection, and potential for flap compromise from pedicle tamponade by a hematoma.<sup>10,19</sup> However, several studies have demonstrated clinical benefit with VTE prophylaxis without an increase in hematoma incidence.<sup>3,10</sup> Similarly, our study did not show a significant difference in reoperative hematoma rate or other complications with the extended enoxaparin regimen.

Chemoprophylaxis has been studied in body contouring, which also carries a high VTE risk.<sup>20–23</sup> In a previous study, body contouring patients designated as “high-risk” were given 7 days total of chemoprophylaxis. There were no patients with DVT.<sup>22</sup> Other investigators found a DVT incidence under 1% after administering enoxaparin for 7 days to 253 patients following abdominoplasty.<sup>20</sup> Patients in our study, postdischarge, received an additional 2 weeks of daily enoxaparin 40 mg subcutaneously for prophylaxis. The risk of VTE remains elevated after discharge, for up to 12 weeks following surgery.<sup>24,25</sup> Extended-duration chemoprophylaxis (up to 35 days) is used in high-risk patients (eg, abdominal/pelvic cancer operations, knee replacements) in other surgical disciplines such as orthopedic surgery, gynecology, general surgery, and urology, with level 1 evidence support.<sup>26–30</sup> When extended VTE prophylaxis is administered in other surgical disciplines, it is most commonly given for 28 days.<sup>2,31,32</sup> A randomized double-blinded study showed decreased VTE risk for patients undergoing abdominal/pelvic cancer operations if daily enoxaparin 40 mg was administered for 27–31 days compared with 6–10 days.<sup>27</sup> Patients undergoing abdominal free flap breast reconstruction may possibly benefit from a longer duration of chemoprophylaxis. However, patient compliance with chemoprophylaxis may decrease as the duration increases. A study analyzing over 1200 patients with hip fractures found only one in five patients were compliant with 28 days of chemoprophylaxis.<sup>33</sup> In addition, patients in our investigation received enoxaparin 40 mg daily subcutaneously. This dosing may be inadequate for some patients based upon antifactor Xa levels.<sup>34</sup> Enoxaparin 30 mg twice daily may be more optimal dosing, but compliance is likely to decrease by doubling the frequency of administration of the subcutaneous injection. Therefore, the duration of our extended regimen (14 days) and daily dosing (enoxaparin 40 mg) may not be optimal for VTE prevention. A larger study duration is necessary to determine antifactor Xa levels. However, prolonging the treatment and increasing the frequency will likely hinder compliance.

Several limitations exist in our study. The patients were retrospectively reviewed. A clinical practice change of a more aggressive home prophylaxis regimen was initiated after a VTE event approximately a year and a half into the collection period rather than prospective randomization. Our limited sample size (total n = 103 patients) did not allow for multivariate regression modeling as VTE was rare. Therefore, confounders are not controlled. A power calculation ( $\beta = 0.2$ ,  $\alpha = 0.05$ ) demonstrates that nearly 2000 patients would be required in each group for 80% power to achieve statistical significance in VTE prevention based on our event rates. Additionally, a screening duplex to evaluate all patients during a specified time frame was not performed. Our results may indicate that the risk reduction in VTE with extended chemoprophylaxis may be clinically irrelevant or underpowered. However, given the morbidity and mortality of a DVT or PE, a larger study would be justified.

### CONCLUSIONS

Patients undergoing an abdominal free flap for breast cancer reconstruction have an increased risk for VTE often because of the length of the operation, elevated BMI, cancer history, and age. Our chemoprophylactic protocol is 5000 units of subcutaneous heparin or 40 mg of enoxaparin given preoperatively, enoxaparin 40 mg daily while inpatient, aspirin 325 mg daily for 30 days, and enoxaparin 40 mg daily postdischarge for an additional 14 days for patients with a Caprini score of 5 or higher. Patients having prophylactic mastectomies, under age 40, and with a BMI less than 25 kg/m<sup>2</sup> may not meet criteria. Imaging is obtained to rule out DVT and PE based on clinical presentation. This pilot study showed that extended-duration chemoprophylaxis can be applied to these patients without an increase in bleeding complications or flap loss. There was no statistical significance between VTE in the extended chemoprophylaxis group (1.5% incidence) and the standard inpatient-only chemoprophylaxis group (2.8%). However, given the high morbidity and mortality of VTE, we continue to use the extended chemoprophylaxis regimen at our institution. Our findings with the small sample size indicate a larger, higher powered study is justified to assess if an extended home chemoprophylaxis protocol should be standard of care post free flap breast reconstruction.

**Aladdin H. Hassanein, MD, MMSc**

Division of Plastic Surgery  
Indiana University School of Medicine  
545 Barnhill Drive Suite 232  
Indianapolis, IN 46202  
E-mail: ahassane@iu.edu

### REFERENCES

1. Beckman MG, Hooper WC, Critchley SE, et al. Venous thromboembolism: a public health concern. *Am J Prev Med.* 2010;38(4 suppl):S495–S501.
2. Gould MK, Garcia DA, Wren SM, et al. Prevention of VTE in nonorthopedic surgical patients: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2 suppl):e227S–e277S.
3. Pannucci CJ, Dreszer G, Wachtman CF, et al. Postoperative enoxaparin prevents symptomatic venous thromboembolism in high-risk plastic surgery patients. *Plast Reconstr Surg.* 2011;128:1093–1103.
4. Pannucci CJ, Bailey SH, Dreszer G, et al. Validation of the Caprini risk assessment model in plastic and reconstructive surgery patients. *J Am Coll Surg.* 2011;212:105–112.
5. Pannucci CJ. Evidence-based recipes for venous thromboembolism prophylaxis: a practical safety guide. *Plast Reconstr Surg.* 2017;139:520e–532e.
6. Murphy RX Jr, Alderman A, Gutowski K, et al. Evidence-based practices for thromboembolism prevention: summary of the ASPS venous thromboembolism task force report. *Plast Reconstr Surg.* 2012;130:168e–175e.
7. Falck-Ytter Y, Francis CW, Johanson NA, et al. Prevention of VTE in orthopedic surgery patients: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2 suppl):e278S–e325S.
8. Kahn SR, Lim W, Dunn AS, et al. Prevention of VTE in non-surgical patients: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2 suppl):e195S–e226S.
9. Pannucci CJ, MacDonald JK, Ariyan S, et al. Benefits and risks of prophylaxis for deep venous thrombosis and pulmonary embolus in plastic surgery: a systematic review and meta-analysis of controlled trials and consensus conference. *Plast Reconstr Surg.* 2016;137:709–730.
10. Pannucci CJ, Wachtman CF, Dreszer G, et al. The effect of postoperative enoxaparin on risk for reoperative hematoma. *Plast Reconstr Surg.* 2012;129:160–168.
11. Modarressi A, Schettini AV, Ruegg EM, et al. Venous thromboembolism events after breast reconstructions with DIEP free flaps in 192 consecutive cases. *Ann Chir Plast Esthet.* 2018;63:11–19.
12. Pannucci CJ, Chang EY, Wilkins EG. Venous thromboembolic disease in autogenous breast reconstruction. *Ann Plast Surg.* 2009;63:34–38.
13. Lemaine V, McCarthy C, Kaplan K, et al. Venous thromboembolism following microsurgical breast reconstruction: an objective analysis in 225 consecutive patients using low-molecular-weight heparin prophylaxis. *Plast Reconstr Surg.* 2011;127:1399–1406.
14. Losken A, Carlson GW, Tyrone JW, et al. The significance of intraabdominal compartment pressure after free versus pedicled TRAM flap breast reconstruction. *Plast Reconstr Surg.* 2005;115:261–263.
15. Losken A, Carlson GW, Jones GE, et al. Significance of intraabdominal compartment pressures following TRAM flap breast reconstruction and the correlation of results. *Plast Reconstr Surg.* 2002;109:2257–2264.
16. Young VL, Watson ME. The need for venous thromboembolism (VTE) prophylaxis in plastic surgery. *Aesthet Surg J.* 2006;26:157–175.
17. Keith JN, Chong TW, Davar D, et al. The timing of preoperative prophylactic low-molecular-weight heparin administration in breast reconstruction. *Plast Reconstr Surg.* 2013;132:279–284.
18. Rochlin DH, Shekter CC, Pannucci C, et al. Venous thromboembolism following microsurgical breast reconstruction: a longitudinal analysis of 12,778 patients. *Plast Reconstr Surg.* 2020;146:465–473.
19. Pannucci CJ, Oppenheimer AJ, Wilkins EG. Practice patterns in venous thromboembolism prophylaxis: a survey of 606 reconstructive breast surgeons. *Ann Plast Surg.* 2010;64:732–737.
20. Marangi GF, Segreto F, Poccia I, et al. Thromboprophylaxis in abdominoplasty: efficacy and safety of a complete perioperative protocol. *Arch Plast Surg.* 2016;43:360–364.

21. Clavijo-Alvarez JA, Pannucci CJ, Oppenheimer AJ, et al. Prevention of venous thromboembolism in body contouring surgery: a national survey of 596 ASPS surgeons. *Ann Plast Surg.* 2011;66:228–232.
22. Reish RG, Damjanovic B, Colwell AS. Deep venous thrombosis prophylaxis in body contouring: 105 consecutive patients. *Ann Plast Surg.* 2012;69:412–414.
23. Venturi ML, Davison SP, Caprini JA. Prevention of venous thromboembolism in the plastic surgery patient: current guidelines and recommendations. *Aesthet Surg J.* 2009;29:421–428.
24. Sweetland S, Green J, Liu B, et al; Million Women Study collaborators. Duration and magnitude of the postoperative risk of venous thromboembolism in middle aged women: prospective cohort study. *BMJ.* 2009;339:b4583.
25. Momeni A, Sorice SC, Li AY, et al. Breast reconstruction with free abdominal flaps is associated with persistent lower extremity venous stasis. *Plast Reconstr Surg.* 2019;143:1144e–1150e.
26. Rausa E, Kelly ME, Asti E, et al. Extended versus conventional thromboprophylaxis after major abdominal and pelvic surgery: systematic review and meta-analysis of randomized clinical trials. *Surgery.* 2018;164:1234–1240.
27. Bergqvist D, Agnelli G, Cohen AT, et al; ENOXACAN II Investigators. Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *N Engl J Med.* 2002;346:975–980.
28. Rasmussen MS, Jørgensen LN, Wille-Jørgensen P, et al; FAME Investigators. Prolonged prophylaxis with dalteparin to prevent late thromboembolic complications in patients undergoing major abdominal surgery: a multicenter randomized open-label study. *J Thromb Haemost.* 2006;4:2384–2390.
29. Kakkar VV, Balibrea JL, Martínez-González J, et al; CANBESURE Study Group. Extended prophylaxis with bemiparin for the prevention of venous thromboembolism after abdominal or pelvic surgery for cancer: the CANBESURE randomized study. *J Thromb Haemost.* 2010;8:1223–1229.
30. Vedovati MC, Becattini C, Rondelli F, et al. A randomized study on 1-week versus 4-week prophylaxis for venous thromboembolism after laparoscopic surgery for colorectal cancer. *Ann Surg.* 2014;259:665–669.
31. Bottaro FJ, Elizondo MC, Doti C, et al. Efficacy of extended thrombo-prophylaxis in major abdominal surgery: what does the evidence show? A meta-analysis. *Thromb Haemost.* 2008;99:1104–1111.
32. Akl EA, Terrenato I, Barba M, et al. Extended perioperative thromboprophylaxis in patients with cancer. A systematic review. *Thromb Haemost.* 2008;100:1176–1180.
33. Gao Y, Long A, Xie Z, et al. The compliance of thromboprophylaxis affects the risk of venous thromboembolism in patients undergoing hip fracture surgery. *Springerplus.* 2016;5:1362.
34. Pannucci CJ, Rockwell WB, Ghanem M, et al. Inadequate enoxaparin dosing predicts 90-day venous thromboembolism risk among plastic surgery inpatients: an examination of enoxaparin pharmacodynamics. *Plast Reconstr Surg.* 2017;139:1009–1020.