



Low diversity and host specificity in the gut microbiome community of *Eciton* army ants (Hymenoptera: Formicidae: Dorylinae) in a Costa Rican rainforest

Bradd MENDOZA-GUIDO*, Natalia RODRÍGUEZ-HERNÁNDEZ*, Aniek B.F. IVENS, Christoph VON BEEREN, Catalina MURILLO-CRUZ, Ibrahim ZUNIGA-CHAVES, Piotr ŁUKASIK, Ethel SANCHEZ, Daniel J.C. KRONAUER & Adrián A. PINTO-TOMÁS

Abstract

Neotropical army ants of the genus *Eciton* are top arthropod predators in tropical rainforests. Microbial symbionts, including Unclassified Firmicutes (UF) and Unclassified Entomoplasmatales (UE), are associated with this genus and likely play a significant role in the biology of these ants. While previous work focused on associations of army ants and gut microbes across large geographic scales, here we report a community survey of the gut microbes colonizing the six sympatric *Eciton* army ant species in a single Costa Rican location. Furthermore, we characterized the gut microbiota associated with different army ant castes in the swarm-raiding species *Eciton burchellii*. We employed a combination of 16S ribosomal RNA (rRNA) amplicon sequencing as well as fluorescence and electron microscopy to identify gut microbes and to verify their presence in ant guts. We also measured the diversity and interaction specificity of the ant-gut microbe interaction network. The two most dominant operational taxonomic unit (OTU) phylotypes in all species were related to UF and UE previously found in army ants, followed by OTUs assigned to the genus *Weissella*. Furthermore, the worker castes of *E. burchellii* shared similar gut microbiota, also dominated by UF and UE phylotypes. Overall, we found a low diversity of gut microbes and a low interaction specificity between army ants and microbes at the community level, mainly because most microbe strains were detected in various *Eciton* species. The fluorescence in-situ hybridization analyses documented the presence of the two dominant phylotypes within ant guts, and electron microscopy located bacterial biofilms in the hindgut near the microvilli. Their morphology suggests that these bacteria probably belong to the dominant phylotypes UF and UE. Taken together, our results confirm that the *Eciton* gut microbiome is consistently dominated by a few species of specialized bacteria that may improve nutrient uptake efficiency of host ants. Further research should employ multi-omics and culture-dependent strategies to fully understand the role of these potential symbionts in ant ecophysiology.

Key words: Symbiosis, microbial diversity, host specificity, Costa Rica, Firmicutes, Entomoplasmatales.

Received 2 May 2022; revision received 12 September 2022; accepted 7 October 2022

Subject Editor: Manuela de Oliveira Ramalho

Bradd Mendoza-Guido & Natalia Rodríguez-Hernández, Centro de Investigación en Estructuras Microscópicas (CIEMic), Universidad de Costa Rica, 11501-2060 San José Costa Rica; Centro de Investigación en Biología Celular y Molecular (CIBCM), Universidad de Costa Rica, 11501-2060 San José, Costa Rica.

E-mail: braddmg@gmail.com; natyrh16@gmail.com

Aniek B.F. Ivens, Laboratory of Social Evolution and Behavior, The Rockefeller University, 1230 York Avenue, New York, NY 10065, USA; Department of Ecological Sciences, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands. E-mail: a.b.f.iven@vu.nl

Christoph von Beeren, Laboratory of Social Evolution and Behavior, The Rockefeller University, 1230 York Avenue, New York, NY 10065, USA; Department of Biology, Ecological Networks, Technical University of Darmstadt, Schnittspanstraße 2, 64287 Darmstadt, Germany. E-mail: cvonbeeren@gmail.com

Catalina Murillo-Cruz, Ibrahim Zuniga-Chaves & Adrián A. Pinto-Tomás (contact author), Centro de Investigación en Estructuras Microscópicas (CIEMic), Universidad de Costa Rica, 11501-2060 San José, Costa Rica; Centro de Investigación en Biología Celular y Molecular (CIBCM), Universidad de Costa Rica, 11501-2060 San José, Costa Rica; Departamento de Bioquímica, Escuela de Medicina, Universidad de Costa Rica, 11501-2060 San José, Costa Rica. E-mail: catalina.murillocruz@ucr.ac.cr; ibrahimzuniga@gmail.com; adrian.pinto@ucr.ac.cr

Piotr Łukasik, Division of Biological Sciences, University of Montana, 32 Campus Dr, Missoula, MT 59812, USA; Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, ul. Gronostajowa 7, 30-387 Kraków, Poland. E-mail: p.lukasik@uj.edu.pl

Ethel Sanchez, Centro de Investigación en Estructuras Microscópicas (CIEMic), Universidad de Costa Rica, 11501-2060 San José, Costa Rica. E-mail: mesanche77@gmail.com

Daniel J.C. Kronauer, Laboratory of Social Evolution and Behavior, The Rockefeller University, 1230 York Avenue, New York, NY 10065, USA; Howard Hughes Medical Institute, New York, NY, USA. E-mail: dkronauer@mail.rockefeller.edu

*These authors contributed equally to this work.

Introduction

Insects are among the most diverse and abundant animal groups on Earth, occupying all types of habitats (BAHRNDORFF & al. 2016). The diversification and evolutionary success of insects depend in part on symbiotic interactions (POULSEN & SAPOUNTZIS 2012), particularly with microorganisms (WEISS & AKSOY 2011). Bacterial symbionts can fulfill different functions in insects, where they can contribute to host nutrition, digestion, reproduction, and defense (OHKUMA 2003, DILLON & DILLON 2004, ENGEL & MORAN 2013).

Bacteria associated with insects commonly inhabit the digestive tract, where they can perform key nutrition-related functions such as nutrient recycling or upgrading (ENGEL & MORAN 2013, MORAN & al. 2019), but also contribute to the defense against pathogens (KOCH & SCHMID-HEMPEL 2011) and other aspects of host health. The gut microbiome composition differs according to host phylogeny (ANDERSON & al. 2012, SANDERS & al. 2014) and environmental factors (AMATO 2013, KALTENPOTH & ENGL 2014). Large differences in microbiome composition and abundance have been described among different taxa and even among individuals within the same taxon, and it has been suggested that certain species may harbor virtually no gut microbes (HAMMER & al. 2017, SANDERS & al. 2017, HAMMER & al. 2019). In some insects, the hindgut harbors large bacterial populations due to favorable pH and redox conditions (DILLON & DILLON 2004, DOUGLAS 2015), while lower bacterial loads are found in the midgut, which tends to be a hostile environment for microorganisms because the epithelium secretes digestive enzymes and is immunologically active. For example, the *Drosophila melanogaster* midgut produces various antimicrobial peptides, digestive enzymes (including lysozymes), and a dual oxidase (DUOX: NADPH oxidase) that generates microbicidal reactive oxygen species (DOUGLAS 2015).

Several ant clades have important symbiotic relationships with their gut microbiome (BOURSAUX-EUDE & GROSS 2000, DOUGLAS 2015, MOREAU 2020). In herbivorous ants, bacteria can help their hosts by synthesizing essential amino acids, unfolding lignocellulosic components, supplying vitamins and cofactors, and detoxifying harmful plant subcomponents (DOUGLAS 2009, RUSSELL & al. 2009, SUEN & al. 2010). A well-known example is the symbiosis between carpenter ants and their endosymbiont, *Blochmannia floridanus*, which recycles nitrogenous

waste and upgrades its host's diet by providing essential amino acids during metamorphosis (ZIENZ & al. 2006, FELDHAAR & al. 2007, KUPPER & al. 2016). Likewise, in turtle ants of the genus *Cephalotes*, abundant, relatively diverse, and highly specific gut microbial communities recycle common nitrogenous waste products such as urea and uric acid to synthesize essential amino acids that are then provided to host workers in substantial quantities (HU & al. 2018, DUPLAIS & al. 2021). Similarly, herbivorous ants of the genus *Dolichoderus* host *Tokpelaia* bacteria that encode largely complete pathways for nitrogen recycling and biosynthesis of all essential amino acids and different vitamins (BISCH & al. 2018).

Recent studies have characterized the gut microbiome of a few clades of predatory ants (FUNARO & al. 2011, NEUVONEN & al. 2016, ŁUKASIK & al. 2017, BISCH & al. 2018, IVENS & al. 2018). The gut microbiome of a conspicuous predatory ant, the bullet ant *Paraponera clavata*, is dominated mainly by two bacterial operational taxonomic units (OTUs) in the genus *Tumebacillus* (Firmicutes), which is known to associate with a diversity of hosts (MOREAU & RUBIN 2017). In at least some instances, the microbial symbionts of predatory ants appear to play significant roles, such as in the case of *Tokpelaia* in the gut of the ant *Harpegnathos saltator*, which seems to import and degrade amino acids such as histidine and arginine, as well as urea, to produce glutamate from these nitrogen sources (NEUVONEN & al. 2016). This contrasts with the role of *Tokpelaia* in the herbivorous ant *Dolichoderus* spp., which harbors genes for the biosynthesis of histidine and arginine. These differences are likely the result of a metabolic adaptation to the diet of their hosts: Protein-rich diets in predatory ants select for microbes involved in amino acid degradation, while carbohydrate-rich diets in herbivorous ants promote symbionts specialized in amino acid biosynthesis (BISCH & al. 2018).

Army ants are considered the ultimate predators in the ant world (SCHNEIRLA 1971, KRONAUER 2020) and are characterized by mass raiding, nomadism, and colony fission, collectively known as the army ant adaptive syndrome (GOTWALD 1995). One of the best-studied species is the swarm-raider *Eciton burchellii*, which is a main predator of arthropods in Neotropical humid forests (FRANKS 1982, RETTENMEYER & al. 1983, GOTWALD 1995, HOENLE & al. 2019) and has an extraordinarily rich fauna of over

300 associated animal species, from mites to birds (RET TENMEYER & al. 2011).

There are several studies focusing on the gut microbiomes of army ants (FUNARO & al. 2011, ANDERSON & al. 2012, ŁUKASIK & al. 2017). In general, these studies have shown that these ants consistently harbor relatively abundant but low-diversity microbial communities. Specifically, microbiomes of all New World army ants seem to be dominated by a few taxa, including Unclassified Entomoplasmatales (UE), Unclassified Firmicutes (UF), Actinomycetales, and *Weissella*. Particularly, UE and UF are the dominant bacteria in the digestive tracts of *Eciton* workers (ŁUKASIK & al. 2017).

Interestingly, there is evidence that different species or even different colonies, as well as different individuals within the same colony, harbor different strains of these bacteria (FUNARO & al. 2011, ANDERSON & al. 2012, ŁUKASIK & al. 2017). Previous studies characterized army ants from different geographic locations, despite the known geographic variation in microbiome composition. Consequently, microbiome diversity and host-microbe specificity patterns across different army ant species living in the same population and across ant castes within the same colony have not yet been addressed systematically.

To bridge this knowledge gap, we systematically evaluated differences in gut microbiome composition between *Eciton* species, colonies, and castes at a single site: La Selva Biological Station (LSBS) in Costa Rica. The goals of the present study were threefold: (1) to characterize the composition and structure of the gut bacterial communities of sympatric *Eciton* species, (2) to test for caste differences in microbiome composition within a single species, *Eciton burchellii*, and (3) to localize bacteria in *E. burchellii* gut tissue to confirm the results from the broader sequencing-based surveys. We addressed these goals by characterizing the gut microbiomes of workers of different colonies from all locally occurring *Eciton* species at La Selva, that is, *E. burchellii*, *E. dulcium*, *E. hamatum*, *E. lucanoides*, *E. mexicanum*, and *E. vagans*, using 16S ribosomal RNA (rRNA) amplicon sequencing. In addition, we compared bacterial communities of different castes from two colonies of *E. burchellii*. Lastly, we employed scanning and transmission electron microscopy (SEM and TEM) and fluorescence in-situ hybridization (FISH) to visualize and localize bacteria within the gut tissue of *E. burchellii* ants. This study shows that *Eciton* army ants harbor a relatively simple and unspecific gut microbiome at the community level, adding evidence to the current general view that this is the case with most predatory ants.

Material and methods

Specimen collection

Army ants of the species *Eciton burchellii*, *Eciton dulcium*, *Eciton hamatum*, *Eciton lucanoides*, *Eciton mexicanum*, and *Eciton vagans*, including regular workers, submajors, majors, and males, were collected from February to April 2013 and March to April 2014 at LSBS

in Costa Rica (10° 25' 50.8188" N, 84° 0' 24.2388" W, altitude 67 m; Tab. S1, as digital supplementary material to this article, at the journal's web pages). Regular workers and submajors can easily be distinguished based on the coloration and the mandible shape, the latter of which are still triangular but more stretched in length in the submajors (RET TENMEYER 1961). Both worker types are easily distinguishable from majors, which possess the *Eciton*-typical hook-shaped mandibles (RET TENMEYER 1961, POWELL & FRANKS 2006). The workers were collected during nocturnal colony emigrations and colony raids (for more information see VON BEEREN & al. 2016, 2018). Ant species were identified employing the taxonomic keys of John T. Longino (LONGINO 2010). Regular workers from one additional *E. burchellii* colony were collected in September 2017 at LSBS. Regular workers were further divided into two size classes, which were previously denoted as minors and medias (FRANKS 1985).

The microbial diversity and distribution were analyzed using pools of 8 specimens for each worker type per *Eciton* colony (1 pooled sample per colony), from 12 colonies of *Eciton burchellii*, 9 of *Eciton dulcium*, 11 of *Eciton hamatum*, 2 of *Eciton lucanoides*, 9 of *Eciton mexicanum* and 6 of *Eciton vagans*. *Eciton* workers of all studied species are deposited at the Technical University Darmstadt Insect Collection and are either mounted on insect cardboards or preserved in absolute ethanol in 50 ml vials and stored in a -30 °C freezer. To compare microbiomes across *E. burchellii* worker size classes and males, three individuals were chosen from each of the following categories: minors, medias, submajors, majors, and males, from two of the colonies characterized in the previous step. Symbionts were localized within ant gut tissues using three individuals for each of the microscopy techniques (SEM, TEM, and FISH).

DNA extraction and sequencing

As a first approximation, the bacterial DNA for gut microbiome profiling was extracted from the whole gaster, the body part that includes most of the ant digestive system. Separating the intestinal tract into different parts would have given additional information about the microbes' preferred locations within the workers, but performing such an analysis was beyond the scope of this study. The aim was to screen for gut microbe differences between sympatric species and between *Eciton burchellii* worker castes and males. To prevent contamination with environmental bacteria, all extractions were performed under sterile conditions in a laminar flow hood. To remove bacteria from the ants' cuticles, all insects were surface-sterilized by immersing them individually for 30 s in 5% bleach and then in 1 × phosphate-buffered saline solution (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, and 1.8 mM KH₂PO₄) for 30 s. A pretreatment was applied to the samples that consisted of placing gasters, previously cut with sterile disposable razors, inside 1.5 ml tubes with stainless steel beads (5 mm) and 180 µl of the enzymatic lysis buffer (20 mM Tris-HCl, pH 8.0, 2 mM sodium EDTA, 1.2% Triton

X100, and 20 mg / ml lysozyme). Then, samples were homogenized in a QIAGEN (Hilden, Germany) TissueLyzer II for 3 min at 30 Hz, and the homogenate was processed, following the manufacturer's modified extraction protocol from the QIAGEN DNeasy Blood & Tissue kit. Each DNA extraction batch had its own extraction control (blank sample) that was used as a negative control throughout the procedure (including sequencing). As a single positive control, a gaster of a *Cephalotes* sp. worker was used.

The ant gut bacterial communities were characterized by sequencing amplicons of the V4 region of bacterial 16S rRNA (CAPORASO & al. 2011, 2012). The V4 region was amplified using universal bacterial primers (515F: 5' GTGCCAGCMGCCGCGGTAA 3', 806R: 5' GGACTACH-VGGGTWTCTAAT 3') (KOZICH & al. 2013) with adapter stubs. The PCR reactions consisted of 2.5 μ l of sample DNA, 0.5 μ l of 10 μ M primers, 12.5 μ l of 2 \times KAPA HiFi Hotstart Readymix (Kapa Biosystems Inc., Wilmington, MA, USA), and water to the total reaction volume of 25 μ l. PCRs ran at 95 $^{\circ}$ C for 3 min, followed by 26 cycles of 95 $^{\circ}$ C for 30 s, 55 $^{\circ}$ C for 30 s and 72 $^{\circ}$ C for 30 s, with a final extension step at 72 $^{\circ}$ C for 5 min. A second PCR reaction was performed to attach Illumina sequencing adapters and barcodes (Illumina Inc., San Diego, CA, USA). The reaction employed 5 μ l of the amplification product of the first PCR, 5 μ l of Nextera Index barcodes (forward and reverse) with 25 μ l of 2 \times KAPA HiFi Hotstart Readymix and water up to a total volume of 50 μ l. The indexing reaction conditions were the same as above, except for fewer cycles: 8 instead of 26. Indexed PCR products were then pooled with eight samples each and purified together using 0.6 \times Agencourt AMPure XP beads (Beckman Coulter, Brea, CA, USA). DNA was quantified in each purified product with QubitTM (ThermoFisher Scientific, Waltham, MA, USA) and equimolarly pooled to create a single sample. Libraries from DNA samples and controls were sequenced on an Illumina MiSeq platform at the Rockefeller University Genomics Resource Center to obtain 2 \times 250bp paired-end sequencing reads.

Sequence processing and bioinformatic analyses

The sequences of both datasets were processed with the software R (R CORE TEAM 2017) using the package DADA2 (version 1.14) through a modified pipeline (CALLAHAN & al. 2016a). First, the paired-end reads were joined into contigs. Then, sequences were clustered into Amplicon Sequence Variants (ASVs) and screened for sequencing errors: ASV sequences of less than 248 bp and those of more than 254 bp, as well as singletons and chimeras, were removed. The remaining sequences were aligned to the SILVA reference database (version 132) (QUAST & al. 2013) with a 97% nucleotide similarity level. Taxonomy assignment was performed with the naive Bayesian classifier method (WANG & al. 2007). Sequences not matching any record in the database (with a bootstrap confidence below 50) were denoted as "Unclassified". Subsequently, to minimize the artifacts created in the ASV designation, all sequences were aligned with the package Biostrings

(PAGÈS & al. 2019), and then clustered into OTUs with 99% sequence similarity applying the neighbor joining clustering method with the package DECIPHER (WRIGHT 2016).

Finally, all the sequences that were not assigned to Bacteria were removed, along with those classified as mitochondrial and chloroplast, employing the package phyloseq (version 1.22.3) (MCMURDIE & HOLMES 2013, CALLAHAN & al. 2016b). Potentially contaminating sequences were identified based on their prevalence in experimental samples and blanks using the decontam R package with a 0.5 threshold (DAVIS & al. 2018). Sequences with higher prevalence in negative control samples than in ant samples were defined as potential contaminants and then removed from the analysis. Moreover, four samples (EB16-Media1, EB16-Male2, EB16-Male3, EB16-Male1) were excluded from the analyses because they had fewer than 1000 sequence reads.

Data availability

Raw data can be accessed at the National Center for Biotechnology Information Sequence Read Archive under Bioproject ID PRJNA818015 and accession numbers SAMN26814244 - SAMN26814318.

Interaction specificity, modularity, and diversity of the ant gut microbe network

A quantitative contingency matrix summarizing the interactions of *Eciton* species (rows) and OTUs (columns) was established to quantify the interaction specificity of the *Eciton*-gut microbe network at LSBS at the species / OTU- and at the network-level. The interaction matrix was based on incidences, which represent the number of times different OTUs were detected in different army ant colonies. Accordingly, link strengths among pairs represent the number of colonies of a given *Eciton* species that contained representatives of a given OTU in ant guts (see also IVENS & al. 2016). The network approach was thus based on spatiotemporally independent collection events, yielding a conservative estimate of microbe specialization because microbes belonging to the same OTU collected from the same colony are only represented by a single incidence (BLÜTHGEN 2010). The species-level specificity was quantified with the standardized Kullback-Leibler distance d' and the network-level specificity with the standardized two-dimensional Shannon entropy H_2' (BLÜTHGEN & al. 2006). Both metrics are normalized relative to minimum and maximum possible values and consequently range from 0 (random, unspecific interactions) to 1 (highly specific interactions) (BLÜTHGEN & al. 2006). The H_2' of the actual ant-microbe network was compared with H_2' values of 10,000 randomized networks to test whether the observed network pattern deviates from random interactions using the null model algorithm for contingency matrices developed by Patefield (PATEFIELD 1981, BLÜTHGEN & al. 2006). Modularity, which describes the extent to which species interactions are clustered into distinct network subgroups, was evaluated using the quantitative modularity metric Q (DORMANN

& STRAUSS 2014). This metric is also normalized and ranges from 0 (minimum modularity) to 1 (maximum modularity).

In addition, several diversity metrics were calculated to assess the diversity of OTUs between *Eciton* species within the network. Species richness S gives the total number of OTUs detected in different *Eciton* colonies. The diversity metric e^H represents the exponential Shannon diversity of network links (JOST 2006, IVENS & al. 2016), serving here as an intuitive measure of microbial diversity per host species. In addition, comparability between S and e^H measurements of different *Eciton* species were improved by randomly sub-setting network incidences per army ant species to the lowest network incidence number of 13 incidences in *Eciton lucanoides*. Rarefaction was performed 100 times and the mean and standard deviation of S and e^H were calculated for the network subsets (S_{rare} and e^H_{rare}). The *Eciton* network of army ants and gut microbes visualizing the host specificity of gut microbes was arranged to minimize the number of crosses. The same network specificity and diversity calculations were performed for different *Eciton* castes / worker types within the species *Eciton burchellii*.

To analyze beta diversity, a non-metric multidimensional scaling plot (NMDS) was created based on Bray-Curtis dissimilarities (BRAY & CURTIS 1957) to compare microbial diversity between the different *Eciton burchellii* castes and sexes, as well as between *Eciton* species. Permutational multivariate analysis of variance (PERMANOVA) and analysis of similarities (ANOSIM) (ANDERSON & WALSH 2013) were performed employing the Bray-Curtis dissimilarity matrix with the R packages phyloseq (version 1.22.3) and vegan (2.5 - 6) (OKSANEN & al. 2019). To complement the analysis, a heat map was created by plotting the relative abundance of OTUs from the two datasets. In this analysis, only OTUs that had a relative abundance $\geq 1\%$ of the total reads in at least one sample were included. The databases were rearranged with the R package tidyverse (WICKHAM & al. 2019), and plots were created using the R packages ggplot2 (WICKHAM 2016), ComplexHeatmap (GU & al. 2016, GU 2022), and phyloseq (version 1.22.3) (MCMURDIE & HOLMES 2013). The heatmap of the *Eciton* species dataset was reordered according to their phylogenetic relationships. Also, a bar plot of the relative abundance of the OTUs of the different *Eciton* species was included as a supplementary material.

Scanning Electron Microscopy

The localization of symbiotic bacteria within the digestive tract of *Eciton burchellii* workers was visualized by SEM, with a special focus on biofilm formations. Digestive tracts of three collected *E. burchellii* media workers were dissected in sterile conditions in $1 \times$ PBS, then fixed in modified Karnovsky solution (2.5% glutaraldehyde and 2% paraformaldehyde) (KARNOVSKY 1965) for at least one hour and stored in fixative at 4°C until processing. To begin processing, samples were washed with a 0.1 M phosphate buffer. Then, samples were post-fixed with 1%

osmium tetroxide (OsO_4) solution and washed with distilled water. Subsequently, samples were dehydrated in an ethanol gradient (30, 50, 70, 80, 90, 95, and $2 \times 100\%$), washed with xylene, and embedded in paraffin (Paraplast, Leica, Wetzlar, Germany) (BANCROFT & al. 2018). Sections ($30 \mu\text{m}$) were obtained using a PT-PC PowerTome ultramicrotome (RMC products, Arizona, USA), mounted on aluminum bases and coated with gold with an Ion Coater (IB-5, Eiko engineering, Tokyo, Japan). Finally, samples were observed under a Hitachi (Okyama, Japan) S-3700N scanning electron microscope.

Transmission Electron Microscopy

The aseptically dissected gut of three *Eciton burchellii* workers were used for in-depth ultrastructural exploration using TEM. The samples were fixed with a modified Karnovsky solution and washed with a 0.1 M phosphate buffer. Subsequently, a 1% solution of OsO_4 was employed for post-fixation, after which the samples were washed with distilled water. For dehydration, an acetone gradient (30, 50, 70, 90, and $3 \times 100\%$) was applied. Samples were placed in Spurr (Electron Microscopy Sciences, Hatfield, PA, USA) medium-viscosity resin to make cuts of different thicknesses (50 to 80 nm) with the Powertome microtome. Finally, samples were contrasted with uranyl acetate and lead and observed in a Hitachi H7100 transmission electron microscope.

Fluorescence in-situ hybridization

The presence of symbiotic bacteria within the digestive tract of ants was determined by fluorescence microscopy. Digestive tracts were dissected from three workers of *Eciton burchellii* as described above and then fixed in 4% formaldehyde. Subsequently, intestinal contents were filtered employing nucleopore polycarbonate filters (Whatman, Maidstone, UK) to avoid, among other issues, auto-fluorescence. FISH was performed over the filters using a combination of eubacterial probes (EUB338 and EUB897, label CY3) and probes specific to UF (UNF16SF1 – 5' GAGTTGTCCTCGTCTTATCGG '3, label CY5) and UE (cute493R – 5' AGAAAGCCACGGCNAACTAT '3, label CY5) (ŁUKASIK & al. 2017).

Filter sections were incubated with 1 ml hybridization buffer (5 M NaCl, 1 M Tris-HCl, 1% (v/v) SDS), formamide (30% FA (v/v) and H_2O), and $3 \mu\text{l}$ (10 μM) probe working solution for 120 min at 46°C in a hybridization oven. The filter sections were then washed in prewarmed buffer (10 M Tris, 5 mM EDTA, a probe dependent amount of 5 M NaCl, H_2O) for 15 min at 48°C . Then, samples were washed in ice cold H_2O for 20s, mounted on glass slides with antifade solution that includes DAPI (ProLong™ Gold Antifade Mountant with DAPI, ThermoFisher Scientific, Waltham, MA, USA), and were finally observed with a confocal microscope Olympus FV1000 (Shinjuku, Japan). For negative controls, probes without their competitors were used to verify the presence of nonspecific labeling, and no fluorescence was observed in either case (Fig. S1).




Fig. 1: OTU abundance and host specificity. (A) Heatmap showing the abundance of OTUs in each colony (x-axis). Order of x-axis is based on phylogenetic relationships between *Eciton* species, which was recovered from WINSTON & al. 2016. (B) Network between *Eciton* army ants and gut microbes visualizing the host specificity of gut microbes. Link widths correspond to the number of times an OTU was detected in different army ant colonies. Link colors depict the microbes' species-level specificity measured as Kullback-Leibler distance (d'). Abbreviations: Ac: *Acinetobacter*, To: *Tokpelaia*, EB: *Eciton burchellii*, ED: *Eciton dulcium*, EH: *Eciton hamatum*, EL: *Eciton lucanoides*, EM: *Eciton mexicanum*, En: Enterobacteriaceae, EV: *Eciton vagans*, RA: Relative abundance, Se: *Serratia*, UE: Unclassified Entomoplasmatales, UF: Unclassified Firmicutes, We: *Weissella*. Images of *Eciton* workers by Christoph von Beeren.

Results

Microbiome composition across *Eciton* species

We obtained a total of 6,823,185 sequencing reads from 49 gut samples of the six *Eciton* species, with a median of 77,225 reads per sample (range: 3134 - 1,117,978 reads per sample, Tab. S2). ASV sequence clustering at 99% identity cutoff resulted in a total of 534 OTUs.

The ant microbial communities were dominated by multiple OTUs associated with UE (GenBank Accession: HM996853.1) and UF taxa (GenBank Accession: KX983349.1). Five bacterial OTUs representing UE and six OTUs belonging to UF were abundant across samples, together accounting for 91.8% of all reads in the dataset (38.41% for UE and 52.77% for UF). In the case of UE OTUs, UE-01 was most abundant in all *Eciton* species except *Eciton mexicanum*, where the host-specific OTU UE-02 dominated instead (Figs. 1A and S2). Regarding UF OTUs, two phylotypes were abundant in all *Eciton* species (UF-01 and UF-03). In contrast, UF-02 was mainly present in *Eciton hamatum* and *Eciton vagans* (Figs. 1A and S2). Besides these two dominant symbionts, other OTUs belonging to the genera *Weissella* (1.96%), *Tokpelaia* (1.29%), *Serratia* (0.67%), and *Acinetobacter* (0.20%) were also present in some of the *Eciton* samples (> 1% of relative abundance in at least one sample).

The final OTU table from this dataset is presented as Table S3.

Microbiome composition in *Eciton burchellii* worker castes and males

Across *E. burchellii* worker castes and males, a total of 1,621,829 sequencing reads were obtained, with a median of 49,714 reads per sample (range: 12,344 - 185,448 reads per sample, Tab. S4), which clustered into a total of 247 OTUs. The gut microbiomes of *E. burchellii* castes and males were dominated by the two bacterial phylotypes UE and UF. Four OTUs belonging to the phylotype UE and two belonging to UF were abundant in different samples, accounting for 89.54% of all reads (62.86% by UE and 26.68% by UF). Additionally, in one male sample and one submajor sample, we observed high relative abundance of *Weissella* OTUs (Fig. 2A). The final OTU table from this dataset is presented as Table S5.

Interaction specificity and diversity of the host-gut microbe network

The network of army ants and gut microbes showed a low degree of specificity ($H_2' = 0.13$), demonstrating that the community was dominated by non-specific associations. Yet, network specificity differed from purely random network models (H_2' tested against null models, $p = 0.001$),




Fig. 2: OTU abundance in digestive tracts of *Eciton burchellii* worker castes and males. (A) Heatmap showing the abundance of different OTUs in each worker caste and male sample. (B) Network between *E. burchellii* males and worker castes and gut microbes visualizing the host specificity of gut microbes. Link widths correspond to the number of times an OTU was detected in each sample type. Link colors depict the microbes' species-level specificity measured as Kullback-Leibler distance (d'). Abbreviations: Cl: *Cloacibacterium*, RA: Relative abundance, Rh: *Rhodanobacter*, UE: Unclassified Entomoplasmatales, UF: Unclassified Firmicutes, We: *Weissella*. Images of *E. burchellii* workers and male were taken from KRONAUER (2020), with permission.




Fig. 3: Alpha and beta diversity plots of the different *Eciton* species. A) Non-metric multidimensional scaling (NMDS) plot based on a Bray Curtis dissimilarity matrix of the gut microbiome of different *Eciton* species. Each point represents the gut microbiome of an individual sample, and the color represents the species to which it belongs. There is a significant difference in the composition of microbial community between the species (analysis of similarities, $p < 0.001$, $R = 0.447$; permutational multivariate analysis of variance, Pseudo- $F = 5.235$, $p < 0.001$; stress = 0.224). The relatively high stress value of the NMDS plot indicates that the plot provides a rather poor representation of the multidimensional data in a 2D space. B) Barplots showing the number of OTUs observed (S) and the exponential form of Shannon diversity (eH) and their standard deviation. S and eH are values obtained from 100 rarefied networks. Order of x-axis is based on phylogenetic relationships between *Eciton* species, which was recovered from WINSTON & al. (2016).




Fig. 4: Alpha and beta diversity plots of different castes and sexes of *Eciton burchellii*. A) Non-metric multidimensional scaling plot based on a Bray Curtis dissimilarity matrix of the gut microbiome of different castes and sexes of *E. burchellii*. Each point represents the gut microbiome of an individual sample, and the color represents its caste or sex. There is no significant difference in the composition of microbial communities between the different castes and sexes (analysis of similarities, $p = 0.257$, $R = 0.040$; permutational multivariate analysis of variance, Pseudo- $F = 0.930$, $p = 0.551$; stress = 0.176). B) Barplots showing the number of OTUs observed (S) and the exponential form of Shannon diversity (eH) and their standard deviation. S and eH are values obtained from 100 rarefied networks.

mainly because few bacterial OTUs showed clear host preferences (see UE-02, UE-05, CT-02, and AC-01 in Fig. 1B). Host specificity at the microbe OTU level was also generally low (mean d' : 0.132; range d' : 0.000 - 0.389). Similarly, interaction specificity was low at the army-ant species level (all d' values ≤ 0.18). Not surprisingly, modularity of the network was also low ($Q = 0.16$), showing that there is little partitioning of hosts between microbial OTUs. Gut microbe diversity was similar in all *Eciton* species (S_{rare} range: 8.35 - 11.00 OTUs, eH_{rare} range: 7.35 - 10.50 OTUs, Fig. 3A). Beta diversity metrics showed significant differences in the composition of the microbial communities of the different *Eciton* species (Fig. 3B; ANOSIM, $p < 0.001$, $R = 0.445$; PERMANOVA, Pseudo $F = 5.264$, $p < 0.001$; stress = 0.212). Mainly, the NMDS plot showed that the microbial communities of *Eciton hamatum* and *Eciton vagans* are more similar to each other than to the other species.









The network between *Eciton burchellii* worker castes / males and gut microbes also showed a low degree of specificity ($H_2' = 0.04$) and it did not differ from random

network models (H_2' tested against null models, $p = 0.156$). As expected then, specificity was also low at the microbe OTU level (mean d' : 0.054; range d' : 0.000 - 0.263, Fig. 2B) and at the *E. burchellii* castes / males level (all d' values ≤ 0.15), and the network showed no sign of modularity ($Q = 0.06$). Gut microbe diversity was similar in all *E. burchellii* worker castes and males (S_{rare} range: 5.91 - 7.54 OTUs, eH_{rare} range: 5.55 - 6.73 OTUs, Fig. 4A). Considering beta diversity, the gut microbial communities in male samples were separated from the worker castes in the NMDS, and they also showed a high dispersion (Fig. 4B). However, no statistically significant difference was detected (ANOSIM, $p = 0.111$, $R = 0.086$; PERMANOVA, Pseudo- $F = 1.2117$, $p = 0.221$; stress = 0.154).

Microscopic analyses of *Eciton burchellii* worker digestive tracts

Microscopic analysis of gut tissues from media workers revealed different types of intestinal cavities, especially in the hindgut. Inside one of these hindgut cavities (Fig. 5A), SEM images showed a possible biofilm formation. This bi-

Fig. 5: Localization of bacteria inside the gut of *Eciton burchellii* media workers. A) Ultrastructure of cavities in the hindgut. B) Close-up of a cavity pictured in A, showing a possible biofilm formation with the presence of coccoid-shaped bacteria. C) Ultrastructure of the hindgut showing biofilm formations (arrows). D) Close view of biofilm formation near gut microvilli. E) Bacteria fission (arrow). Filtered bacterial cells from ant worker digestive tracts employing F) a universal probe for bacteria



(red – EUB338 + EUB897) and DAPI (blue), G) a specific probe for Unclassified Firmicutes bacteria (orange – UNF16SF1) and DAPI (blue), and H) a specific probe for Unclassified Entomoplasmatales (green – cute493R) and DAPI (blue).

ofilm consists mainly of coccoid-shaped bacteria (Fig. 5B), with sizes between 200 - 700 nm. These observations could indicate the presence of UE, one of the most abundant microorganisms detected in our culture-independent analysis. TEM observations showed microorganisms near midgut microvilli (Fig. 5C) with the presence of bacilli bacteria (Fig. 5D). These intracellular bacteria range in size from 0.60 μm to 1 μm , with dense cytoplasm. Some of them can be observed undergoing cell divisions, suggesting that they are metabolically active (Fig. 5E).

FISH results were in accordance with our 16S rRNA sequencing results, showing an abundance of bacteria in the gut (Fig. 5F). The use of specific probes further confirmed the presence of the most abundant bacteria detected in the gut of *Eciton burchellii* in our culture-independent analysis: UF (Fig. 5G) and UE (Fig. 5H). FISH visualizations showed that Firmicutes are bacilliform bacteria, and Entomoplasmatales have a coccoid shape, which is consistent with our SEM observations of the *Eciton* digestive tract.

Discussion

Gut bacterial communities in *Eciton* army ants

We found that the gut bacterial communities of six sympatric *Eciton* species were dominated by a small number of bacterial taxa, which is consistent with previous reports (FUNARO & al. 2011, ANDERSON & al. 2012, ŁUKASIK & al. 2017). As mentioned before, the intestinal microbiome of different predatory ants is often dominated by one to two bacterial taxa (ŁUKASIK & al. 2017, MOREAU & RUBIN 2017), while herbivorous ants may harbor a more diverse microbiome (HU & al. 2014, 2018). However, other herbivorous ants such as *Dolichoderus* and *Pseudomyrmex* also harbor relatively few bacterial OTUs, indicating that diet alone is not sufficient to explain the diversity of gut microbial communities (SANDERS & al. 2017, BISCH & al. 2018).

The dominant phylotypes in the *Eciton* gut community identified in this study, UE and UF, have been detected in other army ant species from both the New World (e.g., *Labidus praedator* and *Nomamyrmex hartigii*) and the Old World (e.g., *Aenictus* spp. and *Dorylus* spp.), and they represent lineages highly specific to army ants (FUNARO & al. 2011, ANDERSON & al. 2012, ŁUKASIK & al. 2017). In general, the Entomoplasmatales order remains poorly understood, with the exceptions of the intracellular human pathogen *Mycoplasma* (DALEY & al. 2014) and plant- and insect-associated *Spiroplasma* (BOVÉ & al. 2003, BALLINGER & PERLMAN 2019). Different lineages of Entomoplasmatales have been associated with diverse insect taxa including bees (MEEUS & al. 2012), beetles (LUNDGREN & al. 2007), fruit flies (ANBUTSU & FUKATSU 2011), and ants (SAPOUNTZIS & al. 2015, DE OLIVEIRA & al. 2016). Bacteria from the phylum Firmicutes have been detected in a wide variety of insects, such as Coleoptera, Hemiptera, Lepidoptera, and Diptera (YUN & al. 2014), as well as in the bullet ant *Paraponera clavata* (see MOREAU & RUBIN 2017) and the edible ant *Liometopum apiculatum* (see GONZÁLEZ-ESCOBAR & al. 2018).

A previous phylogenetic analysis, based on near-full 16S rRNA, placed the UF as a well-supported, divergent, and undescribed clade, representing a potential new order within the phylum Firmicutes that also includes symbionts of other predatory ants (ŁUKASIK & al. 2017). Recently, an unclassified core gut microorganism of the ponerine ant *Diacamma cf. indicum* was included in this clade, adding evidence for a novel evolutionary origin independent from other insect-associated Firmicutes bacteria, such as *Lactobacillus*, *Apilactobacillus*, and *Bombilactobacillus* (SHIMOJI & al. 2021).

Even though these two dominant phylotypes in *Eciton* microbiomes represent lineages with a specific association, their role remains unknown (FUNARO & al. 2011, ANDERSON & al. 2012, ŁUKASIK & al. 2017). It has been proposed that they participate in the degradation of cuticular chitin of the ants' prey, although there is no experimental evidence yet to support this possibility (SAPOUNTZIS & al. 2015). In other predatory ants such as *Harpegnathos saltator*, known microbial symbionts contribute to nitrogen metabolism, including the conversion of histidine, arginine, and urea into glutamine, a more suitable source of energy that could even be involved in insect fecundity by activating cell growth via the target of rapamycin (TOR) pathway (BISCH & al. 2018). Furthermore, it has been speculated that UE bacteria are not essential for growth or development because they are absent in eggs and larvae (FUNARO & al. 2011). Neither UE nor UF bacteria appear to be pathogenic since they are ubiquitous across healthy *Eciton* individuals, consistent with previous reports from healthy *Acromyrmex* leaf-cutter ants and their dominant Entomoplasmatales symbiont (SAPOUNTZIS & al. 2015). Therefore, further research is required to fully elucidate the role of the UE and UF symbionts in army ant physiology.

In addition to UE and UF, the genus *Weissella* (phylum Firmicutes, class Bacillii) was an abundant taxon across army ant samples. This genus has been detected in a wide range of habitats (FUSCO & al. 2015), including the gut of different insects like the lepidopteran *Ostrinia nubilalis*, an important European pest of maize (BELDA & al. 2011), the cockroach *Cryptocercus kyebangensis* (HEO & al. 2019), the ant *Pseudomyrmex ferrugineus* (EILMUS & HEIL 2009), and other army ants such as *Labidus praedator* (ŁUKASIK & al. 2017). In *Eciton burchellii*, workers from the same colony showed variability in the relative abundance of this taxon, and it was suggested that the association with this bacterium is unlikely to be ancient (ŁUKASIK & al. 2017). Members of the genus *Weissella* are facultatively anaerobic bacteria, and several species have an obligate fermentative metabolism, being able to produce lactic acid and acetic acid from different sugar substrates (FUSCO & al. 2015, PRAET & al. 2015). This suggests that *Weissella* could potentially play a fermentative role in army ant guts, processing simple sugars that result from digestion. Nonetheless, several strains of *Weissella ceti* have been reported as pathogens in fish and whales, showing different metabolic adaptations such as

the presence of genes associated with virulence factors and antibiotic resistance (ABRIOUEL & al. 2015). To fully understand the functional interactions of these potential symbionts with their hosts, further experiments employing metagenomics and metatranscriptomics are needed, as well as culture-dependent strategies to isolate *Weissella* and other army ant symbionts (mainly UE and UF) to evaluate their enzymatic and secondary metabolism capabilities by performing different types of bioassays (e.g., MOREAU 2020).

It is currently challenging to contextualize these results with those from other predatory ants, since studies are limited. One possible trend is that Firmicutes are often associated with predatory ants. For example, the bullet ant, *Paraponera clavata*, hosts the Firmicutes taxon *Tumebacillus*, which is consistently present across individuals. However, the role of these microbes is still unknown (MOREAU & RUBIN 2017). Another study of the gut microbiome of predatory ants in the genus *Odontomachus* found a dominant UF bacterium. As in our study, different UF variants (ASVs) appear to be one of the main components of the reported differences between *Odontomachus chelifer* and *Odontomachus hastatus* microbiomes, suggesting species-specific microbe patterns in these species (ROCHA & al. 2022).

Gut bacterial community in worker castes and males of *Eciton burchellii*

It has been proposed that the presence of microorganisms in individual guts might correlate with the host's genotype as well as with its behavior and anatomy (SPOR & al. 2011). All these characteristics can differ between ant worker castes (JAFFÉ & al. 2007). Yet, in our analyses, minors, medias, submajors, majors, and males of *E. burchellii* harbor a similar gut bacterial community, dominated by UF and UE phylotypes. Similar results were recently found in the ponerine ant *Diacamma cf. indicum*, where workers share a similar gut microbiome, dominated by a UF bacterium, although this bacterium has very low abundance in males and queens (SHIMOJI & al. 2021). Furthermore, other adult workers from social insects, such as bees, have a relatively stable set of intestinal bacterial species, but it is distinct from the queen's microbiome (KWONG & MORAN 2015, MILLER & al. 2019).

Notwithstanding that we found a low diversity microbiome with similar dominant OTUs across *Eciton burchellii* worker types and males, a previous study employing more diverse genetic markers showed that even individuals from the same colony can harbor different UF and UE genotypes (ŁUKASIK & al. 2017). This supports the need for further phylogenomic and population genetic analyses to obtain a better understanding of the distribution patterns of these potential symbionts across army ant individuals, castes, sexes, and species.

In addition, the low gut bacterial abundance detected in the removed male samples (EB16-Male1, EB16-Male2, EB16-Male3) might be attributed to the fact that these males must have emerged from their pupae just a few days

before their collection, during the first nomadic colony emigrations (C. von Beeren, unpubl.; see SCHNEIRLA 1971, GOTWALD 1995, KRONAUER 2020). This pattern can be the result of limited social interactions between males and workers, decreasing the opportunities for symbiont transmission. This is consistent with previous results regarding bee-associated microbes, where newly hatched individuals (callow workers, males, and queens) harbored no gut bacteria or a reduced community of symbionts (KEŠNEROVÁ & al. 2016). Moreover, this agrees with previous studies of army ants that found a low incidence of gut bacteria across eggs, larvae, and pupae, suggesting that they are not required for growth and development and therefore are unlikely to be maternally transmitted (FUNARO & al. 2011), but rather socially transferred (ŁUKASIK & al. 2017). Microbiomes of army ant queens have never been studied, partly because collecting queens results in the collapse of the colony.

Host-microbe specificity

Our results for both UE and UF suggest that most bacterial OTUs show little host preference and only a few appear to show a certain level of host specificity ($d' > 0.20$), which is in line with previous findings for a larger number of New World army ant species across the Americas (ŁUKASIK & al. 2017). Our experimental setup, with all species originating from the same location, allowed us to separate the effects of phylogenetic relationship from the effect of geographic locations. As expected, we found a relatively homogeneous community of gut microbes, which can be explained by the dominance of host generalists accompanied with host co-occurrence at a single site. A few UE and UF OTUs were nonetheless differentially distributed across *Eciton* species, supporting previous results, and expanding the finding of such differences even in sympatric, closely related species that form a single community. Overall, these findings support the hypothesis of the specialization and stability between army ants and both bacterial phylotypes (UE and UF), which seems to have originated well before the diversification of the genus *Eciton* and perhaps all Dorylinae (ŁUKASIK & al. 2017). Despite this, there does not appear to be a relationship between the phylogeny of *Eciton* ants and the distribution of their symbionts, indicating that other factors are likely to determine the host-microbe associations (Fig. 1A). However, to conclusively resolve the phylogenetic and evolutionary relationships between army ants and their microbial symbionts it is critical to sequence and characterize their genomes.

Bacterial localization in the *Eciton burchellii* digestive tract

The combination of different microscopic techniques allowed us to detect and localize intestinal bacteria. By using FISH microscopy with specific probes, we confirmed the presence of the most abundant OTUs detected in our culture-independent analysis inside *Eciton* ant guts: UE and UF. The morphology and size of the bacteria detected

by FISH is consistent with observations made by electron microscopy: UE cells have a coccoid shape, like the bacteria observed with SEM (0.20 μm to 0.70 μm); UF cells are bacilliform, like the intracellular bacteria detected by TEM (0.60 μm to 1.00 μm).

The intestinal bacteria in *Eciton burchellii* observed by SEM and TEM are compartmentalized in certain regions of the gut, such as hindgut cavities or near midgut microvilli. The bacterial community of *Cephalotes* ants has been studied in detail in recent years. These ants feed mainly on extrafloral nectar, fungi, pollen, and their guts are divided into sections: the foregut (esophagus and crop), the midgut, and the hindgut (ileum and rectum). These sections show anatomical and physiological differences that likely influence the composition and abundance of the microbial communities, contributing to the function of each section in the digestion process (LANAN & al. 2016, FLYNN & al. 2021). Furthermore, employing TEM, we observed unknown intracellular bacteria near the microvilli in healthy ants, which reinforces the idea that they do not have a direct pathological impact on the health of their host (SAPOUNTZIS & al. 2015), and their function could be related to nutrition.

Bacteria in the order *Entomoplasmatales* (Mollicutes) have also been reported as one of the dominant groups in the gut community of another Neotropical ant genus, *Acromyrmex*, where they are described as cocci with an approximate diameter of 0.7 μm (SAPOUNTZIS & al. 2015). This is consistent with our findings in army ants. However, in *Acromyrmex*, these microorganisms are found extracellularly as well as intracellularly in the fat body tissues, with the presence of an extra plasma membrane that helps them survive in the host's cytoplasm (SAPOUNTZIS & al. 2015, 2018, 2019). In contrast, in army ants, these bacteria appear to live in the gut lumen, specifically within the pylorus and ileum (ŁUKASIK & al. 2017). On the other hand, Firmicutes bacteria have been widely reported in insect gut microbiomes (YUN & al. 2014, MOREAU & RUBIN 2017, GONZÁLEZ-ESCOBAR & al. 2018) and some of them share morphology and size range with UF (AUDISIO & al. 2011, TEGTMEIER & al. 2016).

Different clades of ants, and especially other insects, differ dramatically in abundance and taxonomic and functional diversity of their microbiomes (SANDERS & al. 2017). Many ant clades host very low bacterial numbers, while the microbial communities of others show nutritional or facultative endosymbionts in high numbers. Two clades of Hymenoptera whose microbiomes have been characterized quite comprehensively, herbivorous *Cephalotes* ants and social bees, host abundant, specialized, socially transmitted gut microbiota comprising multiple bacterial taxa and diverse strains, an association dating back at least 46 million years in *Cephalotes* ants (HU & al. 2018) and about 80 million years in social bees (KWONG & al. 2017). In contrast, the microbiome of army ants comprises fewer microbial clades, and given dietary differences, likely fulfills different functions. As noted by ŁUKASIK & al. (2017), another interesting difference among these hymenopteran

insects is how strain-level diversity is distributed: In *Eciton* species, the pool of strains present at the colony level is partitioned so that different workers tend to harbor distinct strains, whereas in bees, many distinct strains are present in each worker. Therefore, more systematic research on other ants and other social insects is required to elucidate the role of colony characteristics (size, nutrition, dependent or independent colony founding) in the composition of the microbial community across and within species, populations, and colonies. However, to address these questions, we require studies including phylogenetic markers with higher taxonomic resolution than the short fragments of the conserved 16S rRNA gene. Simultaneously, we should address another critical question: how the genotype-level differences among symbionts correspond to functions, and ultimately, what the role of the microbiome in the biology of the hosts is.

In the case of *Eciton* ants, these two categories of information can be provided by the rapidly improving “omics” approaches, specifically metagenomics and metatranscriptomics, coupled with cultivation efforts of UF and UE bacteria to sequence and annotate their genomes to characterize their metabolic potential and thus identify their specific functions within the host. This strategy has been applied successfully in turtle ants (CHANSON & al. 2021), fungus-growing ants (SAPOUNTZIS & al. 2015, 2018, 2019), and honeybees (ZHENG & al. 2016), and thus has the potential to allow us to fully understand the biological importance and the evolution of the microbiota associated with army ants and how it influences the host.

Conclusions

We combined microbiome composition analyses with imaging techniques to supplement information on host-microbe associations in army ants. Our results confirmed that gut communities of army ants are composed of few bacterial OTUs and are similar between the sympatric species studied, as well as between *Eciton burchellii* worker castes and males, with a clear dominance of two bacterial phylotypes: UF and UE. Microscopy techniques confirmed our sequencing results, demonstrating the presence of UF and UE bacteria in *E. burchellii* guts. Our community-based evaluation of host-microbe associations unveiled a low degree of specificity, with only a few bacterial OTUs showing clear host preferences. Possible functions of these microorganisms in army ant guts remain unknown and offer plenty of opportunities for future work.

Acknowledgements

Collecting permits were granted by the “Comisión Institucional de Biodiversidad” (Institutional Biodiversity Committee, University of Costa Rica; Resolution Number 023, 048, and 094) and authorized by La Selva Biological Station. We are grateful to the staff of the Research Center for Microscopic Structures (CIEMic) of the University of Costa Rica (UCR), for their support in electron microscopy, especially to Manuel Jiménez. Finally, we thank Dagmar Woekben, Max Nepel, and Florian Strasser for

their assistance implementing FISH techniques at UCR. A.B.F.I. was supported by the European Union Seventh Framework Programme (FP7 / 2007 - 2013) under Grant Agreement 624145. This research was supported by University of Costa Rica projects 810-B3-273 and 810-C1-080. We thank the staff members of La Selva Biological Station for their generous support during field work.

References

- ABRIOUËL, H., LERMA, L.L., CASADO MUÑOZ, M.D.C., MONTORO, B.P., KABISCH, J., PICHNER, R. & BENOMAR, N. 2015: The controversial nature of the *Weissella* genus: technological and functional aspects versus whole genome analysis-based pathogenic potential for their application in food and health. – *Frontiers in Microbiology* 6: art. 1197.
- AMATO, K.R. 2013: Co-evolution in context: The importance of studying gut microbiomes in wild animals. – *Microbiome Science and Medicine* 1: 10-29.
- ANBUTSU, H. & FUKATSU, T. 2011: *Spiroplasma* as a model insect endosymbiont. – *Environmental Microbiology Reports* 3: 144-153.
- ANDERSON, K.E., RUSSELL, J.A., MOREAU, C.S., KAUTZ, S., SULLAM, K.E., HU, Y., BASINGER, U., MOTT, B.M., BUCK, N. & WHEELER, D.E. 2012: Highly similar microbial communities are shared among related and trophically similar ant species. – *Molecular Ecology* 21: 2282-2296.
- ANDERSON, M.J. & WALSH, D.C. 2013: PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous dispersions: What null hypothesis are you testing? – *Ecological Monographs* 83: 557-574.
- AUDISIO, M.C., TORRES, M.J., SABATÉ, D.C., IBARGUREN, C. & APELLA, M.C. 2011: Properties of different lactic acid bacteria isolated from *Apis mellifera* L. bee-gut. – *Microbiological Research* 166: 1-13.
- BAHRNDORFF, S., ALEMU, T., ALEMNEH, T. & LUND, N.J. 2016: The microbiome of animals: implications for conservation biology. – *International Journal of Genomics* 2016: art. 5304028.
- BALLINGER, M.J. & PERLMAN, S.J. 2019: The defensive *Spiroplasma*. – *Current Opinion in Insect Science* 32: 36-41.
- BANCROFT, J.D., LAYTON, C. & SUVARNA, S.K. 2018: Bancroft's theory and practice of histological techniques. – Elsevier Health Sciences, Amsterdam, 557 pp.
- BELDA, E., PEDROLA, L., PERETÓ, J., MARTÍNEZ-BLANCH, J.F., MONTAGUD, A., NAVARRO, E. & PORCAR, M. 2011: Microbial diversity in the midguts of field and lab-reared populations of the European corn borer *Ostrinia nubilalis*. – *Public Library of Science One* 6: art. e21751.
- BISCH, G., NEUVONEN, M.M., PIERCE, N.E., RUSSELL, J.A., KOGA, R., SANDERS, J.G. & ANDERSSON, S.G. 2018: Genome evolution of Bartonellaceae symbionts of ants at the opposite ends of the trophic scale. – *Genome Biology and Evolution* 10: 1687-1704.
- BLÜTHGEN, N. 2010: Why network analysis is often disconnected from community ecology: a critique and an ecologist's guide. – *Basic and Applied Ecology* 11: 185-195.
- BLÜTHGEN, N., MENZEL, F. & BLÜTHGEN, N. 2006: Measuring specialization in species interaction networks. – *BioMed Central Ecology* 6: art. 9.
- BOURSAUX-EUDE, C. & GROSS, R. 2000: New insights into symbiotic associations between ants and bacteria. – *Research in Microbiology* 151: 513-519.
- BOVÉ, J.M., RENAUDIN, J., SAILLARD, C., FOISSAC, X. & GARNIER, M. 2003: *Spiroplasma citri*, a plant pathogenic mollicute: relationships with its two hosts, the plant and the leafhopper vector. – *Annual Review of Phytopathology* 41: 483-500.
- BRAY, R.J. & CURTIS, J.T. 1957: An ordination of the upland forest communities of southern Wisconsin. – *Ecological Monographs* 27: 325-349.
- CALLAHAN, B.J., MCMURDIE, P.J., ROSEN, M.J., HAN, A.W., JOHNSON, A.J.A. & HOLMES, S.P. 2016a: DADA2: High-resolution sample inference from Illumina amplicon data. – *Nature Methods* 13: 581-583.
- CALLAHAN, B.J., SANKARAN, K., FUKUYAMA, J.A., MCMURDIE, P.J. & HOLMES, S.P. 2016b: Bioconductor workflow for microbiome data analysis: from raw reads to community analyses. – *F1000Research* 5: art. 1492.
- CAPORASO, J.G., LAUBER, C.L., WALTERS, W.A., BERG-LYONS, D., HUNTLEY, J., FIERER, N. & KNIGHT, R. 2012: Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. – *The International Society for Microbial Ecology Journal* 6: 1621-1624.
- CAPORASO, J.G., LAUBER, C.L., WALTERS, W.A., BERG-LYONS, D., LOZUPONE, C.A., TURNBAUGH, P.J. & KNIGHT, R. 2011: Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. – *Proceedings of the National Academy of Sciences of the United States of America* 108: 4516-4522.
- CHANSON, A., MOREAU, C.S. & DUPLAIS, C. 2021: Assessing biosynthetic gene cluster diversity of specialized metabolites in the conserved gut symbionts of herbivorous turtle ants. – *Frontiers in Microbiology* 12: art. 678100.
- DALEY, G.M., RUSSELL, D.B., TABRIZI, S.N. & MCBRIDE, J. 2014: *Mycoplasma genitalium*: a review. – *International Journal of STD & AIDS* 25: 475-487.
- DAVIS, N.M., PROCTOR, D.M., HOLMES, S.P., RELMAN, D.A. & CALLAHAN, B.J. 2018: Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. – *Microbiome* 6: art. 226.
- DE OLIVEIRA, T.B., FERRO, M., BACCI, M., DE SOUZA, D.J., FONTANA, R., DELABIE, J.H.C. & SILVA, A. 2016: Bacterial communities in the midgut of ponerine ants (Hymenoptera: Formicidae: Ponerinae). – *Sociobiology* 63: 637-644.
- DILLON, R.J. & DILLON, V.M. 2004: The gut bacteria of insects: nonpathogenic interactions. – *Annual Review of Entomology* 49: 71-92.
- DORMANN, C.F. & STRAUSS, R. 2014: A method for detecting modules in quantitative bipartite networks. – *Methods in Ecology and Evolution* 5: 90-98.
- DOUGLAS, A.E. 2009: The microbial dimension in insect nutritional ecology. – *Functional Ecology* 23: 38-47.
- DOUGLAS, A.E. 2015: Multiorganismal insects: diversity and function of resident microorganisms. – *Annual Review of Entomology* 60: 17-34.
- DUPLAIS, C., SAROU-KANIAN, V., MASSIOT, D., HASSAN, A., PERRONE, B., ESTEVEZ, Y. & MOREAU, C.S. 2021: Gut bacteria are essential for normal cuticle development in herbivorous turtle ants. – *Nature Communications* 12: art. 676.
- EILMUS, S. & HEIL, M. 2009: Bacterial associates of arboreal ants and their putative functions in an obligate ant-plant mutualism. – *Applied and Environmental Microbiology* 75: 4324-4332.
- ENGEL, P. & MORAN, N.A. 2013: The gut microbiota of insects – diversity in structure and function. – *Federation of European Microbiological Societies Microbiology Review* 37: 699-735.
- FELDHAAR, H., STRAKA, J., KRISCHKE, M., BERTHOLD, K., STOLL, S., MUELLER, M.J. & GROSS, R. 2007: Nutritional upgrading for omnivorous carpenter ants by the endosymbiont *Blochmannia*. – *BioMed Central Biology* 5: art. 48.

- FLYNN, P.J., D'AMELIO, C.L., SANDERS, J.G., RUSSELL, J.A. & MOREAU, C.S. 2021: Localization of bacterial communities within gut compartments across *Cephalotes* turtle ants. – *Applied and Environmental Microbiology* 87: art. e02803-20.
- FRANKS, N.R. 1982: Ecology and population regulation in the army ant *Eciton burchelli*. In: LEIGH, E.G., RAND, A.S. & WINDSOR, D.M. (Eds.): *The ecology of a tropical forest: seasonal rhythms and long-term changes*. – Oxford University Press, Oxford, UK, pp. 389-395.
- FRANKS, N.R. 1985: Reproduction, foraging efficiency and worker polymorphism in army ants. – *Fortschritte der Zoologie (Stuttgart)* 31: 91-107.
- FUNARO, C.F., KRONAUER, D.J., MOREAU, C.S., GOLDMAN-HURTAS, B., PIERCE, N.E. & RUSSELL, J.A. 2011: Army ants harbor a host-specific clade of Entomoplasmatales bacteria. – *Applied and Environmental Microbiology* 77: 346-350.
- FUSCO, V., QUERO, G.M., CHO, G.S., KABISCH, J., MESKE, D., NEVE, H. & FRANZ, C.M. 2015: The genus *Weissella*: taxonomy, ecology, and biotechnological potential. – *Frontiers in Microbiology* 6: art. 155.
- GONZÁLEZ-ESCOBAR, J.L., GRAJALES-LAGUNES, A., SMOLIŃSKI, A., CHAGOLLA-LÓPEZ, A., DE LÉON-RODRÍGUEZ, A. & DE LA ROSA, A.P.B. 2018: Microbiota of edible *Liometopum apiculatum* ant larvae reveals potential functions related to their nutritional value. – *Food Research International* 109: 497-505.
- GOTWALD, W.H. 1995: *Army ants: the biology of social predation*. – Cornell University Press, New York, NY, 302 pp.
- GU, Z. 2022: Complex heatmap visualization. – *iMeta* 1: art. e43.
- GU, Z., EILS, R. & SCHLESNER, M. 2016: Complex heatmaps reveal patterns and correlations in multidimensional genomic data. – *Bioinformatics* 32: 2847-2849.
- HAMMER, T.J., JANZEN, D.H., HALLWACHS, W., JAFFE, S.P. & FIERER, N. 2017: Caterpillars lack a resident gut microbiome. – *Proceedings of the National Academy of Sciences of the United States of America* 114: 9641-9646.
- HAMMER, T.J., SANDERS, J.G. & FIERER, N. 2019: Not all animals need a microbiome. – *Federation of European Microbiological Societies Microbiology Letters* 366: art. fnz117.
- HEO, J., HAMADA, M., CHO, H., WEON, H.Y., KIM, J.S., HONG, S.B. & KWON, S.W. 2019: *Weissella cryptocerci* sp. nov., isolated from gut of the insect *Cryptocercus kyebangensis*. – *International Journal of Systematic and Evolutionary Microbiology* 69: 2801-2806.
- HOENLE, P.O., BLÜTHGEN, N., BRÜCKNER, A., KRONAUER, D.J., FIALA, B., DONOSO, D.A. & VON BEEREN, C. 2019: Species-level predation network uncovers high prey specificity in a Neotropical army ant community. – *Molecular Ecology* 28: 2423-2440.
- HU, Y., ŁUKASIK, P., MOREAU, C.S. & RUSSELL, J.A. 2014: Correlates of gut community composition across an ant species (*Cephalotes varians*) elucidate causes and consequences of symbiotic variability. – *Molecular Ecology* 23: 1284-1300.
- HU, Y., SANDERS, J.G., ŁUKASIK, P., D'AMELIO, C.L., MILLAR, J.S., VANN, D.R. & RUSSELL, J.A. 2018: Herbivorous turtle ants obtain essential nutrients from a conserved nitrogen-recycling gut microbiome. – *Nature Communications* 9: art. 964.
- IVENS, A.B.F., GADAU, A., KIERS, E.T. & KRONAUER, D.J.C. 2018: Can social partnerships influence the microbiome? Insights from ant farmers and their trophobiont mutualists. – *Molecular Ecology* 27: 1898-1914.
- IVENS, A.B.F., VON BEEREN, C., BLÜTHGEN, N. & KRONAUER, D.J.C. 2016: Studying the complex communities of ants and their symbionts using ecological network analysis. – *Annual Review of Entomology* 61: 353-371.
- JAFFÉ, R., KRONAUER, D.J.C., BERNHARD KRAUS, F., BOOMSMA, J.J. & MORITZ, R.F. 2007: Worker caste determination in the army ant *Eciton burchellii*. – *Biology Letters* 3: 513-516.
- JOST, L. 2006: Entropy and diversity. – *Oikos* 113: 363-375.
- KALTENPOTH, M. & ENGL, T. 2014: Defensive microbial symbionts in Hymenoptera. – *Functional Ecology* 28: 315-327.
- KARNOVSKY, M.J. 1965: A formaldehyde-glutaraldehyde fixative of high osmolality for use in electron microscopy. – *Journal of Cell Biology* 27: 137A-138A.
- KEŠNEROVÁ, L., MORITZ, R. & ENGEL, P. 2016: *Bartonella apis* sp. nov., a honey bee gut symbiont of the class Alphaproteobacteria. – *International Journal of Systematic and Evolutionary Microbiology* 66: 414-421.
- KOCH, H. & SCHMID-HEMPEL, P. 2011: Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. – *Proceedings of the National Academy of Sciences of the United States of America* 108: 19288-19292.
- KOZICH, J.J., WESTCOTT, S.L., BAXTER, N.T., HIGHLANDER, S.K. & SCHLOSS, P.D. 2013: Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. – *Applied and Environmental Microbiology* 79: 5112-5120.
- KRONAUER, D.J.C. 2020: *Army ants: nature's ultimate social hunters*. – Harvard University Press, Cambridge, MA, 336 pp.
- KUPPER, M., STIGLOHER, C., FELDHAAR, H. & GROSS, R. 2016: Distribution of the obligate endosymbiont *Blochmannia floridanus* and expression analysis of putative immune genes in ovaries of the carpenter ant *Camponotus floridanus*. – *Arthropod Structure & Development* 45: 475-487.
- KWONG, W.K., MEDINA, L.A., KOCH, H., SING, K-W., SOH, E. J.Y., ASCHER, J.S., JAFFÉ, R. & MORAN, N.A. 2017: Dynamic microbiome evolution in social bees. – *Science Advances* 3: art. e1600513.
- KWONG, W.K. & MORAN, N.A. 2015: Evolution of host specialization in gut microbes: the bee gut as a model. – *Gut Microbes* 6: 214-220.
- LANAN, M., RODRIGUES, P., AGELLON, A., JANSMA, P. & WHEELER, D.E. 2016: A bacterial filter protects and structures the gut microbiome of an insect. – *The International Society for Microbial Ecology Journal* 10: 1866-1876.
- LONGINO, J. 2010: *Ants of Costa Rica*. – <<http://ants.biology.utah.edu/home.htm>>, retrieved on 14 January 2021.
- ŁUKASIK, P., NEWTON, J.A., SANDERS, J.G., HU, Y., MOREAU, C.S., KRONAUER, D.J.C. & RUSSELL, J.A. 2017: The structured diversity of specialized gut symbionts of the New World army ants. – *Molecular Ecology* 26: 3808-3825.
- LUNDGREN, J.G., LEHMAN, R.M. & CHEE-SANFORD, J. 2007: Bacterial communities within digestive tracts of ground beetles (Coleoptera: Carabidae). – *Annals of the Entomological Society of America* 100: 275-282.
- MCMURDIE, P.J. & HOLMES, S. 2013: phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. – *Public Library of Science One* 8: art. e61217.
- MEEUS, I., VERCRUYSE, V. & SMAGGHE, G. 2012: Molecular detection of *Spiroplasma apis* and *Spiroplasma melliferum* in bees. – *Journal of Invertebrate Pathology* 109: 172-174.
- MILLER, D.L., PARISH, A.J. & NEWTON, I.L. 2019: Transitions and transmission: Behavior and physiology as drivers of honey bee-associated microbial communities. – *Current Opinion in Microbiology* 50: 1-7.
- MORAN, N.A., OCHMAN, H. & HAMMER, T.J. 2019: Evolutionary and ecological consequences of gut microbial communities. – *Annual Review of Ecology, Evolution, and Systematics* 50: 451-475.

- MOREAU, C.S. 2020: Symbioses among ants and microbes. – *Current Opinion in Insect Science* 39: 1-5.
- MOREAU, C.S. & RUBIN, B.E. 2017: Diversity and persistence of the gut microbiome of the giant neotropical bullet ant. – *Integrative and Comparative Biology* 57: 682-689.
- NEUVONEN, M.M., TAMARIT, D., NÄSLUND, K., LIEBIG, J., FELDBAAR, H., MORAN, N.A. & ANDERSSON, S.G. 2016: The genome of Rhizobiales bacteria in predatory ants reveals urease gene functions but no genes for nitrogen fixation. – *Scientific Reports* 6: art. 39197.
- OHKUMA, M. 2003: Termite symbiotic systems: efficient bio-recycling of lignocellulose. – *Applied Microbiology and Biotechnology* 61: 1-9.
- OKSANEN, J., GUILLAUME BLANCHET, F., FRIENDLY, M., KINDT, R., LEGENDRE, P., MCGLINN, D., MINCHIN, P.R., O'HARA, R.B., SIMPSON, G.L., SOLYMOS, P., HENRY, M., STEVENS, H., SZOEC, E. & WAGNER, H. 2019: vegan: Community Ecology Package. R Package Version 2.5-6. – <https://www.researchgate.net/publication/346579465_vegan_community_ecology_package_version_25-6_November_2020>, retrieved on 12 February 2020.
- PAGÈS, H., ABOYOUN, P., GENTLEMAN, R. & DEBROY, S. 2019: Biostrings: Efficient manipulation of biological strings. R package version 2.54.0. – <<https://rdrr.io/bioc/Biostrings/>>, retrieved on 12 February 2020.
- PATEFIELD, W.M. 1981: Algorithm AS 159: An efficient method of generating random $r \times c$ tables with given row and column totals. – *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 30: 91-97.
- POULSEN, M. & SAPOUNTZIS, P. 2012: Behind every great ant, there is a great gut. – *Molecular Ecology* 21: 2054-2057.
- POWELL, S. & FRANKS, N.R. 2006: Ecology and the evolution of worker morphological diversity: a comparative analysis with *Eciton* army ants. – *Functional Ecology* 20: 1105-1114.
- PRAET, J., MEEUS, I., CNOCKAERT, M., HOUF, K., SMAGGHE, G. & VANDAMME, P. 2015: Novel lactic acid bacteria isolated from the bumble bee gut: *Convivina intestinae* gen. nov., sp. nov., *Lactobacillus bombicola* sp. nov., and *Weissella bombi* sp. nov. – *Antonie van Leeuwenhoek* 107: 1337-1349.
- QUAST, C., PRUESSE, E., YILMAZ, P., GERKEN, J., SCHWEER, T., YARZA, P. & GLÖCKNER, F.O. 2013: The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. – *Nucleic Acids Research* 41: D590-D596.
- R CORE TEAM 2017: R: A language and environment for statistical computing. R Foundation for Statistical Computing. – <<https://www.R-project.org/>>, retrieved on 18 January 2021.
- RETTENMEYER, C.W. 1961: Arthropods associated with neotropical army ants with a review of the behavior of these ants (Arthropoda; Formicidae; Dorylinae). – Ph.D. thesis, University of Kansas, Lawrence, KS, 605 pp.
- RETTENMEYER, C.W., CHADAB-CREPET, R., NAUMANN, M.G. & MORALES, L. 1983: Comparative foraging by Neotropical army ants. – *Proceedings on the International Symposium on Social Insects in the Tropics* 2: 59-73.
- RETTENMEYER, C.W., RETTENMEYER, M.E., JOSEPH, J. & BERGHOFF, S.M. 2011: The largest animal association centered on one species: the army ant *Eciton burchellii* and its more than 300 associates. – *Insectes Sociaux* 58: 281-292.
- ROCHA, F., RONQUE, M., LYRA, M., BACCI, M. & OLIVEIRA, P. 2022: Habitat and host species drive the structure of bacterial communities of two Neotropical trap-jaw *Odontomachus* ants. – *Microbial Ecology*; doi: 10.1007/s00248-022-02064-y.
- RUSSELL, J.A., MOREAU, C.S., GOLDMAN-HUERTAS, B., FUJIWARA, M., LOHMAN, D.J. & PIERCE, N.E. 2009: Bacterial gut symbionts are tightly linked with the evolution of herbivory in ants. – *Proceedings of the National Academy of Sciences of the United States of America* 106: 21236-21241.
- SANDERS, J.G., ŁUKASIK, P., FREDERICKSON, M.E., RUSSELL, J.A., KOGA, R., KNIGHT, R. & PIERCE, N.E. 2017: Dramatic differences in gut bacterial densities correlate with diet and habitat in rainforest ants. – *Integrative and Comparative Biology* 57: 705-722.
- SANDERS, J.G., POWELL, S., KRONAUER, D.J.C., VASCONCELOS, H.L., FREDERICKSON, M.E. & PIERCE, N.E. 2014: Stability and phylogenetic correlation in gut microbiota: lessons from ants and apes. – *Molecular Ecology* 23: 1268-1283.
- SAPOUNTZIS, P., NASH, D.R., SCHIÖTT, M. & BOOMSMA, J.J. 2019: The evolution of abdominal microbiomes in fungus-growing ants. – *Molecular Ecology* 28: 879-899.
- SAPOUNTZIS, P., ZHUKOVA, M., HANSEN, L.H., SØRENSEN, S.J., SCHIÖTT, M. & BOOMSMA, J.J. 2015: *Acromyrmex* leaf-cutting ants have simple gut microbiota with nitrogen-fixing potential. – *Applied and Environmental Microbiology* 81: 5527-5537.
- SAPOUNTZIS, P., ZHUKOVA, M., SHIK, J.Z., SCHIÖTT, M. & BOOMSMA, J.J. 2018: Reconstructing the functions of endosymbiotic Mollicutes in fungus-growing ants. – *elife* 7: art. e39209.
- SCHNEIRLA, T. 1971: Army ants; a study in social organization. – W.H. Freeman & Co., San Francisco, CA, 349 pp.
- SHIMOJI, H., ITOH, H., MATSUURA, Y., YAMASHITA, R., HORI, T., HOJO, M.K. & KIKUCHI, Y. 2021: Worker-dependent gut symbiosis in an ant. – *International Society for Microbial Ecology Communications* 1: art. 60.
- SPOR, A., KOREN, O. & LEY, R. 2011: Unravelling the effects of the environment and host genotype on the gut microbiome. – *Nature Reviews Microbiology* 9: 279-290.
- SUEN, G., SCOTT, J.J., AYLWARD, F.O., ADAMS, S.M., TRINGE, S.G., PINTO-TOMÁS, A.A. & CURRIE, C.R. 2010: An insect herbivore microbiome with high plant biomass-degrading capacity. – *Public Library of Science Genetics* 6: art. e1001129.
- TEGTMEIER, D., RIESE, C., GEISSINGER, O., RADEK, R. & BRUNE, A. 2016: *Breznakia blatticola* gen. nov. sp. nov. and *Breznakia pachnodae* sp. nov., two fermenting bacteria isolated from insect guts, and emended description of the family Erysipelotrichaceae. – *Systematic and Applied Microbiology* 39: 319-329.
- VON BEEREN, C., BRÜCKNER, A., MARUYAMA, M., BURKE, G., WIESCHOLLEK, J. & KRONAUER, D.J.C. 2018: Chemical and behavioral integration of army ant-associated rove beetles – a comparison between specialists and generalists. – *Frontiers in Zoology* 15: art. 8.
- VON BEEREN, C., MARUYAMA, M. & KRONAUER, D.J.C. 2016: Cryptic diversity, high host specificity and reproductive synchronization in army ant-associated *Vatesus* beetles. – *Molecular Ecology* 25: 990-1005.
- WANG, Q., GARRITY, G.M., TIEDJE, J.M. & COLE, J.R. 2007: Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. – *Applied and Environmental Microbiology* 73: 5261-5267.
- WEISS, B. & AKSOY, S. 2011: Microbiome influences on insect host vector competence. – *Trends in Parasitology* 27: 514-522.
- WICKHAM, H. 2016: ggplot2: elegant graphics for data analysis. 2nd edn. – Springer-Verlag, New York, NY, 276 pp.

- WICKHAM, H., AVERICK, M., BRYAN, J., CHANG, W., MCGOWAN, L.D.A., FRANÇOIS, R. & YUTANI, H. 2019: Welcome to the tidyverse. – *Journal of Open Source Software* 4: art. 1686.
- WINSTON, M.E., KRONAUER, D.J.C. & MOREAU, C.S. 2016: Early and dynamic colonization of Central America drives speciation in Neotropical army ants. – *Molecular Ecology* 26: 859-870.
- WRIGHT, E.S. 2016: Using DECIPHER v2.0 to analyze big biological sequence data in R. – *R Journal* 8: 352-359.
- YUN, J.H., ROH, S.W., WHON, T.W., JUNG, M.J., KIM, M.S., PARK, D.S. & BAE, J.W. 2014: Insect gut bacterial diversity determined by environmental habitat, diet, developmental stage, and phylogeny of host. – *Applied and Environmental Microbiology* 80: 5254-5264.
- ZHENG, H., NISHIDA, A., KWONG, W.K., KOCH, H., ENGEL, P., STEELE, M.I. & MORAN, N.A. 2016: Metabolism of toxic sugars by strains of the bee gut symbiont *Gilliamella apicola*. – *MBio* 7: art. e01326-16.
- ZIENTZ, E., BEYAERT, I., GROSS, R. & FELDHAAR, H. 2006: Relevance of the endosymbiosis of *Blochmannia floridanus* and carpenter ants at different stages of the life cycle of the host. – *Applied and Environmental Microbiology* 72: 6027-6033.