

# Development and evaluation of a risk algorithm predicting alcohol dependence after early onset of regular alcohol use

Chrianna Bharat<sup>1</sup>  | Meyer D. Glantz<sup>2</sup> | Sergio Aguilar-Gaxiola<sup>3</sup> |  
 Jordi Alonso<sup>4,5,6</sup> | Ronny Bruffaerts<sup>7</sup> | Brendan Bunting<sup>8</sup> |  
 José Miguel Caldas-de-Almeida<sup>9</sup> | Graça Cardoso<sup>9</sup> | Stephanie Chardoul<sup>10</sup> |  
 Peter de Jonge<sup>11</sup> | Oye Gureje<sup>12</sup> | Josep Maria Haro<sup>13</sup> | Meredith G. Harris<sup>14,15</sup> |  
 Elie G. Karam<sup>16</sup> | Norito Kawakami<sup>17</sup> | Andrzej Kiejna<sup>18</sup> |  
 Viviane Kovess-Masfety<sup>19</sup> | Sing Lee<sup>20</sup> | John J. McGrath<sup>15,21</sup>  |  
 Jacek Moskalewicz<sup>22</sup> | Fernando Navarro-Mateu<sup>5,23</sup>  | Charlene Rapsey<sup>24</sup> |  
 Nancy A. Sampson<sup>25</sup> | Kate M. Scott<sup>24</sup> | Hisateru Tachimori<sup>26</sup> |  
 Margreet ten Have<sup>27</sup> | Gemma Vilagut<sup>4,5</sup> | Bogdan Wojtyniak<sup>28</sup> |  
 Miguel Xavier<sup>9</sup> | Ronald C. Kessler<sup>25</sup> | Louisa Degenhardt<sup>1</sup> 

## Correspondence

Chrianna Bharat, National Drug and Alcohol Research Centre, UNSW Sydney, Sydney, NSW, Australia.

Email: [c.bharat@student.unsw.edu.au](mailto:c.bharat@student.unsw.edu.au)

## Funding information

American Health Organization; Australian National Health and Medical Research Council; Bristol-Myers Squibb; Eli Lilly and Company; GlaxoSmithKline; John D. and Catherine T. MacArthur Foundation; National Institute of Health; National Institute of Mental Health; Ortho-McNeil Pharmaceutical Inc.; Pfizer Foundation; United States Public Health Service; UNSW Sydney

## Abstract

**Aims:** Likelihood of alcohol dependence (AD) is increased among people who transition to greater levels of alcohol involvement at a younger age. Indicated interventions delivered early may be effective in reducing risk, but could be costly. One way to increase cost-effectiveness would be to develop a prediction model that targeted interventions to the subset of youth with early alcohol use who are at highest risk of subsequent AD.

**Design:** A prediction model was developed for DSM-IV AD onset by age 25 years using an ensemble machine-learning algorithm known as ‘Super Learner’. Shapley additive explanations (SHAP) assessed variable importance.

**Setting and Participants:** Respondents reporting early onset of regular alcohol use (i.e. by 17 years of age) who were aged 25 years or older at interview from 14 representative community surveys conducted in 13 countries as part of WHO’s World Mental Health Surveys.

**Measurements:** The primary outcome to be predicted was onset of life-time DSM-IV AD by age 25 as measured using the Composite International Diagnostic Interview, a fully structured diagnostic interview.

**Findings:** AD prevalence by age 25 was 5.1% among the 10 687 individuals who reported drinking alcohol regularly by age 17. The prediction model achieved an

For affiliations refer to page 11

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Addiction* published by John Wiley & Sons Ltd on behalf of Society for the Study of Addiction.

external area under the curve [0.78; 95% confidence interval (CI) = 0.74–0.81] higher than any individual candidate risk model (0.73–0.77) and an area under the precision-recall curve of 0.22. Overall calibration was good [integrated calibration index (ICI) = 1.05%]; however, miscalibration was observed at the extreme ends of the distribution of predicted probabilities. Interventions provided to the 20% of people with highest risk would identify 49% of AD cases and require treating four people without AD to reach one with AD. Important predictors of increased risk included younger onset of alcohol use, males, higher cohort alcohol use and more mental disorders.

**Conclusions:** A risk algorithm can be created using data collected at the onset of regular alcohol use to target youth at highest risk of alcohol dependence by early adulthood. Important considerations remain for advancing the development and practical implementation of such models.

#### KEYWORDS

Adolescence, alcohol use, calibration, dependence, discrimination, machine learning

## INTRODUCTION

Alcohol use is a leading global risk factor for population health, estimated to be responsible for 3.7% of the total burden of disease and injuries world-wide. The interval of risk for starting alcohol use begins typically during adolescence and often progresses from experimentation to regular use [1–3]. Early use is known to be a strong predictor of later progression into alcohol use disorder [4, 5]. Therefore, the World Health Organization recommends that screening and brief interventions be part of routine health care for adolescents [6].

Screening involves the use of brief, easy-to-administer tools that can help to determine an individual's level of involvement with alcohol, detect risky and harmful drinking patterns and assist clinicians to effectively monitor an individual's progress over time [7]. Item responses are scored and compared to a scale which conveys a certain risk level to guide the clinician's response [7, 8]. Good evidence is available regarding the validity of certain instruments when used with young people [9, 10]; however, these risk evaluations are often indicative of current or short-, rather than long-term, outcomes.

There has been recent interest in the development of risk algorithms to predict, among youth, substance use disorder onset over time-periods extending into adulthood [11, 12]. Results indicate that substance use disorder onset by age 30 years can be predicted with fair to good accuracy using sociodemographic, clinical and environmental information obtained prior to the age of 18 years. While promising, the generalizability of these findings is unclear, given that models to date have been developed/applied in unrepresentative samples or their performance evaluated using the same data used in model development [13]. Further, as these models predict the outcome of any substance use disorder diagnosis rather than for a specific substance, the indicative risk for specific substance(s) are unknown, and this may limit the usefulness of the

results in guiding interventions. For these reasons, the extent to which alcohol dependence (AD) prognoses over the long-term can be evaluated among youth remains unclear.

Using data collected retrospectively from cross-sectional, general population surveys in 13 high-income countries, this study aimed to develop and internally validate preliminary prognostic models to predict AD onset by early adulthood among youth with an early onset of regular alcohol use using socio-demographic, mental health and contextual/environmental variables. If good performance measures are obtained, the algorithm could subsequently be validated in other settings, and the results used to inform data collection in future prospective trials. In addition, the use of the risk-evaluation strategies could be used to support the recruitment of target groups for studies evaluating brief interventions and treatment.

## METHOD

### Study design and setting

Data from 14 surveys (13 countries) in the World Mental Health (WMH) Survey Initiative which were defined as high-income at the time of data collection were used in developing and internally validating the prognostic models. The WMH Survey Initiative is a series of community epidemiological household surveys which include retrospective assessments of alcohol use and use disorders among other mental and substance use disorders. All surveys were based on national household samples, the exceptions being a survey of urban areas in Argentina, a specific region in Spain (Murcia region) and a series of metropolitan areas in Japan. Surveys were carried out between 2001 and 2015 (depending on the survey; see Table 1) in respondents' homes face-to-face with trained lay interviewers. Informed consent was obtained prior to all interviews using procedures approved by local ethics committees.

**TABLE 1** World Mental Health sample characteristics of high-income countries.<sup>a</sup>

Country	Sampling	Field dates	Age range at time of interview <sup>b</sup>	Sample size		Response rate <sup>c</sup>	Mean (SD) age of regular use onset among study cohort <sup>d</sup>
				Part 1	Part 2		
Argentina	Eight largest urban areas of the country (approx. 50% of the total national population)	2015	18–98	3927	2116	77.3%	15.2 (2.0)
Australia	Nationally representative	2007	18–85	8463	8463	60.0%	15.7 (1.5)
Belgium	Nationally representative	2001–02	18–95	2419	1043	50.6%	15.4 (1.7)
France	Nationally representative	2001–02	18–97	2894	1436	45.9%	15.1 (2.0)
Germany	Nationally representative	2002–03	19–95	3555	1323	57.8%	15.7 (1.4)
Japan	11 metropolitan areas	2002–06	20–98	4129	1682	55.1%	15.7 (1.7)
The Netherlands	Nationally representative	2002–03	18–95	2372	1094	56.4%	15.4 (1.3)
New Zealand	Nationally representative	2004–05	18–98	12 790	7312	73.3%	15.0 (1.9)
Northern Ireland	Nationally representative	2005–08	18–97	4340	1986	68.4%	15.6 (1.5)
Poland	Nationally representative	2010–11	18–65	10 081	4000	50.4%	15.8 (2.0)
Spain	Nationally representative	2001–02	18–98	5473	2121	78.6%	15.2 (1.9)
Spain, Murcia	Murcia region	2010–12	18–96	2621	1459	67.4%	15.5 (1.6)
United States	Nationally representative	2001–03	18–99	9282	5692	70.9%	15.1 (2.0)
Total				76 195	41 787	62.2%	15.3 (1.8)

<sup>a</sup>The World Bank (2018) data available at: <http://data.worldbank.org/country>. Some of the World Mental Health (WMH) countries have moved into new income categories since the surveys were conducted. The income groupings above reflect the status of each country at the time of data collection. The current income category of each country is available at the preceding URL.

<sup>b</sup>For the purposes of cross-national comparisons, we limit the sample to those 18+.

<sup>c</sup>The response rate is calculated as the ratio of the number of households in which an interview was completed to the number of households originally sampled, excluding from the denominator households known not to be eligible either because of being vacant at the time of initial contact or because the residents were unable to speak the designated languages of the survey.

<sup>d</sup>Unweighted mean [standard deviation (SD)] among the 10 687 individuals who were 25 years or older at time of interview and reported drinking alcohol regularly by 17 years of age.

## Outcome measure

The primary outcome to be predicted was onset of life-time DSM-IV AD by age 25 years as measured using the Composite International Diagnostic Interview [14], a fully structured diagnostic interview. The cut-off of 25 years was selected because it captured the interval of time with the highest acceleration of onset of AD relative to other periods. AD onset was evaluated as the age the respondent reported first experiencing at least three disorder-related symptoms in the same year. The outcome measure was a binary variable representing whether AD criteria were met between the onset of regular alcohol use and age 25 years.

## Sample

The master sample was defined as all individuals who, at the time of interview, were aged 25 years or older and retrospectively reported onset of regular use by age 17 years. The cut-off of 17 years was selected because it indicated an early and accelerated involvement with alcohol, was associated with an increased risk of AD relative to a later onset of regular use and fell below the legal drinking age in most included countries (see Supporting information, Appendix S1). The

analysis sample was drawn from this master sample in a variation of the case-control approach.

The cases were individuals in this master sample who reported an AD onset by age 25. Cases were separated based on serpentine sampling [15, 16] of person-level weights into a training set of 70% and a holdout set with the remaining 30%. Cases in the training sample were further subdivided into 10 training folds based on serpentine sampling of person-level weights [15, 16].

The controls were a stratified probability sample of other individuals in the master sample. Controls were sampled into a training set with 70% of controls and a holdout set, with the remaining 30% based on a serpentine type of person-level weights with implicit stratification by sex, age and survey. Controls in the training sample were further divided into 10 training folds based on serpentine sampling of person-level weights. Sampling fractions within folds were set to generate a sample of controls approximately five times the number of cases, with probability of selection equal to person-level weight.

Cases and controls were weighted by the inverse of their probabilities of selection for purposes of analysis and population projection. The 70% of cases combined with the subsampled controls created the training sample from which the prediction model was developed. The remaining 30% of cases and controls were held out to

validate the model. For model development, weights for cases in the training sample were multiplied by five times the average weight of cases in the same fold. Controls in the training folds were assigned a weight of one. This created a balanced weighted case-control ratio within each fold for model development.

## Predictor classes

All survey questions that could be used to operationalize risk factors identified in the literature review and which were collected in all surveys were identified. A total of 59 predictors (including 14 survey indicators) were evaluated as at the age the respondent reported onset of regular alcohol use across four categories of predictor classes.

### Socio-demographics

There were two variables, including a dichotomous indicator for sex and one continuous variable representing age at prediction (i.e. age of onset of regular alcohol use, range = 6–17).

### Pattern of alcohol use

There were four variables relating to history of alcohol use. Continuous variables were defined to represent the age of onset of first alcohol use (range = 6–17) and the number of years from onset of alcohol use to regular use (range = 0–12). The cumulative life-time prevalence of regular alcohol use among 25–35-year-olds in the respondent's country as of the calendar year when the respondent reported their own onset of regular alcohol use was evaluated, and categorized into units of width five (range = 15–20% to 90–95%, or 16 levels).

### History of prior mental disorders

There were six history of prior mental disorder variables. These included dichotomous indicators of generalized anxiety disorders, panic disorder, major depression and broad bipolar disorder. An ordered categorical variable representing the highest level of drug use (no use, use with no abuse, abuse with no dependence, dependence) was created and standardized. A continuous measure for the total number of prior mental disorders was also created (range = 0–5).

### Traumatic experiences

There were 34 traumatic experience variables. These included dichotomous indicators of 26 different traumatic experiences [17].

Continuous measures of the total number of any traumatic experience types (range = 0–14; entered as both standardized and stabilized variables), as well as the number from each trauma domain were evaluated. Trauma domains included exposure to organized violence (range = 0–3), participated in organized violence (range = 0–5), interpersonal violence (range = 0–4), sexual relationship violence (range = 0–6), other life-threatening trauma (range = 0–6) and secondary trauma (range = 0–2) (see Supporting information, Appendix S2).

Missing age-of-onset values were imputed using regression-based imputation controlling for key socio-demographic and clinical variables at the smallest level possible (within-country otherwise within the high-income group). A skip error in a subset of earlier surveys meant that symptoms of AD were only assessed among respondents with a history of alcohol abuse. Missing values were imputed using regression-based imputation models developed and validated on data from later surveys without the skip error (among which 13.5% of AD cases had not met criteria for alcohol abuse). Validation results suggest that the imputation models produce consistent prevalence estimates, strong individual-level classification accuracy metrics and similar distributions of important correlates; full details are described elsewhere [18].

## Statistical analysis

### Model selection

To predict AD, we used the Super Learner (SL) ensemble method to combine predicted probabilities of AD across a large number of different machine learning algorithms (the 'library') [19]. The library included a generalized linear model with a logistic link function, a series of penalized generalized linear models with different mixing models parameters, random forests, neural networks, a series of gradient boosted decision trees, Bayesian adaptive regression trees and a series of adaptive regression splines (see Supporting information, Appendix S3 for details).

A number of the algorithms in our SL library require hyperparameter tuning for optimal performance. We addressed this in simple cases by including a series of models for a single algorithm with different hyperparameter values in the SL ensemble (e.g. 10 penalized regression classifiers that differed in values of the mixing parameter, several adaptive regression splines that differed in maximum degrees). In more complex cases (random forests, gradient-boosted decision trees), we used cross-validated penalized generalized linear models to select optimal combinations of hyperparameters.

Tenfold cross-validation was used for internal SL cross-validation both to build optimal models with each classifier and to determine the optimal cross-validated weight for each classifier in the ensemble. Model performance was assessed in terms of discrimination (ability to distinguish between cases and controls) and calibration (agreement between observed and predicted probabilities) in the holdout sample.

## Discrimination

Discrimination was estimated by means of generating the receiver operating characteristic (ROC) and precision–recall (PR) curves. Both ROC and PR curves visualize trade-offs in different aspects of performance as the threshold applied to a model's prediction varies. A ROC curve shows the sensitivity (the proportion of cases that were above a given prediction threshold) as a function of the false positive rate (the proportion of controls that were above the same prediction threshold), with the area under the ROC curve (AUC) providing a measure of discriminative ability. In comparison, the PR curve shows the positive predictive value (PPV; the probability of AD among respondents above a given prediction threshold) as a function of the sensitivity, with the area under the PR curve called the 'average precision'. The PR curve is of interest because it does not incorporate correctly predicted controls and is therefore less prone to exaggerate model performance for imbalanced data sets [20]. While the baseline for ROC curves is fixed at 0.5, the baseline of a PR curve is determined by the ratio of the minority class (case) and majority class (controls) [20]. Because the PR curve is prevalence-dependent, the ROC curve should be used when comparing model performance between data sets with different event prevalence.

## Calibration

Calibration was estimated by means of calibration plots and the median ( $E_{50}$ ), 90th percentile ( $E_{90}$ ), maximum ( $E_{max}$ ) [21] and weighted mean (integrated calibration index; ICI) of the absolute difference between observed and predicted probabilities [22]. The ICI provides a measure of central tendency for summarizing the absolute differences, with weights determined by the empirical distribution of predicted probabilities.

Operating characteristics, including sensitivity, specificity and PPV, were calculated for a variety of thresholds. The 5, 10, 15 and 20% of observations with highest predicted probabilities of AD were defined by rank ordering predicted probabilities from the SL in the training sample and using the cut-points to create risk tiers of predicted probabilities in the holdout sample.

## Variable importance

Predictor variables of greatest overall importance were investigated using the model-agnostic Shapley additive explanations (SHAP) method [23], providing a summary measure of contribution to the model prediction from each variable.

## Fairness

Predictive models can be subject to biases stemming from the original data or through model development which, when applied,

may not provide equal benefit to every population subgroup [24]. To evaluate fairness, Poisson regression analyses were used to determine whether the association of observed AD with predicted probabilities were independent of key covariates. Main effects included the key covariate (evaluated at the time of regular alcohol use onset), the SL predicted probability and the interaction of these two variables. Key covariates were selected to identify groups at risk of marginalization for which data were available, and included gender, age (pre-teen: < 13 years versus teenager aged 13–17 years) and predominant country culture (western version non-western).

## Sensitivity analysis

To examine the impact of including other potential predictors of SUD onset on the findings, a sensitivity analysis was conducted which included all variables in the main analysis as well as variables capturing childhood adversities, social phobia and specific phobia disorder. Surveys which did not assess these additional variables, including Australia, New Zealand and Israel, were excluded from the sensitivity analysis. For comparison, the original model was re-run with the original set of predictors and excluding the subset of surveys which did not assess the additional predictors.

All analyses were conducted in R [25] using the SuperLearner [26], glmnet [27], xgboost [28], ranger [29], nnet [30] and earth [31] packages. The analysis was not pre-registered and results should be considered exploratory.

## RESULTS

A total of 41 787 respondents participated in the two-part survey and completed the alcohol use sections. Among the 10 687 individuals who were aged 25 years or older at the time of interview and reported drinking alcohol regularly by age 17 years, 791 [5.1%, standard error (SE) = 0.2] reported an onset of AD by age 25. Age at onset of regular alcohol use in the cohort ranged from 6 to 17 years, with a median of 15.5 (SE = 0.1).

The training sample, containing 553 cases (i.e. individuals with AD onset by age 25) and 2765 sampled controls (i.e. other individuals), was used to develop the prediction model. The remaining 30% of cases ( $n = 238$ ) and 2970 other sampled controls were held out to validate the model.

## Library weighting

The cross-validated weight that defines the relative importance of each classifier in the SL ensemble is shown in Table 2. Extreme gradient boosting was the best classifier for predicting AD onset by age 25 ( $\beta = 0.464$ ). Other weighed classifiers included a generalized linear model ( $\beta = 0.332$ ), neural network ( $\beta = 0.107$ ), multivariate adaptive


**TABLE 2** Super Learner coefficients and area under the receiver operating characteristic curve (AUC) for all candidate learners assigned non-zero coefficients in the Super Learner model as evaluated in the training sample.<sup>a</sup>

Model	Super Learner coefficient ( $\beta$ )	AUC (95% CI)	
		Training AUC	Holdout AUC <sup>b</sup>
Cross-validated SuperLearner	--	0.81 (0.80, 0.82)	--
SuperLearner	--	0.86 (0.85, 0.87)	0.78 (0.74, 0.81)
Extreme gradient boosting	0.48	0.85 (0.84, 0.86)	0.77 (0.73, 0.81)
Generalized linear model	0.33	0.84 (0.83, 0.85)	0.76 (0.72, 0.80)
Neural network	0.12	0.84 (0.83, 0.85)	0.73 (0.69, 0.77)
Adaptive polynomial spline regression	0.04	0.83 (0.84, 0.86)	0.77 (0.72, 0.80)
Adaptive regression splines (d = 1)	0.03	0.82 (0.81, 0.83)	0.77 (0.73, 0.81)
Adaptive regression splines (d = 4)	0.01	0.86 (0.85, 0.87)	0.74 (0.69, 0.78)

AUC = area under the curve; d = maximum degree of interaction; CI = confidence interval.

<sup>a</sup>The Super Learner model was developed using the training sample ( $n = 3318$ ).

<sup>b</sup>Performance of the Super Learner model was evaluated using the holdout sample ( $n = 3208$ ).



**FIGURE 1** Receiver operating characteristic curve (left) and precision-recall curve (right) for the Super Learner model predicting alcohol dependence (AD) among individuals with an onset of regular alcohol use by age 17 years in the holdout sample ( $n = 3208$ )

polynomial spline regression ( $\beta = 0.062$ ) and multivariate adaptive regression splines (1 degree,  $\beta = 0.027$ ; 4 degrees,  $\beta = 0.008$ ).

## Discrimination

The AUC of the SL model when applied to the holdout sample was 0.78 [95% confidence interval (CI) = 0.74–0.81] (see Table 2 and Figure 1). This was higher than the AUCs of the six library algorithms weighted by the SL, which were in the range of 0.73–0.77. However, a comparison of the performance of all candidate algorithms in the ensemble found four classifiers assigned weights of zero in the fitted SL model outperformed the SL model in the holdout sample. AUC was calculated as 0.78 (95% CI = 0.74–0.82) for two random forest models, 0.78 (95% CI = 0.74–0.82) for a differently parameterized multivariate adaptive regression spline fit (compared to the multivariate adaptive regression spline assigned a non-zero weight in the SL

model) and 0.78 (95% CI = 0.74–0.82) for a different extreme gradient-boosted model (see Supporting information, Appendix S4). The average precision for the SL was 0.220 which, when compared to the baseline precision of 0.052 (determined by 5.2% prevalence in the original sample), indicates a significant improvement in classification over chance (see Figure 1).


## Calibration

Calibration was satisfactory for the SL model when applied to the holdout sample with an ICI of 1.05% and  $E_{50}$  of 0.55%.  $E_{90}$  and  $E_{max}$  values for the SL model were 3.02 and 18.62%, respectively, which indicates some variation for calibration-in-the-large. The calibration plot reveals that this miscalibration occurred at the extreme ends of the predicted probability distributions, with high risks underestimated and low risks overestimated (see Figure 2).



## Operating characteristics

Inspection of the ROC curves for the SL model and the weighted classifiers in the holdout sample showed that the slope was steepest for 1-specificity in the range 0–0.05, which corresponds approximately to the 5% of respondents with highest predicted AD risk in the models (see Table 3). Visual inspection of the PR curve also shows that the precision of the model was highest over this interval. Another inflection point in the slope was at the 20th percentile. The sensitivity at these two thresholds show that these respondents



**FIGURE 2** Calibration plot for Super Learner model predicting alcohol dependence (AD) among individuals with an onset of regular alcohol use by age 17 years in the holdout sample ( $n = 3208$ ). ALD, alcohol dependence.

**TABLE 3** Cross-validated performance metric estimates for predicted probability of alcohol dependence 10 years after early onset of regular alcohol use in the holdout sample.

% Highest risk <sup>a</sup>	Holdout sample in risk threshold group		Within risk threshold group			Cumulative <sup>b</sup>		
	$n^d$	% of holdout sample(95% CI)	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV <sup>c</sup> % (95% CI)	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV % (95% CI)
5	81	2.0 (1.2, 2.7)	15.2 (9.6, 20.7)	1.3 (0.6, 2.0)	40.0 (23.7, 56.2)	15.2 (9.6, 20.7)	98.7 (98.0, 99.4)	40.0 (23.7, 56.2)
10	127	3.4 (2.7, 4.2)	12.2 (6.6, 17.9)	2.9 (2.2, 3.6)	18.9 (10.0, 27.8)	27.4 (20.5, 34.3)	95.8 (94.8, 96.8)	26.6 (19.1, 34.2)
15	144	3.4 (2.6, 4.1)	9.7 (6.7, 12.6)	3.0 (2.3, 3.7)	15.1 (9.0, 21.3)	37.0 (30.0, 44.1)	92.8 (91.6, 94.0)	22.2 (17.2, 27.3)
20	166	4.7 (3.6, 5.7)	12.0 (8.8, 15.1)	4.2 (3.2, 5.3)	13.5 (8.1, 18.9)	49.0 (42.0, 56.0)	88.6 (87.1, 90.0)	19.2 (15.4, 23.0)

PPV = positive predictive value; SE = standard error; CI = confidence interval; AD = alcohol dependence.

<sup>a</sup>Risk thresholds were created by rank ordering the Super Learner predicted probabilities in the training sample and creating 20 equally sized groups.

<sup>b</sup>Lower cut-point of risk used as clinical threshold for defining someone as being at risk of developing alcohol dependence.

<sup>c</sup>Equal to the AD prevalence within the given risk threshold group.

<sup>d</sup>Unweighted number of individuals from the holdout sample in the specific risk strata.

accounted for 15.2 and 49.0% of all cases of probable AD, respectively. This means that an intervention delivered only to the 5 or 20% of respondents with highest AD risk would capture 15.2 or 49.0%, respectively, of people who would otherwise develop AD by age 25 years.


The highest PPV for the SL model is 40.0% for the 5% threshold, indicating that this is the proportion of respondents above that threshold who would be expected to have an AD onset by age 25 in the absence of any interventions. PPV decreases to 19.2% at the most liberal threshold considered (20%). That is, we would expect one in five respondents above the 20% threshold to have an AD onset by age 25 years. Specificity in the top 5 and 20% were 98.7 and 88.6%, respectively.

## Predictor importance

The nine top predictors included four indicators of individual or cohort alcohol involvement, two relating to traumatic experiences and one each for the number of mental disorders, level of drug use involvement and gender (see Figure 3). Specifically, increased risk was associated with younger onset of both first and regular alcohol use, males, higher cohort alcohol use, fewer years from first to regular alcohol use, more secondary traumas, more mental disorders, higher levels of drug use and having experienced the unexpected death of a loved one.

## Fairness

Prevalence of AD by age 25 following early onset of regular alcohol use was higher for males (5.7 versus females, 4.2%), pre-teens (< 13 years = 6.6 versus teenagers = 4.6%) and respondents in western countries (5.5 versus non-western countries 0.9%). However, there was no substantial difference in the association between predicted probabilities from the SL model and the observed event



**FIGURE 3** Shapley additive explanations (SHAP) summary plot for the nine top predictors in the Super Learner model as evaluated in the holdout sample. DUD, drug use disorder.

rates for these variables (rate ratios ranged from 0.44 to 1.2; see Appendix S5).

### Sensitivity analysis

Results of the sensitivity analyses are summarized in Supporting information, Appendix S6. Compared to the main analysis including all high-income countries, both the subset analysis (Supporting information, Appendix S6.1) and the sensitivity analysis (Supporting information, Appendix S6.2) generated better measures of discrimination and sensitivity, similar measures of specificity and PPV and worse calibration. Top-ranked predictors similar to both the main and sensitivity analyses included age of onset of drinking, age of onset of regular drinking, prevalence of regular use among persons aged 25–35 years in the same country, gender and drug involvement. For each variable added to the sensitivity analysis, including childhood adversities, social phobia and specific phobia, the relative contribution to the model was less than 1%.

### DISCUSSION

Reflecting the need for innovative approaches to improve the delivery of alcohol prevention programmes [32], this study developed a risk model to identify, among youth who are drinking alcohol regularly, those at highest risk of developing alcohol dependence by early adulthood. In addition to evaluating the predictive performance of a preliminary prognostic algorithm for this purpose, our findings illustrate some of the standard processes and considerations in developing prognostic models as well as indicate potential

approaches to advancing their development and practical implementation.

First, the prevalence of AD by age 25 among the subgroup with the highest quintile of risk assigned by the algorithm was 19.2%; this is approximately three times as high as in the sample of people with an early onset of regular alcohol use (5.1%) and eight times as high as in the full sample of survey respondents (2.4%). These results show promise that a fairly useful risk algorithm could be used to identify a more precise subgroup of individuals at highest risk, providing a tool which could assist health professionals in implementing timely, indicated interventions, especially when endeavouring to maximize cost-effectiveness and resource allocation. However, these findings must be considered alongside the observed miscalibration at the extreme ends of the risk distribution, and the relatively low measures of sensitivity and PPV. If the algorithm presented in this study was implemented in practice, these findings imply that the model would misclassify some AD cases and that most people identified as being at risk would not develop AD by age 25. The implications and consequences of inaccurate identification from a predictive algorithm will vary depending on the context; even where an indicated intervention is warranted, stigma of the risk-label may contribute to a range of negative psychosocial factors [33, 34]. Procedures for averting such risks need to be carefully considered prior to model implementation and in the design of potential interventions.

Secondly, investigating each predictor in the algorithm and disaggregating its contribution to the risk profiles helps in understanding which factors provide the greatest signal for the outcome of interest. The most important predictors for developing AD were related to the individuals' and cohort's use of alcohol, traumatic experiences, comorbid mental disorders, other drug use and gender. Although these features are known risk factors of problematic drinking during



adulthood, these results provide confidence in terms of the contextual validity of the model based on existing domain knowledge. For outcomes or settings where there is a less established understanding of the mechanisms impacting the outcome of interest, evaluations of variable importance may lead to the discovery of new predictors. More generally, variable importance measures may overcome barriers to transparency by providing 'human-intelligible explanations' of decisions formed by otherwise uninterpretable 'black-box' algorithms [35].

Thirdly, while the algorithm in the main analysis showed fairly good internal and external validation performance, the sensitivity analyses incorporating additional information regarding childhood adversities and phobia-related diagnoses resulted in improved discrimination but worse calibration. However, this improvement in discrimination was shown to result primarily from the exclusion of select surveys, rather than from the inclusion of additional predictors. These findings suggest that there may be considerable variability in the extent to which AD can be predicted across geographical sites, even when accounting for site and/or geographical variation in the predictive algorithm. The availability of an expansive list of potentially important variables might help to identify other important predictors across settings and potentially improve model accuracy. Ideally, subsequent studies developing predictive models would focus upon conducting external validation of algorithms and endeavour to identify, from a wide range of predictors, the smallest number from which optimal accuracy can be achieved. This would advance the development and potential practical implementation of future prognostic models globally.

The current findings should be interpreted within the context of several limitations which may provide opportunities for subsequent studies. First, AD and other mental disorders were evaluated based on retrospective reporting, which may be subject to recall bias. These measures, however, were collected using well-validated instruments and have shown high diagnostic concordance with clinical diagnoses [36]. Survival bias may also contribute to a downward trend in predictive performance. Secondly, as data come from 13 high-income countries there is not full representation of all regions, income levels and other country characteristics. There was variation in response rates, the years in which the surveys were conducted and cross-national differences in the legal age of drinking and potential willingness to disclose personal information about alcohol use and other mental health and life experiences, which may be found in external validation studies to either positively or negatively impact model generalizability. Thirdly, our sensitivity test considered an expanded list of variables in model development, but other variables that the literature suggests would be useful in predicting AD, such as behavioural disorders and family history of alcohol use disorder [37], were not collected in all surveys, so could not be included in the analyses.

Finally, this study aimed to determine the potential of a risk algorithm to identify a sufficiently high concentration of at-risk individuals to support the design and implementation of indicated interventions. This has been achieved by showing that nearly half of

all AD cases occur among the 20% of patients classified by our model as having highest risk. Whether this threshold is optimal for implementing indicated interventions for AD, or if any intervention would be cost-effective at any threshold, is beyond the scope of this report. Given that a much more extensive set of predictor variables was available to build this model than exists in most routinely collected data sets, these findings provide evidence to justify a more focused effort to collect expanded information during routine health screening. If youth were prospectively followed-up to age 25 to observe which of them develop AD, longitudinal analysis could be used to see how well the predictors collected at onset of regular alcohol use help to predict which ones will be at risk of AD. This would advance the agenda of creating broadly useful AD risk algorithms to target interventions based on the presenting profiles of people with an early onset of regular alcohol drinking, as well as subsequent algorithms to predict other types of substance use disorders and mental disorders.

## Collaborators

The WHO World Mental Health Survey collaborators are Ali Al-Hamzawi MD; Yasmin A. Altwajiri PhD; Laura Helena Andrade MD, PhD; Lukoye Atwoli MD, PhD; Corina Benjet PhD; Guilherme Borges ScD; Evelyn J. Bromet PhD; Somnath Chatterji MD; Alfredo H. Cia MD; Koen Demyttenaere MD, PhD; Silvia Florescu MD, PhD; Giovanni de Girolamo MD; Oye Gureje MD, DSc, FRCPsych; Hristo Hinkov MD, PhD; Chi-yi Hu MD, PhD; Aimee Nasser Karam PhD; Elie G. Karam MD; Georges Karam MD; Jean-Pierre Lepine MD; Maria Elena Medina-Mora PhD; Marina Piazza MPH, ScD; Jose Posada-Villa MD; Tim Slade PhD; Juan Carlos Stagnaro MD, PhD; Dan J. Stein FRCPsych, PhD; Yolanda Torres MPH, Dra.HC; Maria Carmen Viana MD, PhD; Daniel V. Vigo MD, DrPH; Harvey Whiteford MBBS, PhD; and David R. Williams MPH, PhD. The authors and collaborators thank the staff of the WMH Data Collection and Data Analysis Coordination Centres for assistance with instrumentation, fieldwork, and consultation on data analysis.

## ACKNOWLEDGEMENTS

This work was supported by an Australian National Health and Medical Research Council (NHMRC) project grant (no. 1081984). C.B. is supported by a UNSW Scientia PhD scholarship and a National Drug and Alcohol Research Centre (NDARC) scholarship. L.D. is supported by an NHMRC Senior Principal Research Fellowship (1135991) and a USA National Institute of Health (NIH) National Institute on Drug Abuse (NIDA) grant (R01DA1104470). NDARC, UNSW Sydney, is supported by funding from the Australian Government Department of Health under the Drug and Alcohol Program. The World Health Organization World Mental Health (WMH) Survey Initiative is supported by the United States National Institute of Mental Health (NIMH; R01 MH070884), the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the United States Public Health Service (R13-MH066849, R01-MH069864 and R01 DA016558), the

Fogarty International Center (FIRCA R03-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical Inc., GlaxoSmithKline, and Bristol-Myers Squibb. The 2007 Australian National Survey of Mental Health and Wellbeing is funded by the Australian Government Department of Health and Ageing. The Argentina survey—Estudio Argentino de Epidemiología en Salud Mental (EASM)—was supported by a grant from the Argentinian Ministry of Health (Ministerio de Salud de la Nación) (grant number 2002-17270/13-5). The ESEMeD project is funded by the European Commission (Contracts QL6-1999-01042; SANCO 2004123 and EAHC 20081308), the Piedmont Region (Italy), Fondo de Investigación Sanitaria, Instituto de Salud Carlos III, Spain (FIS 00/0028), Ministerio de Ciencia y Tecnología, Spain (SAF 2000-158-CE), Generalitat de Catalunya (2017 SGR 452; 2014 SGR 748), Instituto de Salud Carlos III (CIBER CB06/02/0046, RETICS RD06/0011 REM-TAP) and other local agencies and by an unrestricted educational grant from GlaxoSmithKline. The World Mental Health Japan (WMHJ) Survey is supported by the grant for Research on Psychiatric and Neurological Diseases and Mental Health (H13-SHOGAI-023, H14-TOKUBETSU-026, H16-KOKORO-013, H25-SEISHIN-IPPAN-006) from the Japan Ministry of Health, Labour and Welfare. Te Rau Hinengaro: the New Zealand Mental Health Survey (NZMHS) is supported by the New Zealand Ministry of Health, Alcohol Advisory Council, and the Health Research Council. The Northern Ireland Study of Mental Health was funded by the Health and Social Care Research and Development Division of the Public Health Agency. The Polish project Epidemiology of Mental Health and Access to Care—EZOP Project (PL 0256) was carried out by the Institute of Psychiatry and Neurology in Warsaw in consortium with Department of Psychiatry—Medical University in Wrocław and National Institute of Public Health National Institute of Hygiene in Warsaw and in partnership with Psykiatrist Institut Vinderen—Universitet, Oslo. The project was funded by the European Economic Area Financial Mechanism and the Norwegian Financial Mechanism. The EZOP project was co-financed by the Polish Ministry of Health. The Portuguese Mental Health Study was carried out by the Department of Mental Health, Faculty of Medical Sciences, NOVA University of Lisbon, with collaboration of the Portuguese Catholic University, and was funded by Champalimaud Foundation, Gulbenkian Foundation, Foundation for Science and Technology (FCT) and Ministry of Health. The Psychiatric Enquiry to General Population in Southeast Spain—Murcia (PEGASUS—Murcia) Project has been financed by the Regional Health Authorities of Murcia (Servicio Murciano de Salud and Consejería de Sanidad y Política Social) and Fundación para la Formación e Investigación Sanitarias (FFIS) of Murcia. The US National Comorbidity Survey Replication (NCS-R) is supported by the National Institute of Mental Health (NIMH; U01-MH60220) with supplemental support from the National Institute of Drug Abuse (NIDA), the Substance Abuse and Mental Health Services Administration (SAMHSA), the Robert Wood Johnson Foundation (RWJF; grant 044708) and the John W. Alden Trust. A complete list of all within-country and cross-national WMH publications can be found at <http://www.hcp.med.harvard.edu/wmh/>. None of the funders had

any role in the design, analysis, interpretation of results or preparation of this paper. The views and opinions expressed in this report are those of the authors and should not be construed to represent the views of the World Health Organization, other sponsoring organizations, agencies or governments. Open access publishing facilitated by University of New South Wales, as part of the Wiley - University of New South Wales agreement via the Council of Australian University Librarians.

## DECLARATION OF INTERESTS

In the past 3 years, L.D. has received investigator-initiated untied educational grants for studies of opioid medications in Australia from Indivior, Mundipharma and Seqirus. M.G.H. reports consulting fees from RAND Corporation outside the submitted work. N.K. reports grants and consulting fees outside the submitted work. He received grants from Fujitsu Japan, Ltd and SBAtWork Corporation and consulting fees from Occupational Health Foundation, Japan Dental Association, Sekisui Chemicals, Junpukai Health Care Center and Osaka Chamber of Commerce and Industry. R.C.K. and N.A.S. report research grants from National Institute of Mental Health, USA (Grants: R01 MH070884; U01 MH60220); John D. and Catherine T. MacArthur Foundation; Pfizer Foundation; US Public Health Service (Grants: R13-MH066849, R01-MH069864 and R01 DA016558); Fogarty International Center (Grant: R03-TW006481); Pan American Health Organization; Eli Lilly and Company; Ortho-McNeil Pharmaceutical; GlaxoSmithKline; Bristol-Myers Squibb; National Institute of Drug Abuse; Substance Abuse and Mental Health Services Administration, USA; Robert Wood Johnson Foundation (Grant 044708); and John W. Alden Trust. In the past 3 years, R.C.K. was a consultant for Datastat, Inc., Holmusk, RallyPoint Networks, Inc. and Sage Therapeutics. He has stock options in Mirah, PYM and Roga Sciences. H.T. reports research grants from the Ministry of Health, Labour and Welfare, Japan. All other authors report no conflicts of interest.

## AUTHOR CONTRIBUTIONS

**Meyer D Glantz:** Conceptualization; funding acquisition; methodology; supervision; writing-original draft; writing-review and editing. **Sergio Aguilar-Gaxiola:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Jordi Alonso:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Ronny Bruffaerts:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Brendan Bunting:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **José M. Caldas-de-Almeida:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Graca Cardoso:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Stephanie Chardoul:** Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Peter de Jonge:** Conceptualization; data curation; funding acquisition; project administration; writing-review

and editing. **Oye Gureje**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Josep Maria Haro**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Meredith Harris**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Elie G Karam**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Norito Kawakami**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Andrzej Kiejna**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Viviane Kovess-Masfety**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Sing Lee**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **John J McGrath**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Jacek Moskalewicz**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Fernando Navarro-Mateu**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Charlene Rapsey**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Nancy A Sampson**: Conceptualization; data curation; funding acquisition; project administration; resources; supervision; writing-review and editing. **Kate Scott**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Hisateru Tachimori**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Margreet ten Have**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Gemma Vilagut**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Bogdan Wojtyniak**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Miguel Xavier**: Conceptualization; data curation; funding acquisition; project administration; writing-review and editing. **Ronald Kessler**: Conceptualization; data curation; formal analysis; funding acquisition; project administration; resources; supervision; writing-original draft; writing-review and editing. **Louisa Degenhardt**: Conceptualization; data curation; funding acquisition; project administration; resources; supervision; writing-original draft; writing-review and editing.

## AFFILIATIONS

<sup>1</sup>National Drug and Alcohol Research Centre (NDARC), University of New South Wales Australia, Sydney, NSW, Australia

<sup>2</sup>Department of Epidemiology, Services, and Prevention Research (DESPR), National Institute on Drug Abuse (NIDA), National Institute of Health (NIH), Bethesda, MA, USA

<sup>3</sup>Center for Reducing Health Disparities, UC Davis Health System, Sacramento, CA, USA

<sup>4</sup>Health Services Research Unit, IMIM-Hospital del Mar Medical Research Institute, Barcelona, Spain

<sup>5</sup>Instituto de Salud Carlos III, Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

<sup>6</sup>Department of Life and Health Sciences, Pompeu Fabra University (UPF), Barcelona, Spain

<sup>7</sup>Universitair Psychiatrisch Centrum - Katholieke Universiteit Leuven (UPC-KUL), Campus Gasthuisberg, Leuven, Belgium

<sup>8</sup>School of Psychology, Ulster University, Londonderry, UK

<sup>9</sup>Lisbon Institute of Global Mental Health and Chronic Diseases Research Center (CEDOC), NOVA Medical School|Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisbon, Portugal

<sup>10</sup>Institute for Social Research, University of Michigan, Ann Arbor, MI, USA

<sup>11</sup>Department of Developmental Psychology, University of Groningen, Groningen, The Netherlands

<sup>12</sup>Department of Psychiatry, University College Hospital, Ibadan, Nigeria

<sup>13</sup>Research, Teaching and Innovation Unit, Parc Sanitari Sant Joan de Déu, Sant Boi de Llobregat, Centre for Biomedical Research on Mental Health (CIBERSAM), Madrid, Spain

<sup>14</sup>School of Public Health, The University of Queensland, Herston, QLD, Australia

<sup>15</sup>Queensland Centre for Mental Health Research, The Park Centre for Mental Health, Wacol, QLD, Australia

<sup>16</sup>Department of Psychiatry and Clinical Psychology, Faculty of Medicine, Institute for Development, Research, Advocacy and Applied Care (IDRAAC), St George Hospital University Medical Center, Balamand University, Beirut, Lebanon

<sup>17</sup>Department of Mental Health, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

<sup>18</sup>Institute of Psychology, University of Lower Silesia, Wroclaw, Poland

<sup>19</sup>Ecole des Hautes Etudes en Santé Publique (EHESP), Paris Descartes University, Paris, France

<sup>20</sup>Department of Psychiatry, Chinese University of Hong Kong, Tai Po, Hong Kong

<sup>21</sup>Queensland Brain Institute, The University of Queensland, National Centre for Register-based Research, Aarhus University, Aarhus V, Denmark

<sup>22</sup>Institute of Psychiatry and Neurology, Warsaw, Poland

<sup>23</sup>Department of Basic Psychology and Methodology, University of Murcia, Murcia Biomedical Research Institute (IMIB-Arrixaca), Unidad de Docencia, Investigación y Formación en Salud Mental, Servicio Murciano de Salud, Murcia, Spain

<sup>24</sup>Department of Psychological Medicine, University of Otago, Dunedin, New Zealand

<sup>25</sup>Department of Health Care Policy, Harvard Medical School, Boston, MA, USA

<sup>26</sup>Department of Clinical Data Science, Clinical Research and Education Promotion Division, National Center of Neurology and Psychiatry, Endowed Course for Health System Innovation, Keio University School of Medicine, Tokyo, Japan

<sup>27</sup>Trimbos-Instituut, Netherlands Institute of Mental Health and Addiction, Utrecht, the Netherlands

<sup>28</sup>Centre of Monitoring and Analyses of Population Health, National Institute of Public Health-National Research Institute, Warsaw, Poland

## ORCID

Chrianna Bharat  <https://orcid.org/0000-0002-9778-0386>

John J. McGrath  <https://orcid.org/0000-0002-4792-6068>

Fernando Navarro-Mateu  <https://orcid.org/0000-0002-3228-623X>

Louisa Degenhardt  <https://orcid.org/0000-0002-8513-2218>

## REFERENCES

- Degenhardt L, Chiu W-T, Sampson N, Kessler RC, Anthony JC, Angermeyer M, et al. Toward a global view of alcohol, tobacco, cannabis, and cocaine use: findings from the WHO World Mental Health Surveys. *PLOS Med*. 2008;5:e141.
- Masten AS, Faden VB, Zucker RA, Spear LP. Underage drinking: a developmental framework. *Pediatrics*. 2008;121:S235–51.
- Glantz MD, Bharat C, Degenhardt L, Sampson NA, Scott KM, Lim CCW, et al. The epidemiology of alcohol use disorders cross-nationally: findings from the World Mental Health Surveys. *Addict Behav*. 2020;102:106128.
- Dawson DA, Goldstein RB, Patricia Chou S, June Ruan W, Grant BF. Age at first drink and the first incidence of adult-onset DSM-IV alcohol use disorders. *Alcohol Clin Exp Res*. 2008;32:2149–60.
- Warner LA, White HR, Johnson V. Alcohol initiation experiences and family history of alcoholism as predictors of problem-drinking trajectories. *J Stud Alcohol Drugs*. 2007;68:56–65.
- World Health Organization (WHO). Orientation Programme on Adolescent Health for Health Care Providers. Geneva, Switzerland: WHO; 2006.
- Harris SK, Louis-Jacques J, Knight JR. Screening and brief intervention for alcohol and other abuse. *Adolesc Med State Art Rev*. 2014;25:126–56.
- Pilowsky DJ, Wu L-T. Screening instruments for substance use and brief interventions targeting adolescents in primary care: a literature review. *Addict Behav*. 2013;38:2146–53.
- Toner P, Böhnke JR, Andersen P, Mccambridge J. Alcohol screening and assessment measures for young people: a systematic review and meta-analysis of validation studies. *Drug Alcohol Depend*. 2019;202:39–49.
- Clark DB, Gordon AJ, Ettaro LR, Owens JM, Moss HB. Screening and brief intervention for underage drinkers. *Mayo Clin Proc*. 2010;85:380–91.
- Meier MH, Hall W, Caspi A, Belsky DW, Cerdá M, Harrington HL, et al. Which adolescents develop persistent substance dependence in adulthood? Using population-representative longitudinal data to inform universal risk assessment. *Psychol Med*. 2016;46:877–89.
- Jing Y, Hu Z, Fan P, Xue Y, Wang L, Tarter RE, et al. Analysis of substance use and its outcomes by machine learning I. childhood evaluation of liability to substance use disorder. *Drug Alcohol Depend*. 2020;206:107605.
- Steyerberg EW, Vergouwe Y. Towards better clinical prediction models: seven steps for development and an ABCD for validation. *Eur Heart J*. 2014;35:1925–31.
- Kessler RC, Üstün TB. The World Mental Health (WMH) survey initiative version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int J Methods Psychiatr Res*. 2004;13:93–121.
- Chromy JR. Sequential sample selection methods. *Proc Surv Res Meth Sect Am Stat Assoc* 1979;1979:401–406.
- Williams RL, Chromy JR. SAS Sample Selection Macros. In: *Proceedings of the Fifth Annual SAS Users Group International Conference* SAS Institute Inc: Cary, NC; 1980.
- Kessler RC, Rose S, Koenen KC, Karam EG, Stang PE, Stein DJ, et al. How well can post-traumatic stress disorder be predicted from pre-trauma risk factors? An exploratory study in the WHO World Mental Health Surveys. *World Psychiatry*. 2014;13:265–74.
- Lago L, Glantz MD, Kessler RC, Sampson NA, al-Hamzawi A, Florescu S, et al. Substance dependence among those without symptoms of substance abuse in the World Mental Health Survey. *Int J Methods Psychiatr Res*. 2017;26:e1557.
- Polley EC. *Super Learner* Berkeley, CA: University of California; 2010.
- Saito T, Rehmsmeier M. The precision-recall plot is more informative than the ROC plot when evaluating binary classifiers on imbalanced datasets. *PLOS ONE*. 2015;10:e0118432.
- Harrell FE. *Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal Regression, and Survival Analysis* New York, NY: Springer Publishing; 2015.
- Austin PC, Steyerberg EW. The integrated calibration index (ICI) and related metrics for quantifying the calibration of logistic regression models. *Stat Med*. 2019;38:4051–65.
- Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. *Proceedings of the 31st International Conference on Neural Information Processing Systems*, Long Beach, CA, USA; 2017, pp. 4768–4777.
- Bharat C, Hickman M, Barbieri S, Degenhardt L. Big data and predictive modelling for the opioid crisis: existing research and future potential. *Lancet Digital Health*. 2021;3:e397–407.
- R Core Team. *R: A Language and Environment for Statistical Computing* Vienna, Austria: R Foundation for Statistical Computing; 2019.
- Polley E, LeDell E, Kennedy C, van der Laan M. *SuperLearner: Super Learner Prediction*. R package version 2.0–26. Vienna, Austria: R Foundation for Statistical Computing; 2019.
- Friedman J, Hastie T, Tibshirani R. Regularization paths for generalized linear models via coordinate descent. *J Stat Softw*. 2010;33:1–22.
- Chen T, He T, Benesty M, Khotilovich V, Tang Y, Cho H et al. *xgboost: Extreme Gradient Boosting*. R package version 1.3.2.1. Vienna, Austria: R Foundation for Statistical Computing; 2021.
- Wright MN, Ziegler A. *Ranger: A fast implementation of random forests for high dimensional data in C++ and R*. *J Stat Softw*. 2017;77:1–17.
- Venables WN, Ripley BD. *Modern Applied Statistics with S*. New York: Springer Publishing; 2002.
- Milborrow S. Derived from mda:mars by Trevor Hastie and Rob Tibshirani. Uses Alan Miller's Fortran utilities with Thomas Lumley's leaps wrapper. *earth: Multivariate Adaptive Regression Splines*. R package version 5.3.0. Vienna, Austria: R Foundation for Statistical Computing; 2020.
- Demant J, Schierff LM. Five typologies of alcohol and drug prevention programmes. A qualitative review of the content of alcohol and drug prevention programmes targeting adolescents. *Drugs Educ Prev Policy*. 2019;26:32–9.
- Yang LH, Link BG, Ben-David S, Gill KE, Girgis RR, Brucato G, et al. Stigma related to labels and symptoms in individuals at clinical high-risk for psychosis. *Schizophr Res*. 2015;168:9–15.
- Livingston JD, Boyd JE. Correlates and consequences of internalized stigma for people living with mental illness: a systematic review and meta-analysis. *Soc Sci Med*. 2010;71:2150–61.
- Goodman B, Flaxman S. European Union regulations on algorithmic decision-making and a 'right to explanation'. *AI Mag*. 2017;38:50–7.

36. Haro JM, Arbabzadeh-Bouchez S, Brugha TS, de Girolamo G, Guyer ME, Jin R, et al. Concordance of the composite international diagnostic interview version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health Surveys. *Int J Methods Psychiatr Res.* 2006;15:167–80.
37. Hawkins JD, Catalano RF, Miller JY. Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: implications for substance abuse prevention. *Psychol Bull.* 1992;112:64–105.

### SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Bharat C, Glantz MD, Aguilar-Gaxiola S, Alonso J, Bruffaerts R, Bunting B, et al. Development and evaluation of a risk algorithm predicting alcohol dependence after early onset of regular alcohol use. *Addiction.* 2023. <https://doi.org/10.1111/add.16122>