

## UvA-DARE (Digital Academic Repository)

### Swift and RXTE follow up observations of the transient currently active in the globular cluster Terzan 5

Altamirano, D.; Degenaar, N.D.; Heinke, C.O.; Homan, J.; Pooley, D.; Sivakoff, G.R.; Wijnands, R.

**Publication date**

2011

**Document Version**

Final published version

**Published in**

The astronomer's telegram

**License**

Unspecified

[Link to publication](#)

**Citation for published version (APA):**

Altamirano, D., Degenaar, N. D., Heinke, C. O., Homan, J., Pooley, D., Sivakoff, G. R., & Wijnands, R. (2011). Swift and RXTE follow up observations of the transient currently active in the globular cluster Terzan 5. *The astronomer's telegram*, 3720.  
<https://www.astronomerstelegram.org/?read=3720>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)*

Outside  
GCN  
IAUCs  
ATel on Twitter  
Patreon

# The Astronomer's Telegram

[Post](#) | [Search](#) | [Policies](#)  
[Credential](#) | [Feeds](#) | [Email](#)

14 Oct 2022; 11:07 UT

This space for free for your conference.



**Thanks to Patrons, The Astronomer's Telegram is free to read, free to publish and always will be. Thank you.**

[ [Previous](#) | [Next](#) | [ADS](#) ]

## Swift and RXTE follow up observations of the transient currently active in the globular cluster Terzan 5

ATel #3720; **D. Altamirano (U. Amsterdam), N. Degenaar (U. Amsterdam), C. O. Heinke (U. Alberta), J. Homan (MIT), D. Pooley (Sam Houston State U.), G. R. Sivakoff (U. Alberta), R. Wijnands (U. Amsterdam)**

on 28 Oct 2011; 01:29 UT

Credential Certification: Gregory R Sivakoff ([grs8g@virginia.edu](mailto:grs8g@virginia.edu))

Subjects: X-ray, Binary, Globular Cluster, Neutron Star, Transient

Referred to by ATel #: 3723, 3729, 3743

[Tweet](#)

Following the detection of an X-ray outburst in the direction of Terzan 5 (ATEL #3714), we obtained a Swift observation and additional RXTE observations.

The XRT aboard Swift observed Terzan 5 on Oct. 26, 2011 in imaging mode for a total exposure time of 967 s. The source was detected at high count rates causing significant pile-up (the core is saturated), and a bad column intersects the point-spread function. We estimate (aligning a circle by eye with symmetric parts of the PSF) a position of 17:48:05.43, -24:46:47.9 (J2000) with a 1" (1  $\sigma$ ) error. In addition, the Swift XRT has a systematic positional uncertainty of 3.5" (Goad et al. 2007, A&A, 476, 1401).

Two short (< 30 min) RXTE observations were performed at 23:04 UT on Oct 26th and 00:38 UT on Oct. 27th, i.e., at about 17.5 and 19 hrs after the first discovery (ATEL #3714). Within this time span the source brightened from ~8 mCrab to 83 and 90 mCrab (2-16 keV), respectively. During our observations we detect highly significant broad-band noise in the range 0.007-10 Hz with fractional rms amplitudes of  $13 \pm 1\%$  and  $15 \pm 1\%$  (2-60 keV), respectively. No thermonuclear X-ray bursts were detected and we find no evidence for quasi-periodic oscillations nor coherent pulsations.

We fit the 3-30 keV RXTE/PCA spectrum assuming  $N_H = 1.2 \times 10^{22} \text{ cm}^{-2}$ , finding a good fit including diskbb, powerlaw, and gaussian components. The diskbb has  $kT = 2.55 \pm 0.08$  keV, powerlaw photon index =  $2.25 \pm 0.14$ , and iron line equivalent width (fixed at 6.5 keV) of  $0.39 \pm 0.07$  keV. The diskbb component contributes the majority,  $59 \pm 3\%$ , of the 3-30 keV flux. Replacing the diskbb with a single blackbody produces a worse fit. While adding an additional blackbody component produces a better statistical fit, this drives the

### Related

- 9072 Outburst from low-mass X-ray binary GRS 1747-312 in Terzan 6
- 4915 Confirmation of GRS 1747-312 as the active transient in Terzan 6
- 4302 Chandra Identification of the 2012 Transient in Terzan 5
- 4264 The new X-ray transient Swift J174805.3-244637 in Terzan 5 is a neutron star LMXB
- 4249 Brightening and hardening of new X-ray transient in globular cluster Terzan 5
- 4242 A new X-ray transient in the globular cluster Terzan 5
- 3924 A Candidate Optical Counterpart of IGR J17480-2446 in Terzan 5 identified with the Hubble Space Telescope
- 3892 Sub-arcsecond position of the the transient 11-Hz X-ray pulsar in the Globular Cluster Terzan 5 derived from a Moon occultation observed by RXTE
- 3743 Chandra Identification of the 2011 Transient in Terzan 5: Same as the 2000 Transient
- 3729 MAXI detection of a super-burst from EXO 1745-248 in Terzan 5
- 3723 Improved Swift-XRT position of the transient source in Terzan 5
- 3720 **Swift and RXTE follow up observations of the transient currently active in the globular cluster Terzan 5**
- 3718 INTEGRAL non detection of renewed activity from Terzan 5
- 3714 New X-ray transient outburst in Terzan 5
- 3650 MAXI J0556-332 is a transient neutron-star Z source
- 3264 A possible IR counterpart to the transient X-ray pulsar IGR J17480-2446 in Terzan 5
- 3044 Terzan 5 transient IGR J17480-2446: return of thermonuclear bursts or were they always there?
- 3000 Type-II bursts from the new Terzan 5 transient: a GRO J1744-28 analogue?
- 2974 Chandra Identification of the Transient in Terzan 5: Not the 2000 Transient.
- 2958 Discovery of mHz QPOs and burst rate evolution in the active Terzan 5 neutron star transient
- 2952 D. Altamirano (Amsterdam), J. Homan (MIT), M. Linares (MIT), A. Patruno (Amsterdam), Y. Yang (Amsterdam), A. Watts (Amsterdam), M. Kalamkar

diskbb temperature to an unreasonably high value.

Our position is consistent with only one source among the quiescent X-ray sources listed in Heinke et al. 2006 (ApJ, 651, 1098), the X-ray burster EXO 1745-248, which was active in 2000 (Heinke et al. 2003, ApJ, 590, 809). It is not consistent with the position of the 11 Hz pulsar IGR J17480-2446. Given this Swift position and the fact that we do not detect any evidence of coherent pulsations at 11 Hz, we conclude that the transient source is most likely not the X-ray pulsar IGR J17480-2446. Although the position is consistent with that of EXO 1745-248 (Markwardt & Swank 2000, IAUC # 7454 ; Wijnands et al. 2005, ApJ, 618, 883), the identification is inconclusive given the size of the Swift error circle in combination with the high number of quiescent X-ray binaries in this globular cluster, and the fact that several X-ray binaries are likely fainter than the detection limits of Chandra in this cluster.

MAXI has reported (T. Mihara, transient alert 2010-Oct-27 09:57:39) a source at the location of Terzan 5 with flux of 0.5 Crab (4-10 keV), appearing at MJD 55858.53 and decaying by MJD 55859.53. We confirm this with the Swift/BAT lightcurve, which in the same time range shows a flare to  $0.01 \text{ cts cm}^{-2}\text{s}^{-1}$ . For a 2.5 keV blackbody spectrum at 5.5 kpc, the Swift/BAT count rate corresponds to  $1.2 \times 10^{37} \text{ ergs s}^{-1}$  (15-30 keV), or  $8 \times 10^{37} \text{ ergs a}^{-1}$  (0.5-30 keV), consistent with the MAXI result. Given an approximate 30 ks characteristic timescale; its fluence is on the order of  $10^{42} \text{ erg}$ . Given this flux, timescale, and fluence, we speculate this flare is a superburst on the surface of the binary's neutron star. This superburst occurred before the INTEGRAL observations reported by Atel #3718, and our RXTE observations. This suggests that the superburst may have triggered the current outburst.

We thank the full Swift and RXTE team for their help in promptly scheduling the observations.

(Amsterdam), P. Casella (Southampton), M. Armas-Padilla (Amsterdam), Y. Cavecchi (Amsterdam), N. Degenaar (Amsterdam), D. Russell (Amsterdam), R. Kaur (Amsterdam), M. van der Klis (Amsterdam), N. Rea (CSIC-IEEC), R. Wijnands (Amsterdam)

- 2946 A Refined Orbital Solution and the Transient Pulsar in Terzan 5 is Not Eclipsing
- 2940 INTEGRAL and RXTE spectral analysis of IGR J17480-2446, the new transient in Terzan 5.
- 2939 A preliminary orbital solution for the transient eclipsing pulsar in Terzan 5
- 2937 Eclipsing X-ray Burster in Terzan 5: Improved Swift Localization
- 2935 Discovery of kilohertz quasi-periodic oscillations from EXO 1745-248
- 2933 The Eclipsing X-ray Burster in Terzan 5 is Probably Not the 2000 Transient
- 2932 Discovery of 11 Hz burst oscillations from the 11 Hz Eclipsing Pulsar in Terzan 5
- 2929 EXO 1745-248 is an 11 Hz Eclipsing Pulsar
- 2924 Further INTEGRAL observations of the transient X-ray burster EXO 1745-248
- 2922 Swift follow-up observations of EXO 1745-248
- 2920 Swift-XRT location of the ongoing Terzan 5 outburst
- 2919 A hard X-ray transient in the direction of Terzan 5 detected by INTEGRAL

### [ Telegram Index ]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

[rrutledge@astronomerstelegram.org](mailto:rrutledge@astronomerstelegram.org)

[dfox@astronomerstelegram.org](mailto:dfox@astronomerstelegram.org)