



UvA-DARE (Digital Academic Repository)

Reservoir spectroscopy of 5s5p 3P2-5snd 3D1,2,3 transitions in strontium

Stellmer, S.; Schreck, F.

DOI

[10.1103/PhysRevA.90.022512](https://doi.org/10.1103/PhysRevA.90.022512)

Publication date

2014

Document Version

Final published version

Published in

Physical Review A

[Link to publication](#)

Citation for published version (APA):

Stellmer, S., & Schreck, F. (2014). Reservoir spectroscopy of 5s5p 3P2-5snd 3D1,2,3 transitions in strontium. *Physical Review A*, *90*(2), 22512. <https://doi.org/10.1103/PhysRevA.90.022512>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Reservoir spectroscopy of $5s5p\ ^3P_2$ – $5snd\ ^3D_{1,2,3}$ transitions in strontium

Simon Stellmer^{1,2,3} and Florian Schreck^{1,4}¹*Institut für Quantenoptik und Quanteninformation (IQOQI), Österreichische Akademie der Wissenschaften, 6020 Innsbruck, Austria*²*Institut für Experimentalphysik und Zentrum für Quantenphysik, Universität Innsbruck, 6020 Innsbruck, Austria*³*Vienna Center for Quantum Science and Technology, Atominsttitut, TU Wien, 1020 Vienna, Austria*⁴*Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, The Netherlands*

(Received 3 June 2014; published 20 August 2014)

We perform spectroscopy on the optical dipole transitions $5s5p\ ^3P_2$ – $5snd\ ^3D_{1,2,3}$, $n \in (5,6)$, for all stable isotopes of atomic strontium. We develop a spectroscopy scheme in which atoms in the metastable 3P_2 state are stored in a reservoir before being probed. The method presented here increases the attained precision and accuracy by two orders of magnitude compared to similar experiments performed in a magneto-optical trap or discharge. We show how the state distribution and velocity spread of atoms in the reservoir can be tailored to increase the spectroscopy performance. The absolute transition frequencies are measured with an accuracy of 5 MHz. The isotope shifts are given to within 200 kHz. We calculate the A and Q parameters for the hyperfine structure of the fermionic isotope at the MHz level. Furthermore, we investigate the branching ratios of the 3D_J states into the 3P_J states and discuss immediate implications on schemes of optical pumping and fluorescence detection.

DOI: [10.1103/PhysRevA.90.022512](https://doi.org/10.1103/PhysRevA.90.022512)

PACS number(s): 32.30.–r, 37.10.De

I. INTRODUCTION

The rich electronic structure of elements with two valence electrons, comprising long-lived metastable states and ultra-narrow transitions, is of great interest for precision measurements, as demonstrated by optical clocks [1–5] or gravimeters [6]. Quantum degenerate samples of Yb [7–11] and the alkaline-earth-metals Ca [12,13] and Sr [14–20] have opened new possibilities, such as the study of $SU(N)$ magnetism [21–34], novel schemes to simulate gauge fields [35–40], the engineering of interactions beyond contact interactions [41–44], the creation of driven-dissipative many-body states [45], the simulation of an extra dimension [46], and new ways to perform quantum computation [47–51].

Many of these applications and proposals rely on broad optical transitions originating from the metastable $^3P_{0,1,2}$ triplet states. These transitions are essential for laser cooling of Mg, Ca, and Sr [52–54]. In quantum computation schemes, they can be employed to drive Raman transitions between the metastable states [48]. Transitions originating from the 3P_2 state are particularly important: A suitable 3P_2 – 3D_3 transition could be used for operation of a magneto-optical trap (MOT) [55,56], for fluorescence detection of single atoms, and for absorption imaging. Precise knowledge of these transitions is also required for the calculation of the polarizability of the 3P_2 state, which is important for quantum computation schemes [48,57]. The frequencies and strengths of the transitions originating from the metastable states are needed to calculate the black-body radiation (BBR) shifts of a clock transition, which is particularly important for strontium [57,58]. Atoms in the 3P_2 state can be used to study systems with quadrupolar interactions [42,44]. Furthermore, magnetic Feshbach resonances have been observed between atoms in the 3P_2 and 1S_0 states [59].

So far, spectroscopic data is available for only some of the dipole transitions within the triplet system of strontium; see Ref. [60] for a recent compilation. The $5s5p\ ^3P_0$ – $5s6s\ ^3S_1$ and $5s5p\ ^3P_1$ – $5s6s\ ^3S_1$ transitions, as well

as the isotope shifts and hyperfine splittings involved, have been measured to a precision of a few 100 kHz [61]. The Rydberg series $5snd\ ^3D_{1,2,3}$ for $n > 12$ has been investigated in a discharge [62], and the series $5snd\ ^3D_2$ for $n > 4$ has been studied by multiphoton ionization [63]. The $5s5p\ ^3P_2$ – $5s4d\ ^3D_2$ transition has been investigated using atoms trapped in a MOT [64]; these three experiments reached an accuracy of about 1 GHz and a precision of a few 10 MHz.

In this article, we present a spectroscopy scheme to probe transitions originating from the $5s5p\ ^3P_2$ state in strontium. Spectroscopy is performed for the three stable bosonic isotopes $^{84,86,88}\text{Sr}$ as well as the fermionic isotope ^{87}Sr . While the bosonic isotopes have zero nuclear spin, the fermionic one has a nuclear spin $I = 9/2$, which gives rise to a rich hyperfine structure. The wealth of hyperfine transitions requires a sophisticated spectroscopy scheme. Specifically, we use a magnetic trap as a reservoir to accumulate atoms in the metastable 3P_2 state, hence we name this method reservoir spectroscopy. Atoms in this trap are then probed, and we can populate or deplete certain hyperfine states beforehand to improve measurement performance. This scheme is used to measure the green $5s5p\ ^3P_2$ – $5s5d\ ^3D_{1,2,3}$ transitions at 497 nm and the blue $5s5p\ ^3P_2$ – $5s6d\ ^3D_{1,2,3}$ transitions at 403 nm, all of which have not yet been investigated spectroscopically. We present two different approaches to determine the number of repumped atoms, namely fluorescence detection and recapture into a narrow-line MOT, followed by absorption imaging. Both precision and accuracy are improved by about two orders of magnitude compared to similar spectroscopy measurements in a MOT or discharge. Note that related schemes have recently been used for spectroscopy in Ca [65] and to determine the lifetimes of the $3s3p\ ^3P_2$ state in Mg [66] and the $5s6d\ ^3D_1$ state in Yb [67]. Furthermore, we calculate the branching ratios of all relevant $^3D_J \rightarrow ^3P_J$ decay channels, compare our findings to experimentally determined values, and comment on possible implications for future experiments.

This article is structured as follows: In Sec. II, we give a short review over various repumping transitions in strontium.

We then describe the spectroscopy scheme used in the present work (Sec. III). The spectroscopy data is given in Sec. IV. The calculation of relevant branching ratios is presented together with experimental investigations of the repump efficiency (Sec. V) and the imperfection of quasicycling 3P_2 - 3D_3 transitions (Sec. VI). A short conclusion follows in Sec. VII and a detailed error analysis is given in the Appendix.

II. REPUMPING IN STRONTIUM

The level structure of strontium and the other alkaline-earth-metals naturally suggests the broad 1S_0 - 1P_1 dipole transition for the first laser cooling stage. Cooling on this transition is simplified through the absence of hyperfine structure in the 1S_0 ground state, such that repumping from hyperfine states, very common to alkali MOTs, is not required. There are, however, long-lived electronic states with energies smaller than the 1P_1 state. The performance of a MOT operated on the 1S_0 - 1P_1 transition may therefore be reduced, as an excited atom can decay into one of these states and leave the MOT cycle. This is reminiscent of the cases of Er [68] and Dy [69], but the intermediate states in these cases are short lived.

There are various decay pathways from the 1P_1 state into the long-lived $^3P_{0,1,2}$ states; see Fig. 1. The dominant decay channel is opened up by the $5s4d$ 1D_2 state, which is located below the $5s5p$ 1P_1 state. The branching ratio from the 1P_1 state into the 1D_2 stage is roughly 1 : 50 000 for Ca and Sr, and roughly 1 : 300 for Ba and Ra. The atoms further decay into the $^3P_{1,2}$ metastable triplet states with a branching ratio of 2 : 1. The second decay channel is of the type $^1P_1 \rightarrow ^3D_{1,2} \rightarrow ^3P_{0,1,2}$ and is about two orders of magnitude weaker: for the first step, a branching ratio of 1 : 3.6×10^6 is given in Ref. [70], and we estimate that about one third of the atoms end up in the 3P_0 state. This type of loss channel is the dominant one in 1S_0 - 1P_1 MOTs of Yb [71,72]. The transition probabilities of a third decay channel, the direct decay $^1P_1 \rightarrow ^3P_{0,1,2}$, have not

yet been calculated, but we expect them to be even smaller. We conclude that all three $^3P_{0,1,2}$ states are eventually populated during operation of the MOT, where the relative rates are about < 0.01 : 2 : 1, respectively.

The three metastable 3P_J states have very different lifetimes. In Sr, the 3P_1 state has a comparably short lifetime of 21 μ s, and atoms arriving in this state decay back into the 1S_0 state and into the MOT cycle well before leaving the MOT region. The 3P_0 and 3P_2 states, however, have much longer lifetimes. The lifetime of the 3P_0 state has been calculated to be thousands of years for bosonic atoms, and around 100 s for fermionic atoms [73]. Atoms in this state are lost from the MOT region. The lifetime of the 3P_2 state has been measured to be around 500 s in absence of BBR [74]. The BBR field of the environment, however, can drive transitions $^3P_2 \rightarrow ^3D_J$, from which the atoms decay into the $^3P_{0,1}$ states. This process reduces the lifetime of the 3P_2 state to a few 10 s at room temperature [74,75], which we verify in our experiment. Atoms in the 3P_2 state can naturally be trapped in the quadrupole field of the MOT, provided that they are in a low-field-seeking m_J state and have a kinetic energy smaller than the trap depth [76].

Various transitions can be employed for repumping, and the choice of the transition is guided by both the anticipated repump efficiency and the availability of suitable lasers at the respective wavelengths. A number of transitions have already been used for repumping. An early experiment tried to close the leakage of atoms into the triplet states by pumping them directly from the 1D_2 state into the $5s6p$ 1P_1 state at 717 nm [77]. This approach is inefficient due to a significant branching ratio from the $5s6p$ 1P_1 state into the triplet states. Other experiments use a repump laser at 707 nm to pump atoms from the 3P_2 state via the 3S_1 state into the 3P_1 state, from where they rapidly decay into the 1S_0 ground state [78]. A large fraction of atoms in the intermediate 3S_1 state decays into the 3P_0 state, necessitating a second laser at 679 nm to repump this state as well. This repumping approach rigorously collects atoms from all possible decay paths and facilitates MOT lifetimes of up to ten seconds. A third strategy involves any of the $5snd$ 3D_2 states at 3.01 μ m [64], 497 nm [79], or 403 nm for $n = 4, 5, 6$, respectively. This strategy requires only one laser. Repumping via the $5p^2$ 3P_2 state at 481 nm is also efficient [80], as favorable branching ratios limit the undesired decay into the 3P_0 state to below 0.5%. When repumping only the 3P_2 state via a 3D_2 state, loss through the 3P_0 state persists and limits the lifetime of a continuously repumped MOT to about one second. Loss into the 3P_0 state also originates from leakages in the repump process; see Sec. V.

Atoms in the metastable states can be returned to the ground state either during or after the MOT stage. Generally, continuous repumping allows for a faster accumulation of atoms, as most of the atoms falling into the 3P_2 state appear in nontrapped m_F states and are lost if not quickly repumped. Repumping of the 3P_0 state improves the MOT atom number and lifetime even further. The MOT atom number is determined by the loading rate, which is about 10^9 ^{88}Sr atoms/s for our experiment, and limited by excited-state collisions, where the loss rate coefficient has been determined to $\beta = 4.5 \times 10^{-10}$ cm^3/s [78]. This strategy of continuous repumping is followed in optical clock experiments.




FIG. 1. (Color online) Schematic illustration of the energy levels and transitions in strontium relevant for this work. Atoms in the metastable $5s5p$ 3P_2 state (indicated by the small solid arrow) are repumped using the $5s5p$ 3P_2 - $5snd$ $^3D_{1,2,3}$ transitions around 497 nm ($n = 5$, dotted line) and around 403 nm ($n = 6$, dash-dotted line).

A second scheme accumulates atoms in the metastable reservoir and transfers them into the ground state after the MOT has been extinguished [76,81]. This accumulation strategy is advantageous when large atom numbers are required. The density limitation of the reservoir arises from inelastic two-body collisions, where the loss rate coefficient has been quantified to be $\beta \approx 1 \times 10^{-10} \text{ cm}^3/\text{s}$ [82], but the large volume of the reservoir ($\sim 20 \text{ cm}^3$) ensures that two-body collisions are rare. After saturation, the reservoir will contain roughly 1000 times more atoms than the MOT, where the factor is given by the ratio of the lifetime of the 3P_2 state (roughly 20 s) and the lifetime of the MOT (roughly 20 ms). In our experiment, we can load the reservoir with up to 10^{10} atoms of ^{88}Sr . Accumulation in the reservoir also allows to sequentially load multiple isotopes [79]. Experiments with the fermionic ^{87}Sr isotope will benefit from the storage in the 3P_2 state, as the hyperfine structure impedes both MOT operation and repumping. This strategy is also preferred when repumping of the 3P_0 state is not available [14,15].

The repump transitions discussed above are dipole allowed and have typical linewidths of a few MHz and saturation intensities of a few mW/cm^2 . The isotope shifts are at most 100 MHz. In the case of ^{87}Sr , efficient repumping is complicated by the hyperfine structure of the states involved. We find that all five hyperfine states $F = 5/2$ through $F = 13/2$ of the 3P_2 level are populated during the MOT phase, however at different relative amounts: roughly 80% of the atoms populate the $F = 13/2$ and $F = 11/2$ states. The hyperfine splittings of the 3P_2 , 3S_1 , and 3D_2 states are on the order of a few GHz. In a typical experimental cycle, repumping is performed only on the $F = 11/2 \rightarrow F' = 13/2$ and $F = 13/2 \rightarrow F' = 13/2$ transitions, or the laser is rapidly scanned across all hyperfine transitions.

In our experiment, the repump light illuminates the entire MOT region with an intensity of typically $0.05 I_{\text{sat}}$, corresponding to a few $100 \mu\text{W}$. Here, $I_{\text{sat}} = \pi \hbar c A_{ik} / 3 \lambda^3$ is the saturation intensity of the transition, where A_{ik} is the transition probability between states i and k , and λ is the wavelength of the transition. Repumping of the reservoir typically takes a few milliseconds.

III. SPECTROSCOPY SCHEME

In this section, we will describe how the reservoir of magnetically trapped atoms can be used to improve spectroscopy performance by orders of magnitude compared to other approaches. We will present two complementary detection schemes and characterize our approach.

A. Reservoir spectroscopy

A straightforward and previously used approach to determine the frequency of a repump transition is to monitor the MOT fluorescence while scanning the repump laser frequency [64]. This works well if the repumping increases the MOT atom number N_{MOT} significantly. Typical values of $(N_{\text{rMOT}} - N_{\text{MOT}})/N_{\text{MOT}}$, where N_{rMOT} is the MOT atom number in presence of the repump light, range between 30 for a nonsaturated bosonic MOT and 0.1 for repumping only one hyperfine state of the ^{87}Sr isotope, to below 0.001 for




FIG. 2. (Color online) Burst in MOT fluorescence after repumping of atoms in the 3P_2 reservoir state back into the MOT cycle. The data is approximated by the sum of two exponential functions (solid red line), consisting of a fast growing (τ_{repump} , dotted blue line) and a slowly decreasing contribution (τ_{rMOT} , dashed blue line). The horizontal dashed line indicates the small and stable background signal, caused by residual stray light.

transitions with poor repump efficiency. Considering that the MOT atom number is typically stable only to within 1%, a more sophisticated scheme is needed for weak repump transitions.

A scheme that employs atoms in the reservoir has the potential to drastically improve the signal. The number of atoms in the reservoir, N_{res} , can be 1000 times larger than N_{MOT} , providing a tremendous leverage to spectroscopically resolve transitions with very poor repump efficiency. Illuminating a filled reservoir with repump light in presence of the MOT light induces a burst in fluorescence [65,76] that can exceed the fluorescence of the bare MOT by a factor of 1000; see Fig 2.

After switching the repump light on at time $t_0 = t(0 \text{ s})$, the reservoir atoms are quickly pumped back into the MOT cycle on a timescale τ_{repump} , which is a few milliseconds, leading to a burst in fluorescence. Subsequently the MOT atom number slowly decreases due to the decay of atoms into the 3P_0 state, which is not repumped in our experiments, and possibly other loss processes [78]. This time scale τ_{rMOT} ranges between 10 ms and 1 s, depending on the efficiency of the repump transition used.

We use a simple model of the form

$$N(t) = N_{\text{MOT}} + N_{\text{repump}} \times \left(e^{-\frac{t-t_0}{\tau_{\text{rMOT}}}} - e^{-\frac{t-t_0}{\tau_{\text{repump}}}} \right)$$

to describe the evolution of the MOT atom number for times $t_0 < t < t_0 + \tau_{\text{rMOT}}$. Here, we assume τ_{repump} to be much smaller than τ_{rMOT} . Fitting this approximation to our data allows us to extract all relevant parameters. The number of repumped atoms, N_{repump} , certainly depends on the frequency of the repump light, and ideally approaches N_{res} near the resonance position. We find that $N_{\text{repump}}/N_{\text{MOT}}$ can assume values of up to 1000, clearly showing the advantage of repump spectroscopy.

The entire experimental sequence used for reservoir spectroscopy is as follows. Atoms emerging from an effusive source are Zeeman slowed and cooled in a MOT using the broad $^1S_0 \rightarrow ^1P_1$ transition at 461 nm. The lifetime of the MOT

is about 20 ms and depends on the intensity of the MOT beams, which sums to about $0.1 I_{\text{sat}}$. Atoms accumulate in the low-field-seeking $m_J = +1$ and $m_J = +2$ components of the long-lived 3P_2 state, and we operate the MOT until the reservoir is filled with about 10^7 atoms. This takes between 50 ms and 10 s, depending on the natural abundance of the respective isotope. Afterwards, we turn on the repump light, which illuminates the entire reservoir volume at an intensity of $10^{-3} I_{\text{sat}}$ and returns the atoms back into the ground state within a few milliseconds. While the $^3P_2 \rightarrow ^3D_2$ transition is commonly used for repumping, also the $^3P_2 \rightarrow ^3D_{1,3}$ transitions return atoms into the ground state, however with lower efficiency. Further details of the apparatus used for this experiment can be found in Ref. [20].

B. Fluorescence detection

The MOT fluorescence is measured by imaging the MOT onto a large-area photo diode with a magnification of 0.5, capturing 0.6% of the total fluorescence. For low densities of the MOT, the fluorescence is roughly proportional to the atom number. We vary the frequency of the repump light in consecutive experimental runs and determine N_{repump} for each run. These values are then fitted by a Lorentzian line shape and allow us to determine the resonance position to at best 300 kHz, where typical uncertainties are about 1 MHz. This scheme of fluorescence detection is very robust and has previously been used to determine the lifetime of the 3P_2 state [74]. Note that, for small values of N_{repump} , it is advantageous to remove the background signal N_{MOT} by removing all MOT atoms prior to the flash of repump light. This can easily be done by blocking the atomic beam or extinguishing the Zeeman slower light prior to repumping.

C. Absorption imaging

We developed a more sophisticated detection method to overcome certain drawbacks of fluorescence detection. The amplitude of the fluorescence signal depends directly on the intensity and polarization of the MOT beams, which might fluctuate and add noise to the signal. In addition, electronic noise of the photo detector, as well as residual stray light on the detector, might impede the detection of very weak transitions. These issues eventually limit the performance and led us to develop a scheme based on absorption imaging. After loading the reservoir with 10^7 atoms, we extinguish the MOT light. A mechanical shutter blocks the atomic beam, and a 1-s wait ensures that only atoms in the magnetic trap remain in the probe volume. Upon a short repump flash, the atoms are captured in a narrow-line MOT operated on the $^1S_0 \rightarrow ^3P_1$ intercombination line [54,83]. The MOT light is initially frequency broadened to a few MHz to increase the capture efficiency. Within 500 ms, the frequency broadening is removed, and the intensity is lowered to $1 I_{\text{sat}}$. The atoms are cooled to a temperature below $1 \mu\text{K}$ and compressed to a dense cloud with a size of order $100 \mu\text{m}$. This stage increases the phase-space density by eight orders of magnitude compared to the broad-linewidth $^1S_0 \rightarrow ^1P_1$ MOT.

Recapture into the narrow-line MOT rather than the broad-line MOT has numerous advantages: the narrow-line MOT

is perfectly cycling, excited-state collisions are rare, and the small size and low temperature are ideal for subsequent absorption imaging. Imaging is performed on the $^1S_0 \rightarrow ^1P_1$ transition and allows for a background-free determination of the atom number. This technique allows us to reduce the uncertainty in the fit of the Lorentzian profile to below 10 kHz for typical settings.

D. Characterization

We will now compare the reservoir spectroscopy scheme with the approach of monitoring the increase of the broad-line MOT fluorescence upon repumping [64]. We will also introduce two methods to increase the signal strength. The detection scheme considered in the following is the absorption imaging scheme, but the conclusions also hold for fluorescence detection after recapture into the broad MOT.

Reservoir spectroscopy is advantageous to direct spectroscopy in the MOT in three ways. First, our measurement is intrinsically background-free. Far away from the repump resonance, no atoms are captured in the narrow-line MOT. The small background noise on the absorption images stems from photon shot noise, which leads to noise in the atom number corresponding to at most 1000 atoms. This background noise can be reduced by averaging over multiple images. On the other hand, spectroscopy in a MOT will always incorporate the background of the MOT itself. The superior signal strength is shown in Fig. 3(a). Depending on the MOT loading rate and repump intensity used, the fluorescence of the MOT increases by a factor of up to 30, exhibiting a substantially broadened linewidth. The atom number increase for the fermionic isotope is at most a few 10% on resonance due to the involved hyperfine structure. The reservoir spectroscopy described here can yield a signal without intrinsic background and an amplitude more than three orders of magnitude larger than the background noise. The inset of Fig. 3(a) shows typical fluorescence spectra of a ^{88}Sr MOT at repump intensities of about 1, 3, and $10 \times 10^{-3} I_{\text{sat}}$, starting from below (solid black lines, normalized to the fluorescence of a MOT without repumping). The dashed red curve shows the corresponding measurement using reservoir spectroscopy, taken at an intensity of $5 \times 10^{-5} I_{\text{sat}}$. The amplitude of 5×10^6 atoms is more than three orders of magnitude above the background noise, and the width is smaller by a factor of four compared to the fluorescence curve at highest repump intensity.

The second advantage of reservoir spectroscopy is the potential to reduce the Doppler broadening: the quadrupole field, which confines the metastable atoms, acts as a velocity filter, as it traps only atoms below a certain kinetic energy. We typically set the axial gradient of the quadrupole field to 55 G/cm during the MOT phase. Atoms may roam the magnetic trap up to an axial radius of 11 mm, given by geometric constraints of our vacuum cell. For bosonic atoms in the 3P_2 , $m_J = +1$ state, the trap depth is $U = 4$ mK, which is slightly larger than the Doppler temperature of $T_D = 0.72$ mK. The maximally trappable velocity generally depends on the m_F state, and, for the fermionic isotope, also on the hyperfine state due to differing g_F factors. A deeper trap will thus capture more atoms also at higher velocity classes, while the confinement can be reduced to deliberately select only the low-velocity




FIG. 3. (Color online) Reservoir spectroscopy scheme. (a) The reservoir spectroscopy signal is background free and rises many orders of magnitude above the background noise. By contrast, the fluorescence signal of a MOT (inset) is increased by only a small factor on resonance. A Lorentzian profile is fit to each data set, and the fluorescence curves are taken at a repump laser intensity of $I = 1, 3,$ and $10 \times 10^{-3} I_{\text{sat}}$, from below. (b) Dependence of the width of the transition on the depth of the magnetic trap. (c) Width (black squares) and atom number (red circles) in dependence of repump light intensity. The repump pulse duration is 10 ms. (d) Accumulation of a certain hyperfine state by optical pumping increases the signal. (e) Depletion of a certain hyperfine state allows for a separation of otherwise overlapping hyperfine transitions. Measurements (a) through (c) are taken on the $5s5p^3P_2-5s5d^3D_2$ transition of ^{88}Sr , measurements (d) and (e) employ ^{87}Sr ; see the text for details.

atoms to reduce the Doppler broadening, as shown in Fig. 3(b). In this measurement, the quadrupole field is ramped down after the MOT phase to adiabatically cool the sample and to allow high-energy atoms to escape.

As a third advantage, we can selectively populate the reservoir with the specific states of interest. This is important for the fermionic isotope with its large number of hyperfine levels. In particular, atoms can be pumped into the state of interest during or after the MOT phase. This is illustrated in Fig. 3(d): During the ^{87}Sr MOT, atoms in all five 3P_2 hyperfine states are accumulated in the reservoir, however at different relative amounts. The $F = 7/2$ state has the smallest g_F factor, therefore only very few atoms are trapped. In addition, π transitions into the 3D_2 states tend to be weaker than σ transitions, making the $F = 7/2 \rightarrow F' = 7/2$ transition one of the weakest among all 13 $^3P_2-^3D_2$ transitions. Constantly pumping atoms from the $F = 9/2$ into the $F = 7/2$ state by standard optical pumping can increase the population of this state significantly.

Similarly, atoms from undesired states can be removed from the reservoir. This is shown in Fig. 3(e): The $F = 11/2 \rightarrow F' = 13/2$ and $F = 9/2 \rightarrow F' = 7/2$ hyperfine transitions of the $5s5p^3P_2-5s5d^3D_2$ line in ^{87}Sr overlap entirely. The initial population of the $F = 11/2$ state is about ten times that of the $F = 9/2$ state, therefore the $F = 11/2 \rightarrow F' = 13/2$ transition completely covers the $F = 9/2 \rightarrow F' = 7/2$ transition. To selectively interrogate only the $F = 9/2 \rightarrow F' = 7/2$ transition, we remove all $F = 11/2$ -state atoms during the MOT phase by optical pumping into the $F' = 11/2$ state. As an additional benefit, this also increases the $F = 9/2$ population. We first verify on the $F = 11/2 \rightarrow F' = 11/2$ (or alternatively the $F = 11/2 \rightarrow F' = 13/2$) transition that all $F = 11/2$ atoms have been removed, and then selectively probe the $F = 9/2$ state and determine the frequency difference between the $F = 11/2 \rightarrow F' = 13/2$ and $F = 9/2 \rightarrow F' = 7/2$ transitions to be 800 (200) kHz. This would not be possible by spectroscopy on a continuously operated MOT.

To optimize our spectroscopy approach, we measure the dependence of the transition linewidth and signal amplitude on the repump intensity. A larger repump intensity will yield a larger amplitude of the signal, but also a larger width; see Fig. 3(c). We find an intensity region around $5 \times 10^{-4} I_{\text{sat}}$ where the width is not broadened by the intensity, but the amplitude has almost saturated. In this region, width and amplitude are almost independent of the repump time for times greater than about 10 ms. A distortion of the line shape can be observed only for intensities lower than $2 \times 10^{-4} I_{\text{sat}}$. Even for the highest intensities used in this study, we do not observe any intensity-dependent line shifts.

IV. SPECTROSCOPIC DATA

A. $5s5p\ ^3P_2-5s5d\ ^3D_{1,2,3}$ spectroscopy

We will now present spectroscopic data of the green $5s5p\ ^3P_2-5s5d\ ^3D_{1,2,3}$ transition around 497 nm, using absorption imaging as the detection method. The loading time of the reservoir is adjusted to compensate for the different natural abundances of the bosonic isotopes, and the intensity of the repump light is $5 \times 10^{-4} I_{\text{sat}}$. A typical set of measurements is shown in Fig. 4. Figure 4(a) shows the spectra of the three bosonic isotopes, where the isotope shifts are roughly 20 MHz per mass unit.

Figure 4(b) shows measurements on the fermionic isotope, taken with a typical intensity of $1 \times 10^{-3} I_{\text{sat}}$. Note that the spectrum spans many GHz in frequency, and that the amplitudes of the strongest lines rise more than two orders of magnitude above the background. Note also the two overlapping lines around 603 285 750 MHz, which were discussed in the previous section and in Fig. 3(e).




FIG. 4. (Color online) Reservoir spectroscopy data obtained with the absorption imaging detection method, shown here for the $5s5p\ ^3P_2-5s5d\ ^3D_2$ transition. (a) Transitions of the bosonic isotopes ^{84}Sr (black, left peak), ^{86}Sr (red, center), and ^{88}Sr (blue, right). (b) A scan across the transitions of the fermionic ^{87}Sr isotope. The separate measurement of the two overlapping lines around 5.75 GHz is discussed in Sec. III D.

A compilation of the transition frequencies can be found in Table I. As discussed in the Appendix, the absolute frequencies have an error of at most 5 MHz. Relative frequency differences, such as the isotope shifts, have an uncertainty of 200 kHz. The measurement of the hyperfine structure of the fermionic isotope is more involved, leading to a larger error in the magnetic dipole and electric quadrupole interaction constants A and Q . Compared to spectroscopy in a MOT or discharge, the spectroscopy scheme used here can improve both the precision and accuracy by at least two orders of magnitude.

B. $5s5p\ ^3P_2-5s6d\ ^3D_{1,2,3}$ spectroscopy

We will now introduce a potential technological simplification to strontium experiments that employ the strategy of accumulation and subsequent repumping. Instead of using the 3S_1 state, which requires two lasers for repumping, such experiments might choose a state that requires only one repump laser, e.g., one of the $5snd\ ^3D_2$ states. Experiments so far used the state with $n = 4$, which requires a technically involved laser source at $3.01\ \mu\text{m}$, or the state with $n = 5$ at 497 nm, which requires a frequency-doubled diode laser system. As a potential simplification, we explore the use of blue light at 403 nm, employing the state with $n = 6$ for repumping; see Fig. 1. Diode lasers at this wavelength are readily available thanks to Blu-ray technology.

As a first step to characterize this approach, we perform reservoir spectroscopy on this transition, analogous to the measurements described above. These measurements are plagued by frequency noise of the laser, which is not locked to a stable reference. This frequency noise occurs on the timescale of a few hundred ms, with an amplitude of a few MHz. We expect this noise to be the dominant error source and therefore match our detection scheme to the less stringent requirements in the precision. Specifically, we do not apply the experimentally involved capture into the narrow-line MOT and subsequent absorption imaging, but recapture into the broad-line MOT and use the peak in MOT fluorescence as a measure of the number of repumped atoms; see Sec. III B. A compilation of the resonance positions is given in Table II.

The relevant parameters of the three $5s5p\ ^3P_2-5snd\ ^3D_J$, $n \in (4,5,6)$, repump transitions are listed in Table III.

C. Hyperfine structure

In contrast to the bosonic isotopes with zero nuclear spin, the fermionic strontium isotope has $I = 9/2$ and shows an involved hyperfine structure. For the $^3P_2-^3D_J$ transitions of ^{87}Sr , we can identify each of the resonances by their relative positions, and subsequently calculate the interaction constants A and Q of the $^3D_{1,2}$ states using

$$\Delta E_{\text{hfs}}/h = \frac{A}{2}K + \frac{Q}{2} \frac{\frac{3}{4}K(K+1) - I(I+1)J(J+1)}{I(2I-1)J(2J-1)}.$$

Here, $K = F(F+1) - I(I+1) - J(J+1)$, where $I = 9/2$ is the nuclear spin, J is the total angular momentum, and F denotes the hyperfine state, given by $\vec{F} = \vec{I} + \vec{J}$. The hyperfine structure of the 3P_2 state is known to the kHz level [84] [$A = -212.765(1)$ and $Q = 67.215(15)$], and we

TABLE I. Transition frequencies, given in MHz, of the three $5s5p\ ^3P_2 - 5s5d\ ^3D_{1,2,3}$ green transitions around 497 nm. Dashes indicate transitions that are allowed by selection rules, but were not observed.

	$5s5p\ ^3P_2 - 5s5d\ ^3D_1$	$5s5p\ ^3P_2 - 5s5d\ ^3D_2$	$5s5p\ ^3P_2 - 5s5d\ ^3D_3$
^{84}Sr	602 833 830.9 (5.0)	603 285 766.0 (5.0)	603 976 415.0 (5.0)
^{86}Sr	602 833 876.1 (5.0)	603 285 811.3 (5.0)	603 976 460.0 (5.0)
^{88}Sr	602 833 924.9 (5.0)	603 285 858.8 (5.0)	603 976 506.6 (5.0)
^{87}Sr $5/2 \rightarrow 3/2$			603 976 697.5 (5.0)
^{87}Sr $5/2 \rightarrow 5/2$		603 284 256.9 (5.0)	-
^{87}Sr $5/2 \rightarrow 7/2$	602 830 264.7 (5.0)	603 284 005.0 (5.0)	-
^{87}Sr $7/2 \rightarrow 5/2$		603 285 034.2 (5.0)	-
^{87}Sr $7/2 \rightarrow 7/2$	602 831 041.6 (5.0)	603 284 772.0 (5.0)	-
^{87}Sr $7/2 \rightarrow 9/2$	-	603 284 459.8 (5.0)	-
^{87}Sr $9/2 \rightarrow 7/2$	602 832 017.1 (5.0)	603 285 748.6 (5.0)	-
^{87}Sr $9/2 \rightarrow 9/2$	602 833 045.1 (5.0)	603 285 436.5 (5.0)	-
^{87}Sr $9/2 \rightarrow 11/2$	602 834 291.5 (5.0)	603 285 047.3 (5.0)	-
^{87}Sr $11/2 \rightarrow 9/2$	602 834 207.3 (5.0)	603 286 608.8 (5.0)	-
^{87}Sr $11/2 \rightarrow 11/2$	602 835 459.1 (5.0)	603 286 211.1 (5.0)	-
^{87}Sr $11/2 \rightarrow 13/2$		603 285 749.4 (5.0)	-
^{87}Sr $13/2 \rightarrow 11/2$	602 836 808.7 (5.0)	603 287 563.6 (5.0)	603 978 453 (10)
^{87}Sr $13/2 \rightarrow 13/2$		603 287 099.1 (5.0)	603 977 428 (10)
^{87}Sr $13/2 \rightarrow 15/2$			603 976 258.8 (5.0)

fix these values for our calculation. The hyperfine structure of the $5s4d\ ^3D_J$ states has been determined independently [85]. A qualitative comparison with our values assures that our designation of hyperfine states is correct.

The $^3P_2 - ^3D_3$ transitions have the lowest repumping efficiency, as the probability of atoms in the 3D_3 state to decay into the 3P_1 state is relatively low; see Secs. V and VI. Instead, atoms might decay into another 3P_2 hyperfine state, where they are dark for the repump light. For the green $5s5p\ ^3P_2 - 5s5d\ ^3D_3$ transition, only four out of 15 lines could be detected, despite a careful search over more than 8 GHz. We speculate that the two strongest lines are the cycling $F = 5/2 \rightarrow F' = 3/2$

and $F = 13/2 \rightarrow F' = 15/2$ transitions, which would imply that the hyperfine structure of the 3D_3 state is inverted with an interaction constant $A \approx -150$. Simultaneous repumping from various hyperfine states would be necessary to obtain a complete spectrum of the $^3P_2 - ^3D_3$ hyperfine transitions.

The signal strength on the blue $5s5p\ ^3P_2 - 5s6d\ ^3D_3$ transition is much larger compared to the green transition, allowing us to detect significantly more lines. The reason is an increased branching ratio into the 3P_1 state; see Sec. V. The eight resonances have very different amplitudes, which helps us to identify them through our knowledge the atoms' initial distribution over the 3P_2 hyperfine states.

TABLE II. Transition frequencies, given in MHz, of the three $5s5p\ ^3P_2 - 5s6d\ ^3D_{1,2,3}$ blue transitions around 403 nm. Dashes indicate transitions that are allowed by selection rules, but were not observed.

	$5s5p\ ^3P_2 - 5s5d\ ^3D_1$	$5s5p\ ^3P_2 - 5s5d\ ^3D_2$	$5s5p\ ^3P_2 - 5s5d\ ^3D_3$
^{84}Sr	743 105 100 (5)	743 252 879 (5)	743 621 723 (5)
^{86}Sr	743 105 147 (5)	743 252 911 (5)	743 621 767 (5)
^{88}Sr	743 105 177 (5)	743 252 960 (5)	743 621 808 (5)
^{87}Sr $5/2 \rightarrow 3/2$			743 622 062 (5)
^{87}Sr $5/2 \rightarrow 5/2$		-	743 621 642 (5)
^{87}Sr $5/2 \rightarrow 7/2$	743 101 462 (5)	-	-
^{87}Sr $7/2 \rightarrow 5/2$		-	-
^{87}Sr $7/2 \rightarrow 7/2$	743 102 252 (5)	743 252 566 (5)	-
^{87}Sr $7/2 \rightarrow 9/2$	-	-	-
^{87}Sr $9/2 \rightarrow 7/2$	743 103 222 (5)	743 253 506 (5)	-
^{87}Sr $9/2 \rightarrow 9/2$	743 104 293 (5)	743 252 802 (5)	-
^{87}Sr $9/2 \rightarrow 11/2$	743 105 617 (5)	743 251 903 (5)	-
^{87}Sr $11/2 \rightarrow 9/2$	743 105 456 (5)	743 253 983 (5)	743 623 285 (5)
^{87}Sr $11/2 \rightarrow 11/2$	743 106 788 (5)	743 253 090 (5)	743 622 387 (5)
^{87}Sr $11/2 \rightarrow 13/2$		743 252 000 (5)	743 621 332 (5)
^{87}Sr $13/2 \rightarrow 11/2$	743 108 137 (5)	743 254 431 (5)	743 623 722 (5)
^{87}Sr $13/2 \rightarrow 13/2$		743 253 352 (5)	743 622 677 (5)
^{87}Sr $13/2 \rightarrow 15/2$			743 621 475 (5)

TABLE III. Compilation of relevant spectroscopic data. For each transition, we state the wavelength λ and frequency ν of the most abundant ^{88}Sr isotope, as well as the isotope shifts $\Delta_{84}^{88} = \nu(^{88}\text{Sr}) - \nu(^{84}\text{Sr})$, and $\Delta_{86}^{88} = \nu(^{88}\text{Sr}) - \nu(^{86}\text{Sr})$, and $\Delta_{87}^{88} = \nu(^{88}\text{Sr}) - \nu(^{87}\text{Sr})$. The interaction constants A and Q characterize the hyperfine structure of the 3D_J manifolds in the fermionic isotope ^{87}Sr . Values labeled by ^a are based on a speculative assignment of the transitions; see Sec. IV C.

transition	λ [nm]	$\nu(^{88}\text{Sr})$ [MHz]	Δ_{84}^{88} [MHz]	Δ_{86}^{88} [MHz]	Δ_{87}^{88} [MHz]	A [MHz]	Q [MHz]
$5s5p\ ^3P_2-5s4d\ ^3D_1$	3067.0 [60]	97 747 180 (150) [60]				139.9 (2) [85]	15 (2) [85]
$5s5p\ ^3P_2-5s4d\ ^3D_2$	3011.8 [64]	99 537 870 (75) [64]	600 (50) [64]	270 (40) [64]	110 (30) [64]	-78.08 (5) [85]	18 (1) [85]
$5s5p\ ^3P_2-5s4d\ ^3D_3$	2923.4 [60]	102 550 490 (170) [60]				-115.3 (2) [85]	51 (9) [85]
$5s5p\ ^3P_2-5s5d\ ^3D_1$	497.30	602 833 924.9 (5.0)	94.0 (2)	48.8 (2)	38 (2)	227.3 (7)	0 (10)
$5s5p\ ^3P_2-5s5d\ ^3D_2$	496.93	603 285 858.8 (5.0)	92.8 (2)	47.5 (2)	17 (2)	-71.5 (5)	0 (30)
$5s5p\ ^3P_2-5s5d\ ^3D_3$	496.36	603 976 506.6 (5.0)	91.6 (2)	46.6 (2)	27 (2) ^a	-156.9 (3) ^a	0 (30) ^a
$5s5p\ ^3P_2-5s6d\ ^3D_1$	403.43	743 105 177 (5)	77 (10)	30 (10)	18 (2)	239.7 (5)	5 (20)
$5s5p\ ^3P_2-5s6d\ ^3D_2$	403.35	743 252 960 (5)	81 (2)	49 (2)	20 (5)	-163.2 (9)	30 (20)
$5s5p\ ^3P_2-5s6d\ ^3D_3$	403.15	743 621 808 (5)	85 (10)	41 (10)	53 (5)	-161.8 (5)	20 (20)

V. REPUMP EFFICIENCY

The efficiency of optical pumping can be reduced by decay from the excited state into undesired states. For the specific case studied here, atoms in the 3P_2 state are to be pumped into the 3P_1 state. The undesired state is the 3P_0 state, which is very long lived, not trapped in the magnetic trap, and dark to repump photons through the fine-structure splitting of about 600 cm^{-1} . Clearly, the probability of decay into the 3P_0 state depends on the excited state used for repumping, and might assume any value between zero and close to unity.

The $5s6s\ ^3S_1$ state can decay into the 3P_0 state: the branching ratios into the $^3P_{0,1,2}$ states are $1/9$, $3/9$, and $5/9$, respectively, such that 25% of all repumped atoms end up in the 3P_0 state. To close this leak, strontium experiments operate another repumper at 679 nm to excite these atoms again into the 3S_1 state [78].

The $5s4d\ ^3D_2$ state can only decay into the $5s5p\ ^3P_{1,2}$ states due to selection rules. Decay into the 3P_0 state is possible only through higher-order processes, which will be neglected here. When going up the ladder of 3D_2 states, however, more and more additional states appear into which the 3D_2 state can decay; see Fig. 5. These might have nonzero branching ratios into the 3P_0 state, possibly through a cascade of various other states.

We will now quantify the branching ratio of higher-lying $5snd\ ^3D_{1,2,3}$ states into the three $5s5p\ ^3P_{0,1,2}$ states. We perform a detailed calculation to trace all possible decay channels for the cases of $n \in (5,6)$, as illustrated in Fig. 5. We use the transition probabilities from Ref. [86]. Again, we will consider only dipole-allowed ($E1$) transitions. The branching ratios into the different fine structure states are obtained using the standard Wigner $6-j$ symbols. We will consider only bosonic isotopes of zero nuclear spin here, ignoring the hyperfine structure of the fermionic isotope. Similarly, we have no knowledge about the m_J -state distribution of atoms in the $5snd\ ^3D_{1,2,3}$ states, and therefore neglect a possible m_J -state dependence of the branching ratios. The fine structure splitting of the states involved is a few 100 cm^{-1} for the $5s5p\ ^3P_{0,1,2}$ states and much smaller for all other states. We take these splittings into account by multiplying correction factors to the branching ratios [87]. The correction factors

favor decay into fine structure states of lower energy (lower J quantum number) by typically a few percent. We carefully compare our results with the transition probabilities tabulated in Ref. [60] and find an overall agreement within a few percent.

The results of these calculations are presented in Table IV. We find that repumping on the green $5s5p\ ^3P_2-5s5d\ ^3D_2$ transition induces a decay into the $5s5p\ ^3P_0$ state of about 0.05%. The multitude of higher-lying states increases this value to about 1.8% for the blue $5s5p\ ^3P_2-5s6d\ ^3D_2$ transition. These values, albeit small, are significant for continuous repumping in the MOT.

We will now follow three different experimental approaches to quantify the relative branching ratio from the $5s5d\ ^3D_2$ state into the $5s5p\ ^3P_{0,1}$ states. These approaches boil down to a precise measurement of τ_{rMOT} , N_{rMOT} , and N_{repump} ; see Fig. 2.




FIG. 5. (Color online) Transition probabilities A_{ik} , given in units of 10^6 s^{-1} , between triplet states of Sr relevant for this work. Values are taken from Ref. [86]. The fine structure splitting, indicated here only for the lowest 3P_J states, is otherwise smaller than the thickness of the horizontal lines. Decay paths with negligible contribution are omitted.

TABLE IV. Branching ratios of relevant 3D_2 states into the $5s5p^3P_{0,1,2}$ states, taking into account all possible decay paths. The precision is given by the uncertainty of the transition probabilities taken from Ref. [86]; see the text for details. The right part of the table shows the repump efficiency of each of the six transitions, as calculated from the branching ratios and experimentally determined from the lifetime τ_{MOT} , the number of atoms in the repumped MOT N_{MOT} , and the number of atoms repumped from the reservoir N_{repump} . The values of the last column were normalized to their largest value.

state used for repumping	calculated branching ratios			repump efficiency			
	$5s5p^3P_0$	$5s5p^3P_1$	$5s5p^3P_2$	calculated	via τ_{MOT}	via N_{MOT}	via N_{repump}
$5s5d^3D_1$	56.24%	41.11%	2.644%	42.22%			
$5s5d^3D_2$	0.035%	76.00%	23.96%	99.95%	96.9(5)%	97(1)%	100%
$5s5d^3D_3$	0.012%	0.057%	99.93%	82.61%			
$5s6d^3D_1$	53.20%	41.75%	5.071%	43.98%	40(5)%	40(2)%	
$5s6d^3D_2$	1.363%	72.80%	25.83%	98.16%	95.7(9)%	92(1)%	98.1(5)%
$5s6d^3D_3$	0.467%	2.276%	97.25%	82.97%	84(2)%	79(1)%	82.6(5)%

At first, we measure the lifetime of the bosonic ^{88}Sr MOT, with and without the repumper applied. The MOT lifetime is strongly density-dependent, therefore measurements are taken with a rather dilute MOT far from saturation. We measure a lifetime of 30 (5) ms without repumper. This lifetime increases to $\tau_{\text{MOT}} = 950$ (50) ms with the green repumper applied, and to 700 (50) ms with the blue repumper. The repumpers close the leak into the 3P_2 state, and we assume that residual decay into the 3P_0 state limits the lifetime of the repumped MOTs. The vacuum lifetime, measured with atoms held in an optical dipole trap, is about two minutes. The factor of 32 (5) in lifetime increase for the green transition [factor 23 (5) for the blue transition] directly shows that about 3% of the repump cycles using green light, and about 4% of the repump cycles using blue light lead the atom into the dark 3P_0 state. Corresponding values of further transitions are given in the column labeled “via τ_{MOT} ” in Table IV.

In a second approach, we measure the steady-state fluorescence of a weak MOT at low MOT light intensity, corresponding to N_{MOT} . When applying the green repumper, we measure a fluorescence increase by a factor of 30 (2) [88], and a factor of 12 (1) for the blue repumper. These numbers can be translated into a probability of atoms to decay from the 3D_2 state into the 3P_0 state. This probability is about 3% for the green repumper and about 8% for the blue repumper; see the column labeled “via N_{MOT} ” in Table IV.

All of these experimentally determined values are significantly larger than the ones calculated above. One possible explanation is a different decay channel not considered so far: the pathways $^1P_1 \rightarrow ^3P_0$ and $^1P_1 \rightarrow ^3D_1 \rightarrow ^3P_0$. The estimated branching ratio from the 1P_1 state into the $5s4d^3D_J$ states is 2.8×10^{-7} [70], about two orders of magnitude smaller than the decay into the $5s4d^1D_2$ state [86]. We estimate that about one third of the atoms following this pathway end up in the dark 3P_0 state. The probability of the direct loss channel $^1P_1 \rightarrow ^3P_0$ is not known. Given the uncertainties in the estimated transition probability values, we find that this decay mechanism might very well explain our observed values of τ_{MOT} and N_{MOT} . This hypothesis is strengthened by the fact that experiments employing repumping of both the 3P_0 and 3P_2 states have reported MOT lifetimes of up to 10 seconds [89]. The branching ratio from the 1P_1 state via all pathways into the 3P_0 state would then be $1 : 1.7(3) \times 10^6$.

We perform a third experiment that does not include the direct decay of the 1P_1 state into the metastable states. Here, we directly measure the number N_{repump} of atoms repumped from the reservoir. We assume that repumping on the green $5s5p^3P_2$ - $5s5d^3D_2$ transition returns all atoms into the ground state, and we use this value to normalize the performance of two other transitions; see last column of Table IV. We find the number of atoms repumped on the blue $5s5p^3P_2$ - $5s6p^3D_2$ transition to be reduced by 1.9% compared to repumping on the green $5s5p^3P_2$ - $5s5p^3D_2$ transition, in very good agreement with our calculation.

Two conclusions can be drawn from these measurements: (i) Using the $5s6d^3D_2$ state instead of the $5s5d^3D_2$ is 1.9% less efficient for a single repump cycle, which seems tolerable when repumping previously accumulated atoms from the reservoir. This is very promising, as it offers a substantial simplification of the required laser system. (ii) The direct decay from the 1P_1 state into the metastable states seems to be significantly larger than expected, and sophisticated calculations might help to resolve this issue. A continuously repumped MOT will therefore greatly benefit in atom number from repumping of the 3P_0 state. Usage of the 3S_1 state here might require the least technological effort.

VI. IS THE $5s5p^3P_2$ - $5s5d^3D_3$ TRANSITION CYCLING?

The $5s5p^3P_2$ - $5s5d^3D_3$ transitions are unique in that they are the only (nearly) cycling transitions originating from the 3P_J manifold. There is an infinite ladder of $5snd^3D_3$ states reachable from the 3P_2 state, starting from $n = 4$.

These transitions have been proposed as MOT transitions to cool atoms in the metastable 3P_2 state. Doppler cooling on similar transitions has been performed in calcium [55] and neon [56]. The $5s5p^3P_2$ - $5s4d^3D_3$ transition at $2.92 \mu\text{m}$ is predestined for operation of a MOT due to a combination of a large wavelength and a small linewidth of only 50 kHz, leading to both a low Doppler and a low recoil temperature. In contrast to all higher-lying 3D_3 states, no other states appear in between the two MOT states to spoil cycling operation, and the absorption of two photons does not ionize the atom. Apart from the fact that lasers at this frequency and linewidth are difficult to operate, efficient MOT operation might however be hindered by inelastic losses [82].

The 3P_2 - 3D_3 transitions could also be used for fluorescence and absorption imaging. A recent publication on quantum computation with fermionic ^{87}Sr [48] suggests the use of a 3P_2 - 3D_3 , $|F = 13/2\rangle \rightarrow |F' = 15/2\rangle$ transition for fluorescence readout of atoms in the metastable states. This application would benefit from a short-wavelength transition, which brings about good imaging resolution and large detection efficiency. The $5s5p\ ^3P_2$ - $5s5d\ ^3D_3$ transition at 497 nm seems a promising candidate. Analogous to the discussion in Sec. V, this transition might not be perfectly cycling due to decay paths that lead into the $^3P_{0,1}$ states. These additional channels gain importance when climbing up the ladder of $5snd\ ^3D_3$ states.

In the following, we will investigate the probability of atoms in the 3D_3 state to decay into the $5s5p\ ^3P_{0,1}$ states. A summation of all possible decay paths for two 3D_3 states can be found in Table IV. The branching ratio into the 3P_1 state is 0.057% for the $5s5d\ ^3D_3$ state and 2.28% for the $5s6d\ ^3D_3$ state. Again, we neglected higher-order transitions. These numbers already suggest that fluorescence detection of individual atoms will require optical pumping of the $^3P_{0,1}$ states into the 3P_2 state, as on the order of 10^5 photons need to be cycled for a sufficiently large signal. For the fermionic isotope, the optical pumping scheme will be more involved due to the hyperfine structure and multitude of m_F states.

To verify our calculation, we perform an experiment that allows us to compare the decay rates from the $5s5d\ ^3D_2$ and 3D_3 states into the $5s5p\ ^3P_1$ state. The $5s5p\ ^3P_2$ - $5s5d\ ^3D_3$ transition is certainly not completely cycling, as can be inferred from the spectroscopy described in Sec. IV: obviously, some atoms are transferred into the 3P_1 state and further into the 1S_0 ground state; see Fig. 5. The measurements presented in the following are performed with the bosonic ^{88}Sr isotope. About 10^7 atoms are loaded into the 3P_2 reservoir state, repumped, and recaptured in the narrow-line MOT. As a reference, we first measure the number of atoms repumped on the $5s5p\ ^3P_2$ - $5s5d\ ^3D_2$ transition in dependence of repump time. For low intensities and short repumping times, there is a linear dependence of repumped atoms on repump time until the atom number saturates; see the inset of Fig. 6. We perform this measurement for a wide range of repump light intensities and, for each measurement series, extract the initial repump rate.

We then perform the same experiment on the 3P_2 - 3D_3 transition; see Fig. 6. We find that repumping via the 3D_3 state instead of the 3D_2 state returns 30 (1) times less atoms into the ground state. The saturation intensities are 2.2 mW/cm² for the 3P_2 - 3D_2 and 10.4 mW/cm² for the 3P_2 - 3D_3 transition. From the calculated branching ratios to the 3P_1 state (see Table IV), we expect a ratio of $76/0.057 \approx 1300$ between the repumping rates of the two transitions examined, constituting a clear discrepancy.

The enhanced decay of the 3D_3 state into the 3P_1 state could be explained by the presence of the black-body radiation (BBR) field. Many of the transitions between the higher-lying states have wavelengths comparable to the maximum of the BBR at room temperature, which is around 10 μm . Such transitions can be driven by the BBR field. As a coincidence, the energy difference of 1049 cm⁻¹ between the $5s6p\ ^3P_2$ and $5s5d\ ^3D_3$ states corresponds exactly to the maximum of the room-temperature BBR spectrum. It has been observed in




FIG. 6. (Color online) Repump rate in dependence of light intensity. The number of atoms repumped into the 1S_0 ground state is proportional to the repump time before reaching saturation (inset, taken on the green $5s5p\ ^3P_2$ - $5s5d\ ^3D_3$ transition at $I = 1 \times 10^{-2} I_{\text{sat}}$). This rate is measured for a large range of intensities on both of the green 3P_2 - 3D_2 and 3P_2 - 3D_3 transitions.

other systems that BBR can redistribute population between near-degenerate levels. Such a process was crucial for the operation of a Ra MOT [90], and we speculate about a similar process for our case. Similar processes might also contribute to the unexpectedly low repump efficiencies discussed in Sec. V.

In conclusion, we find that the 3P_2 - 3D_3 transition at 497 nm is not enough cycling to allow for MOT operation or single-atom fluorescence detection. The situation is even worse for the transition at 403 nm.

VII. CONCLUSION

In conclusion, we have performed spectroscopy on $5s5p\ ^3P_2$ - $5snd\ ^3D_{1,2,3}$ transitions in strontium and determined the isotope shifts and hyperfine parameters. We have accessed all relevant $^3D_J \rightarrow ^3P_J$ branching ratios both theoretically and experimentally. The results have immediate implications for the usage of these transitions in protocols of cooling, manipulation, and detection of ultracold strontium atoms.

The general scheme of performing spectroscopy on atoms stored in a metastable reservoir state can be applied to many other species as well. Elements such as magnesium, calcium [91], ytterbium, and mercury have an electronic structure very similar to strontium. More complex atoms, such as the rare-earth species [92] dysprosium [69,93], holmium [94], erbium [68,95], and thulium [96], which are currently under investigation, all have a wealth of metastable states, which can be trapped in the magnetic field of a MOT and potentially be used for reservoir spectroscopy.

ACKNOWLEDGMENTS

We thank R. Grimm for generous support and stimulating discussions. We thank M. Chwalla, C. Roos, and R. Blatt for the loan of the wave meter and for providing us with an absolute frequency reference based on the $^{40}\text{Ca}^+$ ion. We highly appreciate extensive technical assistance by C. Hempel.

We also wish to thank the strontium groups at JILA, LENS, and Rice University for sharing their experience with us. We gratefully acknowledge support from the Austrian Ministry of Science and Research (BMWF) and the Austrian Science Fund (FWF) through a START grant under Project No. Y507-N20. As member of the project iSense, we also acknowledge the financial support of the Future and Emerging Technologies (FET) programme within the Seventh Framework Programme for Research of the European Commission, under FET-Open Grant No. 250072.

APPENDIX: ERROR ANALYSIS

Laser at 497 nm. The spectroscopy light at 497 nm is generated by a frequency-doubled diode laser, which delivers about 40 mW of green light. The infrared laser is locked to a stable reference cavity, which has a drift of typically 100 kHz per hour, mainly due to piezo creep and possibly thermal drift. The green light is passed through a cascade of four acousto-optic modulators (AOMs) in double-pass configuration. The AOMs have a design frequency of 350 MHz and a bandwidth of about 150 MHz, resulting in an overall scan range of 1.2 GHz. The RF used to drive the AOMs is generated by direct digital synthesizers (DDSs), which are referenced to better than 1 Hz to either the global positioning system GPS or a commercial rubidium atomic clock, both of which have negligible error. The laser light itself has a linewidth smaller than 1 MHz. About 1 mW of light is available for the experiment and collimated to a beam of 10 mm diameter to interrogate the atoms.

Laser at 403 nm. The spectroscopy light at 403 nm is generated by a diode laser, emitting about 30 mW. The light is sent through a series of two 350-MHz AOMs in double-pass configuration and delivered via a fiber to the experiment, where the beam is expanded to illuminate the entire MOT volume. This laser is not locked to a high-finesse cavity, and it shows frequency noise on timescales of a few 100 ms, maximum excursions being about 2 MHz. The drift rate of the laser is rather poor, typically 100 MHz per hour.

Absolute frequency. The frequency of the respective laser used for spectroscopy is constantly monitored on a commercial High Finesse WSU/2 wavemeter. The wavemeter is constantly calibrated by a Ti:Saph laser at 729 nm. This laser is locked to a high-finesse cavity with a well-characterized drift rate of about 200 Hz per hour. The absolute frequency is determined by spectroscopy of the $4s^2S_{1/2}-3d^2D_{5/2}$ transition in the $^{40}\text{Ca}^+$ ion at 432 042 129 776 393.2 (1.0) Hz [97]. Calibration of the wavemeter has an accuracy of 500 kHz, which can be reduced substantially by averaging over many calibrations. If not constantly recalibrated, the wavemeter drifts by about 100 kHz per hour. The precision of the wavemeter is specified to be about 1 MHz.

The accuracy of the wavemeter is about 1 MHz if used within a few nm of the calibration wavelength, but might be substantially larger further away from the calibration wavelength. To characterize the calibration fidelity of the wavemeter over a large range of wavelengths, we measure the frequency of various lasers referenced to various well-established optical transitions, of which the absolute frequencies are well known from the literature. More specifically,




FIG. 7. (Color online) Characterization of the wavemeter error. The wavemeter is calibrated at 729 nm (open diamond) and used to measure light at various well-known frequencies (filled circles). We plot the difference between the obtained values and the literature values, where the error bars are dominated by the calibration and measurement uncertainty of the wavemeter. The stars denote wavelengths at which additional consistency checks were performed, however at a smaller precision. The spectroscopy measurements presented in this work were performed at 403 and 497 nm, denoted by arrows.

these are the $4s^2S_{1/2}-4p^2P_{1/2}$ dipole transition at 397 nm in the $^{40}\text{Ca}^+$ ion, which is known to within 1.7 MHz [98]; the $5s^2^1S_0-5s5p^3P_1$ intercombination line at 689 nm in atomic ^{88}Sr , which is known to within 10 kHz [61,99]; and the $5s^2S_{1/2}-5p^2P_{3/2}$ dipole transition at 780 nm in atomic ^{87}Rb , which is known to within 6 kHz [100]. The infrared light at 793 nm, which is used to generate the light at 397 nm, is used as well. The deviations of the wavemeter reading from the real frequency scatter within about ± 5 MHz; see Fig. 7. We use our frequency-doubled lasers at 397/793 nm, 461/922 nm, and 497/994 nm to connect between different wavelength ranges, as the frequency conversion between the fundamental and the doubled frequency is known to be exactly 2 by design. We perform additional consistency checks with light at wavelengths 461 nm, 854 nm, and 866 nm, corresponding to dipole transitions in either $^{40}\text{Ca}^+$ or ^{88}Sr , which are known at a level of order 100 MHz. From this characterization, we conclude that the wavemeter is accurate to within about 5 MHz over the entire range from 400–1000 nm. As a spinoff, we can determine the frequency of the 461 nm transition in ^{88}Sr to be 650 503 775 (5) MHz, corresponding to an energy of the $5s5p^1P_1$ state of 21 698.4703 (2) cm^{-1} . This value is more precise than (but in slight disagreement with) the one given in Ref. [60].

Error in absolute frequencies. The errors given in Tables I and II include the precision of the wavemeter and the uncertainty in the fit to the data points. The error of the absolute frequencies is entirely dominated by the inaccuracy of the wavemeter, which is certainly below 10 MHz; see Fig. 7. The data is fit by a Lorentzian profile, where deviations from a Lorentzian line shape are visible only for extremely

low repump intensities. The error in the determination of the centroid ranges between 50 kHz and a few MHz, depending on the signal strength. This error becomes comparable to the wavemeter uncertainty only for very weak transitions.

Error in relative frequencies. The error in frequency differences, such as isotope shifts, can be substantially smaller, as the inaccuracy of the wavemeter drops out. For the measurement of the isotope shifts between the bosons, the wavemeter is carefully calibrated and then left free running. A slow drift of the wavemeter then constitutes the dominant error source. The complete spectroscopy scan of the bosonic isotopes is performed within less than half an hour, in which the wavemeter drifts by much less than 100 kHz. A complete scan of all the hyperfine transitions of the fermionic isotope takes about a day, and the wavemeter might drift by up to a few MHz in this period. As a conservative estimate, the isotope shifts between the bosonic isotopes can be measured with an uncertainty of about 200 kHz, and the hyperfine splittings can

be determined to about 4 MHz. While the values stated here apply for the green transition, another error source appears for the blue transition: The frequency noise of the laser, which is about 1 MHz on timescales below 1 s, leads to the comparably large errors given in Tab. III.

Stability and reproducibility. We carefully determine the stability of our MOT atom number and find it to be better than 1% on all timescales between a millisecond and many hours. During a measurement campaign, we repetitively measure certain transitions over the course of a few days to exclude any drifts and find the scatter in absolute frequency to be within less than 1 MHz. Selected frequencies were determined in two independent measurement campaigns separated by a time interval of two years and using largely different setups, and the frequency difference was found to be within the 1 MHz uncertainty of the wavemeter, thereby excluding a large number of possible systematic errors.

-
- [1] T. Ido and H. Katori, *Phys. Rev. Lett.* **91**, 053001 (2003).
- [2] M. Takamoto, F.-L. Hong, R. Higashi, and H. Katori, *Nature (London)* **435**, 321 (2005).
- [3] A. Derevianko and H. Katori, *Rev. Mod. Phys.* **83**, 331 (2011).
- [4] N. Hinkley, J. A. Sherman, N. B. Phillips, M. Schioppo, N. D. Lemke, K. Beloy, M. Pizzocaro, C. W. Oates, and A. D. Ludlow, *Science* **341**, 1215 (2013).
- [5] B. J. Bloom, T. L. Nicholson, J. R. Williams, S. L. Campbell, M. Bishof, X. Zhang, W. Zhang, S. L. Bromley, and J. Ye, *Nature (London)* **506**, 71 (2014).
- [6] N. Poli, F.-Y. Wang, M. G. Tarallo, A. Alberti, M. Prevedelli, and G. M. Tino, *Phys. Rev. Lett.* **106**, 038501 (2011).
- [7] Y. Takasu, K. Maki, K. Komori, T. Takano, K. Honda, M. Kumakura, T. Yabuzaki, and Y. Takahashi, *Phys. Rev. Lett.* **91**, 040404 (2003).
- [8] T. Fukuhara, S. Sugawa, and Y. Takahashi, *Phys. Rev. A* **76**, 051604(R) (2007).
- [9] T. Fukuhara, Y. Takasu, M. Kumakura, and Y. Takahashi, *Phys. Rev. Lett.* **98**, 030401 (2007).
- [10] T. Fukuhara, S. Sugawa, Y. Takasu, and Y. Takahashi, *Phys. Rev. A* **79**, 021601(R) (2009).
- [11] S. Sugawa, R. Yamazaki, S. Taie, and Y. Takahashi, *Phys. Rev. A* **84**, 011610 (2011).
- [12] S. Kraft, F. Vogt, O. Appel, F. Riehle, and U. Sterr, *Phys. Rev. Lett.* **103**, 130401 (2009).
- [13] P. Halder, C.-Y. Yang, and A. Hemmerich, *Phys. Rev. A* **85**, 031603 (2012).
- [14] S. Stellmer, M. K. Tey, B. Huang, R. Grimm, and F. Schreck, *Phys. Rev. Lett.* **103**, 200401 (2009).
- [15] Y. N. Martinez de Escobar, P. G. Mickelson, M. Yan, B. J. DeSalvo, S. B. Nagel, and T. C. Killian, *Phys. Rev. Lett.* **103**, 200402 (2009).
- [16] P. G. Mickelson, Y. N. Martinez de Escobar, M. Yan, B. J. DeSalvo, and T. C. Killian, *Phys. Rev. A* **81**, 051601 (2010).
- [17] B. J. DeSalvo, M. Yan, P. G. Mickelson, Y. N. Martinez de Escobar, and T. C. Killian, *Phys. Rev. Lett.* **105**, 030402 (2010).
- [18] M. K. Tey, S. Stellmer, R. Grimm, and F. Schreck, *Phys. Rev. A* **82**, 011608 (2010).
- [19] S. Stellmer, M. K. Tey, R. Grimm, and F. Schreck, *Phys. Rev. A* **82**, 041602 (2010).
- [20] S. Stellmer, R. Grimm, and F. Schreck, *Phys. Rev. A* **87**, 013611 (2013).
- [21] M. A. Cazalilla and A. M. Rey, *arXiv:1403.2792*.
- [22] C. Wu, J.-P. Hu, and S.-C. Zhang, *Phys. Rev. Lett.* **91**, 186402 (2003).
- [23] C. Wu, *Mod. Phys. Lett. B* **20**, 1707 (2006).
- [24] M. A. Cazalilla, A. Ho, and M. Ueda, *New J. Phys.* **11**, 103033 (2009).
- [25] M. Hermele, V. Gurarie, and A. M. Rey, *Phys. Rev. Lett.* **103**, 135301 (2009).
- [26] A. Gorshkov, M. Hermele, V. Gurarie, C. Xu, P. Julienne, J. Ye, P. Zoller, E. Demler, M. D. Lukin, and A. M. Rey, *Nature Phys.* **6**, 289 (2010).
- [27] M. Foss-Feig, M. Hermele, and A. M. Rey, *Phys. Rev. A* **81**, 051603 (2010).
- [28] C. Xu, *Phys. Rev. B* **81**, 144431 (2010).
- [29] H.-H. Hung, Y. Wang, and C. Wu, *Phys. Rev. B* **84**, 054406 (2011).
- [30] S. Taie, Y. Takasu, S. Sugawa, R. Yamazaki, T. Tsujimoto, R. Murakami, and Y. Takahashi, *Phys. Rev. Lett.* **105**, 190401 (2010).
- [31] S. Taie, R. Yamazaki, S. Sugawa, and Y. Takahashi, *Nature Phys.* **8**, 825 (2012).
- [32] M. J. Martin, M. Bishof, M. D. Swallows, X. Zhang, C. Benko, J. von Stecher, A. V. Gorshkov, A. M. Rey, and J. Ye, *Science* **341**, 632 (2013).
- [33] X. Zhang, M. Bishof, S. L. Bromley, C. V. Kraus, M. S. Safronova, P. Zoller, A. M. Rey, and J. Ye, *arXiv:1403.2964*.
- [34] F. Scazza, C. Hofrichter, M. Höfer, P. C. De Groot, I. Bloch, and S. Fölling, *arXiv:1403.4761*.
- [35] F. Gerbier and J. Dalibard, *New J. Phys.* **12**, 033007 (2010).

- [36] N. R. Cooper, *Phys. Rev. Lett.* **106**, 175301 (2011).
- [37] B. Béri and N. R. Cooper, *Phys. Rev. Lett.* **107**, 145301 (2011).
- [38] A. Górecka, B. Grémaud, and C. Miniatura, *Phys. Rev. A* **84**, 023604 (2011).
- [39] J. Dalibard, F. Gerbier, G. Juzeliunas, and P. Öhberg, *Rev. Mod. Phys.* **83**, 1523 (2011).
- [40] D. Banerjee, M. Bögli, M. Dalmonte, E. Rico, P. Stebler, U.-J. Wiese, and P. Zoller, *Phys. Rev. Lett.* **110**, 125303 (2013).
- [41] S. Diehl, W. Yi, A. J. Daley, and P. Zoller, *Phys. Rev. Lett.* **105**, 227001 (2010).
- [42] S. G. Bhongale, L. Mathey, E. Zhao, S. F. Yelin, and M. Lemeshko, *Phys. Rev. Lett.* **110**, 155301 (2013).
- [43] B. Olmos, D. Yu, Y. Singh, F. Schreck, K. Bongs, and I. Lesanovsky, *Phys. Rev. Lett.* **110**, 143602 (2013).
- [44] M. Lahrz, M. Lemeshko, K. Sengstock, C. Becker, and L. Mathey, *Phys. Rev. A* **89**, 043616 (2014).
- [45] W. Yi, S. Diehl, A. J. Daley, and P. Zoller, *New J. Phys.* **14**, 055002 (2012).
- [46] O. Boada, A. Celi, J. I. Latorre, and M. Lewenstein, *Phys. Rev. Lett.* **108**, 133001 (2012).
- [47] R. Stock, N. S. Babcock, M. G. Raizen, and B. C. Sanders, *Phys. Rev. A* **78**, 022301 (2008).
- [48] A. J. Daley, M. M. Boyd, J. Ye, and P. Zoller, *Phys. Rev. Lett.* **101**, 170504 (2008).
- [49] A. V. Gorshkov, A. M. Rey, A. J. Daley, M. M. Boyd, J. Ye, P. Zoller, and M. D. Lukin, *Phys. Rev. Lett.* **102**, 110503 (2009).
- [50] I. Reichenbach, P. S. Julienne, and I. H. Deutsch, *Phys. Rev. A* **80**, 020701 (2009).
- [51] A. J. Daley, *Quantum Inf. Process.* **10**, 865 (2011).
- [52] N. Rehbein, T. E. Mehlstäubler, J. Keupp, K. Moldenhauer, E. M. Rasel, W. Ertmer, A. Douillet, V. Michels, S. G. Porsev, A. Derevianko *et al.*, *Phys. Rev. A* **76**, 043406 (2007).
- [53] T. Binnewies, G. Wilpers, U. Sterr, F. Riehle, J. Helmcke, T. E. Mehlstäubler, E. M. Rasel, and W. Ertmer, *Phys. Rev. Lett.* **87**, 123002 (2001).
- [54] H. Katori, T. Ido, Y. Isoya, and M. Kuwata-Gonokami, *Phys. Rev. Lett.* **82**, 1116 (1999).
- [55] J. Grünert and A. Hemmerich, *Phys. Rev. A* **65**, 041401 (2002).
- [56] T. Feldker, J. Schütz, H. John, and G. Birkl, *Eur. Phys. J. D* **65**, 257 (2011).
- [57] S. G. Porsev, A. D. Ludlow, M. M. Boyd, and J. Ye, *Phys. Rev. A* **78**, 032508 (2008).
- [58] M. S. Safronova, S. G. Porsev, U. I. Safronova, M. G. Kozlov, and C. W. Clark, *Phys. Rev. A* **87**, 012509 (2013).
- [59] S. Kato, S. Sugawa, K. Shibata, R. Yamamoto, and Y. Takahashi, *Phys. Rev. Lett.* **110**, 173201 (2013).
- [60] J. E. Sansonetti and G. Nave, *J. Phys. Chem. Ref. Data* **39**, 033103 (2010).
- [61] I. Courtillot, A. Quessada-Vial, A. Bruschi, D. Kolker, G. D. Rovera, and P. Lemonde, *Eur. Phys. J. D* **33**, 161 (2005).
- [62] R. Beigang, K. Lücke, D. Schmidt, A. Timmermann, and P. J. West, *Physica Scripta* **26**, 183 (1982).
- [63] P. Esherick, *Phys. Rev. A* **15**, 1920 (1977).
- [64] P. G. Mickelson, Y. N. Martinez de Escobar, P. Anzel, B. J. DeSalvo, S. B. Nagel, A. J. Traverso, M. Yan, and T. C. Killian, *J. Phys. B: At. Mol. Opt. Phys.* **42**, 235001 (2009).
- [65] U. Dammalapati, I. Norris, C. Burrows, and E. Riis, *Phys. Rev. A* **83**, 062513 (2011).
- [66] B. B. Jensen, H. Ming, P. G. Westergaard, K. Gunnarsson, M. H. Madsen, A. Bruschi, J. Hald, and J. W. Thomsen, *Phys. Rev. Lett.* **107**, 113001 (2011).
- [67] K. Beloy, J. A. Sherman, N. D. Lemke, N. Hinkley, C. W. Oates, and A. D. Ludlow, *Phys. Rev. A* **86**, 051404 (2012).
- [68] J. J. McClelland and J. L. Hanssen, *Phys. Rev. Lett.* **96**, 143005 (2006).
- [69] M. Lu, S. H. Youn, and B. L. Lev, *Phys. Rev. Lett.* **104**, 063001 (2010).
- [70] Y. Bidel, Ph.D. thesis, Université de Nice, 2002 (unpublished).
- [71] S. G. Porsev, Y. G. Rakhlina, and M. G. Kozlov, *Phys. Rev. A* **60**, 2781 (1999).
- [72] J. W. Cho, H.-G. Lee, S. Lee, J. Ahn, W.-K. Lee, D.-H. Yu, S. K. Lee, and C. Y. Park, *Phys. Rev. A* **85**, 035401 (2012).
- [73] R. Santra, K. V. Christ, and C. H. Greene, *Phys. Rev. A* **69**, 042510 (2004).
- [74] M. Yasuda and H. Katori, *Phys. Rev. Lett.* **92**, 153004 (2004).
- [75] X. Xu, T. H. Loftus, J. L. Hall, A. Gallagher, and J. Ye, *J. Opt. Soc. Am. B* **20**, 968 (2003).
- [76] H. Katori, T. Ido, Y. Isoya, and M. Kuwata-Gonokami, in *Atomic Physics 17*, edited by E. Arimondo, P. DeNatale, and M. Inguscio (American Institute of Physics, Woodbury, 2001), pp. 382–396.
- [77] T. Kurosu and F. Shimizu, *Jpn. J. Appl. Phys.* **31**, 908 (1992).
- [78] T. P. Dinneen, K. R. Vogel, E. Arimondo, J. L. Hall, and A. Gallagher, *Phys. Rev. A* **59**, 1216 (1999).
- [79] N. Poli, R. E. Drullinger, G. Ferrari, J. Léonard, F. Sorrentino, and G. M. Tino, *Phys. Rev. A* **71**, 061403 (2005).
- [80] T. C. Killian (private communication).
- [81] J. Stuhler, P. O. Schmidt, S. Hensler, J. Werner, J. Mlynek, and T. Pfau, *Phys. Rev. A* **64**, 031405(R) (2001).
- [82] A. Traverso, R. Chakraborty, Y. N. Martinez de Escobar, P. G. Mickelson, S. B. Nagel, M. Yan, and T. C. Killian, *Phys. Rev. A* **79**, 060702(R) (2009).
- [83] T. Mukaiyama, H. Katori, T. Ido, Y. Li, and M. Kuwata-Gonokami, *Phys. Rev. Lett.* **90**, 113002 (2003).
- [84] S. M. Heider and G. O. Brink, *Phys. Rev. A* **16**, 1371 (1977).
- [85] B. Bushaw, H.-J. Kluge, J. Lantsch, R. Schwabach, J. Stenner, H. Stevens, K. Wendt, and K. Zimmer, *Zeitschrift für Physik D* **28**, 275 (1993).
- [86] H. G. C. Werij, C. H. Greene, C. E. Theodosiou, and A. Gallagher, *Phys. Rev. A* **46**, 1248 (1992).
- [87] A. D. Ludlow, Ph.D. thesis, University of Colorado, Boulder, 2008.
- [88] F. Sorrentino, G. Ferrari, N. Poli, R. Drullinger, and G. M. Tino, *Mod. Phys. Lett. B* **20**, 1287 (2006).
- [89] M. J. Martin (private communication).
- [90] J. R. Guest, N. D. Scielzo, I. Ahmad, K. Bailey, J. P. Greene, R. J. Holt, Z.-T. Lu, T. P. O'Connor, and D. H. Potterveld, *Phys. Rev. Lett.* **98**, 093001 (2007).
- [91] D. Hansen and A. Hemmerich, *Phys. Rev. A* **72**, 022502 (2005).
- [92] C. I. Hancox, S. C. Doret, M. T. Hummon, L. Luo, and J. M. Doyle, *Nature (London)* **431**, 281 (2004).
- [93] M. Lu, N. Q. Burdick, S. H. Youn, and B. L. Lev, *Phys. Rev. Lett.* **107**, 190401 (2011).
- [94] J. Miao, J. Hostetter, G. Stratis, and M. Saffman, *Phys. Rev. A* **89**, 041401 (2014).

- [95] K. Aikawa, A. Frisch, M. Mark, S. Baier, A. Rietzler, R. Grimm, and F. Ferlaino, *Phys. Rev. Lett.* **108**, 210401 (2012).
- [96] D. Sukachev, A. Sokolov, K. Chebakov, A. Akimov, S. Kanorsky, N. Kolachevsky, and V. Sorokin, *Phys. Rev. A* **82**, 011405 (2010).
- [97] M. Chwalla, J. Benhelm, K. Kim, G. Kirchmair, T. Monz, M. Riebe, P. Schindler, A. S. Villar, W. Hänsel, C. F. Roos *et al.*, *Phys. Rev. Lett.* **102**, 023002 (2009).
- [98] A. L. Wolf, S. A. van den Berg, C. Gohle, E. J. Salumbides, W. Ubachs, and K. S. E. Eikema, *Phys. Rev. A* **78**, 032511 (2008).
- [99] G. Ferrari, P. Cancio, R. Drullinger, G. Giusfredi, N. Poli, M. Prevedelli, C. Toninelli, and G. M. Tino, *Phys. Rev. Lett.* **91**, 243002 (2003).
- [100] J. Ye, S. Swartz, P. Jungner, and J. L. Hall, *Opt. Lett.* **21**, 1280 (1996).