



UvA-DARE (Digital Academic Repository)

The VLT-FLAMES Tarantula survey. XX. The nature of the X-ray bright emission-line star VFTS 399

Clark, J.S.; Bartlett, E.S.; Broos, P.S.; Townsley, L.K.; Taylor, W.D.; Walborn, N.R.; Bird, A.J.; Sana, H.; de Mink, S.E.; Dufton, P.L.; Evans, C.J.; Langer, N.; Maíz Apellániz, J.; Schneider, F.R.N.; Soszyński, I.

DOI

[10.1051/0004-6361/201424427](https://doi.org/10.1051/0004-6361/201424427)

Publication date

2015

Document Version

Final published version

Published in

Astronomy & Astrophysics

[Link to publication](#)

Citation for published version (APA):

Clark, J. S., Bartlett, E. S., Broos, P. S., Townsley, L. K., Taylor, W. D., Walborn, N. R., Bird, A. J., Sana, H., de Mink, S. E., Dufton, P. L., Evans, C. J., Langer, N., Maíz Apellániz, J., Schneider, F. R. N., & Soszyński, I. (2015). The VLT-FLAMES Tarantula survey. XX. The nature of the X-ray bright emission-line star VFTS 399. *Astronomy & Astrophysics*, 579, [A131]. <https://doi.org/10.1051/0004-6361/201424427>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

The VLT-FLAMES Tarantula survey

XX. The nature of the X-ray bright emission-line star VFTS 399[★]

J. S. Clark¹, E. S. Bartlett², P. S. Broos³, L. K. Townsley³, W. D. Taylor⁴, N. R. Walborn⁵, A. J. Bird⁶, H. Sana⁵,
S. E. de Mink^{7,8,9}, P. L. Dufton¹⁰, C. J. Evans⁴, N. Langer¹¹, J. Maíz Apellániz¹²,
F. R. N. Schneider¹³, and I. Soszyński¹⁴

¹ Department of Physics and Astronomy, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK
e-mail: s.clark@open.ac.uk

² Astrophysics, Cosmology and Gravity Centre, Department of Astronomy, University of Cape Town, Rondebosch 7701,
Republic of South Africa

³ Department of Astronomy & Astrophysics, 525 Davey Laboratory, Pennsylvania State University, University Park, PA 16802,
USA

⁴ UK Astronomy Technology Centre, Royal Observatory Edinburgh Blackford Hill, Edinburgh, EH9 3HJ, UK

⁵ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁶ School of Physics & Astronomy, University of Southampton, Highfield, Southampton, S017 1BJ, UK

⁷ Astronomical Institute Anton Pannekoek, University of Amsterdam, 1098 XH Amsterdam, The Netherlands

⁸ Carnegie Institution for Science: The Observatories, 813 Santa Barbara St, Pasadena, CA 91101, USA

⁹ TAPIR institute, California Institute of Technology, Pasadena, CA 91125, USA

¹⁰ Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, UK

¹¹ Argelander Institut für Astronomie, Auf den Hügeln 71, 53121 Bonn, Germany

¹² Departamento de Astrofísica, Centro de Astrobiología (INTA-CSIC), Campus ESA, Apartado Postal 78,
28 691 Villanueva de la Cañada, Madrid, Spain

¹³ Department of Physics, Denys Wilkinson Building, Keble Road, Oxford, OX1 3RH, UK

¹⁴ Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland

Received 18 June 2014 / Accepted 16 February 2015

ABSTRACT

Context. The stellar population of the 30 Doradus star-forming region in the Large Magellanic Cloud contains a subset of apparently single, rapidly rotating O-type stars. The physical processes leading to the formation of this cohort are currently uncertain.

Aims. One member of this group, the late O-type star VFTS 399, is found to be unexpectedly X-ray bright for its bolometric luminosity – in this study we aim to determine its physical nature and the cause of this behaviour.

Methods. To accomplish this we performed a time-resolved analysis of optical, infrared and X-ray observations.

Results. We found VFTS 399 to be an aperiodic photometric variable with an apparent near-IR excess. Its optical spectrum demonstrates complex emission profiles in the lower Balmer series and select He I lines – taken together these suggest an OeBe classification. The highly variable X-ray luminosity is too great to be produced by a single star, while the hard, non-thermal nature suggests the presence of an accreting relativistic companion. Finally, the detection of periodic modulation of the X-ray lightcurve is most naturally explained under the assumption that the accretor is a neutron star.

Conclusions. VFTS 399 appears to be the first high-mass X-ray binary identified within 30 Dor, sharing many observational characteristics with classical Be X-ray binaries. Comparison of the current properties of VFTS 399 to binary-evolution models suggests a progenitor mass $\gtrsim 25 M_{\odot}$ for the putative neutron star, which may host a magnetic field comparable in strength to those of magnetars. VFTS 399 is now the second member of the cohort of rapidly rotating “single” O-type stars in 30 Dor to show evidence of binary interaction resulting in spin-up, suggesting that this may be a viable evolutionary pathway for the formation of a subset of this stellar population.

Key words. stars: emission-line, Be – binaries: general – stars: individual: VFTS399

1. Introduction

Located within the Large Magellanic Cloud (LMC), the 30 Doradus complex is the most luminous H II region in the Local Group. It contains $>10^3$ OB stars, with star formation apparently commencing ~ 25 Myr ago and continuing to the present day (Walborn & Blades 1997; Grebel & Chu 2000; Walborn et al. 2013). Combined with a moderate line-of-sight extinction, 30 Dor therefore provides an exceptional opportunity to

investigate the complete life-cycle of massive stars. In order to exploit this potential, a multi-epoch spectroscopic observing campaign – the VLT-FLAMES Tarantula Survey (VFTS) – was undertaken between 2008 October–2009 February; a full description of this investigation, including target list, observational strategy and data-reduction methods employed was presented by Evans et al. (2011).

Fulfilling a design goal of the programme, the multi-epoch observations enabled the identification of both single and binary OB star populations (Sana et al. 2013). Investigation of the single O-type stars revealed that the distribution of their rotational

[★] Based on observations collected at the European Southern Observatory under program ID 182.D-0222.




Fig. 1. WFI image of the southern reaches of 30 Dor with VFTS 399 indicated, highlighting its location on the periphery of the complex. The field of view is $\sim 135 \text{ pc} \times 125 \text{ pc}$ for a LMC distance modulus of 18.5 mag (e.g. Gibson 2000).

velocities has a two-component structure (Ramírez-Agudelo et al. 2013), comprising a low velocity peak ($\sim 80 \text{ km s}^{-1}$) and a high-velocity tail (extending to $\sim 600 \text{ km s}^{-1}$). The origin of the high velocity cohort is currently uncertain, with the relative contribution from natal and binary-driven channels still ill-constrained (de Mink et al. 2013; Ramírez-Agudelo et al. 2013). With a projected equatorial rotational velocity, $v_e \sin i \sim 334 \pm 18 \text{ km s}^{-1}$ (Ramírez-Agudelo et al. 2013), the O9 III_n star VFTS 399 is a member of this group (Walborn et al. 2014), while its location on the periphery of the 30 Dor complex (Fig. 1) is suggestive of a potential runaway nature (cf. Blaauw 1961). Unexpectedly, cross correlation of the VFTS sample with the X-ray survey of Townsley et al. (2014) revealed it to be both highly variable and, at its peak, one of the most X-ray luminous stars within 30 Dor. Motivated by these observational findings we report the results of a multi-wavelength analysis of VFTS 399 in order to clarify its nature, presenting the data in Sect. 2, utilising these to infer its physical properties in Sect. 3, discuss the implications of these findings in Sect. 4, before summarising our results in Sect. 5.

2. The observational properties of VFTS 399

2.1. Optical and IR properties

2.1.1. Photometry

Evans et al. (2011) provided single-epoch optical and near-IR photometry of VFTS 399, which we present in Table 1. The latter are derived from the survey of the LMC by Kato et al. (2007), which were obtained on 2003 November 16. Earlier 2MASS magnitudes, obtained on 1998 March 19, are also available (Skrutskie et al. 2006)¹ and are suggestive of variability (see below). Finally *Spitzer* SAGE observations (Meixner et al. 2006) obtained between 2005 July 15–26 and October 26–November 2 are also presented in Table 1.

¹ $J = 15.48$, $H = 15.61$ and $K_s = 14.83$, noting the K_s -band magnitude is flagged as uncertain.

Table 1. Optical and IR photometry of VFTS 399.

B	V	J	H	K_s	[3.6]	[4.5]
15.91	15.83	15.40	15.30	15.22	14.95	14.95

VFTS 399 also lies within fields covered by the Optical Gravitational Lensing Experiment (OGLE-III; Udalski 2003) and VISTA-VMC (Cioni et al. 2011) surveys. Unfortunately, neither dataset coincides with the period over which spectroscopic observations were made but both clearly indicate that VFTS 399 is variable at the level of ~ 0.1 – 0.2 mag over timescales of ~ 10 – 10^2 days in the V , I and K_s -bands (Fig. 2). In the extensive I -band dataset the variability appears characterised by significant scatter superimposed on longer-term variability. The K_s -band data are more sparsely sampled, but are suggestive of coherent variability over a $\sim 10^2$ day timescale. Preliminary analyses of the more extensive OGLE-III I -band dataset revealed a number of potential periodicities – e.g. ~ 156 , 370 and 720 days – although the later two appeared to result from the seasonality of the observations. Nevertheless, in order to assess the significance of all the periodicities we followed the methodology employed by Bird et al. (2012)². No coherent statistically-significant periods were identified for the complete dataset, while searches over subsets of these data also returned no convincing evidence for transient periodicity.

2.1.2. Spectroscopy

The VLT/FLAMES spectroscopy of VFTS 399 is summarised in Table 2 and the data-reduction techniques employed are fully described in Evans et al. (2011), noting that sky subtraction was accomplished utilising a median sky spectrum constructed from all the sky fibres of the frame that were uncontaminated by flux from bright stars. Walborn et al. (2014; their Fig. 10) presented the 3960–4750 Å portion of the spectrum from which they assigned an O9 III_n classification; for brevity we do not replicate this here, but do plot the $H\alpha$, $H\beta$ and He I 6678 Å emission line profiles in Fig. 3.

All three lines demonstrate complex, morphologically similar profiles with pronounced high-velocity emission, of comparable intensity in both blue and red line wings and a strong, narrow component centred on the systemic velocity of VFTS 399. Similar, single-peaked emission at low projected velocities is also present in the photospheric profiles of the higher Balmer series and select He I lines (Walborn et al. 2014). Given the presence of spatially-variable nebular emission across 30 Dor we caution against the over interpretation of the profiles of the core components in these lines. Sky subtraction successfully removed the [N II] 6547 Å nebular line, but left small single-peaked residuals in the [N II] 6584 Å and [S II] 6717+6730 Å lines. Consequently we suspect that nebular contamination might also be present at low projected velocities in the $H\alpha$, $H\beta$ and He I 6678 Å transitions. We discuss the physical interpretation of these profiles further in Sect. 3.

² A fast implementation of the Lomb-Scargle periodogram, with periods searched for ranging from 2 days (the Nyquist frequency) to ~ 1000 days (dictated by the length of the lightcurve). A rolling-mean was further calculated for the lightcurve and was used to de-trend the data, with extensive Monte-Carlo simulations employed to determine the significance and associated errors of any periods detected. See Bird et al. (2012) for further details.




Fig. 2. VISTA-VMC K_s -band (upper panel) and OGLE I - (middle panel) and V -band (lower panel) lightcurves for VFTS 399.

Table 2. Summary of spectroscopic observations.

Mode + setting	λ -coverage (\AA)	R	Date	RV (km s^{-1})
Medusa LR02	3960–4564	7000	22/12/08	266.8 ± 13.3
			24/12/08	286.5 ± 9.1
			27/01/09	300.3 ± 17.3
			17/02/09	265.8 ± 16.0
			07/10/09	276.5 ± 19.2
Medusa LR03	4499–5071	8500	20/12/08	304.9 ± 10.6
			21/12/08	317.1 ± 12.7
			22/12/08	297.3 ± 13.4
Medusa HR15N	6442–6817	16000	18/12/08	–
			19/12/08	–

Notes. The He II 4200 \AA , He I 4387 \AA and He II 4541 \AA transitions were employed for RV determination with Medusa LR02 observations, and He I 4713 \AA and He I 4922 \AA with Medusa LR03. Note the LR02 and LR03 observations made on 2008 Dec. 22 were separated by ~ 68 min.

Ramírez-Agudelo et al. (2013) determined $v_e \sin i \sim 334 \pm 18 \text{ km s}^{-1}$ from the LR02 and LR03 spectra, while Sana et al. (2013) utilised the multi-epoch nature of the dataset to search for radial velocity (RV) shifts. They found that VFTS 399 demonstrated epoch-to-epoch RV shifts with a peak-to-peak amplitude of $\sim 50 \text{ km s}^{-1}$. Given the magnitudes of the observational uncertainties (Table 2), this is significant at the $\sim 2.7\sigma$ level, meaning that VFTS 399 fell below the conservative threshold set by




Fig. 3. Montage of the $H\alpha$, β and He I 6678 \AA line profiles, with the latter two plotted with an arbitrary vertical exaggeration and offset in order to ease comparison to $H\alpha$. Note the central component of $H\alpha$ at projected velocity $\sim 0 \text{ km s}^{-1}$ has a maximum peak intensity of $\sim 6\times$ continuum; we suspect nebular contamination of this feature and the comparable peaks in $H\beta$ and He I 6678 \AA .

these authors for classification as a binary. We return to this issue in Sect. 3.1, noting that these observations also yield a mean RV value of $289 \pm 19 \text{ km s}^{-1}$.

2.2. X-ray properties

Fortuitously, VFTS 399 has been the subject of a number of X-ray observations spanning ~ 15 yr. Both *XMM-Newton* (XMMU J053833.9-691157; Shtykovskiy & Gilfanov 2005) and *Chandra* (CXOU J053833.4-691158 = [TBF2006] 27; Townsley et al. 2006) provide X-ray point source detections that may be associated with VFTS 399. Utilising data from 2001 November 19, the former study quotes $L_{2-10 \text{ keV}} \sim 1.1 \times 10^{35} \text{ erg s}^{-1}$, assuming a power law spectrum with photon index, $\Gamma = 1.7$ and $N_{\text{H}} = 6 \times 10^{20} \text{ cm}^{-2}$. The latter study utilised a ~ 21 ks observation from 1999 September and returned $L_{0.5-8 \text{ keV}} \sim 10^{34} \text{ erg s}^{-1}$ following a power law fit with $\Gamma = 1.7$ and $N_{\text{H}} = 1.3 \times 10^{22} \text{ cm}^{-2}$.

Subsequently, further *Chandra* observations were made on 2006 January 21, 22 and 30 (Townsley et al. 2014). VFTS 399 is also located within the target field of the current ~ 2 Ms T-ReX *Chandra* programme (PI: L. Townsley) which runs through 2015; consequently we appraised the multiple additional observations available to us at the time of writing (Table 3), deferring an analysis of the full dataset for a future paper.

VFTS 399 was detected in each observation and was found to be well separated from other X-ray sources. The resultant lightcurve is presented in Fig. 4 and it is immediately apparent that the source is highly variable, even within a single observational segment. In order to screen for variability in the gross morphological properties of the spectra we determined the median event energy for individual observations (Fig. 4). This metric shows no evidence for spectral variability across this observational sequence. In order to consolidate this finding we

Table 3. Summary of *Chandra* X-ray observations of VFTS 399 during 2006 and 2014.

Obs. ID	Start time	Exposure (s)
5906	2006-01-21T19:04	12 317
7263	2006-01-22T16:51	42 528
7264	2006-01-30T15:06	37 593
16192	2014-05-03T04:10	93 760
16193	2014-05-08T10:15	75 993
16612	2014-05-11T02:15	22 672
16194	2014-05-12T20:00	31 335
16615	2014-05-15T08:24	45 170
16195	2014-05-24T14:09	44 405
16196	2014-05-30T00:05	67 109
16617	2014-05-31T01:27	58 860
16616	2014-06-03T22:26	34 529
16197	2014-06-06T12:32	67 802
16198	2014-06-11T20:20	39 465
16621	2014-06-14T14:46	44 399
16200	2014-06-26T20:01	27 360
16201	2014-07-21T22:13	58 393
16640	2014-07-24T11:21	61 679
16202	2014-08-19T15:30	65 128
17312	2014-08-22T06:21	44 895

analysed the first eight epochs of data which, when taken together, appear representative of the whole dataset inasmuch as they fully sample the full dynamical range found for VFTS 399 ($L_{0.5-8 \text{ keV}} \sim 10^{34}-10^{35} \text{ erg s}^{-1}$). Furthermore, excluding the 1999 data due to calibration issues, we also summed all remaining observations to produce a single, high signal-to-noise (S/N) spectrum which we treated in the same manner.

Specifically, we first subtracted a background spectrum (corresponding to $\sim 3\%$ of the extracted 0.5–8.0 keV X-ray counts) before several different models within *xspec* (Arnaud 1996) were employed in an attempt to fit these data. A simple absorbed power-law (`tbvarabs*pow`) with $Z = 0.4 \times Z_{\odot}$ was found to provide the best fit (Fig. 5): parameters returned for this model upon application to the summed spectrum were $\Gamma = 1.0^{+0.1}_{-0.1}$, $N_{\text{H}} = 2.7^{+0.4}_{-0.3} \times 10^{22} \text{ cm}^{-2}$ (corresponding to $A_{\text{V}} \sim 16.9^{+2.6}_{-1.9}$ mag, suggestive of significant obscuration intrinsic to VFTS 399) and an absorption corrected $L_{0.5-8 \text{ keV}} \sim 5.0 \times 10^{34} \text{ erg s}^{-1}$.

The results obtained from analysis of the first eight individual epochs are presented in Table 4; this yielded a range of parameters spanning $0.8 \leq \Gamma \leq 1.1$, $2.4 \times 10^{22} \leq N_{\text{H}} \leq 3.1 \times 10^{22} \text{ cm}^{-2}$ and $1.0 \times 10^{34} \leq L_{0.5-8 \text{ keV}} \leq 12.6 \times 10^{34} \text{ erg s}^{-1}$. For epochs with very few counts N_{H} and Γ were held at the value determined from the summed spectrum, with only the normalisation allowed to vary. We caution that fits are provided for epochs as a whole and hence do not account for the variable behaviour of the source over the course of a given observation (cf. Fig. 4). Moreover, while some datasets are fitted by this simple model, others are not; this likely indicates deficiencies in the model, but the low count rate precludes more sophisticated treatments at this stage. Nevertheless, we conclude that there is no evidence for evolution of the gross morphological properties of the X-ray spectrum of VFTS 399 over the course of the observations.

2.2.1. X-ray timing analysis

Lomb-Scargle (LS; Lomb 1976; Scargle 1982; Press & Rybicki 1989) analysis was performed on each of the single observation lightcurves as well as on a composite lightcurve, made up of all the datasets. LS analysis is ideal for detecting weak, periodic

modulation in unevenly sampled datasets (such as our composite dataset). A “blind” search for periods between 7 s (approximately twice the frame interval of *Chandra*) and 1×10^5 s was performed with a frequency step of 1×10^{-7} . Foreshadowing the discussion in Sects. 3.2 and 4.1, this exceeds the observed range of pulsational periods exhibited by neutron stars in X-ray binaries with OB star primaries.

At the other extreme, long-term modulation of the lightcurve might be expected from the rotation of magnetic stars, or due to orbital modulation in colliding-wind systems or accreting high-mass X-ray binaries; in all cases the resultant period may range from several days to several years. Unfortunately, the durations of the individual observing blocks are <1 day (with only a single exception; Table 3) meaning that none would be expected to sample a full cycle of a system with a period of a few days. Moreover, the overall time span of the observations is such that we would also not expect to fully sample a single cycle of any long-period ($\geq 10^2$ day) modulation. Thus, we conclude that the current dataset is ill-suited to the *robust* detection of any such putative period; a longer (continuous) time base of observations would be required.

Furthermore, orbital modulation of the X-ray lightcurves of such binaries occurs in the subset of eccentric-orbit systems, which may demonstrate high-amplitude flaring during perisatron passage due to strengthened wind/wind interactions in colliding wind systems (e.g. Corcoran 2005) or enhanced accretion rates in X-ray binaries (e.g. Galache et al. 2008). Such repetitive, periodic flaring is not immediately apparent in the lightcurve of VFTS 399, although this does not preclude its future occurrence since, in the case of X-ray binaries, such behaviour is observed to be transient.

With the above constraints in mind, Fig. 6 shows the LS periodogram of the whole dataset. There appears to be evidence for a period at ~ 2567 s at $>4\sigma$ level of significance. Figure 7 shows the pulse profile, which appears asymmetric and single peaked. A peak at this period is present in the periodograms of a number of individual observation blocks, albeit inevitably at a lower level of significance. The strength of the detection of a period in a given observation depends on the S/N of the observation, the pulsed fraction and the period length: it is easier to find shorter rather than longer periods in any given observation since more “cycles” are present. As such it is not surprising that this period is not robustly detected in some observations; for instance our shortest observation, 5906, is only 12.3 ks in duration, which allows for <5 pulsations. Nevertheless, this latter result gives us confidence that the period is not spuriously introduced by our analysis of the composite lightcurve during the process of “stitching together” its constituent parts; this assumption was quantitatively tested as described below³. Finally, a strong period at 707 s detected in the periodogram of observation 16 193 is the *Chandra* dither period, resulting from a bad CCD column falling within the aperture.

In order to test both the robustness and statistical significance of this period a number of simulations of the data were run. LS analysis was first performed on the barycentric-corrected X-ray event timestamps with a variety of sizes to make sure that the 2567 s period was not an artefact of the binning adopted – in

³ Such a methodology is routinely applied in e.g. the case of optical observations of X-ray binaries, where periodic gaps between distinct segments of the lightcurve are introduced by seasonal visibility constraints (e.g. Bird et al. 2012). Moreover, Reig et al. (2009) encountered exactly this issue in their search for X-ray pulsations in the high-mass binary 4U2206+54, due to visibility gaps in their RXTE lightcurve.




Fig. 4. Concatenated *Chandra* observations of VFTS 399 (975 ks combined exposure). Binned flux and associated 1σ errors are shown in black, while binned median energies are shown in red. A sliding-window lightcurve inferred from the individual events (i.e. no binning) is shown by the blue lines, with a 68% pointwise confidence band limit represented by the two lines. Running *left to right*, the start dates of the 20 epochs of observations are given in Table 3.

each case the period was still detected. The data were then bootstrapped to determine how robust the period was. This was accomplished by creating “fake” datasets by using a random number generator to select $\sim 63\%$ of the data points and performing a LS analysis on the resultant lightcurve. If the detected period was being driven by a small number of data points it will not be consistently detected during the bootstrapping. The period at ~ 2567 s was detected in 96.6% of the trials, suggesting that the period is indeed robust.

We also ran a second bootstrapping analysis, this time selecting random observations (as opposed to data points) to build the simulated datasets. Anywhere from 30 to 80% of the total data were used (since there are fewer ways to randomly select 20 lightcurves in contrast to the full set of data points). In this instance the period at ~ 2567 s was recovered in 85% of the trials. The discrepancy between the two analyses almost certainly arises in the low data fraction regime in the second bootstrapping, since recovering the period with only 30% of the original data will be unlikely.

The significance of the detected period was assessed by estimating the probability of the detection under the null hypothesis that the lightcurve is uniform. Over 15 000 such lightcurves were

generated by randomizing the X-ray event timestamps. This allows us to create simulated lightcurves with the same statistical properties as the original dataset. LS analysis was performed on each of these lightcurves and the maximum LS power achieved recorded. This is shown as the dashed line in Fig. 6. The results of the Monte-Carlo analysis suggest that our ~ 2567 s period, with a LS power of 42.8 has a less than 1 in 15787 (i.e. $<0.006334\%$) probability of being a chance occurrence due to white noise. Although a second peak (at 8718 s) in the periodogram exceeds this significance threshold, it is not robust to the above bootstrapping analyses; specifically, the peak disappears into the noise if the three observing blocks from 2006 are excluded, suggesting that it is most likely an artefact.

We therefore conclude that the Monte-Carlo and bootstrapping analyses both suggest there is a periodic modulation in our dataset although, trivially, it cannot reveal the origin of this periodicity. The ~ 2567 s period does not correspond to the *Chandra* frame interval (3.241040 s), dither period (707 s) or the orbital period (198 ks), nor does it correspond to an alias (the beat frequency) of any of these three potential periods with each other (which corresponds to 3.241093, 3.255966 s and 707.5335 s; for a comprehensive discussion on aliasing and periods see

Table 4. Summary of indicative fits to the *Chandra* X-ray observations of 2006 and 2014.

ObsID	Date	Exposure (s)	Net Counts	N_{H} ($\times 10^{22}$ cm $^{-2}$)	Γ	Normalisation ($\times 10^{-5}$ photons cm $^{-2}$ s $^{-1}$ keV $^{-1}$)	$\chi^2/\text{d.o.f.}$	$\log L_{0.5-8}$ keV (erg s $^{-1}$)
Summed	–	975 400	5397	2.7 $^{+0.4}_{-0.3}$	1.0 $^{+0.1}_{-0.1}$	1.5 $^{+0.2}_{-0.2}$	125.5/115	34.7
5906	21/01/06	12 320	175	2.6 $^{+1.9}_{-1.5}$	0.8 $^{+0.5}_{-0.3}$	2.9 $^{+3.0}_{-1.4}$	4.8/8	35.1
7263	22/01/06	42 528	407	3.0 $^{+1.4}_{-1.0}$	1.0 $^{+0.3}_{-0.3}$	2.4 $^{+1.4}_{-0.8}$	24.1/23	34.9
7264	30/01/06	37 593	40	2.7	1.0	0.2 $^{+0.1}_{-0.1}$	6.7/9	34.0
16192	03/05/14	93 760	419	3.1 $^{+1.8}_{-1.3}$	1.1 $^{+0.4}_{-0.3}$	1.5 $^{+1.2}_{-0.6}$	29.6/24	34.7
16193	08/05/14	75 996	657	3.0 $^{+1.4}_{-1.0}$	1.1 $^{+0.3}_{-0.3}$	3.3 $^{+1.9}_{-1.1}$	38.6/38	35.0
16612	11/05/14	22 678	54	2.7	1.0	0.6 $^{+0.2}_{-0.2}$	9.8/7	34.4
16194	12/05/14	31 335	238	2.7	1.0 $^{+0.3}_{-0.3}$	1.8 $^{+0.6}_{-0.5}$	30.6/13	34.8
16615	15/05/14	45 170	237	2.4 $^{+1.8}_{-1.3}$	0.8 $^{+0.4}_{-0.4}$	1.2 $^{+1.0}_{-0.5}$	6.4/12	34.7

Notes. Summary of the X-ray model parameters for the summed spectrum (*top row*) and the individual spectra derived from the first eight representative, constituent spectra. Parameters given in *italics* have been fixed to the properties derived from the summed spectrum due to low count rate, note also the comparatively poor fit to the spectrum of 2014 May 12. In all cases errors quoted are 90% confidence limits. The X-ray luminosities are derived after correction for extinction and for a distance to the LMC of 50 kpc.




Fig. 5. *Top panel:* absorbed power-law fit to the summed spectrum constructed from the 20 epochs of *Chandra* observations shown in Table 3 (5397 net counts; 975 ks of combined exposure). *Bottom panel:* residuals for this fit.




Fig. 6. LS periodogram for the combined dataset, the dashed line representing the 4σ confidence interval, which the peak corresponding to the ~ 2567 s period clearly exceeds.

Bird et al. 2012). For the remainder of the paper we therefore proceed under the assumption that the period is astrophysical in origin.

Typically, in determining the error on a period one assumes that the period derivative is zero throughout the observation and hence that the dominant source of errors is the Poisson error on the counts themselves⁴. However, given the time span of our observations we may not a priori make this assumption. This in turn prevents a robust quantitative estimate of the error on the period since we may not distinguish between random error and a systematic evolution of the period during the course of the observation.

However, this turns out to be of secondary concern as it is possible that the ~ 2567 s period we detect could be an alias of the true astrophysical period and the *Chandra* frame interval (3.241040 s; note that aliasing is a common problem in the

⁴ In such circumstances one may vary the lightcurve within the errors on the individual data points in order to produce a large number (i.e. $\sim 10^5$) of synthetic lightcurves that are statistically identical to the original and subject each to a LS search. A histogram of the resultant distribution of periods is typically well approximated by a Gaussian, from which the width (i.e. standard deviation) yields a quantitative estimate of the period error.

analysis of e.g. optical lightcurves of X-ray binaries, Bird et al. 2012). This would require the true period of VFTS 399 to be ~ 3.24515 s. Unfortunately following the above discussion such a period would not be *directly* detectable in our current dataset; observations by a different observatory such as XMM would be necessary to distinguish between these possibilities. It is, however, noteworthy that our simulations indicate the ~ 2567 s period is robust to evolution of the period, such as those commonly seen in X-ray binaries due to accretion torques and doppler motion, but *not* so if it is due to an alias. Given the nature of the system (Sects. 3.2 and 4.1) this might be taken as evidence to disfavour the alias hypothesis.

Nevertheless, it is important to emphasise that either interpretation still requires a real periodicity in the X-ray lightcurve to be present. However, we are left in the uncomfortable position of having identified a statistically-significant period of apparently astrophysical origin in our dataset but not being able to *conclusively* determine whether the ~ 2567 s periodicity is an alias or not. As a consequence for the remainder of the paper (e.g. Sects. 3.2 and 4) we are forced to consider the physical implications of both the “short” and “long” periods in parallel. Clearly, independent observations of VFTS 399 with observatories other than *Chandra* would be highly desirable, since one




Fig. 7. X-ray data folded on the ~ 2567 s period. The pulsed fraction, calculated by integrating over the profile and computing the ratio between the pulsed section of the profile and the total flux, is 0.17 ± 0.04 .

requires a much shorter frame time in order to test the alias hypothesis.

3. The nature of VFTS339

VFTS 399 is displaced from the locus occupied by the majority of OB stars in both near-IR ($(J - H)/(H - K_s)$; Dougherty et al. 1994) and mid-IR ($[3.6]/([3.6] - [4.5])$; Bonanos et al. 2009) colour/colour and colour/magnitude diagrams. It is in a region populated by, amongst others, early-type stars exhibiting the classical Be phenomenon. Furthermore, photometric variability, such as that demonstrated by VFTS 399, is a well-known property of classical OeBe stars (e.g. Mennickent et al. 2002).

Emission in the Balmer series is another key classification criterion for classical OeBe stars. Under such a scenario, the emission at large projected velocities in the line wings of $H\alpha$, $H\beta$ and $\text{HeI } 6678 \text{ \AA}$ (Fig. 3) would result from the presence of a compact, quasi-Keplerian decretion disc; during the early stages of disc-reformation, time resolved $H\alpha$ spectra of α And revealed emission components with a comparable peak-to-peak separation (Clark et al. 2003). Similar behaviour is also observed in both the $H\alpha$ and $\text{HeI } 6678 \text{ \AA}$ lines of the B0 Ve primary of the high mass X-ray binary X Per. Comparison of X Per to VFTS 399 appears particularly apposite, given that both stars demonstrate additional emission features interior to those present in the line wings. In X Per this is attributed to the presence of a gaseous circumstellar ring exterior to, and rotating more slowly than, the inner disc that is responsible for the high-velocity emission component of the line profile (Tarasov & Roche 1995; Clark et al. 2001)⁵.

The OeBe phenomenon extends to stars with spectral types as early as that of VFTS399 (e.g. Negueruela et al. 2004), while such stars are also known to be rapid rotators (e.g. Townsend et al. 2004). In light of these observational properties, VFTS 399

appears to be a classical OeBe star and hence we slightly revise its formal classification to O9 IIIne.

3.1. Stellar parameters

The broad-band photometry presented in Table 1 was analysed using the Bayesian code CHORIZOS in order to determine the interstellar reddening (Maíz Apellániz 2004; Maíz Apellániz et al. 2014), from which Walborn et al. (2014) inferred $M_v \sim -4.41$, $T_{\text{eff}} \sim 32.8 \text{ kK}$ and $\log(L_{\text{bol}}/L_{\odot}) \sim 4.9 \pm 0.1$ for VFTS 399. As part of a wider programme (Ramírez-Agudelo, in prep.), the LR02 and LR03 spectroscopic data were subsequently analysed with the non-local thermodynamic equilibrium code FASTWIND (Puls et al. 2005), yielding $T_{\text{eff}} \sim 31.2^{+1.0}_{-0.5} \text{ kK}$, $\log g \sim 3.6^{+0.15}_{-0.1}$ and $\log(L_{\text{bol}}/L_{\odot}) \sim 4.8 \pm 0.1$.

We have also attempted to estimate the nitrogen abundance in VFTS 399. Due to its large projected rotational velocity, the N II spectrum could not be analysed but the N III multiplet near 4515 \AA was observed as a blend in the LR02 and LR03 spectra. In both spectra, this region lies close to either the upper or lower wavelength limits making normalisation of the spectra difficult. We have therefore chosen to treat the spectra separately, with a simple subjective normalisation. These spectra are shown in Fig. 8, together with predictions from model atmosphere calculations. The latter were taken from the O-type star grid of Lanz & Hubeny (2003) and have been convolved with an appropriate broadening function. The weaker theoretical N III spectrum has been interpolated to the FASTWIND atmospheric parameters (and a microturbulence of 10 km s^{-1}) and is for a nitrogen abundance of 7.62 dex and a LMC metallicity. The stronger spectrum is for a nitrogen abundance of 7.92 dex but a galactic metallicity; this leads to more line blanketing, a hotter temperature structure and, for example, enhanced He II absorption. Inspection of grid models with different effective temperatures implies that this will lead to an overestimation of the strength of the N III spectra compared with an LMC metallicity.

Both the LR02 and LR03 spectra imply a nitrogen abundance of approximately 7.8 dex. However we emphasise that this value and the atmospheric parameters from the FASTWIND analysis must be treated with caution due to the limitations of classical non-LTE techniques. Both the $H\alpha$ and $H\beta$ profiles are subject to infilling, with some evidence for similar behaviour in the $\text{H}\gamma$ profile, although a definitive conclusion is complicated by possible nebular contamination. Moreover we have assumed that an isolated stellar model atmosphere is appropriate; the possible systematic errors introduced by the complexity of the system e.g. the presence of an accreting secondary (see Sect. 3.2) are difficult to quantify but could be significant. Nevertheless, we are confident that our conclusion that VFTS 399 demonstrates a pronounced nitrogen enhancement is robust.

Likewise, contamination of the broad-band photometry by continuum emission from the circumstellar disc inferred for VFTS 399 has the potential to seriously compromise reddening, and hence luminosity, estimates. This is evident upon consideration of the long term photometric behaviour of the BeX-ray binary A0535+26. Over a ~ 16 yr period Clark et al. (1999) report $\Delta V \sim 0.4 \text{ mag}$ and $\Delta E(B - V) \sim 0.13 \text{ mag}$ as a result of disc variability (in the sense the system became redder as it brightened). As a consequence if we fail to account for the presence of the circumstellar disc, we risk overestimating the bolometric luminosity of VFTS399 because (i) we will not subtract the excess disc continuum emission and (ii) because we will overestimate

⁵ In this regard we are mindful of the likelihood of nebular contamination in VFTS 399. However even if all emission interior to the high velocity wings were nebular in origin – which appears unlikely upon comparison of the relative widths of the nebular lines to this component – the global consideration that VFTS 399 is a classical OeBe star would not be affected. However, potential unresolved nebular contamination does preclude the measurement of the line equivalent widths.




Fig. 8. LR02 (upper) and LR03 (lower) spectra for VFTS 399 encompassing the N III multiplet near 4515 Å. Also shown are theoretical spectra with nitrogen abundances of 7.62 dex (red, dash) and 7.92 (blue, dot-dash), convolved with an appropriate broadening function. The absorption near 4541 Å is due to He II.

the interstellar reddening component and hence over correct for this.

Riquelme et al. (2012) investigated this problem for Be X-ray binaries in the Galaxy and Magellanic Clouds, finding it to be significant for the latter objects, where contributions from circumstellar and interstellar reddening are comparable. In the absence of contemporaneous spectroscopic and photometric observations of VFTS 399 in a discless state we may not determine the interstellar reddening. Moreover, nebular contamination of the H α line prevents an equivalent width measurement and from this an estimate of the circumstellar reddening via the relationship defined by Riquelme et al. (2012). As a consequence we are forced to adopt a “worst-case” approach, correcting for the maximum contribution from circumstellar reddening reported by Riquelme et al. (2012) of $E^{cs}(B - V) \sim 0.17$ mag. Utilising this value, the photometry from Table 1 and the intrinsic colours of an O9 III star (Martins & Pez 2006) we may infer an interstellar reddening. Then employing the relation from Riquelme et al. (2012) between total optical extinction and interstellar reddening $-A_V^{\text{total}} = 3.1E^{is}(B - V) - 0.3$ mag – for Be X-ray binaries and an appropriate bolometric correction (from Martins et al. 2005) we determine $\log(L_{\text{bol}}/L_{\odot}) \sim 4.4$ for VFTS 399.

This value appears surprisingly low in comparison to either $\log(L_{\text{bol}}/L_{\odot}) \sim 5.17$ or 4.72 expected for, respectively, O9 giants and main sequence stars (Martins et al. 2005). It is difficult to envisage any physical process, such as binary interaction, that would leave a star this under luminous for its spectral

Table 5. Bonnsai predictions for the current mass and lower limit for the age of VFTS 399.

$\log(L_{\text{bol}}/L_{\odot})$	M_{curr} (M_{\odot})	Age (Myr)
–	$21.4^{+2.7}_{-3.7}$	$>5.8^{+0.6}_{-0.7}$
4.7 ± 0.1	$18.0^{+1.2}_{-0.9}$	$>6.1^{+0.5}_{-0.6}$
4.9 ± 0.1	$20.0^{+1.6}_{-1.2}$	$>6.0^{+0.4}_{-0.5}$
5.2 ± 0.1	$24.0^{+2.2}_{-1.7}$	$>5.3^{+0.5}_{-0.5}$

Notes. All four models utilise stellar temperature, surface gravity and projected equatorial rotational velocity as input parameters, while rows 2–4 present the results of including the stellar luminosity estimates (Col. 1) as an additional constraint. The quoted errors are 1σ uncertainties.

classification, suggesting that we have over corrected for circumstellar contamination. Therefore we favour the intrinsic luminosity of an O9 V star as an appropriate lower bound for VFTS 399⁶ and the upper bound from Walborn et al. (2014), leading to $\log(L_{\text{bol}}/L_{\odot}) \sim 4.7$ –4.9.

We employed the Bayesian code BONNSAI⁷ (Schneider et al. 2014b) to determine the current mass (M_{curr}) and age of VFTS 399 (Table 5). Firstly, we neglected the luminosity estimate and simultaneously matched the observed effective temperature, $T_{\text{eff}} = 31.2^{+1.0}_{-0.5}$ kK, surface gravity, $\log g = 3.60^{+0.15}_{-0.10}$ and projected surface rotational velocity, $v_e \sin i = 334 \pm 33$ km s⁻¹ to the rotating, single star LMC models of Brott et al. (2011) and Köhler et al. (2014). We used a Salpeter initial mass function (Salpeter 1955) and the observed distribution of rotational velocities of the 30 Doradus O-type stars (Ramírez-Agudelo et al. 2013) as appropriate priors, together with a uniform distribution of stellar ages and a random orientation of rotational axes. The stellar models predicted a luminosity of $\log(L_{\text{bol}}/L_{\odot}) = 5.0^{+0.25}_{-0.16}$, in agreement with the luminosity estimate of Walborn et al. (2014) and the expectations from the O9 classification. A luminosity of $\log(L_{\text{bol}}/L_{\odot}) \sim 4.4$ was excluded at more than 3σ .

Secondly, we took the luminosity estimates for VFTS 399 into account. Following from above we assumed $\log(L_{\text{bol}}/L_{\odot}) \sim 4.7$ –4.9 but, for completeness, also undertake calculations assuming both $\log(L_{\text{bol}}/L_{\odot}) \sim 4.4$ – appropriate for the maximum expected contamination by a circumstellar disc – and $\log(L_{\text{bol}}/L_{\odot}) \sim 5.2$ – the expected luminosity for an O9 III star (Martins et al. 2005). The results are presented in Table 5. For $\log(L_{\text{bol}}/L_{\odot}) \sim 4.4$, the stellar models could not reproduce the observables simultaneously, likely due to the low luminosity. For our favoured range of bolometric luminosities we determined $M_{\text{curr}} \sim 18.0$ –20.0 M_{\odot} . Finally, we inferred a current age of ~ 6 Myr for VFTS 399, assuming single-star evolution. However, pre-empting the discussion in Sect. 3.2, we consider it likely that VFTS 399 has accreted mass from a binary

⁶ Foreshadowing the discussion in Sect. 4.1, Negueruela & Reig (2001) reported that the emission-line mass-donor BD+53°2790 (classified as O9.5 V) in the high-mass X-ray binary 4U 2206+54 demonstrates variable emission, infilling the He II 4686 Å photospheric line that forms part of the luminosity classification criteria for late O-type stars (Walborn et al. 2014). If such an effect is also present here then it could erroneously lead to the assignment of a giant rather than dwarf luminosity class for VFTS 399.

⁷ The BONNSAI web-service is available at <http://www.astro.uni-bonn.de/stars/bonnsai>

companion in the past, rejuvenating the star (e.g. Schneider et al. 2014a); therefore we consider this value as a lower limit for its true age.

Under this assumption it is of interest to determine whether the observed nitrogen enhancement in VFTS 399 is the result of binary mass transfer. Langer (2012) showed that for Galactic stars such interaction may result in an enrichment of ~ 0.5 – 0.8 dex. Since the initial C/N ratio is ~ 0.2 – 0.3 dex larger in the Galaxy in comparison to the LMC we would expect the nitrogen enrichment in the LMC to be larger by this amount; for a baseline nitrogen abundance of ~ 6.9 dex in the LMC we would predict a post-interaction abundance of ~ 7.6 – 8.0 dex. Unfortunately, while this is consistent with our findings for VFTS 399, it is also consistent with the predictions for rotational mixing in a single, rapidly rotating star, with BONNSAI predicting $\sim 7.76^{+0.24}_{-0.12}$ – $7.83^{+0.22}_{-0.15}$ dex for our favoured range of luminosities. Therefore we are currently unable to utilise the chemical abundance of VFTS 399 in isolation to distinguish between single and binary star evolution.

3.2. Understanding the X-ray emission

Single OB stars are known to be (thermal) X-ray emitters, likely due to shocks propagating through their winds. However, this emission scales as $\log(L_x/L_{\text{bol}}) \sim 10^{-7}$ (e.g. Berghoefer et al. 1997; Sana et al. 2006), while VFTS 399 is orders of magnitude brighter, with $\log(L_x/L_{\text{bol}}) \sim 0.3$ – 4×10^{-4} in the observations reported above, while the emission is of a non-thermal nature. These properties also exclude membership of the class of X-ray bright and variable γ Cas-like OeBe stars, for which a high temperature thermal spectrum and $\log(L_x/L_{\text{bol}}) \sim 10^{-6}$ are expected (Smith et al. 2012; Rauw et al. 2013). Likewise, the peak X-ray luminosity of VFTS 399 is two orders of magnitude greater than that found for Of?p stars and the subset of strongly magnetic O-type stars which, in turn, are X-ray over-luminous with respect to the wider population of massive stars (cf. Nazé et al. 2008; Clark et al. 2009, and refs. therein). Additionally, the X-ray emission in such stars is once again thermal in origin, in contrast to VFTS 399.

At its peak luminosity, VFTS 399 is an order of magnitude brighter than any O+O colliding wind binary, although seven binaries with luminous blue variable or Wolf-Rayet primaries demonstrate $\log L_x \geq 10^{35}$ erg s⁻¹ (Gagné et al. 2012)⁸. However in each of these systems the primaries support significantly more powerful winds than are expected for VFTS 399, suggesting that its X-ray flux is unlikely to arise via wind collisions (indeed no signature of any secondary is visible in our optical spectroscopic observations). Moreover the X-ray emission from such systems is thermal in origin, with $kT \sim 0.9$ – 4.4 keV for those massive binaries with comparable X-ray fluxes.

We are then left with the possibility that VFTS 399 contains an accreting relativistic object and, given the nature of the primary, an obvious identification is with a Be X-ray binary. These systems are overwhelmingly comprised of OeBe stars (\sim O9-B2 V-III; Negueruela & Coe 2002; McBride et al. 2008) orbited by neutron stars and have periods 20 days $\leq P_{\text{orb}} \leq$ 400 days and eccentricities $0.1 \lesssim e \lesssim 0.9$ (Cheng et al. 2014). The interplay between orbital phase and the intrinsically variable circumstellar disc leads to complex, variable X-ray emission (e.g.

Okazaki & Negueruela 2001)⁹. Haberl et al. (2008) determined the X-ray properties of a representative sample of SMC systems from single-epoch observations, finding the majority of spectra to be best fit by power laws ($\Gamma \sim 0.93^{+0.34}_{-0.22}$) with a wide range of absorbing columns ($\sim 10^{20}$ – 10^{22} cm⁻²) and luminosities spanning $\sim 1.5 \times 10^{35}$ – 5.5×10^{36} erg s⁻¹. The X-ray properties of VFTS 399 therefore appear entirely consistent with its classification as a Be X-ray binary.

In such a scenario a neutron star accretor would be strongly favoured on both theoretical and observational grounds. Only one Be+black hole X-ray binary has been identified from a Galactic population of ~ 80 such systems (Casares et al. 2014), while the same study shows it to have a quiescent X-ray luminosity some three orders of magnitude lower than VFTS 399.

Mindful of the caveats in Sect. 2.2.1, the detection of periodic modulation of the X-ray lightcurve provides support for such a conclusion, being most naturally explained as the pulsational period of a rotating, accreting neutron star. Importantly, both “short” (~ 3.245 s) and “long” (~ 2567 s) potential periodicities lie within the range of pulsational periods exhibited by neutron stars within Be X-ray binaries (e.g. Knigge et al. 2011; Reig et al. 2009) and we expand upon this further in Sect. 4.1.

4. VFTS 399 in context

An immediate question to ask is whether we may constrain the orbital parameters of the system from the extant data? Adopting $M_{\text{curr}} \sim 18.0$ – $20.0 M_{\odot}$ for the system primary (Sect. 3.1), the observed RV variability (Table 2 and Sect. 2.1.2) implies an unexpectedly short orbital period of $P_{\text{orb}} \sim 4$ days for a canonical $1.4 M_{\odot}$ neutron star companion. For $P_{\text{orb}} \sim 20$ days, consistent with the commonly found lower bound for Be X-ray binaries (Cheng et al. 2014), we would infer a companion mass of $\sim 2.5 M_{\odot}$, with any longer period strongly favouring a black hole, apparently excluded by the detection of X-ray pulsations¹⁰. Preliminary Monte-Carlo analysis following the methodology of Sana et al. (2013) appears to favour a lower-mass companion with a short period. We caution that this finding should be regarded as provisional since it is primarily driven by the discrepancy in RV between two epochs separated by 0.8 days (Table 2; 2008 December 21, 22). Clearly further observations in order to build an orbital RV curve for the system are required to address this issue.

4.1. The putative neutron star accretor

Is the range of orbital periods suggested by the preceding analysis consistent with one or both of the potential pulsational periods? Following the relationship between the pulsational (P_{spin})

⁹ Most Be X-ray binaries are transient systems, with outbursts typically $>10\times$ their quiescent X-ray luminosities. Following the nomenclature of Stella et al. (1986), Type I outbursts occur periodically during periastron passage and typically yield fluxes of $L_x \sim 10^{36}$ – 10^{37} erg s⁻¹. Type II outbursts are much rarer, aperiodic, have a longer duration and reach $L_x \geq 10^{37}$ erg s⁻¹. In contrast, a subset of systems with wide ($P_{\text{orb}} \geq 100$ days), low-eccentricity orbits are persistent sources with moderate luminosity ($L_x \sim 10^{34}$ – 10^{35} erg s⁻¹) and comparatively low-level variability ($\leq 10\times$ quiescent flux), of which X Per is an exemplar (Haberl et al. 1998; Reig & Roche 2007).

¹⁰ For comparison Antoniadis et al. (2013) determine $\sim 2.01 \pm 0.04 M_{\odot}$ for the pulsar PSR J0348+0432, while Clark et al. (2002) report $\sim 2.44 \pm 0.27 M_{\odot}$ for the non-pulsating relativistic accretor in 4U1700-37 and Casares et al. (2014) find ~ 3.8 – $6.9 M_{\odot}$ for the black hole orbiting the Be star MWC656.

⁸ WR48a, Mk34, η Car, R140a, WR25, R136c and CXO J1745-28.

and orbital (P_{orb}) periods in Be X-ray binaries delineated in the Corbet diagram (Corbet 1986; Knigge et al. 2011) we would infer $P_{\text{orb}} \lesssim 20$ d for $P_{\text{spin}} \sim 3.245$ s and $P_{\text{orb}} \sim 500$ days for $P_{\text{spin}} \sim 2567$ s. While the first combination appears internally consistent, the long orbital period implied by the second would appear to be in tension with our RV data.

However, with $P_{\text{orb}} \lesssim 19.25$ days, and $P_{\text{spin}} \sim 5560$ s (Reig et al. 2009, 2012), the O9.5 Ve+neutron star binary 4U 2206+54 demonstrates that such a combination of long pulsation but short orbital period is viable, although this would require VFTS 399 to host the third most slowly rotating neutron star of any high mass X-ray binary, behind 4U 2206+54 and 2S 0114+650 ($P_{\text{orb}} \sim 11.59$ day and $P_{\text{spin}} \sim 2.7$ h; Crampston et al. 1985; Corbet et al. 1999; Farrell et al. 2008). In this regard we note that both the X-ray luminosity and aperiodic variability of VFTS 399 are replicated in 4U 2206+54 – which is assumed to be powered by direct wind-fed accretion (Negueruela & Reig 2001) – while the primaries in both systems resemble one another (Blay et al. 2006; Walborn et al. 2014).

In each of the systems named above the exceptionally long pulsational periods of the neutron stars are explicable under the assumption that they had been significantly spun down, first via magnetic dipole radiation and subsequently the propellor mechanism (Li & van den Heuvel 1999; Reig et al. 2009, 2012; Popov & Turolla 2012). Operating on comparatively short timescales ($\sim 10^4$ – 10^5 yr; Popov & Turolla 2012), this would also be consistent with the relatively unevolved nature of VFTS 399 (cf. footnote 6), although it would require extremely high magnetic fields, $\geq 10^{14}$ G, in order to facilitate it (cf. Li & van den Heuvel 1999 and Appendix A).

Nevertheless, under either scenario, it is of interest that VFTS 399 appears to host the second young pulsar within the 30 Dor complex after the isolated neutron star PSR J0537-6910 (~ 5 arcmin/ ~ 70 pc distant in projection). With $P_{\text{spin}} \sim 16$ ms (Marshall et al. 1998), PSR J0537-6910 rotates much more rapidly than the putative pulsar within VFTS 399, naturally explicable because of the lack of accretion driven spin-down in the former. However, if the VFTS 399 pulsar is exceptionally slowly rotating due to the effects of an extreme B -field one would be forced to explain their different field strengths ($B \geq 10^{14}$ G versus $B \sim 10^{12}$ G for PSR J0537-6910; Marshall et al. 1998), despite the likelihood of both pulsars having formed from the same underlying stellar population (cf. Dufton et al. 2011).

4.2. The O-type star mass donor

VFTS 399 demonstrates notable physical similarities with VFTS 102, another member of the high rotational-velocity O-type star cohort of 30 Dor (Ramírez-Agudelo et al. 2013). Both are emission-line stars, share the same spectral classification and their temperature, luminosity and surface gravity are identical within the errors quoted (Sect. 3.1 and Table 1 of Dufton et al. 2011). Based on the proximity of VFTS 102 to the isolated pulsar PSR J0537-6910 (Marshall et al. 1998), Dufton et al. (2011) explored a common binary evolution scenario for both objects, with the $16 + 15 M_{\odot}$ mass-transfer model of Cantiello et al. (2007) yielding a rapidly rotating secondary well matched to VFTS 102 at the point of supernova.

While it is tempting to adopt a comparable pre-SN evolutionary scenario for VFTS 399 – albeit with the binary remaining bound at SN – this would be hard to reconcile with the extant RV data. Specifically, a short orbital period would require the binary system to lose a large fraction of the mass and angular momentum released in the pre-SN mass-transfer process

(Petrovic et al. 2005), since conservative evolution leads to orbital periods in the range ~ 50 – 300 day (Wellstein et al. 2001). In other words, this observational constraint implies that VFTS399 accreted comparatively little material, noting that the accretion of only $\sim 2 M_{\odot}$ is sufficient to spin it up to critical rotation. Had the initial mass ratio been close to one – as inferred for VFTS 102 – then the mass transfer would be expected to remain close to conservative (Langer 2012; de Mink et al. 2013). Therefore the properties of the VFTS399 binary system are best understood if the initial mass of the NS progenitor was significantly higher than the current mass of the Oe component.

Given the current absence of a full orbital solution for VFTS 399 the calculation of a tailored evolutionary model is premature, although based on previous analysis we might expect an initial mass of $\geq 25 M_{\odot}$ for the NS progenitor; as a consequence of this we may speculate that VFTS 399 might host a high-mass NS. Under the mass-transfer scenario we favour the mass gainer is expected to quickly rejuvenate, and then behave just as a single star of its new mass, with the possible exception of rapid rotation and surface abundance anomalies. Therefore, we might expect VFTS 399 to have been born with broadly similar properties to those we infer for it now (Table 5 and Sect. 3.1), while our current age estimate (~ 6 Myr; Table 5) would be only slightly lower than expected. Our hypothetical $\sim 25 M_{\odot}$ primary would be expected to explode after ~ 7 Myr (Brott et al. 2011); fully consistent with such a scenario and implying a rather recent ($\lesssim 1$ Myr) SN.

As with VFTS 102, we might therefore expect the systemic velocity of VFTS 399 to have been influenced by the SN. Somewhat surprisingly, there is no indication of it being a radial velocity runaway, with $v_{\text{rad}} \sim 289.4 \pm 18.6$ km s $^{-1}$ in comparison to a cluster mean of ~ 271 km s $^{-1}$ and a velocity dispersion $\sigma \sim 10$ km s $^{-1}$ (Sana et al. 2012, 2013). Nevertheless, the location of VFTS 399 – on the periphery of the complex and displaced from any recognised cluster (Fig. 1; Walborn et al. 2014) – suggests a runaway nature; our prediction of a recent SN would suggest a displacement of ≤ 50 pc from its birth site. Ongoing *Hubble* Space Telescope proper motions studies (PI: D. J. Lennon; see Sabbi et al. 2013) should provide a complementary determination of the transverse velocity in order to test this prediction.

Both VFTS 102 and VFTS 399 have been associated with a population of apparently single, rapidly-rotating O-type star runaways within the 30 Dor complex (Walborn et al. 2014). Of these, 15 stars are of spectral type O8–O9.5 III–V¹¹ and hence might be expected to exhibit the OeBe phenomenon (Negueruela et al. 2004). However, at the time of our spectral observations, only VFTS 102 and 399 were found to support the gaseous circumstellar discs from which a putative neutron star/black hole might accrete. Therefore the remaining stars from this grouping could also harbour relativistic companions which would remain undetectable at X-ray energies until they too formed circumstellar discs. Eldridge et al. (2011) examined the pre- and post-SN evolution of massive binaries and concluded that they remain bound in $\sim 20\%$ of cases, with a most likely systemic velocity of ~ 50 km s $^{-1}$. Given this prediction, long term optical and X-ray observations of this cohort would appear warranted in order to determine if a subset of these stars are indeed quiescent X-ray binaries and hence received their anomalous rotational and/or radial velocities via binary interaction and subsequent SN kick.

¹¹ VFTS 5 (O8 Vn), 74 (O9 Vn), 91 (O9.5 IIIIn), 102 (O9: Vnnne+), 138 (O9 Vn), 249 (O8 Vn), 399 (O9 IIIIn), 530 (O9.5III:nn), 531(O9.5III:nn), 574 (O9.5 IIIIn), 592 (O9.5 Vn), 654 (O9 Vnn), 660 (O9.5 Vnn), 768 (O8 Vn) and 843 (O9.5IIIIn).

5. Concluding remarks

The multi-wavelength properties of the outlying, rapidly rotating O-type star VFTS 399 appear difficult to reconcile with those expected for a single star, instead favouring a binary hypothesis. The X-ray properties strongly favour the presence of a relativistic accretor, with the optical and near-IR properties consistent with classification as a Be X-ray binary, being the seventeenth such system within the LMC at the time of writing (Negueruela & Coe 2002; Vasilopoulos et al. 2013) and the first identified within 30 Dor.

The discovery of an apparently robust, statistically-significant periodicity in the X-ray lightcurve provides strong evidence for a neutron star in the system. Unfortunately we are currently unable to determine *conclusively* whether the ~ 2567 s period reflects P_{spin} or instead is an alias of the true $P_{\text{spin}} \sim 3.245$ s, although the robustness of the former period to the secular evolution common in X-ray binaries argues that it is astrophysical in origin. Breaking this degeneracy via observations with an alternative observatory such as XMM would be of considerable interest, since the former scenario would imply that the neutron star supported a magnetic field of comparable strength to those exhibited by magnetars in order to permit the dramatic spin down required to yield its current P_{spin} .

These findings have a number of ramifications:

- (i) Identification of individual (high-mass) X-ray binaries with their natal stellar aggregates is surprisingly rare – LS I +63 235 lies within the halo of the cluster NGC 663, while PSR B1259-63 and IGR J00370+6122 are likely members of the Cen OB1 and Cas OB8 associations respectively (Negueruela et al. 2011; González-Galán et al. 2014). The location of VFTS 399 within the confines of the 30 Dor complex could therefore aid the observational assessment of the predictions of current population-synthesis models. These include the production (and retention) rates of high mass X-ray binaries as a function of star-formation rate and the duration of such activity. This would also inform studies of e.g. the global X-ray luminosity/star-formation rate relationship for non-active galaxies (e.g. Mineo et al. 2012).
- (ii) In conjunction with determinations of both radial and transverse velocity components, characterisation of the orbital period and eccentricity of VFTS 399 should permit the dynamics of the SN explosion to be reconstructed. Given that Pfahl et al. (2002) and Podsiadlowski et al. (2004) suggest that the magnitude of kick is dependent on the pre-SN binary evolution, the fact that we may place observational constraints on this for VFTS 399 via comparison to the stellar population(s) of 30 Dor is of considerable interest.
- (iii) If the inference of an extreme B -field for the putative neutron star is confirmed, reconstruction of the pre-SN evolutionary pathway and SN dynamics will help constrain the formation requirements for such objects. Moreover, it would support the hypothesis that binarity is a common feature in the formation of magnetars, which in turn would arise from progenitors with a wide mass range from about $17 M_{\odot}$ up to $40 M_{\odot}$ or more (Sect. 4.2 and Clark et al. 2014).
- (iv) After VFTS102, VFTS 399 is the second object drawn from the population of rapidly rotating apparently single O-type stars for which historical/current binarity has subsequently been inferred. Such a finding is consistent with the hypothesis that binary interactions may play a key role in the formation of this cohort (de Mink et al. 2013; Ramírez-Agudelo et al. 2013). If confirmed for further examples it will have considerable import for the production of the

rapidly-rotating, chemically-homogeneous stars inferred to be the progenitors of long gamma-ray bursts.

Acknowledgements. We thank the referee and Danny Lennon and Ignacio Negueruela for their insightful comments which have greatly improved the paper. S.d.M. acknowledges support by NASA through and Einstein Fellowship grant, PF3-140105. STScI is operated by AURA, Inc. under NASA contract NASA 5-26555. L.K.T. and P.S.B. were supported by *Chandra* X-ray Observatory general observer grants GO4-15131X and GO5-6080X and by the Penn State ACIS Instrument Team Contract SV4-74018, issued by the *Chandra* X-ray Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS8-03060. The Guaranteed Time Observations included here were selected by the ACIS Instrument Principal Investigator, Gordon P. Garmire, of the Huntingdon Institute for X-ray Astronomy, LLC, which is under contract to the Smithsonian Astrophysical Observatory; Contract SV2-82024.

Appendix A: Spin-down in the presence of extreme magnetic fields

Following the analysis of Li & van den Heuvel (1999), the evolution of the spin period of the putative neutron star companion to VFTS 399 depends on the accretion rate/torque – a function of the (currently unknown) orbital separation and wind properties of the primary – and its magnetic field. For this mechanism to yield $P_{\text{spin}} \sim 2567$ s it must have been born with an (external) magnetic field of comparable strength to those exhibited by magnetars ($B \gtrsim 10^{14}$ G). If we make the assumption that the neutron star in VFTS 399 is rotating at its equilibrium period we may estimate the dipolar field strength from:

$$P_{\text{spin}} \simeq (20 \text{ s}) B_{12}^{6/7} \dot{M}_{15}^{-3/7} R_6^{18/7} M_{1.4}^{-5/7}$$

(Li & van den Heuvel 1999; Reig et al. 2009) where $B = 10^{12} B_{12}$ G is the dipolar field strength, $\dot{M} = 10^{15} \dot{M}_{15} \text{ g s}^{-1}$ the mass accretion rate and $R = 10^6 R_6$ cm and $M = 1.4 M_{1.4} M_{\odot}$ the neutron star radius and mass. We adopt $P_{\text{spin}} \sim 2567$ s, $R_6 = M_{1.4} = 1$ and $\dot{M}_{15} \approx L_X R / GM = 0.25$ for $L_X \sim 0.5 \times 10^{35} \text{ erg s}^{-1}$ (Reig et al. 2009; Sect. 2.2), which leads to an order-of-magnitude estimate of $B \sim 10^{14}$ G for the neutron star. One might assume that a significantly lower accretion rate may alleviate the requirement for such a B -field. However, following Li & van den Heuvel (1999) e.g. $\dot{M} = 10^{12} \text{ g s}^{-1}$ (leading to $B = 10^{13}$ G) results in a spin-down timescale comparable to the total lifetime of VFTS 399 (cf. Dufton et al. 2011), in contrast to its current relatively unevolved nature.

Recently, Shakura et al. (2012) proposed a new theoretical model for quasi-spherical accretion onto slowly rotating neutron stars (see also Popov & Turolla 2012). Reig et al. (2012) demonstrate that the accretor in 4U2206+54 still supports an extreme ($\sim 10^{14}$ G) B -field under this formulation, although this requires an anomalously slow primary wind (which is observed in this system; Ribó et al. 2006). Clearly, an orbital solution for VFTS 399 and a determination of the wind parameters of the primary (mass loss rate and velocity at the orbital radius) in order to determine \dot{M} would provide an empirical test of the extreme B -field hypothesis, as would a measure of the long term spin-period history of the putative neutron star, from which an estimate of B may also be obtained.

References

- Antoniadis, J., Freire, P. C. C., Wex, N., et al. 2013, *Science*, **340**, 448
 Arnaud, K. A. 1996, in *Astronomical Data Analysis Software and Systems V*, eds. G. H. Jacoby, & J. Barnes, *ASP Conf. Ser.*, **101**, 17
 Berghoefter, T. W., Schmitt, J. H. M. M., Danner, R., & Cassinelli, J. P. 1997, *A&A*, **322**, 167

- Bird, A. J., Coe, M. J., McBride, V. A., & Udalski, A. 2012, *MNRAS*, **423**, 3663
- Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, **15**, 265
- Blay, P., Negueruela, I., Reig, P., et al. 2006, *A&A*, **446**, 1095
- Bonanos, A. Z., Massa, D. L., Sewilo, M., et al. 2009, *AJ*, **138**, 1003
- Brott, I., de Mink, S. E., Cantiello, M., et al. 2011, *A&A*, **530**, A115
- Cantiello, M., Yoon, S.-C., Langer, N., & Livio, M. 2007, *A&A*, **465**, L29
- Casares, J., Negueruela, I., Ribó, M., et al. 2014, *Nature*, **505**, 378
- Cheng, Z.-Q., Shao, Y., & Li, X.-D. 2014, *ApJ*, **786**, 128
- Cioni, M.-R. L., Clementini, G., Girardi, L., et al. 2011, *A&A*, **527**, A116
- Clark, J. S., Lyuty, V. M., Zaitseva, G. V., et al. 1999, *MNRAS*, **302**, 167
- Clark, J. S., Tarasov, A. E., Okazaki, A. T., Roche, P., & Lyuty, V. M. 2001, *A&A*, **380**, 615
- Clark, J. S., Goodwin, S. P., Crowther, P. A., et al. 2002, *A&A*, **392**, 909
- Clark, J. S., Tarasov, A. E., & Panko, E. A. 2003, *A&A*, **403**, 239
- Clark, J. S., Crowther, P. A., & Mikles, V. J. 2009, *A&A*, **507**, 1567
- Clark, J. S., Ritchie, B. W., Najarro, F., Langer, N., & Negueruela, I. 2014, *A&A*, **565**, A90
- Corbet, R. H. D. 1986, *MNRAS*, **220**, 1047
- Corbet, R. H. D., Finley, J. P., & Peele, A. G. 1999, *ApJ*, **511**, 876
- Corcoran, M. F. 2005, *AJ*, **129**, 218
- Crampton, D., Hutchings, J. B., & Cowley, A. P. 1985, *ApJ*, **299**, 839
- de Mink, S. E., Langer, N., Izzard, R. G., Sana, H., & de Koter, A. 2013, *ApJ*, **764**, 166
- Dougherty, S. M., Waters, L. B. F. M., Burki, G., et al. 1994, *A&A*, **290**, 609
- Dufton, P. L., Dunstall, P. R., Evans, C. J., et al. 2011, *ApJ*, **743**, L22
- Eldridge, J. J., Langer, N., & Tout, C. A. 2011, *MNRAS*, **414**, 3501
- Evans, C. J., Taylor, W. D., Hénauld-Brunet, V., et al. 2011, *A&A*, **530**, A108
- Farrell, S. A., Sood, R. K., O'Neill, P. M., & Dieters, S. 2008, *MNRAS*, **389**, 608
- Gagné, M., Fehon, G., Savoy, M. R., et al. 2012, in Proc. Four Decades of Massive Star Research – A Scientific Meeting in Honour of Anthony F. J. Moffat (San Francisco: ASP), *ASP Conf. Ser.*, **465**, 301
- Galache, J. L., Corbet, R. H. D., Coe, M. J., et al. 2008, *ApJS*, **177**, 189
- Gibson, B. K. 2000, *Mem. Soc. Astron. Ital.*, **71**, 693
- González-Galán, A., Negueruela, I., Castro, N., et al. 2014, *A&A*, **566**, A131
- Grebel, E. K., & Chu, Y.-H. 2000, *AJ*, **119**, 787
- Haberl, F., Angelini, L., Motch, C., & White, N. E. 1998, *A&A*, **330**, 189
- Haberl, F., Eger, P., & Pietsch, W. 2008, *A&A*, **489**, 327
- Kato, D., Nagashima, C., Nagayama, T., et al. 2007, *PASJ*, **59**, 615
- Knigge, C., Coe, M. J., & Podsiadlowski, P. 2011, *Nature*, **479**, 372
- Köhler, K., Langer, N., de Koter, A., et al. 2015, *A&A*, **573**, A71
- Langer, N. 2012, *ARA&A*, **50**, 107
- Lanz, T., & Hubeny, I., 2003, *ApJS*, **146**, 417
- Li, X.-D., & van den Heuvel, E. P. J. 1999, *ApJ*, **513**, L45
- Lomb N. R. 1976, *Ap&SS*, **39**, 447
- Maíz Apellániz, J. 2004, *PASP*, **116**, 859
- Maíz Apellániz, J., Evans, C. J., Barbá, R. H., et al. 2014, *A&A*, **564**, A63
- Marshall, F. E., Gotthelf, E. V., Zhang, W., Middleditch, J., & Wang, Q. D. 1998, *ApJ*, **499**, L179
- Martins, F., & Plez, B. 2006, *A&A*, **457**, 637
- Martins, F., Schaerer, D., & Hillier, D. J. 2005, *A&A*, **436**, 1065
- McBride, V. A., Coe, M. J., Negueruela, I., Schurch, M. P. E., & McGowan, K. E. 2008, *MNRAS*, **388**, 1198
- Meixner, M., Gordon, K. D., Indebetouw, R., et al. 2006, *AJ*, **132**, 2268
- Mennickent, R. E., Pietrzyński, G., Geiren, W., & Szcwzyk, O. 2002, *A&A*, **393**, 887
- Mineo, S., Gilfanov, M., & Sunyaev, R. 2012, *MNRAS*, **419**, 2095
- Nazé, Y., Walborn, N. R., & Martins, F. 2008, *Rev. Mex. Astron. Astrofis*, **44**, 331
- Negueruela, I., & Coe, M. J. 2002, *A&A*, **385**, 517
- Negueruela, I., & Reig, P. 2001, *A&A*, **371**, 1056
- Negueruela, I., Steele, I. A., & Bernabeu G. 2004, *Astron. Nachr.*, **325**, 749
- Negueruela, I., Ribó, M., Herrero, A., et al. 2011, *ApJ*, **732**, L11
- Okazaki, A. T., & Negueruela, I. 2001, *A&A*, **377**, 161
- Petrovic, J., Langer, N., & van der Hucht, K. A. 2005, *A&A*, **435**, 1013
- Pfahl, E., Rappaport, S., Podsiadlowski, P., & Spruit, H. 2002, *ApJ*, **574**, 364
- Podsiadlowski, Ph., Langer, N., Poelerends, A. J. T., et al. 2004, *ApJ*, **612**, 1044
- Popov, S. B., & Turolla, R. 2012, *MNRAS*, **421**, L127
- Press, W. H., & Rybicki, G. B. 1989, *ApJ*, **338**, 277
- Puls, J., Urbaneja, M. A., Venero, R., et al. 2005, *A&A*, **435**, 669
- Ramírez-Agudelo, O. H., Simón-Díaz, S., Sana, H., et al. 2013, *A&A*, **560**, A29
- Rauw, G., Nazé, Y., Spano, M., Morel, T., Ud-Doula, A. 2013, *A&A*, **555**, L9
- Reig, P., & Roche, P. 1999, *MNRAS*, **306**, 100
- Reis, J., Torrejón, J. M., Negueruela, I., et al. 2009, *A&A*, **494**, 1073
- Reig, P., Torrejón, J. M., & Blay, P. 2012, *MNRAS*, **425**, 595
- Ribó, M., Negueruela, I., Blay, P., Torrejón, J. M., & Reig, P. 2006, *A&A*, **449**, 687
- Riquelme, M. S., Torrejón, J. M., & Negueruela, I. 2012, *A&A*, **539**, A114
- Sabbi, E., Anderson, J., Lennon, D. J., et al. 2013, *AJ*, **146**, 53
- Salpeter, E. E., 1955, *ApJ*, **121**, 161
- Sana, H., Rauw, G., Nazé, Y., Goset, E., & Vreux, J.-M. 2006, *MNRAS*, **372**, 601
- Sana, H., Dunstall, P. R., Hénauld-Brunet, V., et al. 2012, in Proc. Four Decades of Massive Star Research – A Scientific Meeting in Honour of Anthony F. J. Moffat (San Francisco: ASP), *ASP Conf. Ser.*, **465**, 284
- Sana, H., de Koter, A., de Mink, S. E., et al. 2013, *A&A*, **550**, A107
- Scargle, J. D. 1982, *ApJ*, **263**, 835
- Schneider, F. R. N., Izzard, R. G., de Mink, S. E., et al. 2014a, *ApJ*, **780**, 117
- Schneider, F. R. N., Langer, N., de Koter, A., et al. 2014b, *A&A*, **570**, A66
- Shakura, N., Postnov, K., Kochetkova, A., & Hjalmarsdotter, L. 2012, *MNRAS*, **420**, 216
- Shtykovskiy, P., & Gilfanov, M. 2005, *A&A*, **431**, 597
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, **131**, 1163
- Smith, M. A., Lopes de Oliveira, R., Motch, C., et al. 2012, *A&A*, **540**, A53
- Stella, L., White, N. E., & Rosner, R. 1986, *ApJ*, **308**, 669
- Tarasov, A. E., & Roche, P. 1995, *MNRAS*, **276**, L19
- Townsend, R. H. D., Owocki, S. P., & Howarth, I. D. 2004, *MNRAS*, **350**, 189
- Townsley, L. K., Broos, P. S., Feigelson, E. D., Garmire, G. P., & Getman, K. V. 2006, *AJ*, **131**, 2164
- Townsley, L. K., Broos, P. S., Garmire, G. P., et al. 2014, *ApJS*, **213**, 1
- Udalski, A. 2003, *AcA*, **53**, 291
- Vasilopoulos, G., Maggi, P., Haberl, F., et al. 2013, *A&A*, **558**, A3
- Walborn, N. R., & Blades, J. C. 1997, *ApJS*, **112**, 457
- Walborn, N. R., Barbá, R. H., & Sewilo, M. M. 2013, *AJ*, **145**, 98
- Walborn, N. R., Sana, H., Simón-Díaz, S., et al. 2014, *A&A*, **564**, A40
- Wellstein, S., Langer, N., & Braun, H. 2001, *A&A*, **369**, 939