



UvA-DARE (Digital Academic Repository)

Unravelling the potential mechanisms behind hospitalization-associated disability in older patients

The Hospital-Associated Disability and impact on daily Life (Hospital-ADL) cohort study protocol

Reichardt, L.A.; Aarden, J.J.; van Seben, R.; van der Schaaf, M.; Engelbert, R.H.H.; Bosch, J.A.; Buurman, B.M.; Hospital-ADL study group

DOI

[10.1186/s12877-016-0232-3](https://doi.org/10.1186/s12877-016-0232-3)

Publication date

2016

Document Version

Final published version

Published in

BMC geriatrics

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Reichardt, L. A., Aarden, J. J., van Seben, R., van der Schaaf, M., Engelbert, R. H. H., Bosch, J. A., Buurman, B. M., & Hospital-ADL study group (2016). Unravelling the potential mechanisms behind hospitalization-associated disability in older patients: The Hospital-Associated Disability and impact on daily Life (Hospital-ADL) cohort study protocol. *BMC geriatrics*, 16, [59]. <https://doi.org/10.1186/s12877-016-0232-3>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

STUDY PROTOCOL

Open Access



Unravelling the potential mechanisms behind hospitalization-associated disability in older patients; the Hospital-Associated Disability and impact on daily Life (Hospital-ADL) cohort study protocol

Lucienne A. Reichardt¹, Jesse J. Aarden^{2,3}, Rosanne van Seben¹, Marike van der Schaaf^{2,3}, Raoul H. H. Engelbert^{2,3}, Jos A. Bosch⁴, Bianca M. Buurman^{1,3*} and on behalf of the Hospital-ADL study group

Abstract

Background: Over 30 % of older patients experience hospitalization-associated disability (HAD) (i.e., loss of independence in Activities of Daily Living (ADLs)) after an acute hospitalization. Despite its high prevalence, the mechanisms that underlie HAD remain elusive. This paper describes the protocol for the Hospital-Associated Disability and impact on daily Life (Hospital-ADL) study, which aims to unravel the potential mechanisms behind HAD from admission to three months post-discharge.

Methods/design: The Hospital-ADL study is a multicenter, observational, prospective cohort study aiming to recruit 400 patients aged ≥ 70 years that are acutely hospitalized at departments of Internal Medicine, Cardiology or Geriatrics, involving six hospitals in the Netherlands. Eligible are patients hospitalized for at least 48 h, without major cognitive impairment (Mini Mental State Examination score ≥ 15), who have a life expectancy of more than three months, and without disablement in all six ADLs. The study will assess possible cognitive, behavioral, psychosocial, physical, and biological factors of HAD. Data will be collected through: 1] medical and demographical data; 2] personal interviews, which includes assessment of cognitive impairment, behavioral and psychosocial functioning, physical functioning, and health care utilization; 3] physical performance tests, which includes gait speed, hand grip strength, balance, bioelectrical impedance analysis (BIA), and an activity tracker (Fitbit Flex), and; 4] analyses of blood samples to assess inflammatory and metabolic markers. The primary endpoint is additional disabilities in ADLs three months post-hospital discharge compared to ADL function two weeks prior to hospital admission. Secondary outcomes are health care utilization, health-related quality of life (HRQoL), physical performance tests, and mortality. There will be at least five data collection points; within 48 h after admission (H_1), at discharge (H_3), and at one (P_1 ; home visit), two (P_2 ; by telephone) and three months (P_3 ; home visit) post-discharge. If the patient is admitted for more than five days, additional measurements will be planned during hospitalization on Monday, Wednesday, and Friday (H_2).

Discussion: The Hospital-ADL study will provide information on cognitive, behavioral, psychosocial, physical, and biological factors associated with HAD and will be collected during and following hospitalization. These data may inform new interventions to prevent or restore hospitalization-associated disability.

Keywords: Acute hospitalization, Hospitalization-associated disability, Functional decline, Older patients

* Correspondence: b.m.vanes@amc.uva.nl

¹Department of Internal Medicine, Section of Geriatric Medicine, Academic Medical Center, PO Box 226001100 DD Amsterdam, The Netherlands

³Amsterdam Center for Innovative Health Practice (ACHIEVE), Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands

Full list of author information is available at the end of the article



Background

Studies have observed that at least 30 % of older patients hospitalized with an acute medical illness show a persistent decline in their ability to maintain Activities of Daily Living (ADLs) [1–5]. Such activities are prerequisites to self-care and independent living and include bathing, dressing, transferring out of bed, eating, toileting, and being mobile in and around the house. [1, 3–7]. This decline has been denoted hospitalization-associated disability (HAD), and is defined as the loss of ability to perform one or more of the basic ADLs [6].

HAD is an important problem; it is the leading cause of loss of independence at older age [4] and it is a complex and highly dynamic process with possible recurrent disability episodes in older patients [6, 8]. Research shows that older persons who have been hospitalized have a 60-fold increased risk to develop permanent disabilities [9]. The first month after hospital discharge has been identified as a critical period for recovery, after which disabilities have a high risk of becoming permanent [3]. Moreover, patients with new disabilities are at high risk for other adverse outcomes within three months post discharge: 20 % have readmissions [10], and post-discharge mortality is 25 % [1, 3, 11, 12]. In light of the high prevalence in older patients, and the rapid aging of western societies with a concomitant rise in hospitalizations, a better understanding of HAD is urgently needed.

Previous research has been able to identify a number of generic risk factors for hospitalization-associated disability such as older age [5], the severity of acute illness, geriatric conditions, cognitive impairment and delirium [1, 6, 13, 14]. However, a more fine-grained analyses and characterization of potentially modifiable risk factors is virtually absent from the literature. Little is known, for example, about: 1] the course of loss of muscle mass and strength, and the amount of physical activity older patients perform; 2] the association of cognitive, (psycho-)somatic, behavioral, and psychological restrictive symptoms with the onset and recovery from HAD within the critical period of three months post-discharge such as cognitive impairment, fatigue, pain, sleep quality, shortness of breath, dizziness, fear of falling, diminished self-efficacy, apathy, depression, and anxiety and; 3] the association of the inflammatory activity and related sickness behaviors with the onset and recovery from HAD. Moreover, most previous studies have utilized relatively long follow-up intervals (e.g., from admission to three months or more) [1–3, 15]. Thus information is lacking on events and processes that take place during the weeks after discharge, which are thought to be critical to recovery.

The current study – Hospital-Associated Disability and impact on daily Life (Hospital-ADL study) – aims to investigate cognitive, behavioral, psychosocial, physical, and biological factors that may be associated with HAD in acutely

hospitalized older adults, performing frequent assessments to capture their dynamic development from hospital admission to three months post-discharge. This overall aim can be separated into the following five subordinate aims:

- (1) To study the temporal profile of HAD (i.e., loss of ADL) from hospitalization to three months post-discharge.
- (2) To investigate the course of physical functions that are essential to support ADL, such as muscle mass, muscle strength, and physical performance and spontaneous activity, and test its predictive value for the onset and recovery from HAD, health care utilization, and health-related quality of life (HRQoL) at three months post-discharge.
- (3) To study the prevalence, incidence and course of cognitive, (psycho-)somatic, behavioral, and psychological problems older patients experience from hospitalization up to three months post-discharge that might be restrictive in recovery from HAD post-discharge such as cognitive impairment, fatigue, pain, sleep quality, shortness of breath, dizziness, fear of falling, diminished self-efficacy, apathy, depression, and anxiety.
- (4) To study the association of aforementioned symptoms with HAD, health care utilization, and HRQoL.
- (5) To study the association of metabolic and proinflammatory factors, and physical and behavioral concomitants (e.g., sickness behaviors, loss of muscle mass) with the onset of and recovery from HAD, health care utilization, and HRQoL.

Methods/Design

Study design and setting

The Hospital-ADL study is a multicenter, observational, prospective cohort study designed by an interdisciplinary team of researchers in the field of geriatrics, nursing, psychology, physical therapy and rehabilitation. Six hospitals will participate: 1] the Academic Medical Center in Amsterdam (AMC), a 1002-bed university teaching hospital; 2] the Isala in Zwolle, a 994-bed regional teaching hospital; 3] the Tergooi in Blaricum, a regional teaching hospital (633-bed spread over two sites: Hilversum and Blaricum); 4] the Slotervaart Hospital in Amsterdam, a 310-bed regional teaching hospital; 5] the BovenIJ Hospital in Amsterdam, a 313-bed regional teaching hospital, and; 6] the Meander Medical Center in Amersfoort, a 543-bed regional teaching hospital. The study has started October 1, 2015 and will end after the last patient has been followed up for three months post-discharge. We expect the recruitment phase to be completed late 2016.

Patients

We aim to recruit 400 non-fully disabled adults aged ≥ 70 years. The following inclusion criteria apply: 1] acutely admitted at departments of Internal Medicine, Cardiology or Geriatrics for 48 h or more in one of the above mentioned hospitals; 2] 70 years and older; 3] have approval from the attending Medical Doctor for inclusion; 4] score of 15 or higher on the Mini-Mental State Examination; 5] Dutch language proficiency sufficient to complete questionnaires. Patients will be excluded if they: 1] have a life expectancy of three months or less as assessed by the attending Medical Doctor, or; 2] are disabled in all six basic ADL as determined by the Katz-ADL index [16].

Procedures

Eligible patients will be contacted, and the patient will be informed about the objectives of this study and the study procedures, upon which written informed consent is obtained. Furthermore, a legal representative of the patient will be contacted if the patient has a MMSE score between 15 and 20. Two mobile geriatric assessment teams will visit all six hospitals and will be present on Monday, Wednesday and Friday for consenting and to perform assessments. The mobile geriatric assessment teams consist of a psychologist, physical therapist, and/or a health scientist. The teams are trained in the study procedures of obtaining informed consent, to perform assessments and physical performance tests with adequate inter- and intra-rater reliability (>0.8), and completing the electronic case report form (eCRF).

Table 1 provides an overview of the location, content of assessment and duration of data collection per time point. There will be at least five data collection points; within 48 h after admission (H_1), at discharge (H_3), and at one (P_1 ; home visit), two (P_2 ; by telephone) and three months (P_3 ; home visit) post-discharge. If the patient is admitted for more than five days, additional measurements will be planned during hospitalization on Monday, Wednesday, and Friday (i.e., the days that the mobile geriatric assessment team is present) (H_2).

Data will be collected through: 1] medical and demographical data (e.g., socio-demographic characteristics, severity of acute illness, and geriatric-, and chronic conditions); 2] personal interviews (including cognitive, behavioral, psychosocial, and physical parameters, and health care utilization, see description of information collected below); 3] physical performance tests (e.g., gait speed, muscle strength, muscle mass, mobility and physical functioning, see below) and; 4] blood samples (e.g., to assess markers of inflammation).

The personal interviews will take place during hospitalization (H_1 , H_2 , and H_3), at the participant's home or residence (P_1 and P_3 ; one and three months

post-discharge), and by telephone (P_2 ; two months post-discharge). Physical performance data will be collected within 48 h after admission (H_1), during hospitalization on Monday, Wednesday and Friday (H_2), at discharge (H_3), and at one and three months post-discharge (P_1 and P_3).

Primary outcome

The primary outcome is the level of ADL functioning three months post-discharge compared to premorbid functioning, which are measured with the 6-item Katz-ADL index score of the modified Katz-ADL index [17]. The Katz-ADL index score assesses the degree of independence in bathing, dressing, toileting, use of incontinence materials, transfer from bed-chair and eating [16].

Secondary outcomes

Secondary outcomes include:

- (1) Health care utilization (extension of the Minimal Dataset (MDS) [18] and Comprehensive Geriatric Assessment of the Transitional Care Bridge (TCB) [19], see below).
- (2) Quality of life as measured with the EuroQol-5D [20] and the three items of the MDS [18] (see description below).
- (3) Physical performance tests (see below for description of included tests).
- (4) Mortality.

Scales and assessments

Table 2 gives a detailed overview of the primary and secondary outcomes at each time point.

(1) Medical and demographical data

Socio-demographic characteristics. Socio-demographic data include age, gender, date and time of admission, highest level of education, ethnicity, marital status and living arrangement.

Geriatric conditions. A comprehensive geriatric assessment (CGA) will be collected, which will provide insight in the pre-illness determinants such as polypharmacy, substance use, incontinence, and vision- and hearing impairments.

Chronic conditions. The number and severity of comorbidities will be scored with the Charlson Comorbidity Index [21]. Depending on the risk of mortality, each condition is assigned a score of 1, 2, 3, or 6. Higher scores indicate a greater risk of mortality.

Severity of acute illness. The severity of the acute illness will be measured with the Modified Early Warning Score (MEWS). The MEWS is based on 1] respiratory rate; 2] heart rate; 3] systolic and

Table 1 Time, location, content of assessment and duration of the Hospital-ADL study

| Time | Location | Content of assessment | Duration (minutes) |
|--|---|---|--------------------|
| H ₁ (Within 48 h after admission) | Hospital  | Medical & demographical data Socio-demographic characteristics Medical comorbidity Geriatric conditions Severity of acute illness (medical record) Personal interview/self-report data Cognitive functioning ADL/physical functioning Behavioral & psychosocial functioning Health care utilization (medical record) Physical performance tests Blood parameters | 60 |
| H ₂ (During hospital stay on Monday-Wednesday-Friday) | Hospital  | Medical & demographical data Severity of acute illness (medical record) Short personal interview/self-report data Physical performance tests Blood parameters | 20–30 |
| H ₃ (At hospital discharge) | Hospital  | Personal interview/self-report data Cognitive functioning ADL/Physical functioning Behavioral and psychosocial functioning Physical performance tests Blood parameters | 40 |
| P ₁ (One month post-discharge) | Home visit  | Medical & demographical data Socio-demographic characteristics Geriatric conditions Personal interview/self-report data Cognitive functioning ADL/physical functioning Behavioral & psychosocial functioning Health care utilization Physical performance tests | 60 |
| P ₂ (Two months post-discharge) | By telephone  | Personal interview/self-report data ADL/physical functioning Behavioral and psychosocial functioning Health care utilization | 20 |
| P ₃ (Three months post-discharge) | Home visit  | Medical & demographical data Socio-demographic characteristics Geriatric conditions Personal interview/self-report data Cognitive functioning ADL/physical functioning Behavioral & psychosocial functioning Health care utilization Physical performance tests Mortality (medical record) | 60 |

Table 2 Summary of outcome measures and time points of assessment in Hospital-ADL study

| | Question or instrument | H ₁ | H ₂ | H ₃ | P ₁ | P ₂ | P ₃ |
|---|--|----------------|----------------|----------------|----------------|----------------|----------------|
| 1. Medical & demographical data | | | | | | | |
| Age | Date of birth | × | | | | | |
| Gender | | × | | | | | |
| Postal code | | × | | | | | |
| Date and time of admission | | × | | | | | |
| Education | (In accordance with Verhage, 1966 [57]) | × | | | | | |
| Ethnicity | Country of birth patient and parents | × | | | | | |
| Marital status [18] | | × | | | | | |
| Living arrangement [18, 19] | | × | | | × | | × |
| Medical comorbidity | CCI [21] | × | | | | | |
| Severity of acute illness | MEWS [22] | × | × | × | | | |
| Admission diagnosis | | × | | | | | |
| 2. Personal interviews/self-report data | | | | | | | |
| 2.1 Cognitive functioning | | | | | | | |
| Cognitive impairment | MMSE [23] | × | | × | × | | × |
| Delirium | CAM [24, 58] | × | | | | | |
| | Assessing whether: 1) the patient needs help with self-care; 2) the patient has previously undergone a delirium and; 3) the patient has a cognitive impairment [25] | × | | | | | |
| 2.2 Behavioral & psychosocial functioning | | | | | | | |
| Fear of falling | NRS fear of falling | × | × | × | × | × | × |
| Anxiety | STAI-6 [31] | × | | × | × | × | × |
| Apathy | GDS-15 [29] | × | | × | × | × | × |
| General self-efficacy | ALCOS-12 [34] | | | × | × | × | |
| Quality of life | 1) In general, how is your quality of life?; 2) How would you grade your life at this moment, with a range between 0 and 10? and; 3) Compared to one year ago, how would you rate your health in general now? [18] | × | | × | × | × | × |
| | EQ-5D [20] | × | | × | × | × | × |
| 2.3 ADL/Physical functioning | | | | | | | |
| Disability in ADLs | Modified Katz Index Scale [16, 17] | × | | × | × | × | × |
| Independency in walking | FAC [42] | × | × | × | × | × | × |
| Mobility | Could you walk outside for 5 minutes two weeks before admission/currently? And how often did/do you do physical activity two weeks before admission/currently? [19] | × | | × | × | × | × |
| Falls | Have you fallen once or more in the past (six) month(s)? If yes, how many times? [25] | × | | × | × | × | × |
| Pain | NRS pain [35] | × | × | × | × | × | × |
| Fatigue | NRS fatigue [37] | × | × | × | × | × | × |
| Impact of fatigue | MFIS-5 [38] | | | | × | × | × |
| Sleep quality | PSQI [39] | × | | × | × | × | × |
| Sleep medication | PSQI [39] | × | | × | × | × | × |
| Daytime sleepiness | Do you currently suffer from daytime sleepiness? If yes, does this affect your daily living? | × | × | × | × | × | × |
| Polynocturia | Do you currently suffer from polynocturia? If yes, does this affect your daily living? | × | × | × | × | × | × |
| Dizziness | Do you currently suffer from dizziness? If yes, does this affect your daily living? | × | × | × | × | × | × |

Table 2 Summary of outcome measures and time points of assessment in Hospital-ADL study (Continued)

| | | | | | | | | |
|---|---|----------------|--------------------------------|----------------|----------------|----------------|---|---|
| Shortness of breath | Do you currently suffer from shortness of breath? If yes, does this affect your daily living? | × | × | × | × | × | × | × |
| Hearing impairment | Do you experience difficulties with hearing, despite the use of a hearing aid? | × | | | × | | | × |
| Vision impairment | Do you experience difficulties with your vision, despite the use of glasses? | × | | | × | | | × |
| Nutrition | SNAQ [25, 41] | × | | × | × | × | × | × |
| Dependency | Do you smoke? Do you use alcohol [19]? | × | | | × | | | × |
| Polypharmacy | Do you use five or more different medications [19]? | × | | | × | | | × |
| 2.4 Health care utilization | | | | | | | | |
| Readmission | Have you been hospitalized in the last (six) month(s)? If yes, for how many days? [18] | × | * | | | × | × | × |
| Nursing home admission | Have you had a nursing home admission in the last month? If yes, for how many weeks totally? [18] | | | | | × | × | × |
| Consult physiotherapist and/or occupational therapist | Have you had a consultation with your physiotherapist and/or occupational therapist in the last month? If yes, how many times? | | | | | × | × | × |
| Consult general practitioner | Have you had a consultation with your general practitioner in the last month? If yes, in the evening, night or weekend and how many times totally? [19] | | | | | × | × | × |
| Home care | Do you use home care? If yes, care assistance and/or domestic help and how many hours per week [19] | | | | | × | × | × |
| 3. Physical performance tests | | | | | | | | |
| Handgrip strength | Jamar® [59–61] | × | × | × | × | × | × | × |
| Mobility | DEMMI [45] | × | × | × | × | × | × | × |
| Agility | CSR [47] | × | × | × | × | × | × | × |
| Balance, strength, and gait | SPPB [46] | × | × | × | × | × | × | × |
| Walking distance | 2MWT [49] | × | × | × | × | × | × | × |
| Body composition | BIA (Bodystat Quadscan 4000) [50] | × | × | × | × | × | × | × |
| Activity tracker | Fitbit Flex [51] | × | × | × | × | × | × | × |
| | Question or instrument | H ₁ | H ₂ /H ₃ | P ₁ | P ₂ | P ₃ | | |
| 4. Blood parameters | | | | | | | | |
| Inflammation markers | CRP [52] | × | × | | | | | |
| | WBC diff | × | × | | | | | |
| | TNF-α [53–55] | × | × | | | | | |
| | IL-6 [53–55] | × | × | | | | | |
| | IL-8 [55] | × | × | | | | | |
| Mortality | Date of death | | | | | | | × |

Note: H₁ = within 48 h after admission; H₂ = during hospitalization on Monday, Wednesday, and/or Friday; H₃ = at discharge; P₁ = one month post-discharge (home visit); P₂ = two months post-discharge (by telephone); P₃ = three months post-discharge (home visit);

×* = Data will be obtained from medical record;

CCI Charlson Comorbidity Index, MEWS Modified Early Warning Score, MMSE Mini Mental State Examination, CAM Confusion Assessment Method, NRS Numeric Rating Scale, STAI-6 State Trait Anxiety Inventory-6, GDS-15 Geriatric Depression Scale-15, ALCOS-12 Algemene Competentie Schaal-12 (General Self-Efficacy Scale), EQ-5D EuroQol-5D, FAC Functional Ambulation Categories, MFIS-5 Modified Fatigue Impact Scale-5, PSQI Pittsburgh Sleep Quality Index, SNAQ Short Nutritional Assessment, DEMMI De Morton Mobility Index, CSR Chair Sit and Reach test, SPPB Short Physical Performance Battery, 2MWT 2 Minute Walking Test, BIA Bioelectrical Impedance Analysis, CRP C-Reactive Protein, WBC diff White Blood Cell Differential, TNF-α Tumor Necrosis Factor-α, IL-6 Interleukin-6, IL-8 Interleukin-8

diastolic blood pressure; 4] level of consciousness; 5] temperature, and; 6] oxygen saturation [22].

Personal interviews/self-report data

(2.1) Cognitive functioning

Cognitive impairments. The most commonly used Mini Mental State Examination (MMSE) will be

applied to classify the severity of a cognitive impairment. It is a validated 23-item screening of cognitive impairment. The MMSE consists of a series of questions and tests, which assess different mental abilities, including memory, attention, language, and planning. Cognitive impairment is

defined as a score of 23 or less on the MMSE [23].

Delirium. The Confusion Assessment Method (CAM) will be used to identify the presence of delirium. The CAM consists of four features: 1] acute onset and fluctuating course; 2] inattention; 3] disorganized thinking, and 4] altered level of consciousness. The diagnosis of delirium requires the presence of both features 1 and 2, and the presence of either feature 3 or 4 [24]. Furthermore, we want to assess the risk for developing delirium with the following statements of the Dutch Safety Management Programme (*Veiligheidsmanagementsysteem* (VMS)): 1] the patient needs help with self-care, 2] the patient has previously undergone a delirium, and 3] the patient has a cognitive impairment such as dementia [25, 26].

(2.2) Behavioral and psychosocial functioning
Fear of falling. A Numeric Rating Scale (NRS) will be applied to measure fear of falling, in which a participant selects a whole number (0–10 integers). Zero represents no fear of falling and ten the worst possible fear of falling.

Depression. The Geriatric Depression Scale-15 (GDS-15) will be used to measure symptoms of depression (Cronbach's $\alpha = 0.75$ [27]). The GDS-15 is a self-report scale of 15 items on a binary (yes/no) scale and assesses symptoms over the preceding week. The total score is the sum of the 15 items (range 0–15 points, higher scores indicating more depression). The following categories of the GDS-15 will be used: a score of 0 to 4 will be considered 'normal', a score of 5 to 8 a 'mild depression', 9 to 11 a 'moderate depression', and 12 to 15 a 'severe depression' [28].

Apathy. Three items of the GDS-15 will be used to measure apathy (sensitivity of 69 % and specificity of 85 % [29]). The three apathy items include the following questions: 1] "Do you prefer to stay at home, rather than going out and doing new things?"; 2] "Have you dropped many of your activities and interests?" and; 3] "Do you feel full of energy? Higher scores indicate more apathy. A score of ≥ 2 points is indicative for apathy [29].

Anxiety. The State-Trait Anxiety Inventory-6 (STAI-6) will be used to measure anxiety symptoms (Cronbach's $\alpha = 0.79-0.81$ [30]). The STAI-6 is a short-form of the 20-item state scale of the Spielberger State-Trait Anxiety Inventory (STAI) [31], that maintains results that are comparable with this full-form [30]. It consists of six items on a 4-point Likert scale (1] not at all/almost never; 2] somewhat/sometimes; 3] moderately so/often, and; 4] very much so/almost always). Furthermore, it remains sensitive to different levels of anxiety.

Perceived self-efficacy. The General Self Efficacy Scale (In Dutch: *Algemene Competentie Schaal* (ALCOS-12)) will be used to measure general perceived self-efficacy (Cronbach's $\alpha = 0.78$ [32]). It is based on the Self-Efficacy Scale [33] and is a Dutch translated self-report rating scale of 12 items on a 5 point Likert scale (1] strongly disagree; 2] disagree; 3] no disagreement/agreement; 4] agree and; 5] strongly agree). The ALCOS-12 includes three subscales: competence (Cronbach's $\alpha = 0.72$), perseverance in adversity (Cronbach's $\alpha = 0.67$), and taking initiative (Cronbach's $\alpha = 0.74$) [32]. The total score is the sum of the 12 items (range 12–60), whereby the following categories of the ALCOS-12 will be used: a score of 12 to 38 will be defined as a 'low competence level', a score of 39 to 54 as 'average' and 55 to 60 as 'high' [34].

Health-Related Quality of life. The EuroQol-5D (EQ-5D), a widely used preference based generic health-related quality of life (HRQoL) instrument with well-established psychometric properties will be administered [20]. The EQ-5D consists of five dimensions: 1] mobility; 2] self-care; 3] usual activities; 4] pain/discomfort and; 5] anxiety/depression. These dimensions have three response choices (no problems; some problems or; severe problems). Moreover, the following questions will be used to measure quality of life: 1] "In general, how is your quality of life (participants answer the item with one of five possible responses: excellent; very good; good; moderate or; bad)?"; 2] "How would you grade your life at this moment, with a range between 0 and 10?" and; 3] "Compared to one year ago, how would you rate your health in general now (five response choices: much better; slightly better; much the same; slightly worse or; much worse)?" [18].

(2.3) Physical functioning

Dizziness, polyuria and shortness of breath. Symptoms of dizziness and shortness of breath will be assessed by asking: "Do you suffer from polyuria/dizziness/shortness of breath at this moment? If yes, does this affect your daily functioning?"

Pain. A gold standard of pain intensity measurements, the Numeric Rating Scale (NRS), will be applied to measure pain. The NRS for pain is a validated continuous scale with a score range between 0 and 10 (0 represents no pain and 10 the worst possible pain) [35, 36].

Fatigue. The Numeric Rating Scale (NRS), will be used to measure fatigue. The NRS for fatigue is a continuous scale with a score range between zero and ten (zero represents no pain and ten the worst possible fatigue) [37].

Impact of fatigue. The abbreviated version of the 21-item Modified Fatigue Impact Scale (MFIS) will be used to quantify the impact of fatigue. The short version consists of five items that are divided into three subscales: physical- (2 items), cognitive- (2 items), and psychosocial functioning (1 item) subscale. An example of a MFIS-5 statement is: "Because of my fatigue during the past four week, I have been less alert." The total score of the MFIS-5 is the sum of the raw scores on a 5-point Likert scale (0) never; 1] rarely; 2] sometimes; 3] often, and; 4] almost always). Higher scores indicate greater fatigue [38].

Sleep. The Pittsburgh Sleep Quality Index (PSQI) will be utilized to measure two components of sleep: sleep quality and sleep medication. Sleep quality will be quantified by asking: "During the past month, how would you rate your sleep quality overall?" Sleep medication will be measured by asking: "During the past month, how often have you taken medicine (prescribed or "over the counter") to help you sleep?" The score of sleep quality and sleep medication have a range of 0 (better) to 3 (worse) [39]. In addition, we measure daily sleepiness on a binary scale (yes/no) with the following question: "Do you currently suffer from daytime sleepiness? If yes, does this affect your daily living?"

Nutrition. The widely used Short Nutritional Assessment Questionnaire (SNAQ) will be applied to identify malnourished hospital patients (Cronbach's alpha = 0.58 [40]) [25, 26]. The total score of the SNAQ is the sum of the raw scores, whereby the following categories of the SNAQ will be used: a score of 0 to 1 will be defined as 'no malnutrition', a score of 2 as 'moderate malnutrition' and a score of 3 as 'severe malnutrition' [41].

ADL functioning. The 15 items modified Katz-ADL index will be used to measure physical functioning [16, 17]. The modified Katz-ADL index consists of statements of their independency in performing basic Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) (formulated in two versions on a binary (yes/no) scale: two weeks before admission or currently).

Mobility. The Functional Ambulation Categories (FAC) will be used to classify mobility, using six categories: a category of 1 will be defined as 'independent unlimited', a category of 2 as 'independent limited' and categories 3 to 5 as 'dependent'. Allocation to these last categories is based on levels of assistance and supervision needed [42]. Furthermore, we will measure mobility with two questions in according to the Comprehensive Geriatric Assessment (CGA) of the Dutch Society of Clinical Geriatrics (NVKG, 2012): 1] "Were you able to walk outside the house for five

minutes (formulated in two versions: two weeks before admission or currently)?", and; 2] "How often did/do you perform physical activity two weeks before admission/currently [19]?"

Falls. To measure the number of falls in the past (six) month(s) the following question of the VMS will be used: "Have you fallen once or more in the past (six) month(s)? If yes, how many times [25, 26]?" (2.4) Health care utilization

(Re)admission(s). Any (re)admission(s) to the hospital will be measured. We will search the medical record for (re)admission(s) in the same hospital six months before hospitalization and during three months post-discharge, and we will also retrieve this information by self-report at P₁-P₃ with the following self-report question: "Have you been hospitalized in the last month? If yes, for how many days [18]?" Data that will be collected out of the hospital system are: date of admission and discharge for any readmission, whether the admission was planned or unplanned and the reason for the readmission.

Nursing home admission(s). The amount of nursing home admission or whether they were admitted to the nursing home and the length of stay will be measured with the subsequent question: "Have you had a nursing home admission in the last month? If yes, for how many weeks totally [18]?"

Consult of physical therapist and/or occupational therapist. The amount of consults of a physiotherapist and/or occupational therapist will be measured by asking: "Have you had a consultation with your physical therapist and/or occupational therapist in the last month? If yes, how many times?"

Consult general practitioner. The amount of consults of a general practitioner will be measured by asking: "Have you had a consultation with your general practitioner in the last month? If yes, in the evening, night or weekend and how many times in total [19]?"

Home care. The use of home care will be measured with the subsequent question: "Do you use home care? If yes, care assistance and/or domestic help and how many hours per week [19]?" A distinction will be made between household help from a nursing aid, and help from a registered nurse.

(2) Physical performance tests

Handgrip strength. The hand grip strength will be measured with the widely used Jamar® grip strength dynamometer (Lafayette Instrument Company, USA). The handgrip strength test is used to provide an objective index of general upper body strength. Handgrip strength is a reliable instrument (good to excellent test-retest reproducibility and excellent inter-rater reliability) to indicate skeletal muscle mass [43]. Participants will perform the task thrice with

each hand. The highest score from either hand will be used and registered in the eCRF. Normative values of adults are described in a study of Mathiowetz [44].

Mobility. To measure the mobility we will use the 15-item Morton Mobility Index (DEMMI). Subjects will be asked to perform several mobility tasks, in the order of bed, chair, stand, and walking activities to maximize patient safety, which will result in an ordinal raw score (range: 0–19). The ordinal raw score will be converted into a total interval DEMMI score (range: 0 to 100 points). Moreover, the DEMMI has a hierarchical structure, and thus each assessed participant can be evaluated. Higher scores indicate a better mobility performance [45].

Balance, strength, and gait speed. The Short Physical Performance Battery (SPPB) will be applied to measure the balance, strength and gait speed. Participants will be asked to stand with their feet in various balance positions, walk a distance of four meter and to rise from a chair and return to the seated position five times as quickly as possible. Higher scores indicate a better performance [46].

Back and hamstring flexibility. The Chair Sit and Reach (CSR) test will be used as a measure of flexibility. Participant will be asked to extend one leg as straight as possible, hands on top of each other, and then to reach to his/her foot as far as possible. Lower distances between the tip of his/her toes and their extended fingers indicating a higher back and hamstring flexibility [47, 48].

Walking distance. The 2 Minute Walking Test (2MWT) will be applied to measure the maximal walking distance in meters. Participants will be asked to walk back and forth along a premeasured corridor of 15 meter in two minutes. Longer walking distances indicating a better walking capacity [49].

Body composition. The Bioelectrical Impedance Analysis (BIA) (Bodystat Quadscan 4000) will be used as method for estimating body composition, in particular fat-free mass (FFM) and high fat mass (FM). Electrodes will be attached to the ankle and wrist. A small electric signal will circulate, which measures the resistance and reactance of this electrical signal in the human body [50].

Activity level. The Fitbit Flex will be applied to monitor the sleep quality, measure motion patterns, determine the calories burned, distance traveled, and steps taken [51]. Participants will be asked to wear the Fitbit Flex from hospital admission up to one and a half weeks post-discharge.

(3) Blood parameters

Inflammation markers. Inflammation markers, such as C-Reactive Protein (CRP) [52], Tumor Necrosis

Factor- α (TNF- α), the interleukins IL-6 [53–55] and IL-8) [55], and White Blood Cell Differential (WBC diff), will be determined from blood plasma and serum. Blood will be collected during the customary laboratory rounds during hospitalization. Venous blood will be collected in 4.5 ml EDTA and serum vacutainers. Samples will be centrifuged and stored at $-80\text{ }^{\circ}\text{C}$ until analysis. Sample handling and analyses will be performed according to ISO standards.

Planned statistical analyses

Data will be analyzed in accordance with the research questions outlined in the introduction, applying appropriate General Linear Models (e.g., linear regression, repeated measures ANOVA/ANCOVA) as well as log-linear models (e.g., logistic regression in case of binary outcomes). Mortality, a (censored) numerical outcome, will be tested using survival analysis. The global α level will be set at 0.05 with hypothesis-wise adjustment for multiple testing. All analyses will be performed using SPSS version 22.0 [56]. Castor Electronic Data Capture (EDC) will be used to build electronic Case Report Forms (eCRFs) for save and valid data collection.

Primary endpoint in the study will be HAD as measured with the Katz-ADL index score. For multivariable analyses (General Linear models and log-linear models) a custom 10:1 case-to-outcome ratio is utilized as a maximum. Utilizing a repeated measures design, power calculations imputing a conservative α level of 0.01 yielded a power of 95 % for associations of a small effect-size (Cohen's $f = 0.069$), whereas a power of 80 % was established for associations with an effect-size of 0.058 (Cohen's f).

Discussion

More than 30 % of the older patients experience hospitalization-associated disability (HAD) after acute hospitalization [1, 3, 4], which implies the loss of ability to perform one or more of the basic ADLs [6]. HAD is the leading cause of functional decline at older age [4]. With a higher number of older persons and an increasing life expectancy, there is an urgent need to unravel the potential mechanisms behind HAD as well as how the mechanisms can be influenced. To our knowledge, the Hospital-ADL study is the first study that investigates cognitive, behavioral, psychosocial, physical, and biological factors simultaneously. The current study will provide novel information regarding possible underlying mechanisms behind HAD within the critical period of three months post hospitalization, which is expected to lead to the development of interventions that can prevent or restore HAD.

Ethics approval and consent to participate

The study is approved by the Institutional Review board of the Academic Medical Center (AMC) in The Netherlands (Protocol ID: AMC2015_150). Written informed consent is obtained from all participants before inclusion. The research is performed according to the Dutch Medical Research Involving Human Subjects Act and principles of the Declaration of Helsinki (1964).

Abbreviations

2MWT: 2 min walking test; ADL: activities of daily living; ALCOS-12: *Algemene Competentie Schaal-12* (Perceived Self-Efficacy); BIA: bioelectrical impedance analysis; CAM: confusion assessment method; CCI: Charlson comorbidity index; CGA: comprehensive geriatric assessment; CRF: case report form; CRP: C-Reactive Protein; CSR: Chair Sit an Reach test; DEMMI: De Morton Mobility Index; EQ-5D: EuroQoL-5D; FAC: functional ambulation categories; GDS-15: geriatric depression scale-15; HAD: hospitalization-associated disability; Hospital-ADL study: Hospital-Associated Disability and impact on daily Life study; IADL: instrumental activities of daily living; IL-8: interleukin-8; IL-6: interleukin-6; MDS: minimal dataset; MEWS: modified early warning score; MFIS-5: modified fatigue impact scale-5; MMSE: mini mental state examination; NRS: numeric rating scale; PSQI: Pittsburgh Sleep Quality Index; SNAQ: short nutritional assessment; SPPB: short physical performance battery; STAI-6: state trait anxiety inventory-6; TCB: transitional care bridge; TNF- α : tumor necrosis factor- α ; VMS: safety management programme (In Dutch: *Veiligheidsmanagementsysteem*); WBC diff: white blood cell differential.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

BMB serves as study Principal Investigator (PI), designed the study, and helped draft the manuscript. BMB wrote and reviewed the research proposal that was sent to the funding organization. LAR drafted the manuscript and wrote the protocol for the Medical Ethics Committee. All authors critically reviewed the protocol for the Medical Ethics Committee. All authors are responsible for study management and co-ordination. BMB, LAR, JJA and RS are responsible for the data collection. LAR and JAB will contribute to the storage and processing of blood samples. All authors approved the final manuscript and critically revised the manuscript.

Acknowledgements

This work was supported by the Netherlands Organization for Health Research and Development (NWO-ZonMw) [grant number 16156071].

Author details

¹Department of Internal Medicine, Section of Geriatric Medicine, Academic Medical Center, PO Box 226001100 DD Amsterdam, The Netherlands. ²Department of Rehabilitation, Academic Medical Center, Amsterdam, The Netherlands. ³Amsterdam Center for Innovative Health Practice (ACHIEVE), Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands. ⁴Department of Clinical Psychology, University of Amsterdam, Amsterdam, The Netherlands.

Received: 5 October 2015 Accepted: 25 February 2016

Published online: 05 March 2016

References

- Buurman BM, Hoogerduijn JG, de Haan RJ, Abu-Hanna A, Lagaay AM, Verhaar HJ, Schuurmans MJ, Levi M, de Rooij SE. Geriatric conditions in acutely hospitalized older patients: prevalence and one-year survival and functional decline. *PLoS ONE*. 2011;6(11), e26951.
- Boyd CM, Ricks M, Fried LP, Guralnik JM, Xue QL, Bandeen-Roche K. Functional Decline and Recovery of Activities of Daily Living among Hospitalized, Disabled Older Women: The Women's Health and Aging Study I. *J Am Geriatr Soc*. 2009;57(10):1757–66.
- Boyd CM, Landefeld CS, Counsell SR, Palmer RM, Fortinsky RH, Kresevic D, Burant C, Covinsky KE. Recovery of activities of daily living in older adults after hospitalization for acute medical illness. *J Am Geriatr Soc*. 2008;56(12):2171–9.
- Gill TM, Allore HG, Gahbauer EA, Murphy TE. Change in disability after hospitalization or restricted activity in older persons. *JAMA*. 2010;304(17):1919–28.
- Covinsky KE, Palmer RM, Fortinsky RH, Counsell SR, Stewart AL, Kresevic D, Burant CJ, Landefeld CS. Loss of independence in activities of daily living in older adults hospitalized with medical illnesses: increased vulnerability with age. *J Am Geriatr Soc*. 2003;51(4):451–8.
- Covinsky KE, Pierluissi E, Johnston CB. Hospitalization-associated disability: "She was probably able to ambulate, but I'm not sure". *JAMA*. 2011;306(16):1782–93.
- Buurman BM, De Rooij SE. Functieverlies ouderen bij acute opname in ziekenhuis. *Ned Tijdschr Geneesk*. 2015;159:A8185.
- Hardy SE, Dubin JA, Holford TR, Gill TM. Transitions between states of disability and independence among older persons. *Am J Epidemiol*. 2005;161(6):575–84.
- Gill TM, Allore HG, Holford TR, Guo Z. Hospitalization, restricted activity, and the development of disability among older persons. *JAMA*. 2004;292(17):2115–24.
- Kansagara D, Englander H, Salanitro A, Kagen D, Theobald C, Freeman M, Kripalani S. Risk prediction models for hospital readmission: a systematic review. *JAMA*. 2011;306(15):1688–98.
- Buurman BM, van Munster BC, Korevaar JC, Abu-Hanna A, Levi M, de Rooij SE. Prognostication in acutely admitted older patients by nurses and physicians. *J Gen Intern Med*. 2008;23(11):1883–9.
- Walter LC, Brand RJ, Counsell SR, Palmer RM, Landefeld CS, Fortinsky RH, Covinsky KE. Development and validation of a prognostic index for 1-year mortality in older adults after hospitalization. *JAMA*. 2001;285(23):2987–94.
- Inouye SK, Charpentier PA. Precipitating factors for delirium in hospitalized elderly persons. Predictive model and interrelationship with baseline vulnerability. *JAMA*. 1996;275(11):852–7.
- Sands LP, Yaffe K, Covinsky K, Chren MM, Counsell S, Palmer R, Landefeld CS. Cognitive screening predicts magnitude of functional recovery from admission to 3 months after discharge in hospitalized elders. *J Gerontol Ser A Biol Med Sci*. 2003;58(1):37–45.
- Sager MA, Franke T, Inouye SK, Landefeld CS, Morgan TM, Rudberg MA, Sebens H, Winograd CH. Functional outcomes of acute medical illness and hospitalization in older persons. *Arch Intern Med*. 1996;156(6):645–52.
- Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. *The Gerontologist*. 1970;10(1):20–30.
- Laan W, Zuithoff NP, Drubbel I, Bleijenberg N, Numans ME, de Wit NJ, Schuurmans MJ. Validity and reliability of the Katz-15 scale to measure unfavorable health outcomes in community-dwelling older people. *J Nutr Health Aging*. 2014;18(9):848–54.
- Minimale Data Set Zorgvrager Basismeting [http://topics-mds.eu/?page_id=366].
- Compleet Geriatrisch Assessment TZB [<http://www.effectieveouderenzorg.nl/zorgmodel/downloadformulieren.aspx>].
- Group EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy*. 1990;16(3):199–208.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–83.
- Subbe CP, Kruger M, Rutherford P, Gemmel L. Validation of a modified Early Warning Score in medical admissions. *QJM*. 2001;94(10):521–6.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189–98.
- Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegal AP, Horwitz RI. Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med*. 1990;113(12):941–8.
- VMS. Praktijkijds 'Kwetsbare Ouderen', VMS veiligheidsprogramma. 2009. 978-94-90101-04-6.
- Heim N, van Fenema EM, Weverling-Rijnsburger AW, Tuijl JP, Jue P, Oleksik AM, Verschuur MJ, Haverkamp JS, Blauw GJ, van der Mast RC et al. Optimal screening for increased risk for adverse outcomes in hospitalised older adults. *Age Ageing*. 2015;44(2):239–44.
- Friedman B, Heisel MJ, Delavan RL. Psychometric properties of the 15-item geriatric depression scale in functionally impaired, cognitively intact, community-dwelling elderly primary care patients. *J Am Geriatr Soc*. 2005;53(9):1570–6.
- Kok RM, Heeren TJ, van Hemert AM. De Geriatric Depression Scale. *Tijdschrift voor de Psychiatrie*. 1993;35:416–21.

29. van der Mast RC, Vinkers DJ, Stek ML, Bek MC, Westendorp RG, Gussekloo J, de Craen AJ. Vascular disease and apathy in old age. The Leiden 85-Plus Study. *Int J Geriatr Psychiatry*. 2008;23(3):266–71.
30. Tluczek A, Henriques JB, Brown RL. Support for the Reliability and Validity of a Six-Item State Anxiety Scale Derived From the State-Trait Anxiety Inventory. *J Nurs Meas*. 2009;17(1):19–28.
31. Chlan L, Savik K, Weinert C. Development of a shortened state anxiety scale from the Spielberger State-Trait Anxiety Inventory (STAI) for patients receiving mechanical ventilatory support. *J Nurs Meas*. 2003;11(3):283–93.
32. Bosscher RJ, Smit JH, Kempen GJM. Algemene competentieverwachtingen bij ouderen: Een onderzoek naar de psychometrische kenmerken van de Algemene Competentieschaal (ALCOS)/Global expectations of self-efficacy in the elderly: An investigation of psychometric characteristics of the General Self-Efficacy Scale. *Ned Tijdschr Psychol*. 1997;52:239–48.
33. Sherer M, Maddux JE, Mercandante B, Prentice-Dunn S, Jacobs B, Rogers RW. The self-efficacy scale : Construction and validation. *Psychol Rep*. 1982; 51:663–71.
34. Bosscher RJ, Laurijssen L, Boer E. Competence at later age: An explorative study. (Competentie op latere leeftijd: Een exploratieve studie). *Bewegen Hulpverlening*. 1992;9:225–65.
35. McCaffery M, Beebe A. Pain: clinical manual for nursing practice. C.V. Mosby: St. Louis; 1989.
36. Puntillo KA, Neighbor ML. Two methods of assessing pain intensity in English-speaking and Spanish-speaking emergency department patients. *J Emerg Nurs*. 1997;23(6):597–601.
37. Hwang SS, Chang VT, Cogswell J, Kasimis BS. Clinical relevance of fatigue levels in cancer patients at a Veterans Administration Medical Center. *Cancer*. 2002;94(9):2481–9.
38. Fisk JD, Ritvo PG, Ross L, Haase DA, Marrie TJ, Schlech WF. Measuring the functional impact of fatigue: initial validation of the fatigue impact scale. *Clin Infect Dis*. 1994;18 Suppl 1:S79–83.
39. Buysse DJ, Reynolds 3rd CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193–213.
40. Hanisah R, Suzana S, Lee FS. Validation of screening tools to assess appetite among geriatric patients. *J Nutr Health Aging*. 2012;16(7):660–5.
41. Kruijenga HM, Seidell JC, de Vet HC, Wierdsma NJ, van Bokhorst-de van der Schueren MA. Development and validation of a hospital screening tool for malnutrition: the short nutritional assessment questionnaire (SNAQ). *Clin Nutr (Edinburgh, Scotland)*. 2005;24(1):75–82.
42. Holden MK, Gill KM, Magliozzi MR, Nathan J, Piehl-Baker L. Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. *Phys Ther*. 1984;64(1):35–40.
43. Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, Sayer AA. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age Ageing*. 2011;40(4):423–9.
44. Mathiowetz V, Kashman N, Volland G, Weber K, Dowe M, Rogers S. Grip and pinch strength: normative data for adults. *Arch Phys Med Rehabil*. 1985; 66(2):69–74.
45. de Morton NA, Davidson M, Keating JL. The de Morton Mobility Index (DEMMI): an essential health index for an ageing world. *Health Qual Life Outcomes*. 2008;6:63.
46. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, Scherr PA, Wallace RB. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol*. 1994;49(2):M85–94.
47. Jones CJ, Rikli RE, Max J, Noffal G. The reliability and validity of a chair sit-and-reach test as a measure of hamstring flexibility in older adults. *Res Q Exerc Sport*. 1998;69(4):338–43.
48. Dewhurst S, Bampouras TM. Intraday reliability and sensitivity of four functional ability tests in older women. *Am J Phys Med Rehabil*. 2014;93(8):703–7.
49. Butland RJ, Pang J, Gross ER, Woodcock AA, Geddes DM. Two-, six-, and 12-minute walking tests in respiratory disease. *Br Med J (Clin Res Ed)*. 1982; 284(6329):1607–8.
50. Haverkort EB, Binnekade JM, de van der Schueren MA, Gouma DJ, de Haan RJ. Estimation of body composition depends on applied device in patients undergoing major abdominal surgery. *Nutr Clin Pract*. 2015; 30(2):249–56.
51. Bai Y, Welk GJ, Nam YH, Lee JA, Lee JM, Kim Y, Meier NF. Comparison of Consumer and Research Monitors under Semistructured Settings. *Medicine and science in sports and exercise*: Dixon PM; 2015.
52. Wium-Andersen M, Ørsted D, Nielsen S, Nordestgaard B. Elevated c-reactive protein levels, psychological distress, and depression in 73 131 individuals. *JAMA Psychiatry*. 2013;70(2):176–84.
53. Dantzer R. Cytokine, Sickness Behavior, and Depression. *Immunol Allergy Clin N Am*. 2009;29(2):247–64.
54. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci*. 2008;9(1):46–56.
55. Poon DC, Ho YS, Chiu K, Chang RC. Cytokines: how important are they in mediating sickness? *Neurosci Biobehav Rev*. 2013;37(1):1–10.
56. Corp I. IBM SPSS Statistics for Windows. 220th ed. Armonk: IBM Corp; 2013.
57. Verhage F. Intelligentie en leeftijd: Onderzoek bij Nederlanders van twaalf tot zeventenzeventig jaar. Proefschrift. Van Gorcum: Assen; 1964.
58. Heeren TJ, Kat MG, Stek ML. *Handboek ouderenspsychiatrie* (tweede druk). 2002. De Tijdstroom.
59. Timmerman H, de Groot JF, Hulzebos HJ, de Knikker R, Kerkkamp HE, van Meeteren NL. Feasibility and preliminary effectiveness of preoperative therapeutic exercise in patients with cancer: a pragmatic study. *Physiother Theory Pract*. 2011;27(2):117–24.
60. Trutschnigg B, Kilgour RD, Reinglas J, Rosenthal L, Hornby L, Morais JA, Viganò A. Precision and reliability of strength (Jamar vs. Biodex handgrip) and body composition (dual-energy X-ray absorptiometry vs. bioimpedance analysis) measurements in advanced cancer patients. *Appl Physiol Nutr Metab*. 2008;33(6):1232–9.
61. van Waart H, Stuiver MM, van Harten WH, Sonke GS, Aaronson NK. Design of the Physical exercise during Adjuvant Chemotherapy Effectiveness Study (PACES): a randomized controlled trial to evaluate effectiveness and cost-effectiveness of physical exercise in improving physical fitness and reducing fatigue. *BMC Cancer*. 2010;10:673.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

