



UvA-DARE (Digital Academic Repository)

Small RNAs from *Bemisia tabaci* Are Transferred to *Solanum lycopersicum* Phloem during Feeding

van Kleeff, P.J.M.; Galland, M.; Schuurink, R.C.; Bleeker, P.M.

DOI

[10.3389/fpls.2016.01759](https://doi.org/10.3389/fpls.2016.01759)

Publication date

2016

Document Version

Other version

Published in

Frontiers in Plant Science

License

CC BY

[Link to publication](#)

Citation for published version (APA):

van Kleeff, P. J. M., Galland, M., Schuurink, R. C., & Bleeker, P. M. (2016). Small RNAs from *Bemisia tabaci* Are Transferred to *Solanum lycopersicum* Phloem during Feeding. *Frontiers in Plant Science*, 7, [1759]. <https://doi.org/10.3389/fpls.2016.01759>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Table S1. Number of sRNAs at each analytical step. PC: Phloem control, PW: phloem from whitefly infested plants, LC: leaf control, LW: leaf infested with whiteflies, LE: leaf with eggs deposited on, WN: whitefly nymphs

	Redundant sRNAs			Non-redundant sRNAs		
	PC	PW	WN	PC	PW	WN
Original	5,370,176	7,321,768	31,231,948	3,027,013	4,113,001	17,186,095
After trimming (18 nts < sRNA < 40 nts)	1,999,661	2,958,626	-	1,305,216	1,857,261	-
After alignment to plant virus db	1,990,582	2,947,567	-	1,299,654	1,850,598	-
After alignment to RFAM (rRNA, tRNA, snoRNA, etc.)	1,742,553	2,600,385	-	1,161,605	1,658,597	-
After alignment to tomato mRNA	1,664,670	2,512,089	-	1,112,412	1,603,489	-
PW-PC	1,897,658		-	319,030		-
PW-PC intersect WN	4,346,653			144,646 (set #1)		

	Redundant sRNAs				Non-redundant sRNAs			
	LC	LW	LE	WN	LC	LW	LE	WN
Original	36,793,380	33,780,469	32,730,583	31,231,948	21,956,804	19,461,107	19,516,546	17,186,095
After alignment to spike ins	36,045,154	32,985,853	32,033,206	-	21,560,612	19,054,313	19,154,986	-
After trimming (18 nts < sRNA < 40 nts)	31,394,680	25,383,394	26,107,175	-	18,905,543	15,018,386	15,897,282	-
After alignment to plant virus db	31,319,721	25,310,008	26,031,234	-	18,861,008	14,976,128	15,852,319	-
After alignment to RFAM (rRNA, tRNA, snoRNA, etc.)	29,015,282	21,596,689	24,123,438	-	17,659,020	13,056,740	14,842,019	-
After alignment to tomato mRNA	27,286,742	20,247,471	22,661,674	-	16,685,149	12,297,418	14,004,694	-
LW-LC	9,236,781		-	-	1,587,686		-	-
(LW-LC) - LE	7,990,165			-	1,417,504			-
After alignment to tomato genome	912,215				176,462			
(LW-LC) - LE intersect WN	114,320				32,970 (set #2)			

	Non-redundant sRNAs
(PW-PC intersect WN) intersect [(LW-LC) - LE intersect WN] (set #1 intersect set #2)	10,505