



UvA-DARE (Digital Academic Repository)

Suppressor of fusion, a *Fusarium oxysporum* homolog of Ndt80, is required for nutrient-dependent regulation of anastomosis

Shahi, S.; Fokkens, L.; Houterman, P.M.; Rep, M.

DOI

[10.1016/j.fgb.2016.08.005](https://doi.org/10.1016/j.fgb.2016.08.005)

Publication date

2016

Document Version

Final published version

Published in

Fungal Genetics and Biology

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Shahi, S., Fokkens, L., Houterman, P. M., & Rep, M. (2016). Suppressor of fusion, a *Fusarium oxysporum* homolog of Ndt80, is required for nutrient-dependent regulation of anastomosis. *Fungal Genetics and Biology*, 95, 49-57.
<https://doi.org/10.1016/j.fgb.2016.08.005>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)



Regular Articles

Suppressor of fusion, a *Fusarium oxysporum* homolog of Ndt80, is required for nutrient-dependent regulation of anastomosis



Shermineh Shahi, Like Fokkens, Petra M. Houterman, Martijn Rep*

Molecular Plant Pathology, University of Amsterdam, Amsterdam, The Netherlands

ARTICLE INFO

Article history:

Received 31 January 2016

Revised 10 August 2016

Accepted 12 August 2016

Available online 13 August 2016

Keywords:

Anastomosis

Hyphal fusion

CAT fusion

Heterokaryon incompatibility

ABSTRACT

Heterokaryon formation is an essential step in asexual recombination in *Fusarium oxysporum*. Filamentous fungi have an elaborate nonself recognition machinery to prevent formation and proliferation of heterokaryotic cells, called heterokaryon incompatibility (HI). In *F. oxysporum* the regulation of this machinery is not well understood. In *Neurospora crassa*, Vib-1, a putative transcription factor of the p53-like Ndt80 family of transcription factors, has been identified as global regulator of HI. In this study we investigated the role of the *F. oxysporum* homolog of Vib-1, called Suf, in vegetative hyphal and conidial anastomosis tube (CAT) fusion and HI. We identified a novel function for an Ndt80 homolog as a nutrient-dependent regulator of anastomosis. Strains carrying the *SUF* deletion mutation display a hyper-fusion phenotype during vegetative growth as well as germling development. In addition, conidial pairing of incompatible *SUF* deletion strains led to more heterokaryon formation, which is independent of suppression of HI. Our data provides further proof for the divergence in the functions of different members Ndt80 family. We propose that Ndt80 homologs mediate responses to nutrient quality and quantity, with specific responses varying between species.

© 2016 Elsevier Inc. All rights reserved.

1. Introduction

Filamentous fungi grow by hyphal tip expansion and branching. Hyphae at the growing edge of the colony display a growth pattern described as avoidance, in which hyphal tips show negative autotropism towards each other (Prosser, 1995). Behind the growth front ongoing fusions, or anastomoses, between hyphae build a three dimensional network, the mycelium (Glass et al., 2000). In any given habitat different individuals not only of the same species but also of different species meet and can undergo vegetative hyphal fusion to form heterokaryons (Glass et al., 2000; Saupe, 2000).

In heterokaryotic cells one or more genetically distinct nuclei from each individual share a common cytoplasm (Glass et al., 2000). Although there are potential benefits to heterokaryon formation, such as functional polyploidy and mitotic genetic recombination, filamentous ascomycetes display an elaborate nonself recognition system, by which cells with genetically distinct nuclei are compartmentalized and subject to programmed cell death, a process referred to as heterokaryon incompatibility (HI) (Glass and Dementhon, 2006; Glass and Kaneko, 2003; Pontecorvo, 1956). It has been proposed that HI plays a role in restricting the

transmission of pathogenic elements such as double-stranded RNAs and the exploitation by aggressive genotypes (Cortesi et al., 2001; Debets and Griffiths, 1998).

Nonself recognition during HI is genetically regulated by allelic specification at *het* (for heterokaryon) loci and individuals that differ at one or more *het* loci are incompatible with each other (Glass et al., 2000; Saupe, 2000). In *Neurospora crassa* 11 unlinked *het* loci have been described to be involved in nonself recognition and HI, demonstrating the elaborate character of this immune system (Dementhon et al., 2006; Glass et al., 2000). The *N. crassa* *HET-c/PIN-c* system has been studied as a model and it has been shown that incompatible interactions at these loci lead to severe growth reduction, decreased conidiation, and programmed cell death of the fusion cell and surrounding cells (Glass and Dementhon, 2006; Glass and Kaneko, 2003; Glass et al., 2000). *het* loci have been shown to encode a variety of products, however the HET domain (Pfam PF06985) is conserved among proteins involved in HI in *N. crassa* and *Podospira anserina* (Espagne et al., 2002; Smith et al., 2000). Predicted HET domain genes are common in, and specific to, filamentous ascomycete and basidiomycete genomes (Fedorova et al., 2005; Van der Nest et al., 2014).

In the past decade, extensive studies have been carried out in *N. crassa* to elucidate molecular mechanisms and genetic regulation of HI. Vib-1 was first identified as a regulator for *HET-c* mediated HI. In incompatible interactions between strains carrying the

* Corresponding author.

E-mail address: m.rep@uva.nl (M. Rep).

VIB-1 deletion the phenotype associated with *HET-c/PIN-c* HI is suppressed (Xiang and Glass, 2002). Further investigation revealed that *Vib-1* is required for the expression of genes involved in HI, and these results collectively lead to the conclusion that *Vib-1* is a global mediator of HI in *Neurospora* (Dementhon et al., 2006; Xiang and Glass, 2002). Furthermore it was established that *Vib-1*, like its *A. nidulans* homolog *XprG*, is a positive regulator of extracellular protease production under carbon and nitrogen starvation (Dementhon et al., 2006; Katz et al., 2006). *VIB-1* encodes a putative transcription factor of the *p53*-like *Ndt80/PhoG* DNA binding family. It was shown that the *Ndt80/PhoG* DNA binding domain (Pfam PF05224) is conserved among ascomycete fungi (Dementhon et al., 2006; Hutchison and Glass, 2010; Katz et al., 2013; Lamoureux et al., 2002; Montano et al., 2002; Sopko and Stuart, 2004).

The filamentous ascomycete *Fusarium oxysporum* is an important plant pathogen and has been studied as a model organism for plant microbe interactions (for reviews see (de Sain and Rep, 2015; Takken and Rep, 2010). Although no apparent sexual recombination is known, the genomes of *F. oxysporum* species show a high degree of genetic variability (Rep and Kistler, 2010). Additional comparative genome studies have demonstrated that *Fusarium* species share a conserved core genome, however, *F. oxysporum* and *F. solani* also carry lineage-specific chromosomes (Coleman et al., 2009; Ma et al., 2010). In the case of *F. oxysporum* it was demonstrated that small lineage specific chromosomes harboring essential virulence genes, also known as pathogenicity chromosomes, can be horizontally transferred to a non-pathogenic strain, and thereby conferring pathogenicity towards a specific host (Ma et al., 2010). We have recently described a mechanism in *F. oxysporum* that is likely to enable horizontal chromosome transfer under nutrient-limiting conditions (Shahi et al., 2016). Although this study has added to our knowledge of the mechanics of asexual recombination, the genetic regulation as well as the molecular mechanics of nonself recognition and HI in *F. oxysporum* and their suppression during CAT fusion remain unclear. Given the similarity in phenotypes it was suggested that HI might share a common machinery between different fungal systems (Glass and Kaneko, 2003). The reference *F. oxysporum* genome encodes around 80 putative HET domain proteins (<http://www.fungidb.org>), yet, to our knowledge, none has been characterized. In this species vegetative compatibility groups (VCG) are determined using nitrate non-utilizing (NIT) mutants (Correll et al., 1987; Puhalla, 1985).

Here we investigate the *F. oxysporum* *Vib-1* homolog, encoded by *FOXG_01644* (*f. sp. lycopersici*) and *FOMG_05487* (*f. sp. melonis*), for its role in vegetative fusion and nonself recognition. We show that the *F. oxysporum* homolog of *Vib-1* is a nutrient-dependent negative regulator of hyphal and CAT fusion. In addition, conidial pairing of strains carrying a deletion mutant yielded higher numbers of heterokaryotic cells. However, in contrast to *N. crassa*, the deletion mutant does not overcome HI nor is secretion of proteases affected. We suggest that the increase in heterokaryon formation is a result of increased CAT fusion and propose to rename *FOXG_01644/FOMG_05487* suppressor of fusion (*SUF*).

2. Material and methods

2.1. Strains and culture conditions

Fusarium oxysporum f. sp. lycopersici strain 4287 (*Fol4287*, FGSC9935), *Fusarium oxysporum f. sp. melonis* strain 001 (*Fom001*, FGSC10441), and the non-pathogenic *Fusarium oxysporum* strain 47 (*Fo47*, FGSC10445) were used as the parental strains for fungal transformation. They were stored as a monoconidial culture at -80°C and revitalized on potato dextrose agar (PDA, Difco) at

25°C . *Agrobacterium tumefaciens* EHA105 (Hood et al., 1993) was used for *Agrobacterium*-mediated transformation of *F. oxysporum* and was grown in either Luria broth (LB) or 2YT medium (Sambrook and Russel, 2001) containing $20\ \mu\text{g/ml}$ rifampin at 28°C . Introduction of the plasmids into the *Agrobacterium* strain was performed as previously described (Mattanovich et al., 1989). *Escherichia coli* DH5 α (Invitrogen) was used for construction, propagation, and amplification of the plasmid and was grown at 37°C in LB medium containing $50\ \mu\text{g/ml}$ kanamycin.

2.2. Construction of vectors and *Agrobacterium*-mediated *Fusarium* transformation

2.2.1. *SUF* deletion

To generate *SUF* (*FOXG_01644*) deletion (Δ *suf*) mutants, we used plasmids pRW2h-H1-GFP and pRW2p-H1-RFP (Shahi et al., 2016, 2015). These plasmids have either the hygromycin resistance cassette (*hph*) and histone-GFP under the control of the xylanase promoter or the phleomycin resistance cassette (tested with zeocin, *ble*) and histone-RFP under the control of the xylanase promoter as selection marker. We introduced an $\sim 1\ \text{kb}$ sequence upstream (left flank) and downstream (right flank) of *SUF* open reading frame (ORF) for homologous recombination. For this we PCR amplified the left flank with primer combination FP4278 (5'-cggaattcGCCGTGCCGTTCACTCC-3') and FP4279 (5'-ggggtaccGCC TACGAAGTAGAAGATGAGCTT-3') and the right flank with primer combination FP4280 (5'-tgctctagaCGATATACAAAAGTTGCCAT CAAA-3') and FP4281 (5'-cccaagcttGAACAGATCTATTAACAAAGC CATCC-3'). The right flank was cloned in the *Xba*I site upstream the right border and the left flank in the *Pac*I site downstream the left border.

2.2.2. *SUF* complementation

Binary vectors pRW2h (*hph*) and pRW2p (*ble*) were used as a backbone for vector construction (Houterman et al., 2008; Shahi et al., 2016). We introduced an $\sim 1\ \text{kb}$ sequence upstream (left flank) plus the ORF and an $\sim 1\ \text{kb}$ sequence downstream (right flank) of *SUF* for homologous recombination. For this we PCR amplified the left flank plus *SUF* ORF with primer combination FP6455 (5'-ttaattaaGCCGTGCCGTTCACTC-3') and FP6456 (5'-ttaataaCTAGTTCCTCCATTCCAGGAATATTG-3') and the right flank as described before. The right flank was cloned in the *Xba*I site upstream the right border and the left flank plus *SUF* ORF in the *Pac*I site downstream the left border. pRW2h-*SUF* (*hph*) was used to complement H1-RFP Δ *suf* (*ble*) and pRW2p-*SUF* (*ble*) to complement H1-GFP Δ *suf* (*hph*, for plasmid maps see Fig. S1).

2.2.3. *Agrobacterium*-mediated *Fusarium* transformation

The obtained plasmids H1-GFP Δ *suf* (*hph*), H1-RFP Δ *suf* (*ble*), pRW2h-*SUF* (*hph*) and pRW2p-*SUF* (*ble*) were transformed into *Agrobacterium tumefaciens* EHA105 and the transformants were used for subsequent *A. tumefaciens*-mediated *Fusarium* transformation. *Agrobacterium*-mediated transformation of *F. oxysporum* was performed as previously described (Shahi et al., 2015).

2.3. CAT and hyphal fusion assays

For CAT fusion assay, conidia of *Fol4287* H1-RFP Δ *suf*, *Fom001* H1-GFP Δ *suf*, and *Fo47* H1-GFP Δ *suf* were collected from one-week-old PDA plates in 2 ml of the medium to be tested and filtered through one layer of sterile Miracloth (Calbiochem). $200\ \mu\text{l}$ 7.5×10^5 conidia per ml were incubated in an 8-well microscope chamber slide (Nunc) for 15–18 h in CAT medium and observed with the AMG Evos FL digital inverted microscope. 1000–2500 conidia were counted in two biological replicates. CAT fusion

frequency was calculated as the percentage of CAT fusions per germinated conidia.

For hyphal fusion assays, *Fol4287* H1-RFP Δ *suf*, *Fom001* H1-GFP Δ *suf*, and *Fo47* H1-GFP Δ *suf* were grown on PDA, CDA, or CAT medium supplemented with 1.5% agarose for one week at 25 °C. Side-to-side fusion between hyphae behind the growth front were counted per area (0.5 mm²). 15 technical in 3 biological replicates were performed.

2.4. Co-cultivation

Conidia of *Fol4287* wild type (wt), *Fol4287* H1-RFP Δ *suf*, *Fom001* wt, and *Fom001* H1-GFP Δ *suf* were collected from one-week-old PDA plates in 2 ml water, filtered through one layer of sterile Mira cloth (Calbiochem), and washed with water. To select heterokaryotic cells, 1 ml with 10⁶ conidia from each parental strain were incubated in a 1-well microscope chamber slide (Nunc). After two days conidia were collected and 50 μ l was plated on PDA and incubated for additional two days. Again conidia were collected and washed and 10⁴ conidia per ml were incubated on PDA buffered with 0.1 M Tris (pH 8.0) and supplemented with 100 μ g/ml hygromycin (Duchefa) and 100 μ g/ml zeocin (Invivo-gen) for five days. The number of double drug-resistant colonies was counted.

2.5. Selection of Nit mutations and complementation testing

The screen was performed as previously described (Puhalla, 1985). We selected different nitrate non-utilizing (NIT) mutants for each strain (*Fol4287* wt, *Fol4287* H1-RFP Δ *suf*, *Fom001* wt, and *Fom001* H1-GFP Δ *suf*) and tested compatibility in different combinations.

2.6. Analysis of culture supernatant

Conidia of *Fol4287* wt, *Fol4287* H1-RFP Δ *suf*, *Fom001* wt, and *Fom001* H1-GFP Δ *suf* were collected from 3-day old liquid culture, filtered through one layer of sterile Mira cloth (Calbiochem), and washed with water. Conidia concentration was adjusted to 10⁸ conidia/ml. 20 ml NO₃ (0.17% YNB, 100 mM KNO₃, 3% sucrose) were inoculated with 5 \times 10⁷ conidia and grown at 25 °C shaking. After 2 days the mycelium was pelleted and washed with CAT medium. One pellet each was transferred to 20 ml NO₃, minimal medium (MM, 0.17% YNB, 100 mM KNO₃), CAT, and CAT +3% sucrose media, with (induction for protease activity assay) or without (for SDS-PAGE) 1% BSA and incubated for another 2 days. The culture was again filtered through one layer of sterile Mira cloth and the filtrate was centrifuged. The supernatant used for further testing. The mycelial pellet was dried and the dry weight was measured. Protease activity in the culture supernatant was determined by measuring the release of the trichloroacetic acid (TCA)-soluble orange sulfanilamide component of azocasein upon proteolysis, as previously described (Dementhon et al., 2006).

Further, SDS-PAGE was conducted to visualize protein patterns of culture supernatant. For this 20 μ l of culture supernatant was loaded on a 15% sodium dodecyl sulfate (SDS) polyacrylamide gel followed by protein silver staining (Shevchenko et al., 1996).

2.7. Carbon utilization assay

To analyze carbon utilization of the *Fol4287* *SUF* deletion mutant, BIOLOG FF MicroPlates containing in each well a different carbon source were used, as described (Michielse et al., 2009).

2.8. Phylogenetic analysis

Complete proteomes of *F. oxysporum* f. sp. *lycopersici* 4287, *F. oxysporum* f. sp. *melonis* 26406, *F. graminearum* PH1, *Magnaporthe oryzae* 70-15 and *N. crassa* OR74a were obtained via the Broad institute and the genome of *Aspergillus nidulans* was obtained from GenBank. Proteins that have a Ndt80 domain were identified by searching all proteomes with a hmmer model for PF05224 (Finn et al., 2014) using hmmsearch from the Hmmer package (Eddy, 2009). The domain sequences were cut out using a custom python script and a multiple sequence alignment for the domain sequences was constructed using Clustal Omega with default settings (Sievers et al., 2011). The alignment was inspected but no changes were performed. The alignment was trimmed using trimAl (-strictplus) (Capella-Gutiérrez et al., 2009). Finally, PhyML (Guindon et al., 2010) with 4 substitution rate categories and estimated proportion of invariable sites and gamma distribution was used to infer the phylogeny.

3. Results

3.1. *Fusarium oxysporum* has four Ndt80 homologs

It was previously shown that the number of Ndt80 homologs can vary between different filamentous ascomycetes (Dementhon et al., 2006; Katz and Cooper, 2015). The reference genome of *F. oxysporum* encodes four predicted proteins with a Ndt80/PhoG binding domain that fall into three clades described earlier, a Vib-1 (vegetative incompatibility blocked, NCU03725) clade, a Fsd-1 (female sexual development, NCU09915) clade, and a NCU04729 clade (Dementhon et al., 2006). *VIB-1* and *NCU04729* each have one ortholog in *F. oxysporum*. Interestingly, even though a sexual cycle is not known for *F. oxysporum*, there is a duplication of *FSD-1* in *F. oxysporum* but not *F. graminearum* (Fig. 1). With the aim to better understand the molecular underpinnings of vegetative fusion and nonself recognition in *F. oxysporum*, we decided to investigate the role of *N. crassa* Vib-1 homolog *Suf*, encoded by *FOXG_01644* and *FOMG_05487*, in these processes.

3.2. *SUF* deletion strains display a hyper-fusion phenotype under nutrient-limiting conditions

To understand the role of *F. oxysporum* homologs of Vib-1, we first studied the phenotype of *SUF* deletion mutants. We used a construct based on the flanking regions of *F. oxysporum* f. sp. *lycopersici* strain 4287 (*Fol4287*) *FOXG_01644* to obtain deletion mutants in *Fol4287*, *F. oxysporum* f. sp. *melonis* strain 001 (*Fom001*), and *F. oxysporum* strain 47 (*Fo47*). None of the deletion mutants showed altered colony morphology or conidiation (data not shown). With around 6.5 \times 10⁶ conidia/ml the number of conidia produced after three days incubation in NO₃ medium was similar in all strains.

We next investigated the microscopic phenotype of the deletion mutants. For this we grew the wild type strain, two independent *SUF* deletion (Δ *suf*) mutants, and a complemented deletion strain from each background on PDA, CDA, and CAT medium supplemented with 1.5% agarose. PDA is rich in various carbon and nitrogen sources, CDA offers sucrose as the sole carbon and nitrate as the sole nitrogen source, and CAT medium offers no carbon source and limited nitrate (25 mM). Strains grown on PDA did not show any morphological differences between deletion mutant and wild type strains. However, under nutrient-limiting conditions (CDA and CAT medium), hyphae of the *SUF* deletion mutants of *Fol4287* and *Fom001*, but not *Fo47* *SUF* deletion strains nor the wild type, exhibited a strong increase in side-to-side fusions.




Fig. 1. Phylogenetic tree displaying the relationship of *F. oxysporum* Ndt80 homologs to homologs in several other ascomycetes. The tree is based on alignments of Ndt80/PhoG DNA binding domains (PF05224). Each colored circle represents a species. The appearance of the branch illustrates bootstrap values: grey – bootstrap values 50–80, black – bootstrap values 80–90, bold black: bootstrap values >90. Scale bar indicates number of substitutions per amino acid site. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Complemented strains, in which the *SUF* gene was reintroduced in *locus*, displayed a similar phenotype as the wild type strains (Fig. 2). Interestingly, not all hyphae that came into close proximity showed side-to-side fusion nor did we observe a specific pattern, e.g. in the distance between the fusions.

We examined 15 areas of 0.5 mm² in three biological replicates. In PDA, all strains showed very little or no hyphal fusion (Fig. 2a, PDA). In CDA, *Fol4287* *SUF* deletion mutants peaked at 8 and 10 fusions/area, and *Fom001* *SUF* deletion mutants at 6 and 7 fusions/area (Fig. 2a, CDA). In CAT medium the situation was reversed: *Fol4287* mutants showed lower numbers of fusion/area (2 and 4) than *Fom001* mutants (9 and 10 fusions/area, Fig. 2a,

CAT). *Fo47* did not show any differences in number of fusions between the different media – the number was low in all cases. This hyper-fusion phenotype is distinct from what has been described for any of the Ndt80 homologs so far (Chu and Herskowitz, 1998; Dementhon et al., 2006; Katz and Cooper, 2015; Xiong et al., 2014).

We next decided to investigate another type of fusion, CAT (for conidial anastomosis tube) fusion. In contrast to vegetative hyphal fusion that can occur throughout the colony, CAT fusion is restricted to the developmental stage of colony initiation (Roca et al., 2005a, 2005b). We have observed that CAT fusion in *F. oxysporum* is restricted to carbon starvation and nitrogen limitation and obtained the highest frequency of CAT fusion in a medium with no carbon source and 25 mM nitrate, thus naming this medium CAT medium (Shahi et al., 2016). To test whether *SUF* deletion also has an effect on CAT fusion, we incubated spores of each strain (wild type, two independent deletion mutants, and a complemented strain in each background) in CAT medium for 18 h. We observed that *Fol4287* *SUF* deletion mutants exhibit not only an increase CAT fusion, but also an increase in number of conidia that are interconnected (Fig. 3). In the wild type strain we mostly found two conidia and to a much lesser extent three conidia that are connected, whereas in the *SUF* deletion mutant more than five conidia could be part of an interconnected network (Fig. 3b). We calculated the percentage of CAT fusions connecting 2, 3, 4, 5 and >5 conidia per germinated conidia based on two biological replicates and 1000–2500 conidia. Overall, with *Fol4287* *SUF* deletion strains the percentage of CAT fusions doubled compared to wild type. About half of the CAT fusions connected more than 3 conidia in the mutants. In the complemented strain the wild type phenotype was partially restored. The percentage of total CAT fusion was not much reduced, but the number of conidia that were connected decreased to wild type level. In *Fom001* *SUF* deletion mutants the phenotype was less severe. No increase in percentage of total CAT fusion was detected. However, there was a tendency that more than two conidia are connected through CAT fusions (Fig. 3a). *Fo47* did not show any CAT fusions and *SUF* deletion did not have an effect on this (Fig. 3). We decided to continue our further investigations with *Fol4287* and *Fom001*.

3.3. *SUF* deletion strains display increased heterokaryon formation but no suppression of HI-associated cell death

Under nutrient-limiting conditions CAT fusion between incompatible strains of *F. oxysporum* allows heterokaryon formation (Shahi et al., 2016). *N. crassa* Vib-1 has been described as a global regulator of HI (Dementhon et al., 2006). We investigated the effect of *SUF* deletion on heterokaryon formation between *Fol4287* and *Fom001* during conidial pairing. For this we co-cultivated *Fol4287* expressing phleomycin resistance and *Fom001* expressing hygromycin resistance and allowed for conidium formation. 10⁴ conidia emerging from the mixed incubation were plated on double-selection PDA plates (i.e. containing both hygromycin and zeocin) and the percentage of emerging colonies was calculated. Co-cultivation of *Fol4287* and *Fom001* wild type (at the *SUF* locus) yielded no double-selective colonies. We tested four different combinations of *Fol4287* Δ *suf* and *Fom001* Δ *suf* strains. In all combinations double drug-resistant colonies emerged: *Fol4287* Δ *suf* #2 and *Fom001* Δ *suf* #1 (9.4 ± 2.1%), *Fol4287* Δ *suf* #2 and *Fom001* Δ *suf* #5 (2.5 ± 1.3%), *Fol4287* Δ *suf* #4 and *Fom001* Δ *suf* #1 (14.1 ± 3.5%), and *Fol4287* Δ *suf* #4 and *Fom001* Δ *suf* #5 (1.4 ± 0.9%, average and standard deviation based on 10 replicates). This demonstrates that, similar to *N. crassa*, deletion of *SUF* increases heterokaryon formation.




Fig. 2. *SUF* deletion strains of *Fol4287* and *Fom001*, but not *Fo47*, display increased vegetative hyphal fusion. (a) In CDA and CAT medium, *Fol4287* and *Fom001* strains carrying the *SUF* deletion mutation exhibit higher numbers of hyphal fusion. This phenotype is reversed in the respective complementation strains. *Fo47* does not undergo vegetative hyphal fusion. Depicted are average and standard errors of hyphal fusion per 0.5 mm². The calculation is based on 15 areas from 3 biological replicates. An unpaired *t*-test between wild type and mutant strains was performed for each medium; * *p* < 0.05, ** *p* < 0.005, *** *p* < 0.0005. (b) *Fol4287* and *Fom001* phenotypes on CDA. Arrowheads mark side-to-side fusions. Scale bar: 100 μm.




Fig. 3. *Fol4287* *SUF* deletion strains exhibit higher fusion rates during germling development. (a) *Fol4287* *SUF* deletion strains grown in CAT medium show higher conidial anastomosis tube (CAT) fusion rates. In addition, the number of conidia that are connected increase. *Fom001* *SUF* deletion strains do not exhibit a significant difference to the wild type strain, although a tendency towards more connected conidia was observed. *Fo47* does not undergo CAT fusion. Presented are percentages of CAT fusions connecting 1, 2, 3, 4, 5, and >5 conidia per germinated spores. Calculations are based on 1000–2500 conidia and 2 biological replicates. An unpaired *t*-test comparing CAT fusions including more than two conidia between wild type and mutant strains was performed; * $p < 0.05$, ** $p < 0.0001$. (b) CAT fusion in *Fol4287* wild type (wt) and *SUF* deletion strain (Δ suf #2). Arrowheads mark CAT fusions. A letter indicates conidia that are interconnected. Scale bar: 100 μ m.

We next tested whether the increase in heterokaryon formation is caused by a general suppression of HI-mediated cell death, as was described for *N. crassa*. As mentioned earlier, the mechanisms underlying nonself recognition and HI are not well understood in *F. oxysporum*. In species in which allelic interactions of HI are not known, heterokaryon formation can be tested using nitrate non-utilizing (NIT) mutants. NIT mutants display thin growth on media containing nitrate as a sole nitrogen source. Strains with different NIT mutations are able to complement each other only when they are able to form heterokaryons

and are thus compatible with each other (Correll et al., 1987; Puhalla, 1985). Complementation is apparent from aerial hyphae formation where hyphae of the two strains meet. We used the VCG testing system to determine whether *SUF* deletion mutants can overcome HI during vegetative growth. For this we selected at least one NIT mutant per strain (*Fol4287* wt, *Fol4287* Δ suf #2, *Fom001* wt, and *Fom001* Δ suf #1, for phenotype on nitrate medium see Fig. S2). As was expected, the positive control, i.e. interaction between compatible strains or selfing, resulted in formation of aerial hyphae (Fig. 4, upper left panel). Neither




Fig. 4. *SUF* deletion does not overcome heterokaryon incompatibility. In *F. oxysporum* vegetative compatibility is tested using nitrate non-utilizing (NIT) mutants. Compatible strains carrying different NIT mutations can complement the NIT phenotype (thin growth on nitrate medium) resulting in formation of aerial hyphae. Pairing of complementing *Fol4287* and *Fom001* NIT mutants resulted in production of aerial hyphae (compatible interaction), whereas the pairing of *Fol4287* with *Fom001* did not complement the NIT phenotype (incompatible interaction). The same results were obtained in different pairings between *Fol4287* and *Fom001* *SUF* deletion strains (Δ suf interaction). a: *Fol4287* Nit1, b: *Fol4287* NitM, c: *Fol4287* Δ suf Nit1, d: *Fol4287* Δ suf NitM, e: *Fol4287* Δ suf Nit3, f: *Fom001* Nit1, h: *Fom001* Δ suf Nit1, i: *Fom001* Δ suf Nit3, j: *Fom001* Δ suf NitM.

the interaction between *Fol4287* and *Fom001* wild type nor *SUF* deletion strains resulted in formation of aerial hypha, thus marking these strains as incompatible (Fig. 4, upper right and bottom panel). We conclude that the increase in heterokaryon formation observed upon mixing of conidia is caused by the increase in CAT fusion and that, unlike *N. crassa* *Vib-1*, *Suf* is likely not involved in regulation of HI.

3.4. *Suf* is not important for secretion of proteases or for carbon utilization

Another phenotype that has been associated with *Ndt80* homologs *Vib-1* and *XprG* is absence of extracellular proteases in the culture medium in the respective deletion mutants. In *N. crassa* and *A. nidulans* extracellular protease production is induced upon carbon or nitrogen starvation (Dementhon et al., 2006; Katz et al., 2006). We examined culture supernatants for protease activity in different media (NO_3 , minimal medium (MM), CAT, and CAT +3% sucrose). Similar to *N. crassa* and *A. nidulans*, extracellular protease production is strongly induced under nutrient-limiting condition in *F. oxysporum* (MM and CAT). However, *SUF* deletion had no effect on protease production (Fig. 5). We also found no major differences in culture supernatant protein profiles using SDS-PAGE between wild type and *SUF* deletion mutants in any of the media (Fig. S3). We conclude that *Suf* does not play a major role in protease production in *F. oxysporum* in response to carbon or nitrogen starvation.

In another study a link between *Vib-1* and carbon utilization was demonstrated (Xiong et al., 2014). We analyzed carbon source utilization using BIOLOG FF MicroPlates, which contain a different carbon source in each of the 96 wells. The ratios between *Fol4287* *SUF* deletion strain and wild type strain growth rates were calculated. Values higher than 1.5 or lower than 0.5 are considered a substantial difference in growth rate (Michielse et al., 2009). In our assay, the *SUF* deletion strain performed similar to the wild type strain on all tested carbon sources (Fig. S4). Thus, *Suf* is not required for utilization of these carbon sources.

4. Discussion

In this study we present a novel role for the *F. oxysporum* *Ndt80* homolog suppressor of fusion (*Suf*) as a nutrient-dependent negative regulator of vegetative hyphal and CAT fusion. *Fol4287* and *Fom001* strains carrying a *SUF* deletion show increased vegetative hyphal fusion and in case of *Fol4287* also increased CAT fusion (Figs. 2 and 3). Genes negatively regulating fusion have rarely been characterized. To our knowledge only two such genes have been identified, *N. crassa* NCU006362, a predicted GTPase activating protein, and *SPR7*, a secreted serine protease (Palma-Guerrero et al., 2013). In both cases the deletion strain exhibits higher CAT fusion frequencies and higher numbers of connected conidia. It was suggested that both genes are involved in fungal communication prior to fusion (Palma-Guerrero et al., 2013).

Despite (almost) identical protein sequences of the *Suf* homologs in the three *F. oxysporum* strains analyzed in this study, the deletion strains show different phenotypes. We have previously observed that *Fo47* has a fusion defect and does not undergo vegetative fusion in form of hyphal or CAT fusion and confirmed this phenotype in this study (Shahi et al., 2016 and Figs. 2 and 3). *SUF* deletion does not overcome this fusion defect, indicating that either the two processes are unlinked or that the fusion defect of *Fo47* is caused by a protein downstream of *Suf*, maybe even a target of *Suf*. The differences in phenotype between *Fol4287* and *Fom001* *SUF* deletion strains could also be caused by differences in *Suf* targets.

We show that *SUF* deletion increases heterokaryon formation but does not suppress HI. The increase in heterokaryon formation could be caused by the observed increase in CAT fusion frequency. In other ascomycete fungi, *Ndt80* homologs play different roles. *Saccharomyces cerevisiae* *Ndt80*, the founder of this family of p53-like transcription factors, is a transcriptional activator of ~150 genes involved in completion of meiosis (Chu and Herskowitz, 1998). In *Candida albicans* one *Ndt80* homolog has been characterized, which is involved in drug resistance, biofilm formation, and virulence (Chen et al., 2004; Nobile et al., 2012; Sellam et al., 2010; Wang et al., 2006). The filamentous ascomycete *A. nidulans* has two *Ndt80* homologs. The one most similar to *Ndt80* is involved in sexual reproduction. The second, *XprG*, which is more




Fig. 5. In *F. oxysporum* the production of extracellular proteases in response to carbon and nitrogen starvation is independent of *Suf*. *Fol4287* and *Fom001* wild type and *SUF* deletion strains were assessed for extracellular protease activity. Protease activity was determined by measuring the release of the trichloroacetic acid (TCA)-soluble orange sulfanilamide component of azocasein upon proteolysis at 440 nm. Results are displayed as absorbance per gram dry weight. Measurements were done once, precluding calculation of standard deviation. Nevertheless, each pair of strains behaves similarly and it is clear that *F. oxysporum* extracellular protease production is, similar to *N. crassa* and *A. nidulans*, triggered by carbon source starvation (MM and CAT medium) but deletion of the *F. oxysporum* *NDT80* homolog *SUF* does not suppress the induction of protease production.

closely related to *N. crassa* Vib-1, is a transcriptional activator of a large number of genes in response to carbon limiting conditions (Katz and Cooper, 2015). *N. crassa* Vib-1 was characterized as a global regulator of heterokaryon incompatibility (HI) and a positive regulator of extracellular protease production under carbon and nitrogen starvation (Dementhon et al., 2006; Xiong et al., 2014). A second Ndt80 homolog in *N. crassa*, Fsd-1, is involved in female sexual development, a process that is initiated under nitrogen limiting conditions (Hutchison and Glass, 2010). The one apparent commonality across these diverse functions is response to nutrient-limiting conditions. Although this has not been tested for *N. crassa* Vib-1, in *P. anserina* HI is associated with the response to starvation (Pinan-Lucarré et al., 2005). It was suggested that nutrient sensing might represent one of the ancestral roles for Ndt80 family of proteins (Dementhon et al., 2006; Katz and Cooper, 2015; Katz et al., 2013). Our data provides further support for this hypothesis.

Phylogenetic analyses revealed that the Ndt80 DNA-binding domain (DBD) is conserved among ascomycete fungi, but the proteins have diverged outside this domain (Fig. 1). The question that arises is how these different Ndt80 homologs can respond to the same environmental cues. It is possible that the proteins are activated through a conserved mechanism despite sequence divergence. Alternatively, the expression of *NDT80* and its homologs could be triggered by nutritional cues. In *S. cerevisiae*, Ime1 and Ime2 are positive regulators of expression of *NDT80* (Winter, 2012). *N. crassa* and *A. nidulans* both have a homolog of Ime2, which have been shown to be negative regulators of Vib-1 and XprG, respectively (Hutchison and Glass, 2010; Hutchison et al., 2012; Katz and Cooper, 2015). It was suggested that the regulation of *NDT80* homologs by Ime2 homologs might be conserved among ascomycete fungi (Hutchison et al., 2012). *F. oxysporum* also has a *IME2* homolog, *FOXG_13813* (<http://www.fungidb.org>), and it will be interesting to study its function in regulation of *SUF*.

We have shown that the deletion mutant phenotype is triggered in CDA. CDA contains sucrose as sole carbon source and nitrate as sole nitrogen source but both in adequate quantities, thus the medium is not causing starvation. Apparently, then, it is the absence of complex compounds in CDA (present in PDA) that triggers increased hyphal fusion (in the absence of *Suf*). That may also hold true for other Ndt80 homologs, possibly regulated by Ime2 homologs. Depending on the fungal species, different developmental processes are then activated. In addition, conidial pairing of strains with a *SUF* deletion show an increase in heterokaryon formation, the first step towards horizontal chromosome transfer (Shahi et al., 2016).

Continued work will be required to elucidate the role of Ndt80 homologs in fungal development as well as asexual recombination and horizontal chromosome transfer in *F. oxysporum*. This might include the identification of downstream targets of *Suf* by transcriptome analysis, characterization of other *F. oxysporum* Ndt80 homologs and regulators of *Suf*.

Acknowledgements

This work was made possible by an Innovational Research Incentives Scheme Vici grant of The Netherlands Organisation for Scientific Research (NWO) to M.R. We thank Bas Beerens, Martin Bosch and Ido Vlaardingerbroek for their assistance with experiments.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.fgb.2016.08.005>.

References

- Capella-Gutiérrez, S., Silla-Martínez, J.M., Gabaldón, T., 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25, 1972–1973. <http://dx.doi.org/10.1093/bioinformatics/btp348>.
- Chen, C.-G., Yang, Y.-L., Shih, H.-I., Su, C.-L., Lo, H.-J., 2004. CaNdt80 is involved in drug resistance in *Candida albicans* by regulating CDRI1. *Antimicrob. Agents Chemother.* 48, 4505–4512. <http://dx.doi.org/10.1128/AAC.48.12.4505-4512.2004>.
- Chu, S., Herskowitz, I., 1998. Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. *Mol. Cell* 1, 685–696.
- Coleman, J.J., Rounsley, S.D., Rodriguez-Carres, M., Kuo, A., Wasmann, C.C., Grimwood, J., Schmutz, J., Taga, M., White, G.J., Zhou, S., Schwartz, D.C., Freitag, M., Ma, L.J., Danchin, E.G., Henrissat, B., Coutinho, P.M., Nelson, D.R., Straney, D., Napoli, C.A., Barker, B.M., Gribskov, M., Rep, M., Kroken, S., Molnár, L., Rensing, C., Kennell, J.C., Zamora, J., Farman, M.L., Selker, E.U., Salamov, A., Shapiro, H., Pangilinan, J., Lindquist, E., Lamers, C., Grigoriev, I.V., Geiser, D.M., Covert, S.F., Temporini, E., Vanetten, H.D., 2009. The genome of *Nectria haematococca*: contribution of supernumerary chromosomes to gene expansion. *PLoS Genet.* 5, e1000618. <http://dx.doi.org/10.1371/journal.pgen.1000618>.
- Correll, J.C., Klittich, C.J.R., Leslie, J.F., 1987. Nitrate nonutilizing mutants of *Fusarium oxysporum* and their use in vegetative compatibility tests. *Phytopathology* 77, 1640–1646.
- Cortesi, P., McCulloch, C.E., Song, H., Lin, H., Milgroom, M.G., 2001. Genetic control of horizontal virus transmission in the chestnut blight fungus, *Cryphonectria parasitica*. *Genetics* 159, 107–118.
- de Sain, M., Rep, M., 2015. The role of pathogen-secreted proteins in fungal vascular wilt diseases. *Int. J. Mol. Sci.* 16, 23970–23993. <http://dx.doi.org/10.3390/ijms161023970>.
- Debets, A.J.M., Griffiths, A.J.F., 1998. Polymorphism of het-genes prevents resource plundering in *Neurospora crassa*. *Mycol. Res.* 102, 1343–1349. <http://dx.doi.org/10.1017/S095375629800639X>.
- Dementhon, K., Iyer, G., Glass, N.L., 2006. Vib-1 is required for expression of genes necessary for programmed cell death in *Neurospora crassa*. *Eukaryot. Cell* 5, 2161–2173. <http://dx.doi.org/10.1128/EC.00253-06>.
- Eddy, S.R., 2009. A new generation of homology search tools based on probabilistic inference. *Genome Inform.* 23, 205–211.
- Espagne, E., Balhadère, P., Penin, M.-L., Barreau, C., Turcq, B., 2002. HET-E and HET-D belong to a new subfamily of WD40 proteins involved in vegetative incompatibility specificity in the fungus *Podospora anserina*. *Genetics* 161, 71–81.
- Fedorova, N.D., Badger, J.H., Robson, G.D., Wortman, J.R., Nierman, W.C., 2005. Comparative analysis of programmed cell death pathways in filamentous fungi. *BMC Genom.* 6, 177. <http://dx.doi.org/10.1186/1471-2164-6-177>.
- Finn, R.D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Heger, A., Hetherington, K., Holm, L., Mistry, J., Sonnhammer, E.L.L., Tate, J., Punta, M., 2014. Pfam: the protein families database. *Nucleic Acids Res.* 42, D222–D230. <http://dx.doi.org/10.1093/nar/gkt1223>.
- Glass, N.L., Dementhon, K., 2006. Non-self recognition and programmed cell death in filamentous fungi. *Curr. Opin. Microbiol.* 9, 553–558. <http://dx.doi.org/10.1016/j.mib.2006.09.001>.
- Glass, N.L., Jacobson, D.J., Shiu, P.K.T., 2000. The genetics of hyphal fusion and vegetative incompatibility in filamentous ascomycete fungi. *Annu. Rev. Genet.* 34, 165–186. <http://dx.doi.org/10.1146/annurev.genet.34.1.165>.
- Glass, N.L., Kaneko, I., 2003. Fatal attraction: nonself recognition and heterokaryon incompatibility in filamentous fungi. *Eukaryot. Cell* 2, 1–8.
- Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321. <http://dx.doi.org/10.1093/sysbio/syq010>.
- Hood, E.E., Gelvin, S.B., Melchers, S., Hoekema, A., 1993. New *Agrobacterium* helper plasmids for gene transfer to plants (EHA105). *Transgenic Res.* 2, 208–218.
- Houterman, P.M., Cornelissen, B.J., Rep, M., 2008. Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog.* 4, e1000061. <http://dx.doi.org/10.1371/journal.ppat.1000061>.
- Hutchison, E.A., Bueche, J.A., Glass, N.L., 2012. Diversification of a protein kinase cascade: IME-2 is involved in nonself recognition and programmed cell death in *Neurospora crassa*. *Genetics* 192, 467–482. <http://dx.doi.org/10.1534/genetics.112.142612>.
- Hutchison, E.A., Glass, N.L., 2010. Meiotic regulators Ndt80 and ime2 have different roles in *Saccharomyces* and *Neurospora*. *Genetics* 185, 1271–1282. <http://dx.doi.org/10.1534/genetics.110.117184>.
- Katz, M.E., Braunberger, K., Yi, G., Cooper, S., Nonhebel, H.M., Gondro, C., 2013. A p53-like transcription factor similar to Ndt80 controls the response to nutrient stress in the filamentous fungus, *Aspergillus nidulans*. *F1000Research* 2, 72. <http://dx.doi.org/10.12688/f1000research.2-72.v1>.
- Katz, M.E., Cooper, S., 2015. Extreme diversity in the regulation of Ndt80-like transcription factors in fungi. *G3 (Bethesda)* 5, 2783–2792. <http://dx.doi.org/10.1534/g3.115.021378>.
- Katz, M.E., Gray, K.-A., Cheetham, B.F., 2006. The *Aspergillus nidulans* xprG (phoG) gene encodes a putative transcriptional activator involved in the response to nutrient limitation. *Fungal Genet. Biol.* 43, 190–199. <http://dx.doi.org/10.1016/j.fgb.2005.12.001>.

- Lamoureux, J.S., Stuart, D., Tsang, R., Wu, C., Glover, J.N.M., 2002. Structure of the sporulation-specific transcription factor Ndt80 bound to DNA. *EMBO J.* 21, 5721–5732.
- Ma, L.J., van der Does, H.C., Borkovich, K.A., Coleman, J.J., Daboussi, M.J., Di Pietro, A., Dufresne, M., Freitag, M., Grabherr, M., Henrissat, B., Houterman, P.M., Kang, S., Shim, W.B., Woloshuk, C., Xie, X., Xu, J.R., Antoniw, J., Baker, S.E., Bluhm, B.H., Breakspear, A., Brown, D.W., Butchko, R.A., Chapman, S., Coulson, R., Coutinho, P.M., Danchin, E.G., Diener, A., Gale, L.R., Gardiner, D.M., Goff, S., Hammond-Kosack, K.E., Hilburn, K., Hua-Van, A., Jonkers, W., Kazan, K., Kodira, C.D., Koehrsen, M., Kumar, L., Lee, Y.H., Li, L., Manners, J.M., Miranda-Saavedra, D., Mukherjee, M., Park, G., Park, J., Park, S.Y., Proctor, R.H., Regev, A., Ruiz-Roldan, M.C., Sain, D., Sakthikumar, S., Sykes, S., Schwartz, D.C., Turgeon, B.G., Wapinski, I., Yoder, O., Young, S., Zeng, Q., Zhou, S., Galagan, J., Cuomo, C.A., Kistler, H.C., Rep, M., 2010. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464, 367–373. <http://dx.doi.org/10.1038/nature08850>.
- Mattanovich, D., R uker, F., Machado, A.C., Laimer, M., Regner, F., Steinkellner, H., Himmler, G., Katinger, H., 1989. Efficient transformation of *Agrobacterium* spp. by electroporation. *Nucleic Acids Res.* 17, 6747.
- Michielse, C.B., van Wijk, R., Reijnen, L., Manders, E.M.M., Boas, S., Olivain, C., Alabouvette, C., Rep, M., 2009. The nuclear protein Sge1 of *Fusarium oxysporum* is required for parasitic growth. *PLoS Pathog.* 5, e1000637. <http://dx.doi.org/10.1371/journal.ppat.1000637>.
- Montano, S.P., Cot e, M.L., Fingerman, I., Pierce, M., Vershon, A.K., Georgiadis, M.M., 2002. Crystal structure of the DNA-binding domain from Ndt80, a transcriptional activator required for meiosis in yeast. *Proc. Natl. Acad. Sci. USA* 99, 14041–14046. <http://dx.doi.org/10.1073/pnas.222312199>.
- Nobile, C.J., Fox, E.P., Nett, J.E., Sorrells, T.R., Mitrovich, Q.M., Hernday, A.D., Tuch, B. B., Andes, D.R., Johnson, A.D., 2012. A recently evolved transcriptional network controls biofilm development in *Candida albicans*. *Cell* 148, 126–138. <http://dx.doi.org/10.1016/j.cell.2011.10.048>.
- Palma-Guerrero, J., Hall, C.R., Kowbel, D., Welch, J., Taylor, J.W., Brem, R.B., Glass, N. L., 2013. Genome wide association identifies novel loci involved in fungal communication. *PLoS Genet.* 9, e1003669. <http://dx.doi.org/10.1371/journal.pgen.1003669>.
- Pinan-Lucarr e, B., Balguerie, A., Clav e, C., 2005. Accelerated cell death in *Podospora* autophagy mutants. *Eukaryot. Cell* 4, 1765–1774. <http://dx.doi.org/10.1128/EC.4.11.1765-1774.2005>.
- Pontecorvo, G., 1956. The parasexual cycle in fungi. *Annu. Rev. Microbiol.* 10, 393–400. <http://dx.doi.org/10.1146/annurev.mi.10.100156.002141>.
- Prosser, J.L., 1995. Kinetics of filamentous growth and branching. In: Gow, N.A.R., Gadd, G.M. (Eds.), *The Growing Fungus*. Springer, Netherlands, Dordrecht, pp. 301–318. http://dx.doi.org/10.1007/978-0-585-27576-5_14.
- Puhalla, J.E., 1985. Classification of strains of *Fusarium oxysporum* on the basis of vegetative compatibility. *Can. J. Bot.* 63, 179–183. <http://dx.doi.org/10.1139/b85-020>.
- Rep, M., Kistler, H.C., 2010. The genomic organization of plant pathogenicity in *Fusarium* species. *Curr. Opin. Plant Biol.* 13, 420–426. <http://dx.doi.org/10.1016/j.pbi.2010.04.004>.
- Roca, G.M., Arlt, J., Jeffree, C.E., Read, N.D., 2005a. Cell biology of conidial anastomosis tubes in *Neurospora crassa*. *Eukaryot. Cell* 4, 911–919. <http://dx.doi.org/10.1128/EC.4.5.911-919.2005>.
- Roca, G.M., Read, N.D., Wheals, A.E., 2005b. Conidial anastomosis tubes in filamentous fungi. *FEMS Microbiol. Lett.* 249, 191–198. <http://dx.doi.org/10.1016/j.femsle.2005.06.048>.
- Sambrook, J., Russel, D.W., 2001. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
- Saupe, S.J., 2000. Molecular genetics of heterokaryon incompatibility in filamentous ascomycetes. *Microbiol. Mol. Biol. Rev.* 64, 489–502.
- Sellam, A., Askew, C., Epp, E., Tebbji, F., Mullick, A., Whiteway, M., Nantel, A., 2010. Role of transcription factor CaNdt80p in cell separation, hyphal growth, and virulence in *Candida albicans*. *Eukaryot. Cell* 9, 634–644. <http://dx.doi.org/10.1128/EC.00325-09>.
- Shahi, S., Beerens, B., Bosch, M., Linmans, J., Rep, M., 2016. Nuclear dynamics and genetic rearrangement in heterokaryotic colonies of *Fusarium oxysporum*. *Fungal Genet. Biol.* 91, 20–31. <http://dx.doi.org/10.1016/j.fgb.2016.03.003>.
- Shahi, S., Beerens, B., Manders, E.M.M., Rep, M., 2015. Dynamics of the establishment of multinucleate compartments in *Fusarium oxysporum*. *Eukaryot. Cell* 14, 78–85. <http://dx.doi.org/10.1128/EC.00200-14>.
- Shevchenko, A., Wilm, M., Vorm, O., Mann, M., 1996. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.* 68, 850–858.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., S oding, J., Thompson, J.D., Higgins, D.G., 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* 7, 539. <http://dx.doi.org/10.1038/msb.2011.75>.
- Smith, M.L., Micali, O.C., Hubbard, S.P., Mir-Rashed, N., Jacobson, D.J., Glass, N.L., 2000. Vegetative incompatibility in the het-6 region of *Neurospora crassa* is mediated by two linked genes. *Genetics* 155, 1095–1104.
- Sopko, R., Stuart, D.T., 2004. Purification and characterization of the DNA binding domain of *Saccharomyces cerevisiae* meiosis-specific transcription factor Ndt80. *Protein Expr. Purif.* 33, 134–144. <http://dx.doi.org/10.1016/j.pep.2003.08.025>.
- Takken, F., Rep, M., 2010. The arms race between tomato and *Fusarium oxysporum*. *Mol. Plant Pathol.* 11, 309–314. <http://dx.doi.org/10.1111/j.1364-3703.2009.00605.x>.
- Van der Nest, M.A., Olson, A., Lind, M., V el ez, H., Dalman, K., Brandstr om Durling, M., Karlsson, M., Stenlid, J., 2014. Distribution and evolution of het gene homologs in the basidiomycota. *Fungal Genet. Biol.* 64, 45–57. <http://dx.doi.org/10.1016/j.fgb.2013.12.007>.
- Wang, J.-S., Yang, Y.-L., Wu, C.-J., Ouyang, K.J., Tseng, K.-Y., Chen, C.-G., Wang, H., Lo, H.-J., 2006. The DNA-binding domain of CaNdt80p is required to activate CDR1 involved in drug resistance in *Candida albicans*. *J. Med. Microbiol.* 55, 1403–1411. <http://dx.doi.org/10.1099/jmm.0.46650-0>.
- Winter, E., 2012. The Sum1/Ndt80 transcriptional switch and commitment to meiosis in *Saccharomyces cerevisiae*. *Microbiol. Mol. Biol. Rev.* 76, 1–15. <http://dx.doi.org/10.1128/MMBR.05010-11>.
- Xiang, Q., Glass, N.L., 2002. Identification of *vib-1*, a locus involved in vegetative incompatibility mediated by *het-c* in *Neurospora crassa*. *Genetics* 162, 89–101.
- Xiong, Y., Sun, J., Glass, N.L., 2014. VIB1, a link between glucose signaling and carbon catabolite repression, is essential for plant cell wall degradation by *Neurospora crassa*. *PLoS Genet.* 10, e1004500. <http://dx.doi.org/10.1371/journal.pgen.1004500>.