



## UvA-DARE (Digital Academic Repository)

### Effector profiles distinguish *formae speciales* of *Fusarium oxysporum*

van Dam, P.; Fokkens, L.; Schmidt, S.M.; Linmans, J.J.J.; Kistler, H.C.; Ma, L.-J.; Rep, M.

**DOI**

[10.1111/1462-2920.13445](https://doi.org/10.1111/1462-2920.13445)

**Publication date**

2016

**Document Version**

Final published version

**Published in**

Environmental Microbiology

**License**

Article 25fa Dutch Copyright Act

[Link to publication](#)

**Citation for published version (APA):**

van Dam, P., Fokkens, L., Schmidt, S. M., Linmans, J. J. J., Kistler, H. C., Ma, L.-J., & Rep, M. (2016). Effector profiles distinguish *formae speciales* of *Fusarium oxysporum*. *Environmental Microbiology*, 18(11), 4087-4102. <https://doi.org/10.1111/1462-2920.13445>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

# Effector profiles distinguish *formae speciales* of *Fusarium oxysporum*

Peter van Dam,<sup>1</sup> Like Fokkens,<sup>1</sup> Sarah M. Schmidt,<sup>1</sup>  
Jasper H.J. Linmans,<sup>1</sup> H. Corby Kistler,<sup>2</sup>  
Li-Jun Ma<sup>3</sup> and Martijn Rep<sup>1\*</sup>

<sup>1</sup>Molecular Plant Pathology, Swammerdam Institute for Life Sciences, University of Amsterdam, The Netherlands.

<sup>2</sup>United States Department of Agriculture, ARS Cereal Disease Laboratory, University of Minnesota, St. Paul, MN, USA.

<sup>3</sup>Department of Biochemistry and Molecular Biology, University of Massachusetts, Amherst, MA 01003, USA.

## Summary

**Formae speciales (ff.spp.) of the fungus *Fusarium oxysporum* are often polyphyletic within the species complex, making it impossible to identify them on the basis of conserved genes. However, sequences that determine host-specific pathogenicity may be expected to be similar between strains within the same *forma specialis*. Whole genome sequencing was performed on strains from five different ff.spp. (*cucumerinum*, *niveum*, *melonis*, *radicis-cucumerinum* and *lycopersici*). In each genome, genes for putative effectors were identified based on small size, secretion signal, and vicinity to a “miniature impala” transposable element. The candidate effector genes of all genomes were collected and the presence/absence patterns in each individual genome were clustered. Members of the same *forma specialis* turned out to group together, with cucurbit-infecting strains forming a supercluster separate from other ff.spp. Moreover, strains from different clonal lineages within the same *forma specialis* harbour identical effector gene sequences, supporting horizontal transfer of genetic material. These data offer new insight into the genetic basis of host specificity in the *F. oxysporum* species complex and show that**

**(putative) effectors can be used to predict host specificity in *F. oxysporum*.**

## Introduction

The *Fusarium oxysporum* (Fo) species complex (FOSC) comprises an important group of filamentous fungi that includes plant-pathogenic strains. The species complex as a whole has a very wide host range, but individual pathogenic strains are restricted to one or a few host species. Accordingly, such strains are grouped into *formae speciales* (ff.spp.) based on host specificity (Armstrong and Armstrong, 1981; Baayen *et al.*, 2000). They cause vascular wilt (due to xylem colonization) or root, bulb or foot rot in over 120 plant species, including many economically important crops like tomato and cucurbits (Armstrong and Armstrong, 1981; Michielse and Rep, 2009).

Each *forma specialis* (f.sp.) of Fo consists of one or several clonal lineages (O'Donnell *et al.*, 1998; Katan, 1999). Strains belonging to different *formae speciales* may be more related than strains belonging to the same *forma specialis* (Kistler, 1997; Lievens *et al.*, 2009), which implies that the host of a strain of *F. oxysporum* can not reliably be determined based on conserved gene sequences, such as the translation elongation factor 1-alpha gene (*EF-1 $\alpha$* ) alone. Currently, disease assays remain the primary method of discriminating host range and races (defined by the capacity to infect different cultivars of a plant species and often based on presence/absence of or point mutations in effector genes) of a pathogenic Fo strain (Recorbet *et al.*, 2003; Covey *et al.*, 2014; Martyn, 2014).

Because disease assays are laborious and time consuming, a molecular screening method based on *forma specialis*-specific DNA sequences is highly desirable. However, knowledge of the genetic basis of host-specificity is limited.

For successful infection of their host, pathogens often rely on effector proteins—small secreted proteins that facilitate the colonization process (Stergiopoulos and de Wit, 2009; Dodds and Rathjen, 2010; Giraldo and Valent, 2013). Infection of different plant hosts likely requires a different set of effectors, which renders these proteins potentially informative for discriminating *formae speciales*. In this study we aimed to uncover which putative effector

Received 3 March, 2016; accepted 30 June, 2016. \*For correspondence. E-mail: m.rep@uva.nl; Tel. (+0031) 20525 7764.

[Correction added on 19 September 2016, after first online publication: Errors on Table, Figures and References were corrected.]

genes are shared by strains belonging to each of four economically important (Kim *et al.*, 1993b; Vakalounakis *et al.*, 2005) cucurbit-infecting *formae speciales*, notably *cucumerinum* (Foc; cucumber), *melonis* (Fom; musk melon), *niveum* (Fon; watermelon), and *radicis-cucumerinum* (Forc; cucumber and other cucurbits), as well as the tomato-infecting *F. oxysporum* f.sp. *lycopersici* (Fol). We chose these ff.spp. because we wished to know whether strains that infect related plant species (*Cucurbitaceae*) possess similar effector suites.

Of all of the *formae speciales* mentioned above, Fol has been investigated most extensively. A horizontally transferable “pathogenicity” chromosome (chromosome 14 of Fol4287) harbours all but one of the 14 known Fol effector genes, named *SIX* (for Secreted In Xylem; Ma *et al.*, 2010; Schmidt *et al.*, 2013). These genes encode small, cysteine-rich, secreted proteins with no recognizable protein domain and at least some appear to be employed by Fol to manipulate the host’s defense responses, thereby promoting the infection process (Rep *et al.*, 2004; Gawehns *et al.*, 2014; Ma *et al.*, 2015). Together with chromosomes 1B, 2B, 3, 6, and 15, chromosome 14 forms the accessory genome of Fol4287. The accessory genome lacks synteny with other *Fusarium* species such as *F. verticillioides*, is relatively gene poor and harbours many repeats and transposable elements (TEs), characteristics that differentiate it from the core genome (Ma *et al.*, 2010; 2013).

Analysis of the genomic context of *SIX1* – *SIX7* revealed an association of these genes with two TEs. A miniature impala (mimp) was found in all cases in the upstream region and mFot5 was found frequently downstream of the Open Reading Frame (ORF) (Schmidt *et al.*, 2013). Both mimps and mFots are classes of Miniature Inverted-repeat TEs (MITEs). MITEs are short, nonautonomous DNA transposons, thought to be truncated derivatives of autonomous DNA transposons (Feschotte and Pritham, 2007; Lu *et al.*, 2012). They contain 27–30 nucleotide terminal inverted repeats (TIRs). In the case of mimps, the sequence of these TIRs is conserved between different mimp-subfamilies (Bergemann *et al.*, 2008). A total of 103 mimps are present in the genome of Fol4287 (Schmidt *et al.*, 2013). The highest density was found on chromosome 14, the “pathogenicity chromosome,” with 54 mimps (21 mimps/Mb), compared to 45 (3 mimps/Mb) on the other accessory chromosomes and 4 (0.1 mimps/Mb) on the core chromosomes. Although mimp deletion experiments did not result in altered *SIX* gene expression (Schmidt *et al.*, 2013), the consistent presence of a mimp in the promoter region of known Fo effector genes was successfully exploited to identify novel effector candidates (Schmidt *et al.*, 2016). Sixteen candidate effector genes were identified close to a mimp, and the products of fourteen of these (*SIX1* – *SIX14*) were found in the xylem sap of infected tomato plants using mass spectrometry (Houterman *et al.*, 2007; Schmidt *et al.*, 2013).

We made use of the association between mimps and effector genes in Fo genomes to predict the suite of putative effectors present in 59 *F. oxysporum* genomes (45 new assemblies), without the need to rely on genome annotation. We then compared the predicted “effectoromes” of these 59 different strains to determine whether strains that belong to the same f.sp. have similar effector repertoires and whether we could use presence/absence patterns of putative effector genes to predict the host range of a strain. We find that indeed strains cluster into *formae speciales* based on effector presence/absence profiles. Moreover, identification of ff.spp. based on effector genes is further strengthened by taking their sequences into account, since these are identical or highly similar within a f.sp., but often different between ff.spp.

## Results

### Strain selection

In order to make a well-founded evaluation of genome and effector variation within and between *formae speciales* of *F. oxysporum*, we selected divergent strains for our study. For the *formae speciales* Foc, Fom and Fon, a polyphyletic nature has been described (Jacobson and Gordon, 1990; Kim *et al.*, 1992; 1993a; Lievens *et al.*, 2007). Forc has a broader host range and is able to cause root and stem rot in various cucurbit species such as muskmelon and sponge gourd (*Luffa aegyptica*) (Vakalounakis, 1996; Vakalounakis and Fragkiadakis, 1999; Vakalounakis *et al.*, 2005). According to earlier reports it comprises two vegetative compatibility groups (VCGs) but these might constitute a single clonal lineage (Lievens *et al.*, 2007). Fol is also polyphyletic, comprising at least four clonal lineages (Van Der Does *et al.*, 2008; Ma *et al.*, 2010).

Based on diversity in *EF-1 $\alpha$*  sequence (indicative of clonal lineages, Supporting Information Fig. S1), geographical origin, and disease assays to confirm host-specific pathogenicity, 45 strains (9 Foc, 9 Fom, 9 Fon, 3 Forc, 14 Fol, one non-pathogenic, Table 1) were selected for whole genome Illumina paired-end sequencing. Combined with 14 previously generated Fo assemblies belonging to various ff.spp. (Table 2), these genomes formed the basis for our search for putative effector genes within the FOOSC.

### Most cucurbit-infecting strains are host specific

Previous studies have indicated that Fo strains belonging to f.sp. *cucumerinum* (Cafri *et al.*, 2005) and *niveum* (Zhou and Everts, 2007) display mild cross-pathogenicity towards musk melon. To assess the level of (cross-) pathogenicity of the cucurbit-infecting strains in our collection, we conducted disease assays with all sequenced Foc, Forc, Fom, and Fon strains, plus one Fol strain (Fol029) on susceptible cultivars of cucumber,

**Table 1.** Strains of which the genomes were sequenced and assembled in this study.

Strain <sup>a</sup>	Original designation	VCG <sup>b</sup>	Race <sup>b</sup>	Origin of strain	Reference
Foc001	Foc-1	0183	–	Japan	BL <sup>c</sup>
Foc011	9903-1	0186	–	China	Lievens <i>et al.</i> , 2007
Foc013	9904-1	0186	–	China	Lievens <i>et al.</i> , 2007
Foc015	9906-3	0184	–	China	Lievens <i>et al.</i> , 2007
Foc018	Afu-50(B)	0180	–	Crete, Greece	Lievens <i>et al.</i> , 2007
Foc021	ATCC 16416	0180	–	Florida	Lievens <i>et al.</i> , 2007
Foc030	FOCU-22P	0180	–	Israel	Lievens <i>et al.</i> , 2007
Foc035	NETH 11179	0181	–	Netherlands	Lievens <i>et al.</i> , 2007
Foc037	Tf-213	0185	–	Japan	Lievens <i>et al.</i> , 2007
Forc016	33	0260	–	Canada	Lievens <i>et al.</i> , 2007
Forc024	Afu-11(A)	0260	–	Crete, Greece	Lievens <i>et al.</i> , 2007
Forc031	AK-2	0261	–	Crete, Greece	Lievens <i>et al.</i> , 2007
Fon002	CBS 418.90	–	–	Israel	Lievens <i>et al.</i> , 2007
Fon005	TX-471-1	0080	0	Texas	Zhou and Everts, 2007
Fon010	F-016-1	0082	1	Maryland	Zhou and Everts, 2007
Fon013	F-014-2	0082	2	Maryland	Zhou and Everts, 2007
Fon015	F-063-1	0082	2	Maryland	Zhou and Everts, 2007
Fon019	TX-X1D	0082	2	Texas	Zhou and Everts, 2007
Fon020	F-099-1	0083	2	Delaware	Zhou and Everts, 2007
Fon021	MD-ZE622	–	3	Maryland	Zhou and Everts, 2007
Fon037	NRRL 38539	–	–	Israel	Hadar and Katan, 1989
Fom004	Fom 0122	0134	0	Spain	Schmidt <i>et al.</i> , 2016
Fom005	Fom 0123	0134	1	Spain	Schmidt <i>et al.</i> , 2016
Fom006	Fom 0124	0134	2	Spain	Schmidt <i>et al.</i> , 2016
Fom009	–	0135	2	Israel	Schmidt <i>et al.</i> , 2016
Fom010	–	–	1	Israel	Schmidt <i>et al.</i> , 2016
Fom011	–	–	0	Israel	Schmidt <i>et al.</i> , 2016
Fom012	ML2	0134	0	–	Schmidt <i>et al.</i> , 2016
Fom013	–	0134	2	Spain	Schmidt <i>et al.</i> , 2016
Fom016	Fom26	0134	1	–	Schmidt <i>et al.</i> , 2016
Fol002	WCS862/E241	0030	2	Netherlands	Mes <i>et al.</i> , 1999
Fol004	IPO1530/B1	0030	1	Netherlands	Mes <i>et al.</i> , 1999
Fol007	D2	0030	2	France	Mes <i>et al.</i> , 1999
Fol014	LSU-3	0030	1	Louisiana	Mes <i>et al.</i> , 1999
Fol016	BFOL-51	0031	1	Louisiana	Mes <i>et al.</i> , 1999
Fol018	LSU-7	0030	2	Louisiana	Mes <i>et al.</i> , 1999
Fol026	BRIP 14844 (M1943)	0030	3	Australia	Mes <i>et al.</i> , 1999
Fol029	5397	0030	3	Florida	Mes <i>et al.</i> , 1999
Fol038	CA92/95	0030	3	California	Lievens <i>et al.</i> , 2009
Fol069	DF0-23	0035	2	California	Cai <i>et al.</i> , 2003
Fol072	DF0-38	0031	2	California	Cai <i>et al.</i> , 2003
Fol073	DF0-40	0030	2	California	Cai <i>et al.</i> , 2003
Fol074	DF0-41	0030	3	California	Cai <i>et al.</i> , 2003
Fol075	DF0-62	0031	2 <sup>d</sup>	California	Cai <i>et al.</i> , 2003
FoMN14	MN-14	–	N.P.	California	Gale <i>et al.</i> , 2003

<sup>a</sup> Foc: *Fo* f.sp. *cucumerinum*, Forc: *Fo* f.sp. *radicis-cucumerinum*, Fon: *Fo* f.sp. *niveum*, Fom: *Fo* f.sp. *melonis*, Fol: *Fo* f.sp. *lycopersici*.

<sup>b</sup> Race designation and VCG was taken from the corresponding reference. In the case of Fol, inoculation on differential cultivars [differing in their *I* (immunity)-gene genotype: C32/KG52201/PV2002MM (*i*-i2-i3; susceptible); GCR161 (*i*-i2); OT264 (KG324)/341F (*i*-i2); C295 (*i*-i2-i3); E779 (*i*-i2-i3)] was used to confirm the race of each strain (N.P.: nonpathogenic on tomato).

<sup>c</sup> Foc001 (Foc-1) was obtained from Bart Lievens, Scientia Terrae Research Institute, Belgium.

<sup>d</sup> Fol075 (DF0-62) was reported to be non-pathogenic. Based on retesting we now designate it as a (weak) race 2 strain.

musk melon, watermelon, and tomato, as well as on a Foc-resistant cultivar of cucumber. The results are summarized in Fig. 1.

Fom and Fon strains were highly specific to their described host species, whereas several Foc strains showed some degree of cross-pathogenicity, especially towards musk melon. Disease symptoms on musk melon

plants caused by *formae speciales* other than *melonis* (and *radicis-cucumerinum*) were generally not severe, being limited to growth retardation or light wilting symptoms. Within Foc, strains Foc018, 021, 030 (all belonging to VCG0180) and Foc011 (VCG0186) were pathogenic on susceptible, but not on resistant cucumber plants; the other Foc strains also caused symptoms in the resistant

**Table 2.** Genome assemblies collected from GenBank and the Broad Institute of Harvard and MIT used in this study.

Strain	NRRL #	VCG	Race	f.sp.	Host	GenBank assembly accession
4287	34936	0030	2	<i>lycopersici</i>	<i>Solanum lycopersicum</i>	GCA_000149955.2
MN25	54003	0033	3	<i>lycopersici</i>	<i>Solanum lycopersicum</i>	GCA_000259975.2
CL57	26381	0094	–	<i>radicis-lycopersici</i>	<i>Solanum lycopersicum</i>	GCA_000260155.3
HDV247	37622	–	–	<i>pisi</i>	<i>Pisum sp.</i>	GCA_000260075.2
PHW808	54008	0101	2	<i>conglutinans</i>	<i>Brassica sp.</i>	GCA_000260215.2
PHW815	54005	0102	–	<i>raphani</i>	<i>Raphanus sp.</i>	GCA_000260235.2
Fo5176	–	–	–	( <i>Brassica</i> )	<i>Brassica sp.</i>	GCA_000222805.1
Fom001	26406	0136	1	<i>melonis</i>	<i>Cucumis melo</i>	GCA_000260495.2
Fov	25433	–	–	<i>vasinfectum</i>	<i>Gossypium sp.</i>	GCA_000260175.2
II-5	54006	01213	TR4	<i>cubeuse</i>	<i>Musa sp.</i>	GCA_000260195.2
N2	–	–	1	<i>cubeuse</i>	<i>Musa sp.</i>	GCA_000350345.1
B2	–	–	4	<i>cubeuse</i>	<i>Musa sp.</i>	GCA_000350365.1
FOSC 3-a	32931	–	–	(human)	<i>Homo sapiens</i>	GCA_000271745.2
Fo47	54002	–	–	(biocontrol)	Soil	GCA_000271705.2

plants. Forc strains were strongly pathogenic towards cucumber and melon plants and moderately pathogenic towards Fon-susceptible watermelon plants (cv. Black Diamond) as well as Foc-resistant cucumber plants (cv. Melen). Only Fol029 caused disease in tomato and inoculation with this strain did not cause any symptoms in any of the tested cucurbit plants. This shows that although cross-pathogenicity of Foc and Forc towards related cucurbit plants is possible, this does not extend towards tomato, a member of the *Solanaceae*.

To evaluate the extent to which the different Fo strains are able to colonize the vasculature of the various plant lines and species, slices of surface-sterilized hypocotyls of infected plants were placed on Czapek Dox Agar (CDA) plates. Mycelial outgrowth after four days largely correlated with disease symptoms (Fig. 1). However, sometimes outgrowths were observed from symptomless plants, suggestive of an endophytic interaction.

### Genome features

The genomes of the 45 selected strains were sequenced using paired-end Illumina libraries with different insert sizes, as described in the Experimental procedures. The genomes were sequenced to 60–200× coverage (Supporting Information Table S1, trimmed and cleaned Illumina data), resulting in assemblies of 826 (Forc024) to 3867 (Foc018) scaffolds. The smallest *de novo* assembly has a cumulative size of 48.4 megabasepairs (Mbp), including ambiguous bases (Foc037); the largest *de novo* assembly adds up to 57.7 Mbp (Fom009); 97–98% of highly conserved protein-coding genes are estimated by CEGMA in all assemblies (Parra *et al.*, 2009), which is similar to other *F. oxysporum* reference genome assemblies retrieved from Genbank (Supporting Information Table S1). This highlights the completeness of the assemblies, particularly in core regions of the genomes. Repeat-density,


RNA and DNA transposon abundance (Supporting Information Fig. S2) and GC content (Supporting Information Table S1) is largely similar between the assemblies.

The genome of the reference strain Fol4287 is the best-studied and assembled Fo genome to date, with a near-complete chromosome assembly through the use of Sanger sequencing combined with an optical map (Ma *et al.*, 2010). Figure 2 shows the size and the level of fragmentation in each of the assessed assemblies. This analysis indicates that most strains contain a roughly equally sized core genome of about 41 Mb, with some exceptions that seem to have a slightly larger (e.g. Fom001, Fo f.sp. *cubeuse* [Focub] B2, Forl CL57, Fo47, FOSC-3a) or smaller (e.g. Fo5176, Fom010, Foc037) cumulative core genome size. Differences in reported core genome size between strains may be caused by addition of accessory regions to the ends of core chromosomes, as was observed in chromosomes 1 and 2 of Fol4287 (Ma *et al.*, 2010) or by partial or entire loss of chromosomes. The size of the accessory regions varies considerably (between 4 and 19 Mb), but is usually comparable within a clonal lineage. To make an estimation of the complete genome size, we calculated the number of basepairs mapped against the assembly divided by the median coverage of contigs larger than 100kb. This yielded estimated genome sizes that were considerably larger than the assembly (Supporting Information Table S1), indicating that indeed large-scale duplication events are a relevant factor. This also explains Fol4287's assembly sticking out in size (Fig. 2) due to its relatively complete assembly, most notably in the aforementioned duplicated regions.

### Foc, Fom, Fon, and Fol are all polyphyletic within the FOSC

To determine to what extent host specificity is polyphyletic among the selected strains, we inferred a

	cucumber (Paraiso, susceptible)	cucumber (Melen, resistant)	musk melon (Charantais-T, susceptible)	watermelon (B. Diamond, susceptible)	tomato (C32, susceptible)
mock	- (0/5)	- (0/5)	- (0/5)	- (0/3)	- (0/5)
Fol029	- (0/5)	- (2/5)	- (0/5)	- (0/3)	+++ (5/5)
Foc001	++ (5/5)	++ (5/5)	- (0/5)	- (1/5)	-
Foc011	+ (5/5)	- (2/5)	- (1/5)	- (1/5)	-
Foc013	+++ (5/5)	+ (5/5)	- (2/5)	- (2/5)	- (2/5)
Foc015	++ (5/5)	+ (5/5)	+ (3/5)	- (0/3)	- (0/5)
Foc018	+ (5/5)	- (2/5)	- (1/5)	- (2/5)	- (1/5)
Foc021	++ (5/5)	- (0/5)	- (0/5)	- (1/5)	-
Foc030	++ (5/5)	- (1/5)	- (1/5)	- (1/5)	- (0/5)
Foc035	++ (5/5)	++ (5/5)	- (0/5)	- (0/5)	-
Foc037	+++ (5/5)	++ (5/5)	- (3/5)	- (2/5)	- (0/5)
Forc016	++ (7/7)	- (4/5)	+++ (5/5)	++ (3/4)	- (0/5)
Forc024	+++ (5/5)	+ (5/5)	+++ (4/4)	+ (3/5)	- (0/5)
Forc031	+++ (5/5)	++ (5/5)	+++ (5/5)	++ (2/4)	-
Fom001	- (0/5)		+++ (0/5)	- (1/3)	- (1/5)
Fom004	- (0/5)		+++ (5/5)	- (4/5)	-
Fom005	- (0/5)		+++ (5/5)	- (0/5)	- (0/5)
Fom006	- (0/5)		+++ (5/5)	- (1/5)	-
Fom009	- (0/5)		+++ (5/5)	- (2/5)	-
Fom010	- (0/5)		+++ (5/5)	- (2/5)	-
Fom011	- (0/5)		+++ (5/5)	- (1/5)	- (1/5)
Fom012	- (0/5)		+++ (1/5)	- (0/5)	-
Fom013	- (0/5)		+++ (5/5)	- (1/5)	- (0/5)
Fom016	- (0/5)		+++ (0/5)	- (3/5)	-
Fon002	- (0/5)		- (0/5)	+++ (5/5)	-
Fon005	- (0/5)		- (0/5)	+ (0/5)	- (0/5)
Fon010	- (0/5)		- (0/5)	+ (4/5)	-
Fon013	- (0/5)		- (2/5)	+++ (5/5)	- (0/5)
Fon015	- (0/5)		- (3/5)	+++ (4/5)	-
Fon019	- (0/5)		- (0/5)	+++ (4/5)	-
Fon020	- (1/5)		- (1/5)	+++ (5/5)	-
Fon021	- (0/5)		- (0/5)	++ (6/6)	-
Fon037	- (1/5)		- (0/5)	+++ (5/5)	-




**Fig. 1.** Host-specificity of cucurbit-infecting strains used in this study.

Disease index (0–4) was scored in multiple bioassays under controlled conditions in the greenhouse after two weeks (cucurbits) or three weeks (tomato). All compatible interactions were tested at least twice in individual bioassays. Between brackets is the number of surface-sterilized hypocotyls from which *F. oxysporum* outgrowth was observed on CDA after 4 days.

phylogeny based on a concatenated alignment of 1195 conserved core genes (see Experimental procedures for more detail). The resulting tree, depicted in Fig. 2, is congruent with previously published phylogenetic analyses (Baayen *et al.*, 2000; Lievens *et al.*, 2007). We observe three major clades, of which clade 1 corresponds to a separate phylogenetic species (Laurence *et al.*, 2014). We find that *ff.spp.* are generally grouped into distinct clonal lineages and that these lineages are represented in distinct clades or subclades in the tree. One notable exception is *Forc*, of which all three sequenced strains group into a single clonal line. This is in accordance with previous reports in which 68 different *Forc* strains all belonged to a single Random Amplified Polymorphic DNA (RAPD) and Amplified Fragment Length Polymorphism group (Vakalounakis

and Fragkiadakis, 1999; Vakalounakis *et al.*, 2005; Lievens *et al.*, 2007).

We find that *Foc*, *Fom*, and *Focub* are each represented in two different major clades, *Foc* and *Focub* even in two different phylogenetic species. *Foc* strains are distributed over five clonal lineages: one in clade 1 and four in clade 2. *Focub* (poorly represented in this study with only three strains) is grouped into two lineages, one that belongs to clade 1 and one strain that is placed in clade 2. *Fom* strains cluster into three clonal lineages, two that belong to different subclades in clade 2 and one strain (*Fom001*/NRRL24604) that belongs to clade 3. *Fon* and *Fol* are confined to a single major clade, but within this clade, we find that different clonal lines belong to distinct subclades. *Fon* is clustered into three distinct groups within clade 2. *Fol* harbours four



**Fig. 2.** Most strains analyzed in this study belong to polyphyletic *formae speciales* and contain a core genome of roughly the same size. One thousand one hundred and ninety-five genes residing on FoI4287's core genome were selected that have one-to-one orthologs in all *Fusarium* strains, using *F. verticillioides* 7600 as outgroup (outgroup not shown). A concatenated sequence alignment was generated using ClustalO (total length of 2,542,215 nt after trimming, including gaps) and phylogeny was inferred with 100 bootstrap iterations. Branches with most parsimonious bootstrap partitions >90% are indicated in bold; those with a value >70% are bold and shaded red. A coloured circle (wilting), square (root and shoot rot), or triangle (nonpathogenic/endophytic) representing the strain's host range was plotted on the leaves of the tree. Next to the dendrogram, the scaffolds in the genome assembly are plotted for each strain. Each scaffold is represented by a scaled rectangle and coloured blue if it is part of the core genome (based on alignment to the FoI4287 core genome) and orange if it is not (see Experimental procedures for more detail). The size of the core is very similar amongst different strains, in contrast to the size of the accessory genome, which varies even within a clonal line.

clonal lineages in clade 3, with the majority of the sequenced strains belonging to the same clonal lineage as the reference strain FoI4287, corresponding to VCG 30 (VCG 0030). In summary, all ff.spp. except Forc are polyphyletic, consisting of several clonal lines.

#### *Effector prediction pipeline yields 104 effector candidates in 59 Fo genomes*

If host preference of a strain is largely determined by its suite of effectors, we expect strains that infect the same host to have a similar effector repertoire, even if they

belong to different phylogenetic clades. To determine the effector repertoire of different strains, we exploited the association of mimps with effectors in *Fo*. Each genome was scanned for the presence of mimp TIRs. Subsequently, two methods were used for ORF identification: (i) the sequence 2500 bp downstream of the mimp IR was translated in the three possible reading frames and ORFs bigger than 25 codons were extracted; (ii) AUGUSTUS 3.1 gene prediction software was run on the 5000 bp downstream of the TIR. In both cases, the threshold for distance from TIR to ATG was set to 2000bp. Method ii allowed for the prediction of putative effectors with a short first exon, like *SIX10*. Supporting Information Fig. S3 depicts a summary of the method.

To evaluate the influence of assembly fragmentation on the number of effectors found, we compared the number of mimps, mimp-inverted repeats and associated candidate genes identified in two different short read assemblies (based on Illumina HiSeq and Ion Proton sequencing) to the reference assembly of *Fo*4287 (Sanger/optical mapping). We found that the fragmented assemblies performed only slightly worse than the reference assembly (Supporting Information Tables S1 and S3) in terms of effector candidate prediction, although clearly less complete mimps were found in the Ionproton assembly (21 vs. 40).

When we applied the extended pipeline to all 59 genomes, we found in total 2242 ORFs that met the selection criteria (>25 aa; <300 aa; SignalP value >0.550; distance to closest mimp TIR <2500 bp). To reduce redundancy, we grouped these ORFs into gene families based on a self-BLAST search. This resulted in a final set of 201 mimp-associated gene families encoding small secreted proteins. This set includes all previously described *SIX* genes of *Fo*l except *SIX5* (*Six5* returns a SignalP value of 0.444) and *SIX12* (*Six12* does not have a signal peptide). Subsequently, we ran BLAST2GO and InterProScan to find gene ontology (GO) terms and protein domains in our set of genes, which resulted in a functional annotation for 88 out of 201 genes. Fifty-four genes returned no significant BLAST hit and 59 returned a best BLASTX hit with a hypothetical protein (13 of these had at least one associated GO term). For those genes that had a homolog in any of the annotated *Fo* genomes, we also report the corresponding gene id in this strain (Supporting Information Table S2, column L).

Due to the nature of the method, we sometimes picked up fragments of genes, originating from only the first exon or from an internal gene region. We checked for each candidate if the sequence could be aligned to the start site of the corresponding gene sequence from the Broad annotation in order to update the gene model. In 83 out of 201 candidates the reported Broad gene model could be used. For the others, we did not alter the gene model. To groom the list of candidates, we removed duplicate records (5×),

very short protein products (<35aa after signal peptide removal; 23×); homologs or fragments of *SIX* genes (17×; all 14 *SIX* genes with correct gene models from *Fo*l were manually added to the list instead, resulting in 215 records in Supporting Information Table S2) and sequences with an unlikely signal peptide based on visual inspection (67×), despite a positive (>0.550) detection of a signal peptide by SignalP. These included TEs, transcription factors and integral membrane proteins. This resulted in a final set of 104 candidate effectors (Supporting Information Table S2).

Predicted proteins in the set with recognisable domains include FOVG\_19456, which contains a LysM domain, potentially protecting the fungal cell wall or preventing chitin-triggered immunity in plants (de Jonge and Thomma, 2009; Stergiopoulos and de Wit, 2009; Jiang *et al.*, 2014), Rapid Alkalinization Factor-like protein 33 (FOIG\_11494) and several secreted enzymes with predicted peptidase (FOXG\_17323, FOVG\_19376, FOXB\_07727), polygalacturonase (FOTG\_18786, FOWG\_18016), glycoside hydrolase (FOCG\_17303), carbonic anhydrase (FOMG\_18585), or peroxidase (FOVG\_19731) activity that could play a role in nutrient acquisition or suppressing plant defences.

#### *Strains with the same host-specificity cluster together based on effector content*

Having identified the combined putative “effectorome” of the 59 *Fo* strains, we examined which putative effectors are shared amongst members of the same f.sp., multiple ff.spp., all analyzed *Fo* genomes or with other *Fusarium* species. Presence of a candidate effector gene in a genome was defined as having at least one blastn hit with an e-value ≤ 1E-03 and an identity score (number of identical nucleotides divided by the query length) of at least 30%. This gave rise to a binary “effector-barcode” for each genome. Hierarchical clustering of these presence/absence patterns showed clear grouping of strains with the same host-specificity (Fig. 3), with the exception of *Foc* that split into two clades. We conclude that simple scoring presence/absence patterns of candidate effector genes reflects which host can be infected.

Candidate effector genes were then grouped based on their absence/presence in *formae speciales*. Genes in groups A and B are present in virtually all *Fo* genomes and consists mainly of secreted enzymes and hypothetical proteins, or genes without a BLASTX annotation. Group C is composed of genes that are present in most pathogenic, but not nonpathogenic strains, including some *SIX* genes (marked in red at the top of Fig. 3). *SIX8* forms an exception in this group, it is also found as an intact ORF in *Fo*MN14 while it is incomplete in the FOSC-3a genome assembly, situated at the end of a contig. Group D contains genes that are mostly present in cucurbit-infecting strains



and group E represents Fol genes, including previously described Fol-specific effector genes *SIX2*, 3 and 5 (Lievens *et al.* 2009).

The putative effector repertoires for strains infecting the same host are remarkably similar to each other, suggesting a highly related (at least on the level of presence-absence) shared accessory genome. Interestingly, the cucurbit-infecting strains form a supercluster distinct from the other *formae speciales*, implying that a significantly overlapping set of effectors is associated with the ability to infect cucurbits. This might explain why there is some degree of cross-pathogenicity of Foc towards, for example, melon plants but not from Fol to any of the cucurbits tested. Between cucurbit-infecting *formae speciales*, different sets of effectors are present (particularly in groups C and D) that may be involved in specific pathogenicity towards their host plant.

While still grouping with the other cucurbit-infecting *formae speciales*, the three (highly similar) Foc strains lack a number of putative effectors present in the other cucurbit-infecting strains, predominantly in group D. Their effector pattern is most similar to that of Foc strains not belonging to VCG0180.

Both *FolAVR1* (Houterman *et al.*, 2008) and *AVRFOM2* (Schmidt *et al.*, 2016) show the expected presence-absence pattern (absence of *AVR1* in race 2 and 3 Fol strains and absence of *AVRFOM2* in race 2 Fom strains). Recognition of Avr2 in Fol is evaded by sequence difference rather than deletion of the gene and is present in all Fol strains in the figure.

Strikingly, the hierarchical clustering of genomes based on putative effectors also grouped strains belonging to the same clonal lineage within each polyphyletic *forma specialis*, irrespective of race designation (Fig. 3). This is a signature of vertical inheritance, indicating that the common ancestor of such a clonal lineage at a relatively recent point in time obtained the genetic information needed to infect a new host. Similar selection pressures on different clonal lineages likely resulted in the emergence of the same races in different lineages. This is most clear in the cases of Fol and Fom, where strains belonging to races 1, 2, 3, and races 0, 1, 2, respectively, do not cluster according to race, even though the core genome (Fig. 2), as well as the accessory genome (Fig. 3) are strongly similar within each clonal lineage.

Both Foc and Fon are clearly subdivided into two groups, whose accessory genomes are more different from each other compared to other polyphyletic *formae speciales*. Fon strains 005, 010, 013, 020, 021 form one clonal lineage (Fig. 2) and their putative effector profiles are also nearly identical, while the profiles of strains Fon002, 037, 015, 019 are also highly similar, but lack several effector candidates in group F. The same was found in the case of Foc, where Foc018, 021 and 030, belonging to

one clonal lineage, are highly similar but quite different from the other Foc strains. This suggests relatively large differences in the accessory genomes between clonal lineages in Fon and Foc.


Two non-pathogenic strains were included in this study: Fo47, a biocontrol strain (Aimé *et al.*, 2013) and FoMN14 that was isolated from diseased tomato plants from the same field as MN25 but turned out to be nonpathogenic towards tomato (Gale *et al.*, 2003). These two strains group close to each other and contain relatively few candidate effector genes (47 in Fo47 and 46 in FoMN14, compared to an average of 69 in pathogenic strains). The clinical strain FO SC-3a as well as Fo f.sp. *radicis-lycopersici* CL57 were found in the same branch. These strains also have a relatively low number of candidate effector genes. Based on core-genome sequence (Fig. 2), as well as effector content, Focub strains B2 (race 4) and I15 (tropical race 4; TR4) are nearly identical to each other with only two differentials. This is in line with Fo f.sp. *cubense* TR4 being a single clonal lineage (Ordonez *et al.*, 2015). Since these two strains do not differ that much from strain N2 in their effector pattern (race 1; six presence/absence polymorphisms), the Cavendish banana resistance breaking TR4 may have evolved from race 1 (or race 2, for which a time of this study no genome sequence was available).

To see whether clustering of strains belonging to the same *forma specialis* depends on effector gene sequences and not DNA sequences close to mimps *per se*, we scored presence/absence of 2.5 kb regions downstream of a mimp TIR. The resulting figure (Supporting Information Fig. S4) shows that although some *formae speciales* form a single group, much more fragmentation is found and the clustering generally follows the core genome tree (Fig. 2). For example, Fom001 is found among the Fol strains and Foc011 and 013 are close to Focub strains.

For comparison, the pipeline was also run on 29 other *Fusarium* (non-Fo) genome assemblies of strains for which a whole-genome assembly is available. In most cases, 0–4 mimps or mimp-TIRs could be identified and only in the species *F. avenaceum*, *F. fujikuroi*, and *F. nygamai*, 1–4 candidate ORFs with a secretion signal downstream of a mimp TIR were found (Supporting Information Table S4). This illustrates the specificity of this method for the FO SC and suggests that specifically within this species complex, mimps developed their contextual association with effector genes.

#### *Sequence comparison of SIX genes shows evidence of horizontal transfer*

Strong grouping of host-specificity can already be seen based on presence/absence clustering. We next



**Fig. 3.** *Formae speciales* of *F. oxysporum* cluster based on presence/absence of 104 candidate ORFs close to a mimp, identified in 59 genomes.

Presence was defined as detection with BLASTN with  $e\text{-value} \leq 1e\text{-}3$  and identity score (number of identical nucleotides divided by the query length)  $\geq 30\%$ . A color indicates presence, gray indicates absence. Both the genomes (rows) and effector candidates (columns, 104 $\times$ ) in the resulting table were clustered using a Jaccard binary distance matrix and average linkage. Top: secreted enzymes are marked with black, *SIX* genes and *AVRFOM2* with red, and all four described *AVR* genes are tagged with an asterisk. From left to right: *AVRFOM2*<sup>\*</sup>, *SIX4*<sup>\*</sup> (*AVR1*), 14, 2, 3<sup>\*</sup> (*AVR2*), 7, 10, 5, 12, 1<sup>\*</sup> (*AVR3*), 9, 8, 6, 13, 11.

wanted to find out if a higher resolution could be achieved by comparing sequence types of (candidate) effectors. We aligned the homologs of *SIX1-14* and looked at their phylogeny. Remarkably, different clonal lines within the same f.sp. possess identical or highly

similar sequence variants of each of the 14 described *SIX* genes.

The *SIX1* coding sequence (Rep *et al.*, 2004; van der Does *et al.*, 2008; Fig. 4A) was detected in *Fol*, *Fom*, *Focub*, *Focon* and *Fop*, with strains within each f.sp.

possessing a (nearly) identical sequence for this gene. All Fon strains except Fon002 and 037 possess a *SIX1* homolog that is interrupted by a *Hornet1-TE* at the same position, resulting in a pseudogene. Fom009, 010 and 011 have a second copy of *SIX1* that seems to have been duplicated in the ancestor of this clonal lineage, indicative of vertical inheritance. For *SIX6* (Gawehns *et al.*, 2014), five sequence variants were found: Fol, Foc, Forc-Fon-Fom, Fon015/019, and Focub (Fig. 4B). This suggests that the cucurbit-infecting *formae speciales* Forc (single lineage), Fom (strains from multiple lineages), and Fon (one lineage) have acquired this sequence from the same source. *SIX13* (Schmidt *et al.*, 2013) harbours more sequence variation, but again all Fom, Fol, and Focub strains have identical sequences within the respective f.sp. (Fig. 4C). Interestingly, two different *SIX13* sequence variants were found in Foc and Fon, with one of the two variants being similar to the *ForcSIX13* homolog and the other most closely related to *FomSIX13*.

*SIX5*, 7, 10, and 12 were only found in Fol, with no sequence variation. Earlier studies have shown that some of these genes can be used as Fol-markers (Lievens *et al.*, 2009). *SIX3* (*AVR2*) was also only encountered in Fol, albeit with some sequence variation in strains of Fol that evade I-2 recognition, thus becoming race 3 (V41→M, R45→H or R46→P; Takken and Rep, 2010). *SIX2*, 4, 8, 9, 11, and 14 (de Sain and Rep, 2015) show similar distributions of sequence types as described for *SIX1*, 6 and 13. Phylogenetic trees of these genes can be found in Supporting Information Fig. S5. *SIX2* and *SIX14* were also found in *F. verticillioides* 7600 (not shown).

The *SIX* gene sequences evaluated here do not show the same phylogeny as conserved (core) genes (Fig. 2), which strongly suggests horizontal transfer of these genes. However, Foc, Fom, and Fon strains sometimes also have very different *SIX* gene sequence types, suggesting a highly dynamic origin of host-specificity to cucurbits, possibly with multiple horizontal transfer events.

#### *Most effector genes are expressed during invasive growth*

To assess whether the candidates identified were indeed expressed during plant infection, we performed RNA sequencing on RNA extracted from infected plant roots 10 days after inoculation and five-day-old mycelium from an *in vitro* liquid KNO<sub>3</sub> medium culture. We assessed eight strains belonging to the ff.spp. Foc, Fom, Fon, Forc, Fol. On average, 60.6% (± 3.9% S.D.) of the candidates showed evidence for *in planta* transcription, and 36.7% (± 10.7% S.D.) of the candidates qualified as being clearly expressed *in planta*, but with no or little expression *in vitro* (Supporting Information Fig. S6). In Fol4287, 12 out of 13


*SIX* genes fall inside the latter category (Supporting Information Table S5).

## Discussion

Using a bioinformatics pipeline, putative effectors were identified in 45 newly sequenced strains and 14 publicly available Fo genomes. Effector candidates were identified by searching for a small TIR sequence of miniature Impala TEs. In this way, 104 effector candidates were identified across all 59 *F. oxysporum* genomes. Hierarchical clustering of the presence/absence patterns of these sequences led to grouping of strains that largely coincided with host specificity. This indicates that the accessory genome and specifically the effector genes residing in these regions can be used to identify *formae speciales* in the FOSC. Clustering of ff.spp. was observed even without taking sequence divergence or copy number variation into consideration, underlining the robustness of the method.

We have improved an existing method to identify putative and real effector genes in *F. oxysporum* (Schmidt *et al.*, 2013; 2016) by incorporating a gene prediction module into the pipeline. This at least partially solved the issue of effector genes that were not recognized due to a short first exon, so that *SIX10* was now identified, too. The remaining known false negatives were *SIX5* and *SIX12*, which were not identified due to a low SignalP score or the absence of a signal peptide, respectively. False positives in the list, like duplicate records, very short gene fragments, transcription factors, TEs, membrane proteins and hypothetical proteins with an unlikely signal peptide were identified and removed from the list. The effect of fragmentation of the genome assembly on the number of effectors identified was limited, and by combining the effector candidates from multiple short-read Illumina genomes into one large list, a sufficient resolution was achieved. The level of fragmentation in the Illumina assemblies, did, however, prevent an analysis of the degree to which putative effector genes form clusters in the same genomic regions and possible synteny of such regions between strains.

Other methods of computationally recognizing effector genes include the selection of (predicted) proteins that meet some or all of the following requirements: small size, secreted, high number of cysteines, traces of diversifying selection and *in planta* transcriptional induction (Sperschneider *et al.*, 2015a, 2015b). The advantage of the method presented here is that genome annotation is not necessary and that relevant contextual genome location is taken into account. However, while the association between effector genes and mimps has been demonstrated to hold true in many cases (e.g., *SIX* genes, *AVRFOM2*; Schmidt *et al.*, 2013; 2016), it is possible that several effector candidates have been missed.



**Fig. 4.** Identical *SIX* gene sequence types are found in strains belonging to polyphyletic *formae speciales*, suggesting a combination of vertical and horizontal inheritance of these genes.

A MUSCLE alignment was made of the nucleotide sequence of (a) *SIX1* (881 nt), (b) *SIX6* (727 nt), and (c) *SIX13* (943 nt). Phylogeny was inferred using PhyML with 100 bootstrap iterations and plotted with mid-point rooting. Branches with most parsimonious bootstrap partitions below 50% were collapsed; values  $\geq 50\%$  and  $< 100\%$  are indicated in red; 100% are not indicated. A coloured circle (wilting) or square (root and shoot rot) representing the strain's *forma specialis* was plotted on the leaves of the dendrogram.

For *Fom*, *Fon*, *Forc*, and *Fol* a clear and unambiguous clustering could be observed. *Foc* separated into two clades. Previously, RAPD has been applied as a way of distinguishing between cucurbit-infecting *formae speciales* (Vakalounakis and Fragkiadakis, 1999; Wang *et al.*, 2001; Vakalounakis *et al.*, 2004; Lievens *et al.*, 2007). Aside from being laborious, difficult to replicate and time-consuming, the technique focuses on both the core as well as the

accessory genome, which results in different RAPD profiles for each clonal lineage. Vakalounakis and Fragkiadakis (1999) showed that by transforming RAPD patterns into a binary data matrix and calculating the genetic distance value based on presence or absence of RAPD bands, a similar dendrogram could be generated as shown in our study (Fig. 3), again with VCG 0180 of *Foc* (here represented by *Foc*018, 021, and 030) clustering in a

different group than the other Foc clonal lines. Both Forc (68 strains) and Fom (three related strains from Spain) cluster as single clades in the RAPD study. The fact that clustering based on all DNA sequences associated with mimps resulted in a more fragmented clustering showing more traces of the core phylogeny (Supporting Information Fig. S4) illustrates the importance and relevance of clustering based on ORFs for secreted proteins close to a mimp in *F. oxysporum*.

For about 60% of the identified effector genes, evidence for transcription was found *in planta* 10 days post inoculation (Supporting Information Fig. S5), indicating that many of these genes might play a role during infection. Some candidate genes might be expressed earlier in the infection process and are therefore not found to be expressed at this timepoint.

Focub strains II5, N2 and B2 and *Brassicaceae*-infecting strains PHW808 (f.sp. *conglutinans*) and Fo5176 (no f.sp. reported, but likely f.sp. *conglutinans*) also clustered together, but these *formae speciales* were not sampled extensively enough in this dataset to cover the complete variation present. In both Fon and Foc, and more subtly also in the other *formae speciales*, footprints of vertical inheritance could be seen. Strains from a single clonal lineage usually contain (near) identical putative effector presence patterns (Fig. 3) as well as *SIX* gene sequences (Fig. 4 and Supporting Information Fig. S5).

Foc018, 021, 030 as well as Foc011 (all lacking *SIX6*) were pathogenic on *C. sativus* cv. Paraiso, but not on "Foc-resistant" *C. sativus* cv. Melen plants; the other Foc strains also caused symptoms in cv. Melen. Since they are not in the same core genome clade (Fig. 2), the cause of this race differentiation must have developed at least twice independently, or passed on horizontally. Foc013, which is in the same phylogenetic species as Foc011 (clade 1), was aggressive on both cucumber cultivars.

Because strains belonging to the same *forma specialis* cluster together, the presence of certain putative effectors can indicate with high confidence to which *forma specialis* a strain belongs. For instance, a distinct cluster of putative effector genes is associated with cucurbit-infecting strains (Fig. 3, cluster D). Furthermore, all of the ff.spp. represented by multiple strains share sets of differentiating effector genes. In the case of Fol, it was shown in previous studies that presence of effector genes can be indicative of the host; a PCR screen with *SIX1*, *SIX2*, *SIX3*, and *SIX5* primers resulted in a 100% success rate in identifying Fol strains in a large collection containing fifteen other Fo ff.spp. (Van Der Does *et al.*, 2008; Lievens *et al.*, 2009). From the clustering shown in Fig. 3 we conclude that the suite of putative effector genes present in the genomes of cucurbit-infecting strains, as well as other monophyletic or polyphyletic *formae speciales* can be applied diagnostically if the genome sequence of a novel unknown strain is

available. Members of the same *forma specialis* might share parts of their (accessory) genome involved in specific pathogenic interaction with their host. An important requirement is the completeness of sampling to be able to make a well-supported choice of *forma specialis*-specific marker loci (e.g., group E for Fol, genes in groups C and D for Fom and Foc, cluster F for Fon). Such marker loci are essentially the smallest possible set of effectors that is shared by all strains of a f.sp. and absent (at least as a set) in all other strains. This would greatly improve host range identification, something that so far has proved to be a difficult objective to achieve. Applicability to other *formae speciales* will need to be investigated, for example by sequencing genomes of multiple differential strains of the same *forma specialis* in existing culture collections. A similar clustering result would be expected for other host ranges besides tomato and cucurbits. This would allow *forma specialis*-identification for crops where *Fusarium* wilt and root and shoot rot is a pressing problem.

## Experimental procedures

### Plant lines and fungal strains

The following plant cultivars were used: *Cucumis sativus* cv. Paraiso (susceptible to Foc), *C. sativus* cv. Melen (resistant to Foc), *C. melo* cv. Cha-T (susceptible to Fom), *Citrullus lanatus* cv. Black Diamond (susceptible to Fon) and *Solanum lycopersicum* C32 (susceptible to Fol).

Strains of Fo were selected based on core genome divergence, pathogenicity testing and geographical origin (Table 1). *F. oxysporum* was grown at 25°C in the dark on Czepek Dox agar (CDA, Difco) plates containing 100 mg/l penicillin and 200 mg/l streptomycin.

### Pathogenicity testing

Pathogenicity testing was performed using the root dip method (Wellman, 1939). In short, conidia were isolated from five-day-old cultures in NO<sub>3</sub>-medium (0.17% yeast nitrogen base, 3% sucrose, 100 mM KNO<sub>3</sub>) by filtering through miracloth (Merck; pore size of 22–25 µm). Spores were centrifuged, resuspended in sterile MilliQ water, counted and brought to a final concentration of 10<sup>7</sup> spores/ml. When the first true leaves were emerging, seedlings were uprooted, inoculated, individually potted and kept at 25°C in the greenhouse. Two weeks (cucurbits) or three weeks (tomato) after inoculation, disease was scored using a disease index ranging from 0 to 4 (Supporting Information Methods S1) based on published methods (Rep *et al.*, 2004; Vakalounakis *et al.*, 2004; Pavlou and Vakalounakis, 2005).

From each bioassay combination, pieces of hypocotyl were collected randomly from five plants, surface sterilized with 96% ethanol and a slice was cut of each piece to be placed on CDA plates containing 100 mg/l penicillin and 200 mg/l streptomycin. After four days of incubation at 25°C in the dark, *Fusarium* outgrowth was assessed and scored.

### DNA isolation, genome sequencing, and assembly

Genomic DNA was isolated as described in detail in Supporting Information Methods S2. Library preparation of insert size 550 bp and Illumina HiSeq 2000 (Foc001, 011, 013, 015, 021, 035, 037, Forc016, 024, 031) and Illumina HiSeq 2500 (Fon and Foc018, 030) paired-end sequencing was performed at Keygene N.V. (Wageningen, The Netherlands). All Fom genomes, “Fol4287-illumina” and Fol007 were sequenced and assembled as described in Schmidt *et al.* (2016) at the Beijing Genome Institute (BGI, Hong Kong), using multiple insert libraries. “Fol4287-ionproton” was sequenced at the University of Amsterdam (MAD Dutch Genomics Service and Support Provider) using an Ion Proton™ Sequencer (Thermo Fisher Scientific). Fol paired-end genome sequencing of the remaining strains was performed using Illumina HiSeq 2000 180 bp insert libraries at the Broad Institute of Harvard and MIT.

Genome assemblies were made with CLC workbench v8.0, as described in Methods Supporting Information S3. We used nucmer (with `-maxmatch`) from the MUMmer package (Delcher *et al.*, 2002) to align all genome sequences to the reference genome, that of Fol4287. We designated scaffolds as “core” if its best match in terms of bp that could be aligned was a scaffold that is part of a core chromosome in Fol4287 (with a minimum overlap of either query or the reference scaffold of 30%). All scaffolds that did not meet this criterion were designated “accessory.”

In our assemblies, repeats that are longer than the length of our reads will typically be collapsed into a single sequence; the cumulative length of individual contigs in our assemblies is likely to be an underestimate of true genome size. To improve upon this estimate we determine the coverage by taking the median read depth of all contigs > 100 kb, assuming that these large contigs contain relatively few repeats and divide the number of reads that are mapped to our assembly by the coverage to obtain a putative genome size (Supporting Information Table S1).

### RNA isolation and transcriptome sequence analysis

For transcriptome sequencing, 10-day-old melon Cha-T (Fom), cucumber Paraiso (Foc, Forc), watermelon Black Diamond (Fon), and tomato C32 (Fol) seedlings were inoculated with conidia of strain Fom001 by dipping the roots in the spore suspension for 5 min; roots of infected plants were harvested ten days after inoculation and flash-frozen in liquid nitrogen. Additionally, mycelium from five-day-old *in vitro* KNO<sub>3</sub> cultures was harvested. Total RNA was extracted as described previously (Schmidt *et al.*, 2013). cDNA synthesis, library preparation (200 bp inserts) and Illumina sequencing was performed at BGI (Foc013, Fom001, Forc016, Forc031, Fol4287) and Keygene N.V. (Fon019, Fon020, Fom006).

The obtained reads were mapped against the set of candidate effector sequences retrieved from the genome sequence of each of the strains (only sequences starting with “ATG” were used) with CLC workbench v8.0 with both length and similarity settings set to 0.9. The number of unique mapped reads was divided by the gene length and then multiplied by 1000 to find the number of reads per kb (RPK). This value was then used to calculate relative expression (RE) compared with EF1 alpha expression. Genes with five or more reads

were distinguished as showing evidence for transcription. If a RE of >2% was found, the gene was considered to be strongly expressed (Supporting Information Table S5).

### Data access

The Whole-Genome Shotgun projects for the newly sequenced strains of Foc, Forc, Fom, and Fon have been deposited at Genbank under the BioProject PRJNA306247. Raw sequence data have been deposited into the Sequence Read Archive under the accession numbers SRP067515 (Foc, Fon, Forc, Fol007, Fol4287), SRP042982 (Fom), and SRP002087 (Fol). All publically available genome sequences that were used were obtained from Genbank and the Broad Institute of Harvard and MIT (<http://www.broadinstitute.org>). Detailed information on these strains and their respective accession numbers are listed in Table 2.

### Phylogenetic analysis

We searched for homologs of 15,956 Fol4287 core genes (including introns) in the sequences of the other genomes using BLASTN with default parameters. We selected all sequences that overlap > 70% with the query sequence and more than 80% identity to the query. We then selected query genes for which we find only a single hit in each species, leaving us with 1194 genes. We used ClustalO (Sievers *et al.*, 2014) to construct a multiple sequence alignment for each query and a custom python script to concatenate these alignments. This alignment was trimmed using trimAl `-strictplus`. We used RaxML (with `-T2 -N 100 -m GTRGAMMAIX -x 1234567 -p 123 -f a`) to infer a phylogeny with 100 bootstrap replicates (Stamatakis, 2014).

For the phylogenetic analysis of the genes encoding SIX proteins, we extracted the sequences by BLASTN (with `-evalue 0.001 -task = “blastn”`), then manually curated them to extract complete coding sequences. We used MUSCLE (default values) to construct a multiple sequence alignment for each query, followed by PhyML to infer a phylogeny with 100 bootstrap replicates (Guindon and Gascuel, 2003).

### Putative effector identification

Prediction of candidate effectors was performed as described (Schmidt *et al.*, 2013) on each genome using a custom Python script. Briefly, the genome was searched for a consensus sequence of the mimp inverted repeats (IRs), “TT[TA]TTGCNNCCCACTGNN.” Subsequently, two methods were used for ORF identification: (i) the sequence 2500 bp downstream of the mimp IR was translated in the three possible ORFs and ORFs bigger than 25 codons were extracted; (ii) AUGUSTUS 3.1 gene prediction software was run on the 5000 bp downstream of the mimp IR with options “species = fusarium singlestrand = true” (Stanke *et al.*, 2006). In both cases, the threshold for distance from TIR to ATG was set to 2000 bp. Method (ii) allowed for the

prediction of putative effectors with a short first exon, like *SIX10*. The identified putative ORFs were submitted to a local instance of SignalP4.1 with the option “-u 0.550” for stringent signal peptide prediction (Petersen *et al.*, 2011). All ORFs that met the criteria were collected. After concatenating the putative effectors of the 59 genomes, they were BLASTed against themselves in order to identify and remove redundant ORFs. For each set of homologous putative effectors (criteria: e-value: 1E-03; percent identity: >60%; alignment length: >60%), the longest entry was selected. A total of 201 candidates were saved and used for the downstream hierarchical clustering analysis. For functional annotation of the found candidate genes, we performed BLASTX and InterProScan using BLAST2GO v3.1.3 with standard settings (Conesa *et al.*, 2005).

### Hierarchical clustering

Screening for presence of the putative effectors collected from multiple genomes was done by conducting a BLASTN search (-evalue 0.001, -task = “blastn”) on each genome assembly. Presence of a candidate effector gene in a genome was defined as having at least one blast hit with an e-value  $\leq$  1E-03 and an identity score (number of identical nucleotides in the correct position in the alignment divided by the query length) of at least 30%. A binary datamatrix was generated containing presence (“1”) or absence (“0”) of each candidate in each genome. This table was used as input for hierarchical clustering performed in R, using a Jaccard binary distance matrix and average linkage.

### Acknowledgements

The authors are grateful to K. Broz for Fol DNA isolations and P.M. Houterman for Fol DNA isolations and retesting of Fol races. They thank B. Lievens, D.J. Vakalounakis, and K. Everts for providing their cucurbit-infecting *F. oxysporum* strains. Furthermore, we are grateful to ENZA seeds and Rijk Zwaan for kindly providing cucumber, melon, and watermelon seeds. This work was supported by the Horizon programme (project 93512007) of The Netherlands Genomics Initiative (NGI) and the Innovational Research Incentives Scheme Vici of The Netherlands Organisation for Scientific Research (NWO, project 865.10.002) through grants to M.R.

### References

Aimé, S., Alabouvette, C., Steinberg, C., and Olivain, C. (2013) The endophytic strain *Fusarium oxysporum* Fo47: a good candidate for priming the defense responses in tomato roots. *Mol Plant Microbe Interact* **26**: 918–926.

Armstrong, G.M. and Armstrong, J.K. (1981) *Formae speciales* and races of *Fusarium oxysporum* causing wilt diseases. In *Fusarium: Diseases, Biology, and Taxonomy*. Nelson, P.E., Toussoun, T.A., and Cook, R.J. (eds). Pennsylvania State University Press: University Park, PA, pp. 391–399.

Baayen, R.P., O'Donnell, K., Bonants, P.J., Cigelnik, E., Kroon, L.P., Roebroek, E.J., and Waalwijk, C. (2000) Gene genealogies and AFLP analyses in the *Fusarium oxysporum* complex identify monophyletic and nonmonophyletic *formae speciales* causing wilt and rot disease. *Phytopathology* **90**: 891–900.

Bergemann, M., Lespinet, O., M'barek, S., Ben, Daboussi, M.J., and Dufresne, M. (2008) Genome-wide analysis of the *Fusarium oxysporum* mimp family of MITEs and mobilization of both native and *de novo* created mimps. *J Mol Evol* **67**: 631–642.

Cafri, D., Katan, J., and Katan, T. (2005) Cross-pathogenicity between *formae speciales* of *Fusarium oxysporum*, the pathogens of cucumber and melon. *J Phytopathol* **153**: 615–622.

Cai, G., Gale, L.R., Schneider, R.W., Kistler, H.C., Davis, R.M., Elias, K.S., and Miyao, E.M. (2003) Origin of race 3 of *Fusarium oxysporum* f. sp. *lycopersici* at a single site in California. *Phytopathology* **93**: 1014–1022.

Conesa, A., Gotz, S., Garcia-Gomez, J.M., Terol, J., Talon, M., and Robles, M. (2005) Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* **21**: 3674–3676.

Covey, P. A., Kuwitzky, B., Hanson, M., and Webb, K.M. (2014) Multilocus analysis using putative fungal effectors to describe a population of *Fusarium oxysporum* from sugar beet. *Phytopathology* **104**: 886–896.

Delcher, A.L., Phillippy, A., Carlton, J., and Salzberg, S.L. (2002) Fast algorithms for large-scale genome alignment and comparison. *Nucleic Acids Res* **30**: 2478–2483.

Dodds, P.N. and Rathjen, J.P. (2010) Plant immunity: towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* **11**: 539–548.

van der Does, H.C., Duyvesteyn, R.G.E., Goltstein, P.M., van Schie, C.C.N., Manders, E.M.M., Cornelissen, B.J.C., and Rep, M. (2008) Expression of effector gene *SIX1* of *Fusarium oxysporum* requires living plant cells. *Fungal Genet Biol* **45**: 1257–1264.

van Der Does, H.C., Lievens, B., Claes, L., Houterman, P.M., Cornelissen, B.J.C., and Rep, M. (2008) The presence of a virulence locus discriminates *Fusarium oxysporum* isolates causing tomato wilt from other isolates. *Environ Microbiol* **10**: 1475–1485.

Feschotte, C. and Pritham, E.J. (2007) DNA transposons and the evolution of eukaryotic genomes. *Annu Rev Genet* **41**: 331–368.

Gale, L.R., Katan, T., and Kistler, H.C. (2003) The probable center of origin of *Fusarium oxysporum* f. sp. *lycopersici* VCG 0033. *Plant Dis* **87**: 1433–1438.

Gawehns, F., Houterman, P.M., Ichou, F.A., Michielse, C.B., Hijdra, M., Cornelissen, B.J.C., *et al.* (2014) The *Fusarium oxysporum* effector Six6 contributes to virulence and suppresses I-2-mediated cell Death. *Mol Plant Microbe Interact* **27**: 336–348.

Giraldo, M.C. and Valent, B. (2013) Filamentous plant pathogen effectors in action. *Nat Rev Microbiol* **11**: 800–814.

Guindon, S. and Gascuel, O. (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* **52**: 696–704.

- Hadar, E. and Katan, J. (1989) The use of Nitrate-nonutilizing mutants and a selective medium for studies of pathogenic strains of *Fusarium oxysporum*. *Plant Dis* **73**: 800.
- Houterman, P.M., Cornelissen, B.J.C., and Rep, M. (2008) Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* **4**: e1000061.
- Houterman, P.M., Speijer, D., Dekker, H.L., De Koster, C.G., Cornelissen, B.J.C., and Rep, M. (2007) The mixed xylem sap proteome of *Fusarium oxysporum*-infected tomato plants. *Mol Plant Pathol* **8**: 215–221.
- Jacobson, D.J. and Gordon, T.R. (1990) Variability of mitochondrial DNA as an indicator of relationships between populations of *Fusarium oxysporum* f. sp. *melonis*. *Mycol Res* **94**: 734–744.
- Jiang, C., He, B., Huang, R., Huang, M., and Xu, L. (2014) Identification and functional analysis of LysM effectors from *Marssonina brunnea*. *Australas Plant Pathol* **43**: 615–622.
- de Jonge, R. and Thomma, B.P.H.J. (2009) Fungal LysM effectors: extinguishers of host immunity? *Trends Microbiol* **17**: 151–157.
- Katan, T. (1999) Current status of vegetative compatibility groups in *Fusarium oxysporum*. *Phytoparasitica* **27**: 51–64.
- Kim, D.H., Martyn, R.D., and Magill, C.W. (1993a) Chromosomal polymorphism in *Fusarium oxysporum* f. sp. *niveum*. *Phytopathology* **83**: 1209–1216.
- Kim, D.H., Martyn, R.D., and Magill, C.W. (1993b) Mitochondrial DNA (mtDNA)-relatedness among *formae speciales* of *Fusarium oxysporum* in the Cucurbitaceae. *Phytopathology* **83**: 91–97.
- Kim, D.H., Martyn, R.D., and Magill, C.W. (1992) Restriction fragment length polymorphism groups and physical map of mitochondrial DNA from *Fusarium oxysporum* f. sp. *niveum*. *Phytopathology* **82**: 53.
- Kistler, H.C. (1997) Genetic diversity in the Plant-pathogenic fungus *Fusarium oxysporum*. *Phytopathology* **87**: 474–479.
- Laurence, M.H., Summerell, B.A., Burgess, L.W., and Liew, E.C.Y. (2014) Genealogical concordance phylogenetic species recognition in the *Fusarium oxysporum* species complex. *Fungal Biol* **118**: 374–384.
- Lievens, B., Claes, L., Vakalounakis, D.J., Vanachter, A. C.R.C., and Thomma, B.P.H.J. (2007) A robust identification and detection assay to discriminate the cucumber pathogens *Fusarium oxysporum* f. sp. *cucumerinum* and f. sp. *radicis-cucumerinum*. *Environ. Microbiol* **9**: 2145–2161.
- Lievens, B., Houterman, P.M., and Rep, M. (2009) Effector gene screening allows unambiguous identification of *Fusarium oxysporum* f. sp. *lycopersici* races and discrimination from other *formae speciales*. *FEMS Microbiol. Lett* **300**: 201–215.
- Lu, C., Chen, J., Zhang, Y., Hu, Q., Su, W., and Kuang, H. (2012) Miniature Inverted-repeat transposable elements (MITEs) have been accumulated through amplification bursts and play important roles in gene expression and species diversity in *Oryza sativa*. *Mol Biol Evol* **29**: 1005–1017.
- Ma, L., Houterman, P.M., Gawehns, F., Cao, L., Sillo, F., Richter, H., et al. (2015) The *AVR2-SIX5* gene pair is required to activate I-2-mediated immunity in tomato. *New Phytol* **208**: 507–518.
- Ma, L.J., van der Does, H.C., Borkovich, K.A., Coleman, J.J., Daboussi, M.J., Di Pietro, A., et al. (2010) Comparative analysis reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* **464**: 367–373.
- Ma, L.J., Geiser, D.M., Proctor, R.H., Rooney, A.P., O'Donnell, K., Trail, F., et al. (2013) *Fusarium* pathogenomics. *Annu Rev Microbiol* **67**: 399–416.
- Martyn, R.D. (2014) *Fusarium* wilt of watermelon: 120 years of Research. *Hortic Rev* **42**: 349–442. Vol.
- Mes, J.J., Weststeijn, E.A., Herlaar, F., Lambalk, J.J.M., Wijbrandi, J., Haring, M.A., and Cornelissen, B.J.C. (1999) Biological and molecular characterization of *Fusarium oxysporum* f. sp. *lycopersici* divides race 1 isolates into separate virulence groups. *Phytopathology* **89**: 156–160.
- Michielse, C.B. and Rep, M. (2009) Pathogen profile update: *Fusarium oxysporum*. *Mol Plant Pathol* **10**: 311–324.
- O'Donnell, K., Kistler, H.C., Cigelnik, E., and Ploetz, R.C. (1998) Multiple evolutionary origins of the fungus causing panama disease of banana: concordant evidence from nuclear and mitochondrial gene genealogies. *Proc Natl Acad Sci U S A* **95**: 2044–2049.
- Ordonez, N., Seidl, M.F., Waalwijk, C., Drenth, A., Kilian, A., Thomma, B.P.H.J., et al. (2015) Worse comes to worst: bananas and panama disease—when plant and pathogen clones Meet. *PLOS Pathog* **11**: e1005197.
- Parra, G., Bradnam, K., Ning, Z., Keane, T., and Korf, I. (2009) Assessing the gene space in draft genomes. *Nucleic Acids Res* **37**: 289–297.
- Pavlou, G.C. and Vakalounakis, D.J. (2005) Biological control of root and stem rot of greenhouse cucumber, caused by *Fusarium oxysporum* f. sp. *radicis-cucumerinum*, by lettuce soil amendment. *Crop Prot.* **24**: 135–140.
- Petersen, T.N., Brunak, S., von Heijne, G., and Nielsen, H. (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* **8**: 785–786.
- Recorbet, G., Steinberg, C., Olivain, C., Edel, V., Trouvelot, S., Dumas-Gaudot, E., et al. (2003) Wanted: pathogenesis-related marker molecules for *Fusarium oxysporum*. *New Phytol* **159**: 73–92.
- Rep, M., Van Der Does, H.C., Meijer, M., Van Wijk, R., Houterman, P.M., Dekker, H.L., et al. (2004) A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for I-3-mediated resistance in tomato. *Mol Microbiol* **53**: 1373–1383.
- de Sain, M. and Rep, M. (2015) The role of Pathogen-secreted proteins in fungal vascular wilt diseases. *Int J Mol Sci* **16**: 23970–23993.
- Schmidt, S.M., Houterman, P.M., Schreiber, I., Ma, L., Amyotte, S., Chellappan, B., et al. (2013) MITEs in the promoters of effector genes allow prediction of novel virulence genes in *Fusarium oxysporum*. *BMC Genomics* **14**: 119.
- Schmidt, S.M., Lukasiewicz, J., Farrer, R., van Dam, P., Bertoldo, C., and Rep, M. (2016) Comparative genomics of *Fusarium oxysporum* f. sp. *melonis* reveals the secreted protein recognized by the *Fom-2* resistance gene in melon. *New Phytol* **209**: 307–318.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., et al. (2014) Fast, scalable generation of high-quality protein multiple sequence alignments using clustal Omega. *Mol Syst Biol* **7**: 539–539.
- Sperschneider, J., Dodds, P.N., Gardiner, D.M., Manners, J.M., Singh, K.B., and Taylor, J.M. (2015a) Advances and challenges in computational prediction of effectors from plant pathogenic fungi. *PLoS Pathog* **11**: e1004806.



- Sperschneider, J., Gardiner, D.M., Thatcher, L.F., Lyons, R., Singh, K.B., Manners, J.M., and Taylor, J.M. (2015b) Genome-wide analysis in three fusarium pathogens identifies rapidly evolving chromosomes and genes associated with pathogenicity. *Genome. Biol Evol* **7**: 1613–1627.
- Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**: 1312–1313.
- Stanke, M., Schöffmann, O., Morgenstern, B., and Waack, S. (2006) Gene prediction in eukaryotes with a generalized hidden markov model that uses hints from external sources. *BMC Bioinformatics* **7**: 62.
- Stergiopoulos, I. and de Wit, P.J.G.M. (2009) Fungal effector proteins. *Annu Rev Phytopathol* **47**: 233–263.
- Takken, F. and Rep, M. (2010) The arms race between tomato and *Fusarium oxysporum*. *Mol Plant Pathol* **11**: 309–314.
- Vakalounakis, D.J. (1996) Root and stem rot of cucumber caused by *Fusarium oxysporum* f. sp. *radicis-cucumerinum* f. sp. nov. *Plant Dis* **80**: 313–316.
- Vakalounakis, D.J. and Fragkiadakis, G. A. (1999) Genetic diversity of *Fusarium oxysporum* isolates from cucumber: differentiation by pathogenicity, vegetative compatibility, and RAPD Fingerprinting. *Phytopathology* **89**: 161–168.
- Vakalounakis, D.J., Wang, Z., Fragkiadakis, G. A., Skaracis, G.N., and Li, D.B. (2004) Isolates obtained from cucumber in china by pathogenicity, VCG, and RAPD. *Plant Dis* **88**: 645–649.
- Vakalounakis, D.J., Doulis, A. G., and Klironomou, E. (2005) Characterization of *Fusarium oxysporum* f. sp. *radicis-cucumerinum* attacking melon under natural conditions in Greece. *Plant Pathol* **54**: 339–346.
- Wang, P.H., Lo, H.S., and Yeh, Y. (2001) Identification of *F. o. cucumerinum* and *F. o. luffae* by RAPD-generated DNA probes. *Lett Appl Microbiol* **33**: 397–401.
- Wellman, F.L. (1939) A technique for studying host resistance and pathogenicity in tomato fusarium wilt. *Phytopathology* **29**: 945–956.
- Zhou, X.G. and Everts, K.L. (2007) Characterization of a regional population of *Fusarium oxysporum* f. sp. *niveum* by race, cross pathogenicity, and vegetative Compatibility. *Phytopathology* **97**: 461–469.

### Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web-site.