



UvA-DARE (Digital Academic Repository)

Draft genome sequence of *Dethiobacter alkaliphilus* strain AHT1(T), a gram-positive sulfidogenic polyextremophile

Melton, E.D.; Sorokin, D.Y.; Overmars, L.; Lapidus, A.L.; Pillay, M.; Ivanova, N.; del Rio, T.G.; Kyrpides, N.C.; Woyke, T.; Muyzer, G.

DOI

[10.1186/s40793-017-0268-9](https://doi.org/10.1186/s40793-017-0268-9)

Publication date

2017

Document Version

Final published version

Published in

Standards in Genomic Sciences

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Melton, E. D., Sorokin, D. Y., Overmars, L., Lapidus, A. L., Pillay, M., Ivanova, N., del Rio, T. G., Kyrpides, N. C., Woyke, T., & Muyzer, G. (2017). Draft genome sequence of *Dethiobacter alkaliphilus* strain AHT1(T), a gram-positive sulfidogenic polyextremophile. *Standards in Genomic Sciences*, 12, [57]. <https://doi.org/10.1186/s40793-017-0268-9>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.


UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

EXTENDED GENOME REPORT

Open Access



Draft genome sequence of *Dethiobacter alkaliphilus* strain AHT1^T, a gram-positive sulfidogenic polyextremophile

Emily Denise Melton¹, Dimitry Y. Sorokin^{2,3}, Lex Overmars¹, Alla L. Lapidus⁴, Manoj Pillay⁶, Natalia Ivanova⁵, Tijana Glavina del Rio⁵, Nikos C. Kyrpides^{5,6,7}, Tanja Woyke⁵ and Gerard Muyzer^{1*} 

Abstract

Dethiobacter alkaliphilus strain AHT1^T is an anaerobic, sulfidogenic, moderately salt-tolerant alkaliphilic chemolithotroph isolated from hypersaline soda lake sediments in northeastern Mongolia. It is a Gram-positive bacterium with low GC content, within the phylum *Firmicutes*. Here we report its draft genome sequence, which consists of 34 contigs with a total sequence length of 3.12 Mbp. *D. alkaliphilus* strain AHT1^T was sequenced by the Joint Genome Institute (JGI) as part of the Community Science Program due to its relevance to bioremediation and biotechnological applications.

Keywords: Extreme environment, Soda lake, Sediment, Haloalkaliphilic, Gram-positive, *Firmicutes*

Introduction

Soda lakes are formed in environments where high rates of evaporation lead to the accumulation of soluble carbonate salts due to the lack of dissolved divalent cations. Consequently, soda lakes are defined by their high salinity and stable highly alkaline pH conditions, making them dually extreme environments. Soda lakes occur throughout the American, European, African, Asian and Australian continents and host a wide variety of *Archaea* and *Bacteria*, specialized at surviving under such high salt and high pH conditions [1]. These haloalkaliphiles drive a number of biogeochemical cycles essential to their survival, most notably; the sulfur cycle is very active in these unique habitats [2–4]. The most noteworthy taxa associated with the reductive sulfur cycle are the *Deltaproteobacteria* and the *Firmicutes*. Recently, a number of Gram-positive *Firmicutes* genomes have been analyzed and published describing their metabolic potential and environmental adaptations, including the polyextremophile *Natronaerobius thermophilus* [5], and species belonging to the *Desulfotomaculum* spp. [6–8] and the *Desulfosporosinus* spp. [9]. Here we give an extended insight into the first known

genome of a haloalkaliphilic Gram-positive sulfur disproportionator within the phylum *Firmicutes*: *Dethiobacter alkaliphilus* AHT1^T.

Organism information

Classification and features

The haloalkaliphilic anaerobe *D. alkaliphilus* AHT1^T was isolated from hypersaline soda lake sediments in northeastern Mongolia [10]. *D. alkaliphilus* AHT1^T cells are Gram-positive and the motile rod-shaped cells form terminal ellipsoid endospores (Fig. 1). The strain tolerates salt concentrations ranging from 0.2–0.8 M Na⁺ with an optimum at 0.4 M and is an obligate alkaliphile, growing within a pH range from 8.5–10.3 with an optimum at 9.5 [10]. Phylogenetic analysis showed that strain AHT1^T is a member of the phylum *Firmicutes* and the order *Clostridiales* (Fig. 2). Its closest relative is an acetate-oxidizing syntrophic alkaliphile, described as “*Candidatus* Contubernalis alkalaceticum” which was isolated from a soda lake [11] (Fig. 2). The 16S ribosomal RNA of *D. alkaliphilus* AHT1^T (EF422412) is 88% identical to the 16S rRNA of “*Candidatus* Contubernalis alkalaceticum” (DQ124682) [12].

Extended feature descriptions

D. alkaliphilus AHT1^T is an obligate anaerobe that can produce sulfide by using elemental sulfur and polysulfides

* Correspondence: g.muyzer@uva.nl

¹Department of Freshwater and Marine Ecology, Microbial Systems Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands

Full list of author information is available at the end of the article




Fig. 1 Morphology of *D. alkaliphilus* AHT1^T. **a** Phase contrast micrograph of cells. **b** Electron microscope image of a *D. alkaliphilus* AHT1^T cell

as electron acceptor [10]. Additionally, it has been shown to incompletely reduce thiosulfate to sulfide and sulfite with hydrogen or formate as electron donor [10]. Strain AHT1^T is the first representative from the *Firmicutes* with the metabolic capacity to grow by elemental sulfur disproportionation [13] and, therefore, is a very interesting organism to compare to the typical sulfur disproportionators from the *Deltaproteobacteria*. This species may play an important role in the reductive sulfur cycle in soda lake environments [2] and possibly also in other alkaline anaerobic habitats, such as serpentinization “cement springs”, where sequences closely related to *Dethiobacter* have been found [14, 15]. Also, its affiliation with the syntrophic *Clostridia* “*Candidatus* Contubernalis alkalaceticum” (Fig. 2) implies that *D. alkaliphilus* AHT1^T could be involved in syntrophic anaerobic metabolic activity. More classifications and features of this species are listed in Table 1.

Genome sequencing information

Genome project history

This organism was selected for sequencing at the JGI (<http://jgi.doe.gov>) based on its potential for bioremediation and biotechnological applications. It is part of the Community Science Program: Haloalkaliphilic sulfate-, thiosulfate- and sulfur-reducing bacteria (CSP_788492). The project is registered in the Genomes OnLine Database (Ga0028528) [16] and the permanent draft genome sequence is deposited in GenBank (RefSeq: NZ_ACJMO0000000.1). Draft sequencing and assembly were performed at the JGI using state of the art sequencing technology [17]. The project information is summarized in Table 2.

Growth conditions and genomic DNA preparation

Strain AHT1^T was grown anaerobically at 30 °C in Na-carbonate buffered mineral medium (22 g/L Na₂CO₃, 8 g/L NaHCO₃, 6 g/L NaCl, 1 g/L K₂HPO₄) with a pH of 10 and 0.6 M total Na⁺. Additionally, 4 mM NH₄Cl, 1 mM MgCl₂ × 6H₂O and 1 mL⁻¹ trace element solution were added [18]. After sterilization, acetate serving as carbon source (2 mM) and thiosulfate (20 mM) the electron-acceptor, were also added to the medium. The culture (2 L) was grown in a 10 L bottle mounted on a magnetic stirrer whereby the headspace (8 L) was replaced by 100% (v/v) H₂, at 0.5 Bar overpressure, acting as the electron-donor. Half the culture volume (1 L) was centrifuged at 13,000 g for 30 min, the pellet was washed with 1 M NaCl and frozen at -80 °C until further downstream processing. DNA was extracted from the pellet by the phenol-chloroform method after pre-treatment with SDS-proteinase K according to Marmur [19]. The concentration and molecular weight of the DNA were checked by UV spectroscopy and gel electrophoresis, respectively.

Genome sequencing and assembly

The size of the assembled *D. alkaliphilus* AHT1^T genome sequence was 3.12 Mbp. The draft genome was generated at the JGI using a combination of Sanger, Solexa/Illumina [20] and 454 DNA sequencing technologies [21]. An 8 Kb Sanger library was constructed that provided 2.5 x coverage of the genome (15,321 reads generated) and a Solexa shotgun library and a 454 Titanium standard library, which provided 25x genome coverage totalling 110.0 Mbp of 454 data. The 454 Titanium data were assembled with Newbler. The Newbler consensus sequences were computationally shredded into 2 Kb overlapping fake reads (shreds). Illumina sequencing data was assembled with VELVET, version 1.0.13 [22], and the consensus sequences were computationally shredded into 1.5 Kb overlapping fake reads (shreds). We then integrated Sanger reads, the 454 Newbler




Fig. 2 Neighbour-joining tree based on 16S rRNA gene sequences showing the phylogenetic position of *D. alkaliphilus* AHT1^T to other species within the phylum Firmicutes. The Deltaproteobacteria were used as an outgroup, but were pruned from the tree. The dots indicate bootstrap values between 80 and 100%. The scale bar indicates a 2% sequence difference. The tree was constructed with the ARB software package [48] and the SILVA database [29]. The bootstrap values were calculated using MEGA-6 [49]

Table 1 Classification and general features of *D. alkaliphilus* AHT1^T

MIGS ID	Property	Term	Evidence code
	Classification	Domain: <i>Bacteria</i> Phylum: <i>Firmicutes</i> Class: <i>Clostridia</i> Order: <i>Clostridiales</i> Family: <i>Syntrophomonadaceae</i> Genus: <i>Dethiobacter</i> Species: <i>Dethiobacter alkaliphilus</i> Type strain: AHT1 ^T	TAS [51] TAS [52–54] TAS [55, 56] TAS [57, 58] TAS [59] TAS [10, 60] TAS [10, 60] TAS [10]
	Gram stain	positive	TAS [10]
	Cell shape	rod-shaped	TAS [10]
	Motility	motile	TAS [10]
	Sporulation	endospore-forming	TAS [10]
	Temperature range	mesophile	TAS [10]
	Optimum temperature	33	
	pH range; Optimum	8.5-10.3; 9.5	TAS [10]
	Carbon source	CO ₂ , acetate	TAS [10]
MIGS-6	Habitat	hypersaline soda lakes, sediments	
MIGS-6.3	Salinity	moderately salt-tolerant	
MIGS-22	Oxygen requirement	anaerobe	
MIGS-15	Biotic relationship	free-living	
MIGS-14	Pathogenicity	none	
MIGS-4	Geographic location	northeastern Mongolia; lakes Hotontyn and Shar-Burdiin	TAS [2]
MIGS-5	Sample collection	September 1999	
MIGS-4.1	Latitude	48° 19' 40"	TAS [2]
MIGS-4.2	Longitude	114° 30' 16"	TAS [2]
MIGS-4.4	Altitude	1000 m	

Evidence codes - *IDA* Inferred from Direct Assay, *TAS* Traceable Author Statement (i.e., a direct report exists in the literature); *NAS* Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [Cite ontology project]

Table 2 Project information

MIGS ID	Property	Term
MIGS 31	Finishing quality	permanent draft
MIGS 28	Libraries used	Solexa
MIGS 29	Sequencing platforms	454
MIGS 31.2	Fold coverage	33.2
MIGS 30	Assemblers	Newbler, (2.0.00.20-PostRelease-11-05-2008-gcc-3.4.6), PGA [23], VELVET [22]
MIGS 32	Gene calling method	Prodigal [28]
	Locus Tag	DealDRAFT
	Genbank ID	ACJM00000000
	Genbank Date of Release	12.12.2013
	GOLD ID	Gp0001962
	BIOPROJECT	PRJNA30985
	Project relevance	bioremediation, environmental biotechnology

consensus shreds and the Illumina VELVET consensus shreds using the PGA assembler [23], to combine sequence data from all three platforms for a most contiguous assembly. The software Consed [24] was used in the computational finishing process as described previously [25]. The final draft assembly contained 34 contigs in 5 scaffolds.

Genome annotation

The assembled sequence was automatically annotated with the JGI prokaryotic annotation pipeline [26] with additional manual review using the IMG-ER platform [27]. Genes were predicted using Prodigal [28], ribosomal RNAs were detected using models built from SILVA [29] and tRNAs were predicted with tRNAScanSE [30]. The predicted CDs were translated and used to search the NCBI non-redundant database UniProt, TIGRFam, Pfam, KEGG, COG and InterPro databases. The final annotated genome is available from the IMG system [31]. We performed a CheckM analysis [32] and assessed that the genome is 95.8% complete.

Genome properties

The genome is 3,116,746 bp long with a GC content of 48.46%. A total of 3213 genes were found, of which 3163 coded for proteins and 50 genes encoded only RNA. From the total genes, 69.19% was assigned a putative function. The IMG taxon ID is 643,886,183. The different functional gene groups are summarized in Table 3. Furthermore, the number of genes assigned to functional COG categories is displayed in Table 4.

Table 3 Nucleotide content and gene count levels of the genome

Attribute	Value	% of total
Genome size (bp)	3,116,746	100
DNA coding (bp)	2,773,015	88.97
DNA G + C (bp)	1,510,353	48.46
DNA scaffolds	34	100
Total genes	3213	100
Protein coding genes	3163	98.44
RNA genes	50	1.56
Pseudo genes	0	0
Genes in internal clusters	177	not reported
Genes with function prediction	2223	69.19
Genes assigned to COGs	1971	61.34
Genes with Pfam domains	2632	81.92
Genes with signal peptides	170	5.29
Genes with trans-membrane helices	962	29.94
CRISPR repeats	0	0

Insights from the genome sequence

Extended insights: Metabolic potential

Hydrogen metabolism requires a number of hydrogenase operons, including the *hyd* operon, and a Ni-Fe metallo-center assembly (*hyp*) [33]. The first part of the hydrogenase *hyd* operon is the small hydrogenase subunit *hydA* located at DealDRAFT_1217, the closest NCBI BLAST hit [12] of this protein is the *hydA* gene in *Desulfotomaculum gibsoniae* (Desgi_1397) with 70.4% similarity in a pair-wise alignment [34]. Directly adjacent to *hydA*, is the large subunit *hydB* (DealDRAFT_1218) in the *D. alkaliphilus* AHT1^T genome. This subunit is most similar (75.9%) to the *hydB* subunit in *Dehalobacter* sp. UNSWDHB (UNSWDHB_1527) [12, 34]. DealDRAFT_1219 is a cytochrome B561 of 198 amino acids and could therefore be the interacting partner and gamma subunit *hydC* in the *hyd* operon. The 6-gene *hyp* operon *hypABCDEF* is responsible for the assemblage of the Ni-Fe uptake hydrogenases [35]. The last 5 proteins of the *hyp* operon are annotated in the *D. alkaliphilus* AHT1^T genome (DealDRAFT_0838-DealDRAFT_0842) and follow the organization *hypBFCDE*, as has been seen before in *Rhizobium* [36]. The first gene in the operon (DealDRAFT_0843) is a hypothetical protein of 88 nucleotides length and is assigned to pfam01155 *hypA*, which is 42.6% identical to the *hypA* gene in *Moorella thermoacetica*. Therefore, this hypothetical protein is most likely *hypA* in *D. alkaliphilus* AHT1^T. Using hydrogen as electron donor, *D. alkaliphilus* AHT1^T can grow autotrophically by fixing inorganic carbon through the Wood Ljungdahl pathway, the key genes are all present in the genome (Fig. 3a), including the *acs* gene


Table 4 Number of genes associated with general COG functional categories

Code	Value	% of total	Description
J	175	7.89	Translation, ribosomal structure and biogenesis
A	not reported	not reported	RNA processing and modification
K	134	6.04	Transcription
L	83	3.74	Replication, recombination and repair
B	1	0.05	Chromatin structure and dynamics
D	45	2.03	Cell cycle control, cell division, chromosome partitioning
V	58	2.62	Defense mechanisms
T	131	5.91	Signal transduction mechanisms
M	124	5.59	Cell wall/membrane biogenesis
N	52	2.35	Cell motility
U	34	1.53	Intracellular trafficking and secretion
O	90	4.06	Posttranslational modification, protein turnover, chaperones
C	178	8.03	Energy production and conversion
G	81	3.65	Carbohydrate transport and metabolism
E	227	10.24	Amino acid transport and metabolism
F	69	3.11	Nucleotide transport and metabolism
H	149	6.72	Coenzyme transport and metabolism
I	80	3.61	Lipid transport and metabolism
P	133	6.00	Inorganic ion transport and metabolism
Q	24	1.08	Secondary metabolites biosynthesis, transport and catabolism
R	183	8.25	General function prediction only
S	129	5.82	Function unknown
-	1242	38.66	Not in COGs


The total is based on the number of protein coding genes in the genome

cluster (Fig. 3b). Heterotrophic growth by *D. alkaliphilus* AHT1^T can be achieved with glucose and fructose [10], the entire glycolysis pathway is present in the genome (Fig. 4). Carbohydrate metabolism in *D. alkaliphilus* AHT1^T also includes oxidation of short chain organic acids; the tetrameric pyruvate oxidoreductase is present in the conformation *porBADC* (DealDRAFT_1244 – DealDRAFT_1247). Lactate dehydrogenases could not be found, although there is an L-lactate permease (DealDRAFT_0239), an L-lactate transport protein (DealDRAFT_1845) and a large and small subunit acetolactate synthase (DealDRAFT_2169 and 2170). For assimilation of acetate, strain AHT1^T has an acetyl coenzyme A synthetase (DealDRAFT_1887).

D. alkaliphilus AHT1^T might play a role in the reductive sulfur cycle in alkaline habitats since it grows as a thiosulfate and sulfur/polysulfide reducer or by sulfur disproportionation in laboratory cultures [10]. The



genome sequence contains a thiosulfate sulfurtransferase (DealDRAFT_1917), which is located directly adjacent to another sulfur transferase (Rhodanese domain DealDRAFT_1918). Both alpha and beta subunits of the adenylylsulfate reductase *apr* operon were also found (DealDRAFT_1379, DealDRAFT_1380). The *qmo* electron transfer complex, which usually accompanies the




apr operon [37], is not found. Key sulfur reduction genes such as *sat* (sulfate reduction), *dsr* (sulfite reduction) and *psr* (sulfur reduction) were also not found in this draft genome. As *D. alkaliphilus* AHT1^T can reduce and disproportionate elemental sulfur/polysulfide in laboratory

cultures [10, 13], the absence of these genes is surprising. It is conceivable however, that the sequencing quality of the permanent draft is insufficient to recover complete pathways. Indeed, CheckM analysis revealed that the genome was only 95.8% complete. Unfortunately, we can therefore not explain the key dissimilatory disproportionation mechanism from this genomic data. The genome also contains some assimilatory sulfate reduction genes, such as *cysND* (DealDRAFT_1193 and DealDRAFT_1192).

Extended insights: Haloalkaliphilic adaptations

In order to generate ATP, *D. alkaliphilus* AHT1^T has an *ntp* gene operon encoding a vacuolar ATP synthase (V₀V₁-type) (DealDRAFT_1677 – DealDRAFT_1685) (Fig. 5a). This operon structure is conserved among the *Clostridia* (Fig. 5b). The *ntp* operon encodes the ATP synthase for ATP generation and follows the GILEX-FABD organization in the *Deinococcus-Thermus* phylum [38]. In the *Firmicutes*, the gene organization is slightly different at GIKECFABD (Fig. 5a, b). In *D. alkaliphilus* AHT1^T these genes are located from DealDRAFT_1685



(*ntpG*) to DealDRAFT_1677 (*ntpD*). The *ntpD* subunit within the operon is annotated as being of the V-type. In order to confirm that the ATP synthase is indeed V-type [39], we constructed a phylogenetic tree of the transmembrane *c/K* subunits of *Firmicutes* known specifically to be V- or F-type [40] and NCBI annotation] and aligned the *D. alkaliphilus* AHT1^T *ntpC* sequence (DealDRAFT_1683) with these other sequences (Fig. 6a) [41]. As seen before, there was a clear separation between V-type and F-type ATP synthase, where the AHT1^T sequence clustered together with the V-type ATP synthase. In addition, the sequences are tentatively clustered into separate H⁺ or Na⁺ coupled ATPase branches. The AHT1^T sequence was positioned within a Na⁺ coupled V-type ATP synthase group, indicating that this organism's ATP synthase is coupled specifically to Na⁺ translocation across the membrane. In order to explore this further, we looked at specific Na⁺ binding residues and ligands on the transmembrane *c/K* subunit [40], and

created a Weblogo for the Na⁺ specific *Firmicutes* V-type ATP synthase (Fig. 6b) [42, 43]. When we aligned the *ntpC* sequence of *D. alkaliphilus* AHT1^T we found that it contains all the conserved five amino acids (Ser26, Leu57, Thr60, Gln61 and Tyr64) specific for Na⁺ translocation [40] (Fig. 6c). Thus, the *D. alkaliphilus* AHT1^T genome contains a Na⁺ coupled V-type ATP synthase.

In order to import protons to retain the intracellular pH, the genome contains the multi-subunit electrogenic sodium/proton antiporter *mrp* (DealDRAFT_2487–2497), that pumps protons into the cell and sodium out of the cell [44]. To retain osmotic balance, *D. alkaliphilus* AHT1^T has numerous substrate binding regions and transporters for glycine betaine (e.g. DealDRAFT_2378, _2380 and DealDRAFT2842, _2844), leading to the conclusion that osmoprotectants are used to maintain cellular turgor pressure, instead of the salt-in strategy. Another necessity for alkaliphilic bacteria is to prevent




Fig. 6 a Phylogeny of the F- vs. V-type ATPase within the *Firmicutes*. Numbers on the tree nodes indicate bootstrap values (1000 replicates). Scale bar indicates 0.2% sequence difference. **b** Weblogo of conserved region within the *ntpC/K* *Firmicute* subunit [42, 43]. **c** Weblogo of aligned *D. alkaliphilus* AHT1^T subunit *ntpC* (DealDRAFT_1683) where conserved Na⁺ binding regions (in B and C) are indicated with black arrows

proton leakage from cells, which they can achieve through structural membrane adaptations [1]. The genome contains the genes to synthesize the squalene precursors dimethylallyl diphosphate and isopentenylallyl diphosphate through the non-mevalonate pathway [45]. The accompanying locus tags within the KEGG non-mevalonate pathway (M00096) are *dxs* (DealDRAFT_0731), *dxr/ispC* (DealDRAFT_2409), *ispD* (DealDRAFT_2331), *ispE* (DealDRAFT_2584), *ispF* (DealDRAFT_2332), *ispG* (DealDRAFT_2411) and *ispH* (DealDRAFT_0659). However, we did not find genes similar to *hpnCDE*, which function in the formation of squalene from its precursors [46]. Thus, *D. alkaliphilus* AHT1^T does not seem to have this membrane adaptation to haloalkaline environments, although it could also be due to the incompleteness of the genome. Nevertheless, it has been shown that *Bacillus lentus* C-125, also a *Firmicute*, survives in the haloalkaline environment by increased levels of acidic polymers in its cellular membrane resulting in a cell wall negative charge [47]. It is possible that *D. alkaliphilus* AHT1^T supports a similar mechanism to survive the alkaline pH values of its environment.

Conclusions

In this manuscript we globally characterize the genome of *D. alkaliphilus* AHT1^T, which was isolated from hypersaline soda lakes sediment in north-eastern Mongolia. Investigation of the genome of this anaerobic sulfidogen identified genes for the Wood-Ljungdahl pathway (autotrophic growth, Fig. 3) and the Embden-Meyerhof pathway (heterotrophic growth Fig. 4). Thus the carbon metabolism of this microbe is fairly versatile. *D. alkaliphilus* AHT1^T is capable of disproportionation in laboratory cultures, thus future genomic analyses with qPCR may provide insights into the disproportionation of sulfur compounds. *D. alkaliphilus* AHT1^T is well adapted to the haloalkaline environment, we found genes for active energy generation with a sodium V-type ATP synthase (Fig. 6). In addition, transporters for the osmoprotectants glycine and betaine were found to maintain cellular homeostasis and protection from the saline external environment. Further research will extend our knowledge on the ecophysiology of haloalkaliphiles, their role in nutrient cycling in extreme environments and their adaptations to this polyextreme environment. Moreover, insight in the genome sequence and subsequent transcriptomic or proteomic analysis will be helpful to infer the potential role of *D. alkaliphilus* AHT1^T in the biotechnological removal of sulfur compounds from wastewater and gas streams.

Abbreviations

F-type: Phosphorylation factor-type; IMG: Integrated Microbial Genomes; IMG-ER: Integrated Microbial Genomes - Expert Review; JGI: Joint

Genome Institute; NCBI: National Center for Biotechnology Information; THF: tetrahydrofolate; V-type: Vacuole-type

Acknowledgements

Emily Denise Melton, Lex Overmars and Gerard Muyzer are supported by ERC Advanced Grant PARASOL (No. 322551); Dimitry Sorokin is supported by RFBR grant 16-04-00035 and by the Gravitation (SIAM) program (grant 24002002, Dutch Ministry of Education and Science). Alla L. Lapidus is supported by the St. Petersburg State University grant 15.61.951.2015. The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, was supported under Contract No. DE-AC02-05CH11231.

Authors' contributions

EDM drafted and wrote the manuscript. DYS, GM, LO, NCK and ALL contributed to the written manuscript. LO, DYS and GM stimulated critical discussions. DS cultured AHT1 and extracted the DNA. The sequencing and annotation of the genome were performed at the JGI by ALL, MP, NI, TGR, NCK and TW. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interest.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Freshwater and Marine Ecology, Microbial Systems Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands. ²Winogradsky Institute of Microbiology, Research Centre of Biotechnology, RAS, Moscow, Russia. ³Department of Biotechnology, Delft University of Technology, Delft, The Netherlands. ⁴Center for Algorithmic Biotechnology, Institute of Translational Biomedicine, St. Petersburg State University, St. Petersburg, Russia. ⁵Joint Genome Institute, Walnut Creek, CA, USA. ⁶Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. ⁷Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia.

Received: 23 March 2017 Accepted: 8 September 2017

Published online: 21 September 2017

References

- Sorokin DY, Berben T, Melton ED, Overmars L, Vavourakis CD, Muyzer G. Microbial diversity and biogeochemical cycling in soda lakes. *Extremophiles*. 2014;18:791–809.
- Sorokin DY, Gorlenko VM, Namsaraev BB, Namsaraev ZB, Lysenko AM, Eshinimaev BT, Khmelenina VN, Trotsenko YA, Kuenen JG. Prokaryotic communities of the north-eastern Mongolian soda lakes. *Hydrobiologia*. 2004;522:235–48.
- Sorokin DY, Rusanov I, Pimenov NV, Tourouva TP, Abbas B, Muyzer G. Sulfidogenesis under extremely haloalkaline conditions in soda lakes of Kulunda steppe (Altai, Russia). *FEMS Microbiol Ecol*. 2010;73:278–90.
- Sorokin DY, Kuenen JG, Muyzer G. The microbial sulfur cycle at extremely haloalkaline conditions of soda lakes. *Front Microbiol*. 2011;2:44.
- Zhao B, Mesbah NM, Dalin E, Goodwin L, Nolan M, Pitluck S, et al. Complete genome sequence of the anaerobic, halophilic alkalithermophile *Natronaerobius thermophilus* JW/NM-WN-LF. *J Bacteriol*. 2011;193:4023–4.
- Junier P, Junier T, Podell S, Sims DR, Detter JC, Lykidis A, et al. The genome of the gram-positive metal- and sulfate-reducing bacterium *Desulfotomaculum reducens* strain MI-1. *Environ Microbiol*. 2010;12:2738–54.
- Aüillo T, Ranchou-Peyruse A, Ollivier B, Magot M. *Desulfotomaculum* spp. and related gram-positive sulfate-reducing bacteria in deep subsurface environments. *Front Microbiol*. 2013;4:362.
- Visser M, Parshina SN, Alves JJ, Sousa DZ, Pereira IAC, Muyzer G, et al. Genome analyses of the carboxydophilic sulfate-reducers *Desulfotomaculum nigrificans* and *Desulfotomaculum carboxydivorans* and reclassification of *Desulfotomaculum nigrificans*. *Stand Genomic Sci*. 2014;9:655–75.
- Pester M, Brambilla E, Alazard D, Rattei T, Weinmaier T, Han J, et al. Complete genome sequences of *Desulfosporosinus orientis* DSM765T,

- Desulfosporosinus youngiae* DSM17734T, *Desulfosporosinus meridiei* DSM13257T, and *Desulfosporosinus acidiphilus* DSM22704T. *J Bacteriol.* 2012;194:6300–1.
10. Sorokin DY, Tourova TP, Mussmann M, Muyzer G. *Dethiobacter alkaliphilus* gen. nov. sp. nov., and *Desulfurivibrio alkaliphilus* gen. nov. sp. nov.: two novel representatives of reductive sulfur cycle from soda lakes. *Extremophiles.* 2008;12:431–9.
 11. Zhilina TN, Zavarzina DG, Kolganova TV, Tourova TP, Zavarzin GA. “*Candidatus* Contubernalis alkalaceticum,” an obligately syntrophic alkaliphilic bacterium capable of anaerobic acetate oxidation in a coculture with *Desulfonatronum cooperativum*. *Microbiology.* 2005;74:800–9.
 12. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215:403–10.
 13. Poser A, Lohmayer R, Vogt C, Knoeller K, Planer-Friedrich B, Sorokin D, et al. Disproportionation of elemental sulfur by haloalkaliphilic bacteria from soda lakes. *Extremophiles.* 2013;17:1003–12.
 14. Tiago I, Verissimo A. Microbial and functional diversity of a subterrestrial high pH groundwater associated to serpentinization. *Environ Microbiol.* 2012;15:1687–706.
 15. Suzuki S, Ishiia S, Wua A, Cheung A, Tenneya A, Wangera G, et al. Microbial diversity in the cedars, an ultrabasic, ultrareducing, and low salinity serpentinizing ecosystem. *Proc Nat Acad Sci USA.* 2013;110:15336–41.
 16. Reddy TBK, Thomas AD, Stamatis D, Bertsch J, Isbandi M, Jansson J, et al. The Genomes OnLine Database (GOLD) v.5: a metadata management system based on a four level (metal)genome project classification. *Nucleic Acids Res.* 2014;43:1099–106.
 17. Mavromatis K, Land ML, Brettin TS, Quest DJ, Copeland A, Clum A, et al. The fast changing landscape of sequencing technologies and their impact on microbial assemblies and annotations. *PLoS ONE* 2012;doi: <https://doi.org/10.1371/journal.pone.0048837>.
 18. Pfenning N, Lippert KD. Über das Vitamin B12-Bedürfnis phototropher Schwefelbakterien. *Arch Mikrobiol.* 1966;55:245–56.
 19. Marmur J. A procedure for isolation of DNA from microorganisms. *J Mol Biol.* 1961;3:208–14.
 20. Bennett S. Solexa Ltd. *Pharmacogenomics.* 2004;5:433–8.
 21. Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, et al. Genome sequencing in microfabricated high-density picolitre reactors. *Nature.* 2005;437:326–7.
 22. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 2008;18:821–9.
 23. Zhao F, Zhao F, Li T, Bryant DA. A new pheromone trail-based genetic algorithm for comparative genome assembly. *Nucl Acids Res.* 2008;36:3455–62.
 24. Gordon D, Abajian C, Green P. Consed: a graphical tool for sequence finishing. *Genome Res.* 1998;8:195–202.
 25. Sims D, Brettin T, Dettler JC, Han C, Lapidus A, Copeland A, et al. Complete genome sequence of *Kytococcus sedentarius* type strain (541^T). *Stand Genomic Sci.* 2009;1:12–20.
 26. Mavromatis K, Ivanova NN, Chen IM, Szeto E, Markowitz VM, Kyrpides NC. The DOE-JGI standard operating procedure for the annotations of microbial genomes. *Stand Genomic Sci.* 2009;1:63–7.
 27. Markowitz VM, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics.* 2009;25:2271–8.
 28. Hyatt D, Chen G, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinf.* 2010;11:119.
 29. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl Acids Res.* 2013; doi: [10.1093/nar/gks1219](https://doi.org/10.1093/nar/gks1219).
 30. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucl Acids Res.* 1997;25:955–64.
 31. Markowitz VM, Chen I-M A, Palaniappan K, Chu K, Szeto E, Grechkin Y, et al. IMG: the integrated microbial genomes database and comparative analysis system. *Nucleic Acids Res.* 2012;40:D115–22.
 32. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 2015;25:1043–55.
 33. Schwartz E, Fritsch J, Friedrich B. H2-Metabolizing prokaryotes. In: Rosenberg E, DeLong E. F., Lory S., Stackebrandt E., Thompson F., editors. *The Prokaryotes*. Berlin: Springer; Verlag; 2013. p. 119–199.
 34. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Lopez R, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal omega. *Mol Systems Biol.* 2011;7:539.
 35. Agervald Å, Stensjö K, Holmqvist M, Lindblad P. Transcription of the extended *hyp*-operon in *Nostoc* sp. strain PCC 7120. *BMC Microbiol.* 2008;8:69.
 36. Hernando Y, Palacios JM, Imperial J, Ruiz-Argüeso T. The *hypBFCDE* operon from *Rhizobium leguminosarum* biovar viciae is expressed from an Fnr-type promoter that escapes mutagenesis of the *fnrN* gene. *J Bacteriol.* 1995;177:5661–9.
 37. Duarte AG, Santos AA and Pereira IAC electron transfer between the QmoABC membrane complex and adenosine 5′phosphosulfate reductase. *Biochim Biophys Acta.* 1857;2016:380–6.
 38. Yokoyama K, Ohkuma S, Taguchi H, Yasunaga T, Wakabayashi T, Yoshida M. V-type H+ ATPase/synthase from a thermophilic eubacterium, *Thermus thermophilus*. *J Biol Chem.* 2000;275:13955–61.
 39. Hicks DB, Liu J, Fujisawa M, Krulwich TA. F₁F₀-ATP synthases of alkaliphilic bacteria: lessons from their adaptations. *Biochim Biophys Acta.* 2010;1797:1362–77.
 40. Mulikidjanian AY, Galperin MY, Makarova KS, Wolf YI, Koonin EV. Evolutionary primacy of sodium bioenergetics. *Biol Direct.* 2008;3:13.
 41. Li W, Cowley A, Uludag M, Gur T, McWilliam H, Squizzato S, et al. The EMBL-EBI bioinformatics web and programmatic tools framework. *Nucleic Acids Res.* 2015;43:580–4.
 42. Schneider TD, Stephens RM. Sequence logos: a new way to display consensus sequences. *Nucleic Acids Res.* 1990;18:6097–100.
 43. Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: A sequence logo generator. *Genome Res.* 2004;14:1188–90.
 44. Krulwich TA, Sachs G, Padan E. Molecular aspects of bacterial pH sensing and homeostasis. *Nat Rev Microbiol.* 2011;9:330–43.
 45. Eisenreich W, Bacher A, Rigoni D, Rohdich F. Biosynthesis of isoprenoids 544 via the non-mevalonate pathway. *Cell Mol Life Sci.* 2004;61:1401–26.
 46. Pan JJ, Solbiati JO, Ramamoorthy G, Hillerich BS, Seidel RD, Cronan JE, Almo SC, Poulter CD. Biosynthesis of squalene from farnesyl diphosphate in bacteria: three steps catalysed by three enzymes. *ACS Cent Sci.* 2015;1:77–82.
 47. Aono R, Ito M, Joblin KN, Horikoshi K. A high cell wall negative charge is necessary for the growth of the alkaliphile *Bacillus lentus* C-125 at elevated pH. *Microbiology.* 1995;141:2955–64.
 48. Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadukumar, et al. ARB: a software environment for sequence data. *Nucl Acids Res.* 2004;32:1363–71.
 49. Tamura K, Stecher G, Peterson D, Filipiński A, Kumar S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol.* 2013;30:2725–9.
 50. Overmars L, Kerkhoven R, Siezen RJ, Francke C. MGCV: the microbial genomic context viewer for comparative genome analysis. *BMC Genomics.* 2013;14:209.
 51. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, bacteria, and Eucarya. *Proc Natl Acad Sci U S A.* 1990;87:4576–9.
 52. Gibbons NE, Murray RGE. Proposals concerning the higher taxa of bacteria. *Int J Syst Bacteriol.* 1978;28:1–6.
 53. Garrity GM, Holt JG. The road map to the manual. In: Garrity GM, Boone DR, Castenholz RW, editors. *Bergey's Manual of Systematic Bacteriology*, 2nd ed. Volume 1, New York: Springer; 2001. p. 119–169.
 54. Murray RGE. The higher taxa, or, a place for everything...? In: Holt JG editor. *Bergey's Manual of Systematic Bacteriology*, 1st ed. Volume 1, The Williams and Wilkins Co., Baltimore; 1984. p. 31–34.
 55. List of new names and new combinations previously effectively, but not validly, published. List no. 132. *Int J Syst Evol Microbiol.* 2010;60:469–472.
 56. Rainey FA. Class II. *Clostridia* class nov. In: De Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, Schleifer KH, Whitman WB, editors. *Bergey's Manual of Systematic Bacteriology*, 2nd ed. Volume 3, Springer-Verlag, New York; 2009. p. 736.
 57. Skerman VBD, McGowan V, Sneath PHA. Approved lists of bacterial names. *Int J Syst Bacteriol.* 1980;30:225–420.
 58. Prévot AR. In: Hauderoy P, Ehringer G, Guillot G, Magrou J, Prévot AR, Rosset D, Urbain A, editors. *Dictionnaire des Bactéries Pathogènes*. 2nd ed. Paris: Masson et Cie; 1953. p. 1–692.
 59. Jumas-Bilak E, Roudière L, Marchandin H. Description of ‘*Synergistetes*’ phyl. nov. and emended description of the phylum ‘*Deferribacteres*’ and of the family *Syntrophomonadaceae*, phylum ‘*Firmicutes*’. *Int J Syst Bacteriol.* 2009; 59:1028–35.
 60. Euzéby J. Validation list no. 123. *Int J Syst Evol Microbiol.* 2008;58:1993–1994.