



UvA-DARE (Digital Academic Repository)

Testing scientific models using Qualitative Reasoning: Application to cellulose hydrolysis

Kansou, K.; Rémond, C.; Paës, G.; Bonnin, E.; Tayeb, J.; Bredeweg, B.

DOI

[10.1038/s41598-017-14281-4](https://doi.org/10.1038/s41598-017-14281-4)

Publication date

2017

Document Version

Other version

Published in

Scientific Reports

[Link to publication](#)

Citation for published version (APA):

Kansou, K., Rémond, C., Paës, G., Bonnin, E., Tayeb, J., & Bredeweg, B. (2017). Testing scientific models using Qualitative Reasoning: Application to cellulose hydrolysis. *Scientific Reports*, 7, [14122]. <https://doi.org/10.1038/s41598-017-14281-4>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Supplementary Information

For “Testing scientific models using Qualitative Reasoning: Application to cellulose hydrolysis”

Kamal Kansou^{1*}, Caroline Rémond², Gabriel Paës², Estelle Bonnin¹, Jean Tayeb², Bert Bredeweg³

¹INRA, Biopolymères Interactions et Assemblages, BP 71267, 44316 Nantes, France

²FARE laboratory, INRA, University of Reims Champagne-Ardenne, 51100 Reims, France

³Informatics Institute, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

*To whom correspondence should be addressed. Tel: +33 (0)2 40 67 51 49, Email: kamal.kansou@inra.fr

APPENDIX

Model M1 encompassment test information

Reduced amount of Free enzyme. Figure S1 simulation curves are produced from a model built after equations given in Maurer et al.¹, with inputs describing a reduced amount of free enzyme compared to the available surface. In Figure S1, one can notice the consecutive monotonic increase of the amount of adsorbed enzyme (red curve) and then amount of complexed enzyme (blue curve), prior to stabilization. Neither the adsorption rate nor the catalytic rate decrease. Figure S1 depicts a limitative absorption rate compared to complexation and desorption rates. QR Model M1, produces the pathway [1→2→3→4→5] (Fig. 10ab), which encompasses this behaviour.




Figure S1. Simulation from kinetic model described in Maurer et al.,¹, with limitative amount of Free enzyme for 100h hydrolysis

Excess of Free enzyme. Figure S2 simulation curves are produced with an excess of Free enzyme compared to the available surface. The amount of Free enzyme is too high to be shown on the graph. One can notice the peak of adsorbed surface enzyme (red curve), which denotes a bottleneck effect at the adsorption stage due to limitative complexation rate. QR Model M1 produces the pathway [1→2→3→4→6→7→5] (Fig. 10ab), which encompasses this behaviour.






Figure S2. Simulation from kinetic model described in Maurer et al.¹, with excess of Free enzyme for 100h hydrolysis. One can notice the peak of adsorbed enzyme.

Model M1, sufficiency test information

a)



b)



c)




Figure S3. Results for M1 sufficiency test for TB2 and TB2' (restart scenario). Partial state-graph of model M1 (a). Value histories of selected quantities depicting the behaviour path [1→2→3→4] consistent with TB2 (Tab. 3) (b). Value histories of selected quantities depicting the behaviour path [1→2→3→6→10→15→19→21→26→29→30→37→4] consistent with TB2' (Tab. 4) (c).

Model M3, simulation results




Figure S4. Simulation results for M3. State-graph of model M3 of 41 distinct states with one end-state (state 4) (a). Value histories of selected quantities for model M3 depicting the behaviour path [1→2→3→5→6→10→4] (b).

References

- 1 Maurer, S.A., Bedbrook, C.N., & Radke, C.J., Cellulase Adsorption and Reactivity on a Cellulose Surface from Flow Ellipsometry. *Industrial & Engineering Chemistry Research* 51 (35), 11389-11400 (2012).