



UvA-DARE (Digital Academic Repository)

Bayesian estimation of Kendall's τ using a latent normal approach

van Doorn, J.; Ly, A.; Marsman, M.; Wagenmakers, E.-J.

DOI

[10.1016/j.spl.2018.10.004](https://doi.org/10.1016/j.spl.2018.10.004)

Publication date

2019

Document Version

Final published version

Published in

Statistics and Probability Letters

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

van Doorn, J., Ly, A., Marsman, M., & Wagenmakers, E.-J. (2019). Bayesian estimation of Kendall's τ using a latent normal approach. *Statistics and Probability Letters*, 145, 268-272. <https://doi.org/10.1016/j.spl.2018.10.004>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



Bayesian estimation of Kendall's τ using a latent normal approach



Johnny van Doorn^{a,*}, Alexander Ly^{a,b}, Maarten Marsman^a,
Eric-Jan Wagenmakers^a

^a University of Amsterdam, The Netherlands

^b Centrum Wiskunde & Informatica, The Netherlands

ARTICLE INFO

Article history:

Received 6 March 2017

Received in revised form 1 October 2018

Accepted 3 October 2018

Available online 15 October 2018

Keywords:

Semi-parametric inference

Rank correlation

ABSTRACT

The rank-based association between two variables can be modeled by introducing a latent normal level to ordinal data. We demonstrate how this approach yields Bayesian inference for Kendall's τ , improving on a recent Bayesian solution based on its asymptotic properties.

© 2018 Elsevier B.V. All rights reserved.

1. A Bayesian framework for Kendall's τ

Kendall's τ is a popular rank-based correlation coefficient. Compared to Pearson's ρ , Kendall's τ is robust to outliers, invariant under monotonic transformations, and has an intuitive interpretation (Kendall and Gibbons, 1990). Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ be two data vectors each containing ranked measurements of the same n units. For instance, x could be the rank ordered scores on a math exam and y the rank ordered scores on a geography exam, for n test-takers. A *concordant* pair is defined as a pair of subjects (i, j) where subject i has a higher score on x and y compared to subject j , whereas a *discordant* pair is defined as one where i scores higher on y , but j scores higher on x , or the other way around. Kendall's τ is defined as the difference between the number of concordant and discordant pairs, expressed as proportion of the total number of pairs:

$$\tau = \frac{\sum_{1 \leq i < j \leq n} Q((x_i, y_i), (x_j, y_j))}{n(n-1)/2}, \tag{1}$$

where the denominator is the total number of pairs and Q is the concordance indicator function, which is defined by:

$$Q((x_i, y_i), (x_j, y_j)) = \begin{cases} -1 & \text{if } (x_i - x_j)(y_i - y_j) < 0, \\ +1 & \text{if } (x_i - x_j)(y_i - y_j) > 0, \end{cases} \tag{2}$$

which returns -1 if a pair is discordant, and returns $+1$ if a pair is concordant. However, due to the nonparametric nature of Kendall's τ and the lack of a likelihood function for the data, Bayesian inference is not trivial.

An innovative method for overcoming this problem was proposed by Johnson (2005), and involves the modeling of the test statistic itself, rather than the data. This method has been applied to Kendall's τ by Yuan and Johnson (2008), and was

* Correspondence to: University of Amsterdam, Department of Psychological Methods, Valckeniersstraat 59, 1018 XA Amsterdam, The Netherlands.
E-mail address: JohnnyDoorn@gmail.com (J. van Doorn).




Fig. 1. A graphical model of the latent normal method. Here, x and y are observed rank data. The latent level is denoted with z^x and z^y , and ρ_{z^x, z^y} represents the latent correlation.

recently developed by [van Doorn et al. \(2018\)](#). The inferential framework that follows from this work uses the limiting normal distribution of the test statistic T^* ([Hotelling and Pabst, 1936](#); [Noether, 1955](#)), where

$$T^* = \tau \sqrt{\frac{9n(n-1)}{4n+10}}. \tag{3}$$

Under \mathcal{H}_0 , this limiting normal distribution is the standard normal, whereas under \mathcal{H}_1 , this distribution is specified with a non-centrality parameter Δ for the mean, and a sampling variance of 1.

However, the method –henceforth the *original asymptotic method* –might fall short on two counts. Firstly, the asymptotic assumptions only hold for sufficiently large n (i.e., $n \geq 20$, see [van Doorn et al., 2018](#)). Secondly, the variance of the sampling distribution of the test statistic depends on the population value of Kendall’s τ . For $\tau = 0$, the sampling variance equals 1, but as $|\tau| \rightarrow 1$, the variance decreases to 0 ([Kendall and Gibbons, 1990](#); [Hotelling and Pabst, 1936](#)).

In the current article, we will explore two corrections that aim to improve Bayesian inference for Kendall’s τ :

1. Within the asymptotic framework, the observed value of Kendall’s τ can be used to set its sampling variance. We label this the *enhanced asymptotic method*.
2. Within a Bayesian latent normal framework, a latent level correlation is obtained and transformed to Kendall’s τ . We label this the *latent normal method*.

2. Correction using the sample τ

A first correction to consider is to use the sample value of Kendall’s τ , denoted τ_{obs} , to estimate the sampling variance of T^* , denoted $\sigma_{T^*}^2$. A convenient expression for the upper bound of $\sigma_{T^*}^2$ in terms of τ is given in [Kendall and Gibbons \(1990\)](#):

$$\sigma_{T^*}^2 \leq \frac{2.5n(1-\tau^2)}{2n+5}. \tag{4}$$

Using τ_{obs} as an estimate of τ provides a somewhat crude approximation to the sampling distribution of T^* . However, compared to using $\sigma_{T^*}^2 = 1$ as in the original asymptotic method, working with the upper bound will result in a more narrow posterior for cases where $\tau \neq 0$. However, the enhanced asymptotic method still suffers from the use of asymptotic assumptions about the sampling distribution and variance of the test statistic.

3. Correction using the latent normal approach

3.1. Latent normal models

Several latent variable models quantify the association between two ordinal variables. These methods often introduce a latent bivariate normal distribution to the ordinal variables, where the association between variables is modeled through a latent correlation ([Pearson, 1900](#); [Olssen, 1979](#); [Pettitt, 1982](#); [Albert, 1992](#); [Alvo and Yu, 2014](#)). The observed rank data (x, y) can then be seen as the ordinal manifestations of the continuous latent variables (z^x, z^y) , which have a bivariate normal distribution. [Fig. 1](#) offers a graphical representation of such a model. Using this methodology, the nonparametric problem of ordinal analysis is transformed to a parametric data augmentation problem.

3.2. Posterior distribution for the latent correlation

The joint posterior can be decomposed as follows:

$$P(z^x, z^y, \rho_{z^x, z^y} \mid x, y) \propto P(x, y \mid z^x, z^y) \times P(z^x, z^y \mid \rho_{z^x, z^y}) \times P(\rho_{z^x, z^y}). \tag{5}$$

The second factor on the right-hand side is the bivariate normal distribution of the latent scores given the latent correlation:

$$\begin{pmatrix} z^x \\ z^y \end{pmatrix} \sim \mathcal{N} \left[\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \rho_{z^x, z^y} \\ \rho_{z^x, z^y} & 1 \end{pmatrix} \right]. \tag{6}$$

The factor $P(x, y | z^x, z^y)$ consists of a set of indicator functions that map the observed ranks to latent scores, such that the ordinal information is preserved. For the value z_i^x , this means that its range is truncated by the lower and upper thresholds that are respectively defined as:

$$a_i^x = \max_{j: x_j < x_i} (z_j^x) \tag{7}$$

$$b_i^x = \min_{j: x_j > x_i} (z_j^x). \tag{8}$$

The third factor is the prior distribution on the latent correlation. In the remainder of this article, the prior is specified by a uniform distribution on $(-1, 1)$ (but see Berger and Sun, 2008; Ly et al., 2016).

The general Bayesian framework for estimating the latent correlation involves data augmentation through a Gibbs sampling algorithm (Geman and Geman, 1984) for the latent values z^x and z^y , combined with a random walk Metropolis–Hastings sampling algorithm for ρ_{z^x, z^y} . At sampling step s :

1. For each value of z_i^x , sample from a truncated normal distribution:

$$(z_i^x | z_i^x, z_i^y, \rho_{z^x, z^y}) \sim N(z_i^y \rho_{z^x, z^y}, 1 - \rho_{z^x, z^y}^2) \mathbf{1}_{(a_i^x, b_i^x)}(z_i^x),$$

where $\mathbf{1}_{(a_i^x, b_i^x)}(z_i^x)$ indicates truncation between the lower threshold a_i^x given in (7) and the upper threshold b_i^x given in (8).

2. For each value of z_i^y , the sampling procedure is analogous to step 1.
3. Sample a new proposal for ρ_{z^x, z^y} , denoted ρ_{z^x, z^y}^* , from the asymptotic normal approximation to the sampling distribution of Fisher’s z-transform of ρ (Fisher, 1915):

$$\tanh^{-1}(\rho_{z^x, z^y}^*) \sim N \left(\tanh^{-1}(\rho_{z^x, z^y}^{s-1}), \frac{1}{n-3} \right).$$

The acceptance rate α is determined by the likelihood ratio of $(z^x, z^y | \rho_{z^x, z^y}^*)$ and $(z^x, z^y | \rho_{z^x, z^y}^{s-1})$, where each likelihood is determined by the bivariate normal distribution in (6):

$$\alpha = \min \left(1, \frac{P(z^x, z^y | \rho_{z^x, z^y}^*)(1 - (\rho_{z^x, z^y}^*)^2)}{P(z^x, z^y | \rho_{z^x, z^y}^{s-1})(1 - (\rho_{z^x, z^y}^{s-1})^2)} \right),$$

where $(1 - \rho^2)$ is the Jacobian of Fisher’s z-transform.

Repeating the algorithm a sufficient number of times yields samples from ρ_{z^x, z^y} given z^x, z^y , thus, $P(\rho_{z^x, z^y} | z^x, z^y)$, and the posterior of z^x, z^y , that is, $P(z^x, z^y | x, y)$.

3.3. Relation to Kendall’s τ

With the posterior distribution for the latent ρ_{z^x, z^y} in hand, the transition to the posterior distribution for Kendall’s τ can be made using Greiner’s relation (Greiner; Kruskal, 1958). This relation, defined as

$$\tau = G(\rho) = \frac{2}{\pi} \sin^{-1}(\rho) \tag{9}$$

enables the transformation of Pearson’s ρ to Kendall’s τ when the data follow a bivariate normal distribution. The latent normal framework thus models the posterior distribution for Kendall’s τ as

$$P(\tau | x, y) = P(G(\rho) | x, y) = \int \int P(G(\rho) | z^x, z^y) P(z^x, z^y | x, y) dz^x dz^y.$$

Introducing the latent normal level to the observed variables enables the link between Pearson’s ρ and Kendall’s τ , and turns posterior inference for Kendall’s τ into a parametric data augmentation problem that can be solved with the above MCMC-methods. Thus, Greiner’s relation can be applied to the posterior samples of ρ_{z^x, z^y} to yield posterior samples of τ .

Furthermore, the application of Greiner’s relation in this manner implicitly alters the prior from a uniform distribution on the latent correlation to the following distribution on Kendall’s τ :

$$p(\tau) = \frac{\pi}{4} \cos \left(\frac{\pi \tau}{2} \right), \text{ for } \tau \in (-1, 1). \tag{10}$$




Fig. 2. To illustrate the performance of the three methods, quantile averaged posterior distributions for several values of τ and n are shown. Each column corresponds to a value of n , and each row corresponds to a value of τ . The quantile averaged posterior distributions were obtained with 10,000 synthetic datasets per combination of n and τ . The vertical grey line indicates the population value of τ .

4. Results: Simulation study

The performance of the original asymptotic method, the enhanced asymptotic method, and the latent normal method was assessed with a simulation study. For four values of τ (0, 0.2, 0.4, 0.7) and three values of n (10, 20, 50), 10,000 data sets were generated under four copula models: Clayton, Gumbel, Frank, and Gaussian (Sklar, 1959; Nelsen, 2006; Genest and Favre, 2007; Colonijs, 2016). Using Sklar's theorem, copula models decompose a joint distribution into univariate marginal distributions and a dependence structure (i.e., the copula). The aforementioned copulas are governed by Kendall's τ , so the performance of each method can be assessed through a parameter recovery simulation study. Furthermore, the univariate marginal distributions can be transformed to any other distribution using the cumulative distribution function and its inverse. Because these functions are monotonic, this does not affect the copula or ordinal information in the synthetic data and therefore vastly increases the scope of the simulation study.

For each data set, a posterior distribution was obtained using the three methods and the population value of τ was estimated using the posterior median. Per combination of n and τ , this resulted in 10,000 posterior distributions. For an overall view of each method's performance, Fig. 2 shows the quantile averaged posterior distributions, along with a vertical line indicating the population value of τ . The data in Fig. 2 were generated using the Clayton copula; other copula models yielded highly similar results. The quantile averaged posteriors indicate no difference between the inferential methods under \mathcal{H}_0 , which corroborates the assumption of $\sigma_{\tau^*}^2 = 1$ when $\tau = 0$. However, the difference in methods becomes pronounced in the scenario where $n = 10$ and $\tau = 0.7$. Both asymptotic approaches show a degree of underestimation, and yield

a relatively broad posterior distribution. In the panels where $\tau \neq 0$, the misspecification of the sampling variance also becomes clear, as it is overestimated and results in a wider posterior distribution compared to the latent normal method. Although the assumption of latent normality is the price to pay for the Bayesian latent normal methodology, the simulation results indicate robustness of the method to various violations of this assumption.¹

5. Discussion

This article has outlined two methods of improving the Bayesian inferential framework in cases where n is low and/or τ is high. Although an extension of the asymptotic framework performs somewhat better than the original asymptotic framework in van Doorn et al. (2018), both are outperformed by the latent normal approach. Under \mathcal{H}_0 , the methods do not differ from each other, underscoring the validity of the general framework.

The outlined methods are useful for both estimation and hypothesis testing. In the former case, the posterior distribution enables point estimation through the posterior median, or interval estimation through the credible interval. For hypothesis testing, the Savage–Dickey density ratio (Dickey and Lientz, 1970; Wagenmakers et al., 2010) can be used to obtain Bayes factors (Kass and Raftery, 1995). A concrete example is presented in the online appendix. Because the method uses only the ordinal information in the data, it retains the robust properties of Kendall's τ , such as invariance to monotone transformations, robustness to outliers or violations of normality, and ability to detect nonlinear monotone relations.

Acknowledgments

The project was supported partly by Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO; Grant Number 022.005.022). The authors thank two anonymous reviewers for their helpful comments and suggestions. Centrum Wiskunde & Informatica (CWI) is the national research institute for mathematics and computer science in the Netherlands.

Appendix A. Supplementary data

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.spl.2018.10.004>.

References

- Albert, James H., 1992. Bayesian estimation of the polychoric correlation coefficient. *J. Stat. Comput. Simul.* 44, 47–61.
- Alvo, M., Yu, P.L.H., 2014. Statistical methods for ranking data. In: *Frontiers in Probability and the Statistical Sciences*. Springer New York, New York.
- Berger, J.O., Sun, D., 2008. Objective priors for the bivariate normal model. *Ann. Statist.* 36, 963–982.
- Colonius, Hans, 2016. An invitation to coupling and copulas: With applications to multisensory modeling. *J. Math. Psych.* 74, 2–10.
- Dickey, J.M., Lientz, B.P., 1970. The weighted likelihood ratio, sharp hypotheses about chances, the order of a Markov chain. *Ann. Math. Stat.* 41, 214–226.
- van Doorn, Johnny, Ly, Alexander, Marsman, Maarten, Wagenmakers, Eric-Jan, 2018. Bayesian inference for Kendall's rank correlation coefficient. *Amer. Statist.* 1–6.
- Fisher, Ronald A., 1915. Frequency distribution of the values of the correlation coefficient in samples from an indefinitely large population. *Biometrika* 507–521.
- Geman, S., Geman, D., 1984. Stochastic relaxation, Gibbs distributions and the Bayesian restoration of images. *IEEE Trans. Pattern Anal. Mach. Intell.* 6, 721–741.
- Genest, C., Favre, Anne-Catherine, 2007. Everything you always wanted to know about copula modeling but were afraid to ask. *J. Hydrol. Eng.* 12, 347–368.
- Greiner, R., 1909. Über das fehlersystem der kollektivmasslehre. In: *Zeitschrift für Mathematik und Physik*, vol. 57. pp. 121–158.
- Hotelling, H., Pabst, M.R., 1936. Rank correlation and tests of significance involving no assumption of normality. *Ann. Math. Stat.* 7, 29–43.
- Johnson, Valen E., 2005. Bayes factors based on test statistics. *J. R. Stat. Soc.* 67, 689–701.
- Kass, R.E., Raftery, A.E., 1995. Bayes factors. *J. Amer. Statist. Assoc.* 90, 773–795.
- Kendall, M., Gibbons, J.D., 1990. *Rank Correlation Methods*. Oxford University Press, New York.
- Kruskal, W.H., 1958. Ordinal measures of association. *J. Amer. Statist. Assoc.* 53, 814–861.
- Ly, A., Verhagen, A.J., Wagenmakers, E.J., 2016. Harold Jeffreys's default Bayes factor hypothesis tests: Explanation, extension, and application in psychology. *J. Math. Psych.* 72, 19–32.
- Nelsen, R.B., 2006. *An Introduction to Copulas*, second ed. Springer-Verlag, New York.
- Noether, G.E., 1955. On a theorem of Pitman. *Ann. Math. Stat.* 26, 64–68.
- Olssen, U., 1979. Maximum likelihood estimation of the polychoric correlation coefficient. *Psychometrika* 44, 443–460.
- Pearson, Karl, 1900. Mathematical contributions to the theory of evolution. VII. On the correlation of characters not quantitatively measurable. *Phil. Trans. R. Soc. A* 195, 1–405.
- Pettitt, A.N., 1982. Inference for the linear model using a likelihood based on ranks. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 44, 234–243.
- Sklar, A., 1959. Fonctions de répartition à n dimensions et leurs marges. *Publ. l'Inst. Stat. L'Univ. Paris* 8, 229–231.
- Wagenmakers, Eric-Jan, Lodewyckx, Tom, Kuriyal, Himanshu, Grasman, Raoul, 2010. Bayesian hypothesis testing for psychologists: A tutorial on the savage–dickey method. *Cogn. Psychol.* 60, 158–189.
- Yuan, Ying, Johnson, Valen E., 2008. Bayesian hypothesis tests using nonparametric statistics. *Statist. Sinica* 18, 1185–1200.

¹ R-code, plots, and further details of the simulation study are available at <https://osf.io/u7jj9/>.