



UvA-DARE (Digital Academic Repository)

Drought decreases incorporation of recent plant photosynthate into soil food webs regardless of their trophic complexity

Chomel, M.; Lavallee, J.M.; Alvarez-Segura, N.; de Castro, F.; Rhymes, J.M.; Caruso, T.; de Vries, F.T.; Baggs, E.M.; Emmerson, M.C.; Bardgett, R.D.; Johnson, D.

DOI

[10.1111/gcb.14754](https://doi.org/10.1111/gcb.14754)

Publication date

2019

Document Version

Final published version

Published in

Global Change Biology

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Chomel, M., Lavallee, J. M., Alvarez-Segura, N., de Castro, F., Rhymes, J. M., Caruso, T., de Vries, F. T., Baggs, E. M., Emmerson, M. C., Bardgett, R. D., & Johnson, D. (2019). Drought decreases incorporation of recent plant photosynthate into soil food webs regardless of their trophic complexity. *Global Change Biology*, 25(10), 3549-3561. <https://doi.org/10.1111/gcb.14754>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

PRIMARY RESEARCH ARTICLE

Drought decreases incorporation of recent plant photosynthate into soil food webs regardless of their trophic complexity

Mathilde Chomel¹  | Jocelyn M. Lavelle^{1,2}  | Nil Alvarez-Segura³ | Francisco de Castro⁴ | Jennifer M. Rhymes^{1,5}  | Tancredi Caruso⁶  | Franciska T. de Vries⁷  | Elizabeth M. Baggs⁸  | Mark C. Emmerson⁶ | Richard D. Bardgett¹ | David Johnson¹

¹School of Earth and Environmental Sciences, The University of Manchester, Manchester, UK

²Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA

³Marine and Continental Waters Program, IRTA, Sant Carles de la Ràpita, Catalonia, Spain

⁴AgriFood & Biosciences Institute, Belfast, UK

⁵School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, UK

⁶School of Biological Sciences and Institute for Global Food Security, Queen's University of Belfast, Belfast, UK

⁷Institute of Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, the Netherlands

⁸Global Academy of Agriculture and Food Security, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Midlothian, UK

Correspondence

Mathilde Chomel, School of Earth and Environmental Sciences, The University of Manchester, Manchester, UK.
Email: mathilde.chomel@manchester.ac.uk

Funding information

Natural Environment Research Council, Grant/Award Number: NE/M017028/1

Abstract

Theory suggests that more complex food webs promote stability and can buffer the effects of perturbations, such as drought, on soil organisms and ecosystem functions. Here, we tested experimentally how soil food web trophic complexity modulates the response to drought of soil functions related to carbon cycling and the capture and transfer below-ground of recent photosynthate by plants. We constructed experimental systems comprising soil communities with one, two or three trophic levels (microorganisms, detritivores and predators) and subjected them to drought. We investigated how food web trophic complexity in interaction with drought influenced litter decomposition, soil CO₂ efflux, mycorrhizal colonization, fungal production, microbial communities and soil fauna biomass. Plants were pulse-labelled after the drought with ¹³C-CO₂ to quantify the capture of recent photosynthate and its transfer below-ground. Overall, our results show that drought and soil food web trophic complexity do not interact to affect soil functions and microbial community composition, but act independently, with an overall stronger effect of drought. After drought, the net uptake of ¹³C by plants was reduced and its retention in plant biomass was greater, leading to a strong decrease in carbon transfer below-ground. Although food web trophic complexity influenced the biomass of Collembola and fungal hyphal length, ¹³C enrichment and the net transfer of carbon from plant shoots to microbes and soil CO₂ efflux were not affected significantly by varying the number of trophic groups. Our results indicate that drought has a strong effect on above-ground–below-ground linkages by reducing the flow of recent photosynthate. Our results emphasize the sensitivity of the critical pathway of recent photosynthate transfer from plants to soil organisms to a drought perturbation, and show that these effects may not be mitigated by the trophic complexity of soil communities, at least at the level manipulated in this experiment.

KEYWORDS

Acari, Collembola, microorganisms, plant–soil interaction, pulse-labelling, root-derived C, soil biodiversity, stable isotope

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Global Change Biology* published by John Wiley & Sons Ltd

1 | INTRODUCTION

Drought events are a recurring phenomenon in many ecosystems and are predicted to increase in frequency and intensity in the coming decades (IPCC, 2017; Reichstein et al., 2013). Drought has the potential to severely change ecosystem functioning by impacting plant and soil faunal communities. For example, drought has been shown to alter the composition of grassland plant communities (Cleland et al., 2013; Kardol, Cregger, Company, & Classen, 2010; Liu et al., 2018) and decrease primary productivity (Reichstein et al., 2013; Schwalm et al., 2010). Furthermore, drought can have considerable effects on the composition and activity of soil microbial communities (de Vries et al., 2018; de Vries & Shade, 2013; Hawkes et al., 2011; Meisner, Deyn, Boer, & van der Putten, 2013), and structure, diversity and activity of soil arthropods (Bardgett & Wardle, 2010; Brose et al., 2012; Kardol, Reynolds, Norby, & Classen, 2011; Santonja et al., 2017; Siebert et al., 2019). These effects on soil organisms generally lead to reductions in ecosystem processes, including litter decomposition and nutrient mineralization (de Vries et al., 2013; Fuchslueger, Bahn, Fritz, Hasibeder, & Richter, 2014; Hagedorn et al., 2016; Santonja et al., 2017; Siebert et al., 2019). However, the effect of drought on the entire soil food web is less clear and contrasting responses of different groups of organisms are sometimes reported (Siebert et al., 2019). Furthermore, drought can modify species interactions within the soil food web. For example, warmer and drier conditions can affect decomposition mediated by trophic interactions (Lang, Rall, Scheu, & Brose, 2014; Thakur et al., 2018) or alter predator-induced indirect effects on lower trophic levels (Lensing & Wise, 2006), implying that ecological effects of drought will depend on the trophic structure of the food web.

In most terrestrial ecosystems, 80%–90% of photosynthetically fixed carbon (C) ultimately enters the decomposer food web (Gessner et al., 2010). This happens via two main pathways: the decomposition of dead material (above-ground or below-ground) or via the transfer of recent photosynthates to the rhizosphere and to mycorrhizal fungi (Bais, Weir, Perry, Gilroy, & Vivanco, 2006; Bardgett, Bowman, Kaufmann, & Schmidt, 2005). There is increasing evidence that C from recent photosynthate allocated to roots and associated mycorrhizal mycelium is of major importance for soil food webs (de Vries & Caruso, 2016; Eissfeller et al., 2013; Goncharov, Tsurikov, Potapov, & Tiunov, 2016; Kanters, Anderson, & Johnson, 2015; Pollierer, Langel, Koerner, Maraun, & Scheu, 2007; Ruf, Kuzyakov, & Lopatovskaya, 2006). For example, mycorrhizal fungi act as a rapid conduit for energy and nutrient supply between the plant and fungal feeding Collembola, such as *Protaphorura armata* Tullberg (Endlweber, Ruess, & Scheu, 2009; Johnson et al., 2005). However, perturbations such as drought can severely alter below-ground C allocation, which can in turn affect microbial activity and C turnover (Fuchslueger et al., 2014; Hagedorn et al., 2016; Hasibeder, Fuchslueger, Richter, & Bahn, 2015; Ruehr et al., 2009).

Soil communities form complex networks of direct (e.g. feeding interactions) or indirect (e.g. trophic cascades) interactions, and

these networks are increasingly recognized as driving key processes in soils (Bardgett & Wardle, 2010). For example, the biomass and activity of fungi, which are the primary drivers of C and mineral nutrient cycling, are greatly influenced by grazing activities of Collembola or Acari (Bardgett & Wardle, 2010; Mikola, Bardgett, & Hedlund, 2002; Nielsen, Ayres, Wall, & Bardgett, 2011). As a consequence, these grazing activities can alter rates of decomposition, soil respiration and carbon and nitrogen mineralization (Bardgett & Wardle, 2010; Mikola et al., 2002; Scheu & Setälä, 2002). These interactions can also cascade through food webs, for example, via predation of detritivores, predators can in turn affect microbial activity by controlling the detritivore abundance and grazing activity (Bardgett & Wardle, 2010; Mikola et al., 2002). The diversity of soil organisms plays a key role in simultaneously mediating many ecosystem processes and driving ecosystem multifunctionality (Wagg, Bender, Widmer, & Heijden, 2014), which raises the possibility that different food web components maximize different ecosystem processes in space and time (Gamfeldt, Hillebrand, & Jonsson, 2008; Hector & Bagchi, 2007). For example, the diversity within and across trophic groups influences the mineralization and sequestration of C and energy pathways (Gessner et al., 2010). Soil biodiversity is under growing threat from a range of disturbances associated with global change (Handa et al., 2014; Wagg et al., 2014), which is of concern since biodiversity loss is a major driver of changes in ecosystem function (Bardgett & van der Putten, 2014). Recent empirical and theoretical evidence has shown that soil biodiversity, and the structural asymmetry of C fluxes through food web, can modulate the resistance and resilience of terrestrial ecosystems to perturbations (de Vries et al., 2012, 2018; Rooney & McCann, 2012; Rooney, McCann, Gellner, & Moore, 2006; Schwarz Müller, Eisenhauer, & Brose, 2015; Yang, Wagg, Veresoglou, Hempel, & Rillig, 2018). The difficulty of manipulative reconstructions of soil food webs make experimental studies with soil fauna scarce, and often the effects on soil functions are not consistent (Cole, Dromph, Boaglio, & Bardgett, 2004; Cortet, Joffre, Elmholt, & Krogh, 2003; Lang et al., 2014; Liiri, Setälä, Haimi, Pennanen, & Fritze, 2002). Moreover, the scarcity of manipulative studies means that we have poor understanding of how soil food webs impact critical processes related to C cycling, or whether they can buffer the effects of perturbations on soil functions.

Here, our goal was to test how soil food web trophic complexity impacts microbial community structure, soil functions related to C cycling and the capture and transfer below-ground of recent photosynthate C by plants and whether soil food web trophic complexity modulates the response of these functions to drought. We used microcosms containing soil from semi-natural acid grassland and examined how reconstructed food webs with one, two or three trophic groups, with or without drought, influence microbial communities, mycorrhizal fungal abundance, decomposition and soil CO₂ efflux. In addition, we quantified the flux of C from a common temperate grassland plant species, *Agrostis capillaris*, to soil organisms using ¹³C stable isotope labelling. We hypothesized that: (a) greater trophic complexity of soil food webs will increase litter

decomposition due to the complementarity of organisms, leading to a higher nutrient availability in soil, and hence an increase of plant growth and C allocation below-ground; (b) drought decreases plant photosynthesis and C allocation below-ground, reducing the availability of recently fixed plant C for soil organisms; and (c) soil food webs with fewer trophic groups will be less resistant to drought, leading to a stronger effect of the drought on soil functioning and soil fauna biomass.

2 | MATERIALS AND METHODS

2.1 | Construction of the microcosms

Microcosms were constructed using soil collected from a semi-natural temperate grassland with a history of occasional grazing but no fertilizer application, at the Glensaugh Research Farm, Aberdeenshire, Scotland (56°53'38"N 2°32'29"W). The soil is a humus-iron podsol derived from Old Red Sandstone and acid igneous rocks (pH ~ 5.47; 11.4 C%; 0.76 N%), and the vegetation is classified as *Agrostis-Festuca* grassland, which is widespread in upland regions of the United Kingdom (UK National Vegetation Classification U4a; Rodwell, 1998). Before potting, soil was defaunated by two consecutive freezing events at -80°C for 24 hr (Johnson et al., 2005), sieved to 4 mm, homogenized and air-dried. Microcosms were constructed from PVC pots (9 cm internal diameter × 6 cm depth). The base was filled with clay balls to improve drainage, covered with 100 µm mesh to retain animals and 200 g of dry soil was added. Seeds of *A. capillaris* (Emorsgate Seeds), the dominant grass species of these grasslands, were planted in trays filled with a mixture of defaunated soil and chopped fresh roots of mixed species from the same field site to promote colonization of roots by arbuscular mycorrhizal (AM) fungi. After a 20 day initialization period, nine seedlings of *A. capillaris* were transplanted into each pot. Each pot was inoculated with 10 ml of filtered (10 µm) soil slurry (1:2 soil water mix) to promote establishment of native soil microorganisms (Cole, Dromph, et al., 2004). The microcosms were incubated in a controlled environment cabinet (Weiss Technik UK LiAcarid) at 18°C with a light/dark cycle 18/6 hr, and soil moisture was maintained at 60% (w/w) throughout the experiment. Microcosms were preincubated for 14 days under these conditions to allow microorganisms to colonize the soil, and the plants to extend their root systems. After that time, 1 g of grass litter (with a concentration of C of $47 \pm 0.3\%$ and N of $4.8 \pm 0.2\%$) was placed on top of the soil in each microcosm providing both a habitat and resource to organisms. Senescent grass leaves, composed mainly of *A. capillaris*, were collected from the same site as the soil, homogenized and air-dried for 5 days. For the decomposer and the predator trophic groups, we used one species of Collembola, *Protaphorura armata* (Onychiuridae) and one species of gamasid Acari, *Stratiolaelaps scimitus* (formerly *Hypoaspis miles*) respectively. The euedaphic collembolan *P. armata* is an eyeless and nonpigmented species that reproduces parthenogenetically, and is common in UK grasslands (Cole, Buckland, & Bardgett, 2005;

Hopkin, 2007). The species is frequently used as an ecologically relevant model organism to study invertebrate-fungus grazing interactions (Cole, Staddon, Sleep, & Bardgett, 2004; Johnson et al., 2005; Scheu & Simmerling, 2004). The gamasid *S. scimitus* is a widespread and common soil mite and is known to feed on a range of different soil-dwelling prey species including Collembola. Collembolans were extracted from soil collected at the Glensaugh Research Farm; the soil was spread on a tray and individuals of *P. armata* were gathered with pooters and transferred to polypropylene pots with a mixture of plaster of Paris and charcoal (9:1) in the base. Collembolans were kept at 18°C and fed with yeast prior to the experiment. The predatory Acari, *S. scimitus*, was supplied by Koppert UK Ltd. We constructed three food web treatments: only microorganisms; microorganisms + Collembola; microorganisms + Collembola + predatory Acari. We placed 100 individuals of *P. armata* (equivalent to $15,625 \text{ ind/m}^2$) and 20 individuals of *S. scimitus* (equivalent to $3,125 \text{ ind/m}^2$) in the corresponding treatment pots, which is consistent with natural densities and ratios of predator:prey at Glensaugh ($16,092 \text{ Collembolans/m}^2$ and $7,985 \text{ gamasids/m}^2$). The organisms were added to the microcosms and a transparent plastic sheet (20 cm high) was wrapped around the rim of each pot to prevent animals escaping.

The experiment comprised a 3 (trophic groups) × 2 (drought vs. control) factorial design with six replicates per treatment. Six supplementary microcosms with the full food web treatment were constructed and were used to determine ^{13}C natural abundance in major pools and fluxes. All microcosms (36 microcosms + 6 for ^{13}C natural abundance) were randomly interspersed in a controlled environment growth chamber, and locations were swapped randomly and regularly. Each microcosm was wetted with distilled water every 2 days to maintain their original weight. After 68 days of incubation, 7 days of drought was applied to half of the replicates to reach a $25 \pm 2.04\%$ moisture reduction based on average moisture of control microcosms, which reflects the effects of chronic summer drought on UK upland grasslands (e.g. Grime et al., 2008). *A. capillaris* was clipped to a height of 10 cm at 29 and 49 days after their transplantation; the biomass clipped was dried, weighed and summed to obtain the final plant biomass.

2.2 | Isotopic labelling

At the end of the drought, all droughted pots were watered lightly with 20 ml to release the drought and encourage photosynthetic activity; control pots were also watered with 20 ml for consistency. Immediately following, microcosms were randomly assorted into two plastic wooden-framed chambers (60 × 60 × 50 cm) which received a continuous flow of air containing 99 atom% $^{13}\text{C}\text{-CO}_2$ at $400 \text{ cm}^3/\text{m}^3$ for 7 hr into the chambers. Immediately after labelling, and each day for three consecutive days, approximately 20 mg of plant shoots (<2% of total shoot biomass) was harvested from each microcosm. To capture the release of $^{13}\text{C}\text{-CO}_2$ from soil CO_2 efflux, an Eppendorf tube containing 0.5 ml of 2 M NaOH was placed on the soil surface inside a small tube (2 cm diameter × 4 cm height)

inserted 1 cm into the soil, and the tube was closed for 24 hr and replaced periodically for 3 days following the ^{13}C labelling. A 0.1 ml aliquot of NaOH was transferred to hydrogen-flushed exetainers (Labco, UK) and 0.5 ml of 1.3 M H_3PO_4 was used to acidify the solution 1 day before the analysis of the $^{13}\text{C}/^{12}\text{C}$ ratio of the accumulated CO_2 on a 20/20 isotope ratio mass spectrometer (Sercon Ltd coupled to an ANCA TGII gas preparation module). After 3 days, microcosms were harvested destructively. Plant shoots were collected and the experimental litter that was applied to the surface was picked with tweezers and freeze-dried for analysis of nutrient concentrations and mass loss. Litter mass loss was calculated as the difference of the initial and final litter dry weight. A subsample of soil was taken for the analysis of hyphal length, mycorrhizal colonization of roots and phospholipid fatty acid (PLFA) concentrations. From the remaining soil, animals were extracted by Berlese funnel over 5 days. They were counted under a dissecting microscope, dried and transferred into tin cups for weighing and ^{13}C analysis. The six supplementary microcosms that were not ^{13}C -labelled were used to determine the ^{13}C natural abundance signature of each C pool. Shoot material and organisms were dried at 60°C for 48 hr and weighed. All samples for C analysis were ground and analysed for total C content and $\delta^{13}\text{C}$ using an elemental analyser (PDZ Europa ANCA-GSL, Sercon Ltd) coupled to a 20–20 isotope ratio mass spectrometer (Sercon Ltd).

2.3 | Mycorrhizal colonization and hyphal length

Analysis of *A. capillaris* root colonization by AM fungi was determined by calculating the percentage root length colonization (%RLC). After washing, roots were cleared by boiling for 5 min in a 10% KOH solution, rinsed and stained for 3 min in a boiling ink–vinegar solution (5% ink) (Vierheilig, Coughlan, Wyss, & Piche, 1998). Stained roots were then mounted on slides and fixed with 50% glycerol. Observations were done under a microscope at $\times 200$ magnification and per cent root colonization by arbuscules, vesicles and total AM fungal colonization (including hyphae, arbuscules and vesicles) were quantified using the magnified intersect method (McGonigle, Miller, Evans, Fairchild, & Swan, 1990).

Total length of fungal hyphae was measured in soil extracts using the membrane filter technique modified after Hanssen, Thingstad, Goksøyr, and Goksoyr (1974), followed by staining the hyphae with calcofluor white M2R fluorescent brightener (Sigma-Aldrich) (Bloem, Bolhuis, Veninga, & Wieringa, 1995) and quantification with the grid-line intersect method (Brundett, Bougher, Dell, Grove, & Malajczuk, 1996). Briefly, 10 g (fresh weight) of soil in 95 ml of dH_2O was blended, 5 ml of the soil slurry was diluted with 5 ml of dH_2O and then 1 ml of the diluted soil slurry was added to 1 ml of formalin, 1 ml of Calcofluor and 7 ml of dH_2O . Samples were incubated in the dark at room temperature for 2 hr and stored at 4°C until further processed. Two millilitres of these samples was filtered through a 25 mm diameter black 1 μm pore size polycarbonate filter (Osmonics Inc.) and rinsed three times with 3 ml of dH_2O . The filters were placed on slides with

immersion oil and observed with a gridded ocular lens for a total of 50 fields at $\times 100$ magnification under a UV illuminated microscope (Olympus BX61). Hyphal length (H) was calculated using the equation $H = (l\pi A)/(2L)$, where l is the average number of intersections per grid, A is the grid area and L is the total length of the grid lines. Then, the total length of fungal hyphae (F) (m/g soil) was calculated using the equation $F = H10^{-6} (A/B) (1/S)$ where A is the area of the filter, B is the grid area and S is the amount of soil filtered (Bloem et al., 1995).

2.4 | Soil microbial community analysis

The characterization of soil microbial communities was estimated using PLFA analysis, which were extracted using 19:0 phosphatidylcholine (Avanti Polar Lipids) as an internal standard added at the beginning of the extraction procedure for quantitative analysis (Buyer & Sasser, 2012). Fatty acids were extracted from 0.5 g of dry soil in Bligh-Dyer extractant, containing the internal 19:0 standard, for 2 hr rotating end-over-end. The liquid phase was then collected after centrifugation, 1.0 ml chloroform:water (1:1) was added and the lower phase siphoned off and used for lipid separation. Lipids were separated by solid-phase extraction (SPE) using a 96-well SPE plate (100 mg silica, Phenomenex) with chloroform, acetone and 5:5:1 methanol:chloroform:water. Fatty acids were then transesterified and extracted. The $\delta^{13}\text{C}$ values of individual PLFAs and their quantification (Thornton, Zhang, Mayes, Högberg, & Midwood, 2011) were analysed by GC–C–IRMS using a Trace GC Ultra gas chromatograph with combustion column attached via a GC Combustion III to a Delta V Advantage IRMS (Thermo Finnigan). In summary, 36 PLFAs were identified in these samples, of which 20 microbial-specific PLFAs comprising approximately 80% of the total concentration were used in subsequent data analysis. The fatty acids i15:0, a15:0, i16:0 and i17:0 were used as biomarkers for Gram-positive bacteria; 16:1 ω 7, 18:1 ω 7, cy17:0 and cy19:0 were used as biomarkers of Gram-negative bacteria; and 15:0 and 17:0 were used as general bacterial markers (Frostegård, Bååth, & Tunlio, 1993). The fatty acids 10Me17:0 and 10Me18:0 were used as specific biomarkers of actinomycetes, and 17:1 ω 8c and 19:1 ω 8 were used as biomarkers of methane-oxidizing bacteria (Frostegård et al., 1993). Gram-positive, Gram-negative and general bacterial markers were summed to give total bacterial PLFA, and 18:2 ω 6,9 was used as a marker of fungi (Bååth, 2003; Bååth & Anderson, 2003). The $\delta^{13}\text{C}$ value of each PLFA molecule was corrected for the C added during derivatization using the formula:

$$\delta^{13}\text{C}_{\text{PLFA}} = \frac{C_{\text{FAME}} \times \delta^{13}\text{C}_{\text{FAME}} - C_{\text{MeOH}} \times \delta^{13}\text{C}_{\text{MeOH}}}{C_{\text{PLFA}}}$$

where C_{FAME} , C_{MeOH} and C_{PLFA} denote the number of carbon atoms in FAME, methanol and PLFA, respectively, and $\delta^{13}\text{C}_{\text{FAME}}$ and $\delta^{13}\text{C}_{\text{MeOH}}$ are the measured ratio of stable isotopes $^{13}\text{C}/^{12}\text{C}$ (in part per thousand, ‰) of FAME and methanol respectively (methanol $\delta^{13}\text{C} = -29.3\text{‰}$). While the fungi are represented by only one PLFA, the bacterial community is represented by several PLFAs. To calculate an overall ^{13}C enrichment

(atom% excess) of bacterial PLFA, the net ^{13}C of all individual (a) bacterial PLFA were summed and divided by the sum of the net C of all individual bacterial PLFA using the following equation:

$$\%^{13}\text{C}_{\text{BacterialPLFA}} = \frac{\sum (C_{\text{PLFA}_i} \times \text{Atom}\%^{13}\text{C}_{\text{PLFA}_i})}{\sum C_{\text{PLFA}}}$$

2.5 | Calculations

PLFAs were converted to biomass C using the following factors: 363.6 nmol of total bacterial PLFA = 1 mg bacterial C (Frostegård & Bååth, 1996), 11.8 nmol of the PLFA 18:2 ω 6,9 = 1 mg of fungal C (Klamer & Bååth, 2004). All ^{13}C data were converted from $\delta^{13}\text{C}$ values (‰) to ^{13}C values (atom%) and then in ^{13}C atom% excess subtracting the ^{13}C of unlabelled controls from each enriched sample (Johnson, Vachon, Britton, & Helliwell, 2011). The net incorporation of the ^{13}C tracer into the different compartment for each pot was calculated as:

$$\text{Net } ^{13}\text{C}_{\text{pool}} = C_{\text{pool}} \times \text{Atom}\%^{13}\text{C}_{\text{pool}}$$

where C_{pool} is the stock of C in each pool (mg per pot) and $\text{Atom}\%^{13}\text{C}_{\text{pool}}$ is the atom% excess of ^{13}C in each pool.

The C budget was calculated using the net ^{13}C excess, and allowed investigation of the dynamics of ^{13}C allocation within individual C pools. This C budget was expressed as a percentage of the net ^{13}C exported from the plant shoots (via either shoot respiration or allocation to other organs) to the different C pools during the 3 days after labelling calculated using the equation:

$$\%^{13}\text{C}_{\text{transferred}} = \left(\text{Net } ^{13}\text{C}_{\text{pool}} \div \text{Plant Net } ^{13}\text{C}_{\text{exported}} \right) \times 100,$$

where C_{pool} is the stock of C in each pool (mg per pot) and $\text{Plant net } ^{13}\text{C}_{\text{exported}}$ is the stock of ^{13}C exported by the plant.

2.6 | Statistical analyses

The experimental design included two fixed factors: treatment (drought or control) and food web trophic complexity (three levels). For all measured variables, two-way analysis of variance (ANOVA) was used to test the effects of the drought treatment and food web trophic complexity, followed by Tukey post hoc pairwise comparisons. Normality and homoscedasticity of the data were first checked using Anderson Darling and Levene tests respectively. If the assumptions were not met, data were log-transformed to meet the test assumptions before performing statistical tests. Data from the six additional pots with the full food web that were established and used primarily for natural abundance determination were combined with data from the main experiment for analysis of responses not involving analysis of ^{13}C . All statistical analyses were performed using the R software (version 3.4.3, R Core Team, 2017). Statistical tests used $\alpha = 0.05$ to determine statistical significance.

3 | RESULTS

Both Collembola and Acari were present at the final harvest, showing that consumers and predators established populations. The number of individual Collembola ranged from 8 to 234 (median = 66) and Acari ranged from 3 to 27 (median = 5) per pot. Generally, we found significant main effects of the drought and the number of trophic groups rather than their interaction on the different response variables (Table 1).

3.1 | Effects of food web trophic complexity and drought on plant productivity, soil communities and soil functions

Total hyphal length was reduced by 36% in the presence of Collembola only, and 41% in the presence of Collembola and Acari compared to the microorganisms only ($p < .001$, $F = 15.09$, $df = 2$, $n = 42$, Figure 1b, Table 1). The biomass of Collembola was reduced by 41% in the presence of predators ($p = .008$, $F = 8.20$, $df = 1$, $n = 30$, Figure 1c, Table 1). Above-ground plant biomass was not influenced by food web trophic complexity (Figure 1a, Table 1), whereas litter mass loss was slower in mesocosms with three trophic groups compared with one or two trophic groups ($p = .005$, $F = 6.07$, $df = 2$, $n = 42$, Figure 1e, Table 1), and soil CO_2 efflux was less in mesocosms with two trophic groups than with one or three trophic groups ($p = .048$, $F = 3.31$, $df = 2$, $n = 42$, Figure 1c, Table 1). Trophic complexity had no significant impact on fungal or bacterial PLFAs, but there was a trend of decreasing fungal PLFA with two or three trophic groups (Tables 1 and 2). Likewise, the colonization of roots by AM fungi was not influenced by food web trophic complexity (Tables 1 and 2).

Overall, drought had a consistent effect on the response variables regardless of soil food web trophic complexity, as reflected by the absence of any significant interaction between the two factors. Drought treatment reduced plant above-ground biomass by an average of $27 \pm 2\%$, and litter mass loss by $16 \pm 5\%$ compared to the controls (Figure 1a,e, Table 1). Soil CO_2 efflux increased by $63 \pm 8\%$ after the release of the drought compared to controls (Figure 1f, Table 1). Drought had no significant impact on the fungal PLFA 18:2 ω 6,9, but there was a marginal, but not significant increase in bacterial PLFA under drought relative to controls ($F = 3.65$, $df = 1$, $n = 42$, $p = .064$) (Tables 1 and 2); as such, the fungal/bacterial ratio was reduced ($F = 9.22$, $df = 1$, $n = 42$, $p = .005$) by drought (Tables 1 and 2). While the colonization of roots by AM fungi was not impacted by drought, the production of arbuscules was significantly reduced by drought by $21 \pm 8\%$ relative to the control (Tables 1 and 2).

3.2 | Effects of food web trophic complexity and drought on pools and fluxes of ^{13}C

The number of trophic groups had no effect on ^{13}C enrichment of the majority of C pools, except the Collembola C pool under control conditions, which had greater ^{13}C enrichment in the presence of predators compared to without predators (treatment \times trophic groups interaction, $F = 4.698$, $df = 1$, $n = 36$, $p = .0425$, Figure 2e).

Variables	Drought		Trophic groups (TG)		Drought × TG	
	F	p value	F	p value	F	p value
Process						
Litter mass loss	6.74	.014	6.07	.005	1.55	.226
Soil CO ₂ efflux	9.27	.004	3.31	.048	0.03	.976
Soil biota						
Bacterial PLFA	3.65	.060	1.84	.180	0.60	.556
Fungal PLFA	0.63	.433	1.66	.203	0.06	.940
f/b ratio	9.22	.005	2.05	.144	0.38	.687
% root colonized by AM fungi	1.25	.270	0.23	.793	1.12	.339
% arbuscules	4.91	.033	1.21	.311	1.32	.281
Collembola biomass	38.95	.000	8.20	.008	0.48	.496
Acari biomass	1.72	.210	–	–	–	–
Carbon budget						
Plant shoot ¹³ C net (mg)	5.83	.022	0.38	.687	0.10	.903
Plant shoot ¹³ C exported after 3 days (%)	19.235	.000	0.454	.640	0.307	.738
Net ¹³ C transfer from plant export to respiration	0.11	.747	0.294	.748	2.177	.132
Net ¹³ C transfer from plant export to fungal biomass	0.18	.677	0.745	.484	0.147	.864
Net ¹³ C transfer from plant export to bacterial biomass	0.25	.622	0.31	.736	0.241	.787
Net ¹³ C transfer from plant export to Collembola	8.12	.010	1.93	.180	0.01	.934
Net ¹³ C transfer from plant export to Acari	0.19	.674	–	–	–	–

Note: Significant treatment effects ($p < .05$) are in bold.

Abbreviations: AM, arbuscular mycorrhizal; PLFA, phospholipid fatty acid.

At the end of the pulse-labelling period, ¹³C enrichment of droughted plant shoots was on average $27 \pm 4\%$ greater compared to control plants ($F = 35.2$, $df = 1$, $n = 36$, $p < .001$, Figure 2a). During the first 24h after pulse-labelling, ¹³C enrichment of the CO₂ released from droughted soil was on average $49 \pm 4\%$ lower relative to the control, regardless of the food web trophic complexity ($F = 6.9$, $df = 1$, $n = 36$, $p < .001$, Figure 2b). PLFAs were significantly enriched in ¹³C 3 days after the pulse-labelling. The fungal PLFA 18:2 ω 6,9 had the greatest ¹³C enrichment, reaching a maximum of 0.71 atom% ¹³C excess, while maximal enrichment of bacterial PLFA was 0.14 atom% ¹³C excess for the Gram-negative PLFA 16:1 ω 7c. While ¹³C enrichment of fungal PLFA was similar between the droughted and control mesocosms ($F = 0.04$, $df = 1$, $n = 36$, $p = .84$, Figure 2c), ¹³C enrichment of bacterial PLFA was reduced in droughted mesocosms, regardless of food web

TABLE 1 Results of two-way ANOVAs for the effects of the drought (drought or control) and food web trophic complexity (one, two or three trophic groups) and their interaction on the processes, soil biota and ¹³C carbon budget

trophic complexity ($F = 38.7$, $df = 1$, $n = 36$, $p < .001$; treatment × trophic groups interaction, $p = .09$; Figure 2d). The incorporation of ¹³C into Collembola was greater in droughted mesocosms relative to controls, but only in the absence of predators (interaction between treatment × trophic groups, $F = 4.698$, $df = 1$, $n = 36$, $p = .0425$, Figure 2e). There was no difference in ¹³C enrichment of Acari in response to the drought treatment (Figure 2f).

3.3 | Carbon budget and net C incorporation into the food web

Plants that had been subjected to drought had shoots more enriched in ¹³C immediately after labelling than the controls, but they had overall smaller biomass. Consequently, the net ¹³C in the plant

FIGURE 1 Above-ground plant biomass (A), total fungal hyphal length (B), Collembola biomass (C), Acari biomass (D), litter mass loss (E) and soil CO₂ efflux (F) in response to food web trophic complexity (1: microorganisms, 2: microorganisms + Collembola or 3: microorganisms + Collembola + Acari) and drought (control in white and drought in grey). Lines in boxes represent median, top and bottom of boxes represent first and third quartiles, and whiskers represent the largest value no further than 1.5 interquartile range; dots represent single observations. Only the significant terms of the ANOVA are presented (**p* < .05; ***p* < .01; ****p* < .001). Tukey comparison between trophic groups are indicated by different letters, a < b




TABLE 2 Bacterial and fungal PLFA, fungal/bacterial PLFA ratio and AM fungal root colonization

Variables	Drought	ANOVA	Trophic groups		
			1	2	3
Bacterial PLFA (nmol/g of soil)	Control		1,417 ± 260	1,043 ± 241	1,451 ± 150
	Drought		2,222 ± 499	1,461 ± 277	1,647 ± 300
Fungal PLFA (nmol/g of soil)	Control		34.17 ± 10.45	21.55 ± 8	19.37 ± 4.52
	Drought		24.32 ± 6.91	15.13 ± 3.36	15.16 ± 3.04
Fungal/bacterial PLFA ratio	Control (b)	**	0.015 ± 0.002	0.018 ± 0.004	0.013 ± 0.002
	Drought (a)		0.01 ± 0.001	0.01 ± 0.001	0.009 ± 0
% root colonized by AM fungi	Control		59.8 ± 2	65.8 ± 4.88	60.23 ± 2.27
	Drought		57.03 ± 4.74	56.22 ± 3.66	61.39 ± 4.56
% arbuscules	Control (b)	*	28.55 ± 2.88	30.07 ± 3.76	23.2 ± 1.55
	Drought (a)		20.36 ± 3.01	21.58 ± 2.95	22.13 ± 2.76

Note: Values are presented as mean ± SE. Significant differences among drought treatment are indicated by asterisks (**p* < .05; ***p* < .01) and the Tukey comparison with different letter with a < b. Results of the ANOVA are in Table 1.

Abbreviations: AM, arbuscular mycorrhizal; PLFA, phospholipid fatty acid.

biomass C pool was less in droughted mesocosms than in controls, and was on average 10.8 ± 0.29 mg in the controls and 9.7 ± 0.32 mg in the droughted mesocosms at the end of the labelling period

(*F* = 5.829, *df* = 1, *n* = 36, *p* = .02, Figure 3a, Table 1). Three days later, only 1.9 ± 0.32 mg of ¹³C was left in control plant shoots, but 2.5 ± 0.16 mg ¹³C was left in the droughted plant shoots. Therefore,




FIGURE 2 ^{13}C enrichment of the different carbon pools as a function of the food web trophic complexity (one, two or three trophic groups) and drought (control in white and drought in grey): Above-ground plant (A), soil CO_2 efflux (B), bacterial PLFA (C), fungal PLFA (D), Collembola (E) and Acari (F). Lines in boxes represent median, top and bottom of boxes represent first and third quartiles, whiskers represent the largest value no further than 1.5 interquartile range; dots represent single observations. Only the significant terms of the ANOVA are presented ($*p < .05$; $**p < .01$; $***p < .001$). Tukey comparison between trophic groups are indicated by different letters, $a < b$

the relative amount of ^{13}C exported (through reallocation to the roots or from respiration) from the plant shoots during these 3 days was greater under control compared to drought, with losses of $83 \pm 1.19\%$ and $75 \pm 1.28\%$ from the initial ^{13}C amount respectively ($F = 19.2$, $df = 1$, $n = 36$, $p < .001$, Figure 3b, Table 1). This export of ^{13}C was traced into bacteria, fungi, Collembola, Acari and soil CO_2 efflux. On average across all the treatments, the percentage of the plant ^{13}C exported recovered after 3 days in the below-ground pools was $1.5 \pm 0.2\%$ in bacteria, $8.9 \pm 1.5\%$ in fungi, $0.02 \pm 0.004\%$ in Collembola and $0.0006 \pm 0.0002\%$ in Acari (Figure 3c-f). There were no significant differences in the proportion of recent photosynthate ^{13}C transferred to fungi or bacteria in soils with different food web trophic complexities or under drought (Figure 3c,d, Table 1). There was also no significant effect of either drought or food web treatments on the proportion of net ^{13}C recovered in Acari or in soil CO_2 efflux (Figure 3f,g). However, the proportion of recent photosynthate ^{13}C exported from plant shoots and recovered in Collembola was consistently lower after drought, decreasing from $0.025 \pm 0.006\%$ to $0.008 \pm 0.002\%$ respectively ($F = 8.1$, $df = 1$, $n = 36$, $p = .01$, Figure 3e, Table 1). In addition, transfer of ^{13}C to Collembola in communities with three trophic levels was almost half of that measured

in communities with only two trophic levels ($F = 1.9$, $df = 1$, $n = 36$, $p = .18$, Figure 3e, Table 1). Overall, there was no significant interactive effect of food web trophic complexity on the quantities of ^{13}C transferred to the different C pools.

4 | DISCUSSION

The aim of this study was to test how trophic complexity, manipulated here by increasing the number of trophic levels, moderates the response to drought of soil functions related to C cycling and the capture and transfer below-ground of recent plant photosynthate C. We found limited evidence for interactive effects between drought and food web trophic complexity; only the transfer of recent photosynthate to Collembola was affected by drought \times food web interactions. Therefore, our findings suggest that soil faunal trophic complexity, at least at the level manipulated in this study, does not substantially enhance the resistance of soil functions to drought. Our results show that there was a consistently strong effect of drought on most variables regardless of food web trophic complexity.




FIGURE 3 Quantification of the major pools and fluxes of ¹³C labelled recent plant assimilate in response to food web trophic complexity (1, 2 or 3 trophic groups) and drought (control in white and drought in grey). Net ¹³C (mg) in plant shoots immediately after the pulse labelling (a) and ¹³C exported from plant shoots (comprising either allocation to roots or lost as plant respiration) after 3 days expressed as a percentage of the net ¹³C fixed (b). From the amount of ¹³C (mg) exported from plant shoots, we calculate the percentage transfer to different carbon pools comprising bacteria (c), fungi (d), Collembola (e), predatory Acari (f) and soil CO₂ efflux (g). The soil CO₂ efflux includes root respiration and was measured during the first 24 hr after the labelling. Mean ± SE, significant differences among drought are indicated by asterisks (**p* < .05; ***p* < .01; ****p* < .001)

We found that drought increased the residence time of recently assimilated C in the leaves, leading to a smaller flux of C from the plant shoots to roots, soil and the atmosphere. Plant-derived ¹³C was rapidly incorporated into soil microbial communities and mesofauna (Collembolans and predatory Acari), and partially released from the system as soil ¹³C-CO₂ efflux. This finding supports the view that numerically abundant and widespread species, such as *P. armata*, are fuelled with recent plant photosynthate via root exudates or mycorrhizal mycelium (Johnson et al., 2005; Pausch et al., 2016; Pollierer, Dyckmans, Scheu, & Haubert, 2012). Incorporation of ¹³C was greater in fungal PLFA than in bacterial PLFAs, with the former representing a larger total C pool when converted into biomass (Frostegård & Bååth, 1996). In our system, this supports the idea that fungi could be key conduits of recent assimilate flow and are critical for transferring energy through higher trophic groups, as suggested by other studies (e.g. Johnson et al., 2005; Pausch et al., 2016).

The food web treatments used in this experiment were effective because Collembola density decreased in the presence of

the predatory Acari, and fungal hyphal length, a measure of the production of extramatrical hyphae, decreased in the presence of Collembola, showing a strong cascading top-down effect of higher trophic levels. In accordance with our first hypothesis, food web trophic complexity regulated litter decomposition and the soil CO₂ efflux. However, while we expected litter to decompose faster in the presence of more trophic groups, due to complementarity of soil organisms, we observed slower litter decomposition with the three trophic groups compared to only one or two trophic groups. It has been suggested that predation pressure may prevent Collembola from over-exploiting fungal populations (Hasegawa & Takeda, 1995), which could enhance the rate of litter breakdown. Conversely, predation on Collembola could retard decomposition by preventing the positive effects of Collembola on litter breakdown (Berg & Laskowski, 2005; Cortet et al., 2003). Food web trophic complexity did not influence plant performance, measured in terms of biomass and ¹³C uptake by photosynthesis.

This was despite there being significant effects of food web trophic complexity on litter decomposition. There is currently no clear consensus in the literature on plant responses to food web composition. While several correlative studies have shown that the diversity of soil organisms is important to maintain ecosystem functioning, including plant productivity (de Vries et al., 2013; Wagg et al., 2014), other experimental studies have shown little effect of food web composition on plant growth (Bradford et al., 2002; Cole, Dromph, et al., 2004; Liiri et al., 2002). For example, Cole, Dromph, et al. (2004) detected only a small effect of 18 different food web compositions on the growth of *A. capillaris*. The lack of effect of the soil food web trophic complexity on plant growth in our study might be for a number of reasons. For example, it is possible that variation in food web trophic complexity has multiple, but opposing effects on plant growth (e.g. Bradford et al., 2002), and changes in food web trophic complexity might be realized only in the long term.

In contrast to our first hypothesis, ^{13}C enrichment and the total amount of C transferred from plant shoots to the other C pools varied little with food web trophic complexity. This finding suggests that variations in the trophic structure at the scale of our study had no influence on the plant C uptake and transfer to microbial communities, despite having some effect on their biomass (as also seen by Cole, Dromph, et al., 2004). However, in the absence of drought, ^{13}C - CO_2 efflux tended to decrease with the number of trophic groups, suggesting an efficient C transfer through the trophic levels. This transfer of plant-derived C to higher trophic groups leads to greater retention of C below-ground as a result of reduced loss via soil respiration.

According to our second hypothesis, our study revealed large effects of drought on several response variables, and most of these responses were likely mediated indirectly through plants and Collembola given the magnitude by which those responded to drought. The physiological and growth responses of plants to drought were complex. For example, despite observing an overall decrease of plant biomass in response to drought, more ^{13}C was fixed by plant shoots during the pulse-labelling immediately after the cessation of the drought. This observation might be explained by either greater ^{13}C uptake due to higher photosynthetic activity stimulated by the cessation of drought or by reduced loss of ^{13}C from leaf respiration (Atkin & Macherel, 2009) during the pulse-labelling period. Once fixed by shoots, the ^{13}C may be respired, used to produce new shoot material, temporarily stored or allocated below-ground into roots and to soil organisms (Leake, Ostle, Rangel-Castro, & Johnson, 2006). Three days after labelling, we observed greater retention of recent photosynthate in plant shoots subjected to drought. These findings are in line with results from beech saplings where drought stress doubled the residence time of fresh assimilates in foliar biomass due to a decrease in phloem transport velocity (Ruehr et al., 2009), and reduced below-ground C allocation (Fuchslueger et al., 2014; Ruehr et al., 2009).

Under control conditions, soil CO_2 efflux mainly comprised recent photosynthate, while after drought, a greater proportion of CO_2 originated from other sources. This result confirms that

proportionally less recent photosynthate was invested below-ground under drought, which might promote a shift of soil microbial community towards increased decomposition of soil organic C (Bradford, Fierer, & Reynolds, 2008), explaining the reduction in ^{13}C enrichment of soil CO_2 efflux. Collectively, our results indicate that the rate of ^{13}C transport from leaves below-ground is slower after a drought perturbation, and thus, the connection between above- and below-ground processes is reduced. Our results confirm that drought increases the time-lag between photosynthesis and soil CO_2 efflux (Kuzyakov & Gavrichkova, 2010; Ruehr et al., 2009).

In contrast to theory (Rooney & McCann, 2012; Rooney et al., 2006) and our third hypothesis, manipulation of food web trophic complexity had limited impact on C flow to microorganisms and on buffering the effects of drought on ecosystem processes. However, these theoretical analyses also suggest that asymmetry of pathways of C flow below-ground, reflected here by recent photosynthate and litter decomposition, is expected to confer stability on ecosystem function. Our findings that drought may weaken the recent photosynthate pathway of C flow below-ground (i.e. impacting the flux of ^{13}C from plant shoots to below-ground pools and fluxes) suggest it also impacts the asymmetry of C flow through soil food webs. In addition, inclusion of predators reduced the effect of drought on ^{13}C transfer to Collembola, indicating the importance of higher trophic groups on C dynamics through food webs. These results provide the basis for further experimental work to disentangle the mechanisms by which drought affects energy flow through, and asymmetry of, different channels.

We observed less ^{13}C enrichment of bacterial PLFA after drought, suggesting either that drought leads to bacteria using alternative sources of C to recent assimilate, or that they were less efficient in using this pathway of C. However, this difference was not observed for fungi. Fungi have stronger cell walls preventing water loss, and they can redistribute water by mycelial networks, and so are considered to be more resistant to drought than bacteria (Gühr, Borken, Spohn, & Matzner, 2015). Furthermore, bacteria and fungi often occupy different habitats within soil: bacteria often live within water-filled pore spaces while most fungi live in air-filled pore spaces (Ritz 2011). These habitat preferences may help explain why bacteria and fungi were affected by drought differently. In the absence of predators, the ^{13}C enrichment of Collembola was less in control conditions compared to drought. This finding could be related to the reduced biomass of Collembola after the drought that likely decreased the competition for ^{13}C -enriched substrates (such as fungal hyphae). By contrast, in the presence of predators, ^{13}C enrichment of Collembola did not differ between control and drought treatments. Here, there was overall smaller biomass of Collembola and the difference in biomass between control and drought treatments was less pronounced than in the absence of predators, which could have maintained a lower ratio of Collembola to ^{13}C -labelled substrates thus leading to greater ^{13}C enrichment of Collembola. Our results showed no evidence of an impact of drought on the net transfer of C from the plants to microbial communities; however, we observed less net C transfer from plants to Collembola, mainly

due to the decrease in biomass reflecting their sensitivity to drought (Hopkin, 1997; Makkonen et al., 2011).

Overall, our study identifies an important effect of drought on plant–soil C fluxes mainly by acting on plant physiology (respiration and C allocation), that regulates the availability of resources to the below-ground food web, and which may affect food web stability (Rooney et al., 2006). We also showed that an abundant and widespread species of Collembola is highly sensitive to drought, and its feeding activity is moderated by the presence of predators, suggesting the importance of higher trophic groups on C dynamics through below-ground food webs. The recent photosynthate C pathway is critical to ecosystem functioning and the soil food web, and our work demonstrates its sensitivity to perturbations resulting from global change.

ACKNOWLEDGEMENTS

The project was initiated by RB, DJ, ME, TC, FdV and EB, who also secured the funding from the NERC Soil Security Programme (ref. NE/M017028/1).

AUTHOR CONTRIBUTION

MC, DJ and RDB designed the study, with contributions from all the authors and MC and NAS carried out labwork and analysed the data. MC carried out statistical analysis and wrote the manuscript with DJ, and with contributions from all authors. DJ receives partial support from the N8 AgriFood programme. We thank the University of Aberdeen, L. Harrold for isotopic analysis, and the James Hutton Institute, B. Thornton and G. Martin for the PLFA isotopic analysis and access to Glenshagh Farm. We also thank two anonymous reviewers for their helpful suggestions on a previous version of the manuscript.

ORCID

Mathilde Chomel  <https://orcid.org/0000-0001-5110-2355>

Jocelyn M. Lavallee  <https://orcid.org/0000-0002-3028-7087>

Jennifer M. Rhymes  <https://orcid.org/0000-0001-9347-9863>

Tancredi Caruso  <https://orcid.org/0000-0002-3607-9609>

Franciska T. Vries  <https://orcid.org/0000-0002-6822-8883>

Elizabeth M. Baggs  <https://orcid.org/0000-0003-2014-2148>

REFERENCES

- Atkin, O. K., & Macherel, D. (2009). The crucial role of plant mitochondria in orchestrating drought tolerance. *Annals of Botany*, 103(4), 581–597. <https://doi.org/10.1093/aob/mcn094>
- Bååth, E. (2003). The use of neutral lipid fatty acids to indicate the physiological conditions of soil fungi. *Microbial Ecology*, 45(4), 373–383. <https://doi.org/10.1007/s00248-003-2002-y>
- Bååth, E., & Anderson, T.-H. (2003). Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and PLFA-based techniques. *Soil Biology and Biochemistry*, 35(7), 955–963. [https://doi.org/10.1016/S0038-0717\(03\)00154-8](https://doi.org/10.1016/S0038-0717(03)00154-8)
- Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S., & Vivanco, J. M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, 57, 233–266. <https://doi.org/10.1146/annurev.arplant.57.032905.105159>
- Bardgett, R. D., Bowman, W. D., Kaufmann, R., & Schmidt, S. K. (2005). A temporal approach to linking aboveground and belowground ecology. *Trends in Ecology & Evolution*, 20(11), 634–641. <https://doi.org/10.1016/j.tree.2005.08.005>
- Bardgett, R. D., & van der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature*, 515(7528), 505–511. <https://doi.org/10.1038/nature13855>
- Bardgett, R. D., & Wardle, D. A. (2010). *Aboveground-belowground linkages: Biotic interactions, ecosystem processes, and global change*. Oxford, New York: Oxford University Press.
- Berg, B., & Laskowski, R. (2005). Decomposers: Soil microorganisms and animals. In *Litter decomposition: A guide to carbon and nutrient turnover. Advances in ecological research* (Vol. 38, pp. 73–100). San Diego, CA: Academic Press. [https://doi.org/10.1016/S0065-2504\(05\)38003-2](https://doi.org/10.1016/S0065-2504(05)38003-2)
- Bloem, J., Bolhuis, P. R., Veninga, M. R., & Wieringa, J. (1995). Microscopic methods for counting bacteria and fungi in soil. In K. Alef & P. Nannipieri (Eds.), *Methods in applied soil microbiology and biochemistry* (pp. 162–173). Dordrecht, The Netherlands: Academic Press. Retrieved from <http://library.wur.nl/WebQuery/wurpubs/300230>
- Bradford, M. A., Fierer, N., & Reynolds, J. F. (2008). Soil carbon stocks in experimental mesocosms are dependent on the rate of labile carbon, nitrogen and phosphorus inputs to soils. *Functional Ecology*, 22(6), 964–974. <https://doi.org/10.1111/j.1365-2435.2008.01404.x>
- Bradford, M. A., Jones, T. H., Bardgett, R. D., Black, H. I. J., Boag, B., Bonkowski, M., ... Lawton, J. H. (2002). Impacts of soil faunal community composition on model grassland ecosystems. *Science*, 298(5593), 615. <https://doi.org/10.1126/science.1075805>
- Brose, U., Dunne, J. A., Montoya, J. M., Petchey, O. L., Schneider, F. D., & Jacob, U. (2012). Climate change in size-structured ecosystems. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 367(1605), 2903–2912. <https://doi.org/10.1098/rstb.2012.0232>
- Brundett, M., Bougher, N., Dell, B., Grove, T., & Malajczuk, N. (1996). *Working with mycorrhizas in forestry and agriculture*. Canberra: Australian Centre for International Agricultural Research.
- Buyer, J. S., & Sasser, M. (2012). High throughput phospholipid fatty acid analysis of soils. *Microorganisms and the Sustainable Management of Soil*, 61, 127–130. <https://doi.org/10.1016/j.apsoil.2012.06.005>
- Cleland, E. E., Collins, S. L., Dickson, T. L., Farrer, E. C., Gross, K. L., Gherardi, L. A., ... Suding, K. N. (2013). Sensitivity of grassland plant community composition to spatial vs. temporal variation in precipitation. *Ecology*, 94(8), 1687–1696. <https://doi.org/10.1890/12-1006.1>
- Cole, L., Buckland, S. M., & Bardgett, R. D. (2005). Relating microarthropod community structure and diversity to soil fertility manipulations in temperate grassland. *Soil Biology and Biogeochemistry*, 37(9), 1707–1717. <https://doi.org/10.1016/j.soilbio.2005.02.005>
- Cole, L., Dromph, K. M., Boaglio, V., & Bardgett, R. D. (2004). Effect of density and species richness of soil mesofauna on nutrient mineralisation and plant growth. *Biology and Fertility of Soils*, 39(5), 337–343.
- Cole, L., Staddon, P. L., Sleep, D., & Bardgett, R. D. (2004). Soil animals influence microbial abundance, but not plant-microbial competition for soil organic nitrogen. *Functional Ecology*, 18, 631–640. <https://doi.org/10.1111/j.0269-8463.2004.00894.x>
- Cortet, J., Joffre, R., Elmholt, S., & Krogh, P. H. (2003). Increasing species and trophic diversity of mesofauna affects fungal biomass,

- mesofauna community structure and organic matter decomposition processes. *Biology and Fertility of Soils*, 302–312.
- de Vries, F. T., & Caruso, T. (2016). Eating from the same plate? Revisiting the role of labile carbon inputs in the soil food web. *Soil Biology and Biochemistry*, 102, 4–9. <https://doi.org/10.1016/j.soilbio.2016.06.023>
- de Vries, F. T., Griffiths, R. I., Bailey, M., Craig, H., Girlanda, M., Gweon, H. S., ... Bardgett, R. D. (2018). Soil bacterial networks are less stable under drought than fungal networks. *Nature Communications*, 9(1), 3033. <https://doi.org/10.1038/s41467-018-05516-7>
- de Vries, F. T., Liiri, M. E., Bjørnlund, L., Bowker, M. A., Christensen, S., Setälä, H. M., & Bardgett, R. D. (2012). Land use alters the resistance and resilience of soil food webs to drought. *Nature Climate Change*, 2, 276. <https://doi.org/10.1038/nclimate1368>
- de Vries, F. T., & Shade, A. (2013). Controls on soil microbial community stability under climate change. *Frontiers in Microbiology*, 4, 265. <https://doi.org/10.3389/fmicb.2013.00265>
- de Vries, F. T., Thebault, E., Liiri, M., Birkhofer, K., Tsiafouli, M. A., Bjørnlund, L., ... Bardgett, R. D. (2013). Soil food web properties explain ecosystem services across European land use systems. *Proceedings of the National Academy of Sciences of the United States of America*, 110(35), 14296–14301. <https://doi.org/10.1073/pnas.1305198110>
- Eissfeller, V., Beyer, F., Valtanen, K., Hertel, D., Maraun, M., Polle, A., & Scheu, S. (2013). Incorporation of plant carbon and microbial nitrogen into the rhizosphere food web of beech and ash. *Soil Biology & Biochemistry*, 62, 76–81. <https://doi.org/10.1016/j.soilbio.2013.03.002>
- Endlweber, K., Ruess, L., & Scheu, S. (2009). Collembola switch diet in presence of plant roots thereby functioning as herbivores. *Soil Biology and Biochemistry*, 41(6), 1151–1154. <https://doi.org/10.1016/j.soilbio.2009.02.022>
- Frostegård, A., & Bååth, E. (1996). The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biology and Fertility of Soils*, 22(1), 59–65. <https://doi.org/10.1007/BF00384433>
- Frostegård, A., Bååth, E., & Tunlio, A. (1993). Shifts in the structure of soil microbial communities in limed forests as revealed by phospholipid fatty acid analysis. *Soil Biology and Biochemistry*, 25(6), 723–730. [https://doi.org/10.1016/0038-0717\(93\)90113-P](https://doi.org/10.1016/0038-0717(93)90113-P)
- Fuchslueger, L., Bahn, M., Fritz, K., Hasibeder, R., & Richter, A. (2014). Experimental drought reduces the transfer of recently fixed plant carbon to soil microbes and alters the bacterial community composition in a mountain meadow. *New Phytologist*, 201(3), 916–927. <https://doi.org/10.1111/nph.12569>
- Gamfeldt, L., Hillebrand, H., & Jonsson, P. R. (2008). Multiple functions increase the importance of biodiversity for overall ecosystem functioning. *Ecology*, 89(5), 1223–1231. <https://doi.org/10.1890/06-2091.1>
- Gessner, M. O., Swan, C. M., Dang, C. K., McKie, B. G., Bardgett, R. D., Wall, D. H., & Hattenschwiler, S. (2010). Diversity meets decomposition. *Trends in Ecology & Evolution*, 25(6), 372–380. <https://doi.org/10.1016/j.tree.2010.01.010>
- Goncharov, A. A., Tsurikov, S. M., Potapov, A. M., & Tiunov, A. V. (2016). Short-term incorporation of freshly fixed plant carbon into the soil animal food web: Field study in a spruce forest. *Ecological Research*, 31(6), 923–933. <https://doi.org/10.1007/s11284-016-1402-7>
- Grime, J. P., Fridley, J. D., Askew, A. P., Thompson, K., Hodgson, J. G., & Bennett, C. R. (2008). Long-term resistance to simulated climate change in an infertile grassland. *Proceedings of the National Academy of Sciences of the United States of America*, 105(29), 10028–10032. <https://doi.org/10.1073/pnas.0711567105>
- Guhr, A., Borken, W., Spohn, M., & Matzner, E. (2015). Redistribution of soil water by a saprotrophic fungus enhances carbon mineralization. *Proceedings of the National Academy of Sciences of the United States of America*, 112(47), 14647. <https://doi.org/10.1073/pnas.1514435112>
- Hagedorn, F., Joseph, J., Peter, M., Luster, J., Pritsch, K., Geppert, U., ... Arend, M. (2016). Recovery of trees from drought depends on below-ground sink control. *Nature Plants*, 2(8), 5. <https://doi.org/10.1038/nplants.2016.111>
- Handa, I. T., Aerts, R., Berendse, F., Berg, M. P., Bruder, A., Butenschoten, O., ... Hättenschwiler, S. (2014). Consequences of biodiversity loss for litter decomposition across biomes. *Nature*, 509(7499), 218–221. <https://doi.org/10.1038/nature13247>
- Hanssen, J. F., Thingstad, T. F., Goksøyr, J., & Goksoyr, J. (1974). Evaluation of hyphal lengths and fungal biomass in soil by a membrane filter technique. *Oikos*, 25(1), 102–107. <https://doi.org/10.2307/3543552>
- Hasegawa, M., & Takeda, H. (1995). Changes in feeding attributes of four collembolan populations during the decomposition process of pine needles. *Pedobiologia*, 39(2), 155–169.
- Hasibeder, R., Fuchslueger, L., Richter, A., & Bahn, M. (2015). Summer drought alters carbon allocation to roots and root respiration in mountain grassland. *The New Phytologist*, 205(3), 1117–1127. <https://doi.org/10.1111/nph.13146>
- Hawkes, C. V., Kivlin, S. N., Rocca, J. D., Huguet, V., Thomsen, M. A., & Suttle, K. B. (2011). Fungal community responses to precipitation. *Global Change Biology*, 17(4), 1637–1645. <https://doi.org/10.1111/j.1365-2486.2010.02327.x>
- Hector, A., & Bagchi, R. (2007). Biodiversity and ecosystem multifunctionality. *Nature*, 448(7150), 188–190. <https://doi.org/10.1038/nature05947>
- Hopkin, S. P. (1997). *Biology of the springtails: (Insecta: Collembola)*. Oxford: OUP.
- Hopkin, S. P. (2007). *A key to the Collembola (springtails) of Britain and Ireland*. Shrewsbury: Field Studies Council.
- IPCC. (2017). *Fifth assessment report - Climate change 2013*. Retrieved from <http://www.ipcc.ch/report/ar5/wg1/>
- Johnson, D., Krsek, M., Wellington, E. M. H., Stott, A. W., Cole, L., Bardgett, R. D., ... Leake, J. R. (2005). Soil invertebrates disrupt carbon flow through fungal networks. *Science*, 309(5737), 1047–1047. <https://doi.org/10.1126/science.1114769>
- Johnson, D., Vachon, J., Britton, A. J., & Helliwell, R. C. (2011). Drought alters carbon fluxes in alpine snowbed ecosystems through contrasting impacts on graminoids and forbs. *New Phytologist*, 190(3), 740–749. <https://doi.org/10.1111/j.1469-8137.2010.03613.x>
- Kanters, C., Anderson, I. C., & Johnson, D. (2015). Chewing up the wood-wide web: Selective grazing on ectomycorrhizal fungi by collembola. *Forests*, 6(8), 2560–2570. <https://doi.org/10.3390/f6082560>
- Kardol, P., Cregger, M. A., Company, C. E., & Classen, A. T. (2010). Soil ecosystem functioning under climate change: Plant species and community effects. *Ecology*, 91(3), 767–781. <https://doi.org/10.1890/09-0135.1>
- Kardol, P., Reynolds, W. N., Norby, R. J., & Classen, A. T. (2011). Climate change effects on soil microarthropod abundance and community structure. *Applied Soil Ecology*, 47(1), 37–44. <https://doi.org/10.1016/j.apsoil.2010.11.001>
- Klamer, M., & Bååth, E. (2004). Estimation of conversion factors for fungal biomass determination in compost using ergosterol and PLFA 18:2ω6,9. *Soil Biology and Biochemistry*, 36(1), 57–65. <https://doi.org/10.1016/j.soilbio.2003.08.019>
- Kuzyakov, Y., & Gavrichkova, O. (2010). Time lag between photosynthesis and carbon dioxide efflux from soil: A review of mechanisms and controls. *Global Change Biology*, 16(12), 3386–3406. <https://doi.org/10.1111/j.1365-2486.2010.02179.x>
- Lang, B., Rall, B. C., Scheu, S., & Brose, U. (2014). Effects of environmental warming and drought on size-structured soil food webs. *Oikos*, 123(10), 1224–1233. <https://doi.org/10.1111/j.1600-0706.2013.00894.x>
- Leake, J. R., Ostle, N. J., Rangel-Castro, J. I., & Johnson, D. (2006). Carbon fluxes from plants through soil organisms determined by field ¹³C₂ pulse-labelling in an upland grassland.

- Applied Soil Ecology*, 33(2), 152–175. <https://doi.org/10.1016/j.apsoil.2006.03.001>
- Lensing, J. R., & Wise, D. H. (2006). Predicted climate change alters the indirect effect of predators on an ecosystem process. *Proceedings of the National Academy of Sciences of the United States of America*, 103(42), 15502–15505. <https://doi.org/10.1073/pnas.0607064103>
- Liiri, M., Setälä, H., Haimi, J., Pennanen, T., & Fritze, H. (2002). Soil processes are not influenced by the functional complexity of soil decomposer food webs under disturbance. *Soil Biology & Biochemistry*, 34(7), 1009–1020. [https://doi.org/10.1016/S0038-0717\(02\)00034-2](https://doi.org/10.1016/S0038-0717(02)00034-2)
- Liu, H., Mi, Z., Lin, L. I., Wang, Y., Zhang, Z., Zhang, F., ... He, J.-S. (2018). Shifting plant species composition in response to climate change stabilizes grassland primary production. *Proceedings of the National Academy of Sciences of the United States of America*, 115(16), 4051–4056. <https://doi.org/10.1073/pnas.1700299114>
- Makonen, M., Berg, M. P., van Hal, J. R., Callaghan, T. V., Press, M. C., & Aerts, R. (2011). Traits explain the responses of a sub-arctic Collembola community to climate manipulation. *Soil Biology & Biochemistry*, 43(2), 377–384. <https://doi.org/10.1016/j.soilbio.2010.11.004>
- McGonigle, T. P., Miller, M. H., Evans, D. G., Fairchild, G. L., & Swan, J. A. (1990). A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. *New Phytologist*, 115(3), 495–501. <https://doi.org/10.1111/j.1469-8137.1990.tb00476.x>
- Meisner, A., Deyn, G. B. D., de Boer, W., & van der Putten, W. H. (2013). Soil biotic legacy effects of extreme weather events influence plant invasiveness. *Proceedings of the National Academy of Sciences of the United States of America*, 110(24), 9835–9838. <https://doi.org/10.1073/pnas.1300922110>
- Mikola, J., Bardgett, R. D., & Hedlund, K. (2002). Biodiversity, ecosystem functioning and soil decomposer food webs. In M. Loreau, S. Naeem, & P. Inchausti (Eds.), *Biodiversity and ecosystem functioning: Synthesis and perspectives* (pp. 169–180). Oxford, UK: Oxford University Press.
- Nielsen, U. N., Ayres, E., Wall, D. H., & Bardgett, R. D. (2011). Soil biodiversity and carbon cycling: A review and synthesis of studies examining diversity-function relationships. *European Journal of Soil Science*, 62(1), 105–116. <https://doi.org/10.1111/j.1365-2389.2010.01314.x>
- Pausch, J., Kramer, S., Scharroba, A., Scheunemann, N., Butenschon, O., Kandeler, E., ... Ruess, L. (2016). Small but active – Pool size does not matter for carbon incorporation in below-ground food webs. *Functional Ecology*, 30(3), 479–489. <https://doi.org/10.1111/1365-2435.12512>
- Pollierer, M. M., Dyckmans, J., Scheu, S., & Haubert, D. (2012). Carbon flux through fungi and bacteria into the forest soil animal food web as indicated by compound-specific ¹³C fatty acid analysis. *Functional Ecology*, 26(4), 978–990. <https://doi.org/10.1111/j.1365-2435.2012.02005.x>
- Pollierer, M. M., Langel, R., Koerner, C., Maraun, M., & Scheu, S. (2007). The underestimated importance of belowground carbon input for forest soil animal food webs. *Ecology Letters*, 10(8), 729–736. <https://doi.org/10.1111/j.1461-0248.2007.01064.x>
- R Core Team. (2017). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org>
- Reichstein, M., Bahn, M., Ciais, P., Frank, D., Mahecha, M. D., Seneviratne, S. I., ... Wattenbach, M. (2013). Climate extremes and the carbon cycle. *Nature*, 500(7462), 287–295. <https://doi.org/10.1038/nature12350>
- Ritz, K. (2011). Microbes, habitat space, and transport in soil. In J. Gliński, J. Horabik, & J. Lipiec (Eds.), *Encyclopedia of agrophysics* (pp. 472–475). Encyclopedia of Earth Sciences Series. Dordrecht: Springer.
- Rodwell, J. S. (1998). *British plant communities: Volume 3. Grasslands and montane communities*. Cambridge: Cambridge University Press.
- Rooney, N., & McCann, K. S. (2012). Integrating food web diversity, structure and stability. *Trends in Ecology & Evolution*, 27(1), 40–46. <https://doi.org/10.1016/j.tree.2011.09.001>
- Rooney, N., McCann, K., Gellner, G., & Moore, J. C. (2006). Structural asymmetry and the stability of diverse food webs. *Nature*, 442, 265. <https://doi.org/10.1038/nature04887>
- Ruehr, N. K., Offermann, C. A., Gessler, A., Winkler, J. B., Ferrio, J. P., Buchmann, N., & Barnard, R. L. (2009). Drought effects on allocation of recent carbon: From beech leaves to soil CO₂ efflux. *The New Phytologist*, 184(4), 950–961. <https://doi.org/10.1111/j.1469-8137.2009.03044.x>
- Ruf, A., Kuzyakov, Y., & Lopatovskaya, O. (2006). Carbon fluxes in soil food webs of increasing complexity revealed by C-14 labelling and C-13 natural abundance. *Soil Biology & Biochemistry*, 38(8), 2390–2400. <https://doi.org/10.1016/j.soilbio.03.008>
- Santonja, M., Fernandez, C., Proffit, M., Gers, C., Gauquelin, T., Reiter, I. M., ... Baldy, V. (2017). Plant litter mixture partly mitigates the negative effects of extended drought on soil biota and litter decomposition in a Mediterranean oak forest. *Journal of Ecology*, 105(3), 801–815. <https://doi.org/10.1111/1365-2745.12711>
- Scheu, S., & Setälä, H. (2002). Multitrophic interactions in decomposer food webs. In T. Scharntke & B. A. Hawkins (Eds.), *Multitrophic level interactions* (pp. 223–264). Cambridge: Cambridge University Press.
- Scheu, S., & Simmerling, F. (2004). Growth and reproduction of fungal feeding Collembola as affected by fungal species, melanin and mixed diets. *Oecologia*, 139(3), 347–353. <https://doi.org/10.1007/s00442-004-1513-7>
- Schwalm, C. R., Williams, C. A., Schaefer, K., Arneith, A., Bonal, D., Buchmann, N., ... Richardson, A. D. (2010). Assimilation exceeds respiration sensitivity to drought: A FLUXNET synthesis. *Global Change Biology*, 16(2), 657–670. <https://doi.org/10.1111/j.1365-2486.2009.01991.x>
- Schwarz Müller, F., Eisenhauer, N., & Brose, U. (2015). “Trophic whales” as biotic buffers: Weak interactions stabilize ecosystems against nutrient enrichment. *The Journal of Animal Ecology*, 84(3), 680–691. <https://doi.org/10.1111/1365-2656.12324>
- Siebert, J., Sünemann, M., Auge, H., Berger, S., Cesarz, S., Ciobanu, M., ... Eisenhauer, N. (2019). The effects of drought and nutrient addition on soil organisms vary across taxonomic groups, but are constant across seasons. *Scientific Reports*, 9(1), 639. <https://doi.org/10.1038/s41598-018-36777-3>
- Thakur, M. P., Reich, P. B., Hobbie, S. E., Stefanski, A., Rich, R., Rice, K. E., ... Eisenhauer, N. (2018). Reduced feeding activity of soil detritivores under warmer and drier conditions. *Nature Climate Change*, 8(1), 75–78. <https://doi.org/10.1038/s41558-017-0032-6>
- Thornton, B., Zhang, Z., Mayes, R. W., Högberg, M. N., & Midwood, A. J. (2011). Can gas chromatography combustion isotope ratio mass spectrometry be used to quantify organic compound abundance? *Rapid Communications in Mass Spectrometry: RCM*, 25(17), 2433–2438. <https://doi.org/10.1002/rcm.5148>
- Vierheilig, H., Coughlan, A. P., Wyss, U., & Piche, Y. (1998). Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology*, 64(12), 5004–5007.
- Wagg, C., Bender, S. F., Widmer, F., & van der Heijden, M. G. A. (2014). Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Sciences of the United States of America*, 111(14), 5266–5270. <https://doi.org/10.1073/pnas.1320054111>
- Yang, G., Wagg, C., Veresoglou, S. D., Hempel, S., & Rillig, M. C. (2018). How Soil Biota Drive Ecosystem Stability. *Trends in Plant Science*, 23(12), 1057–1067. <https://doi.org/10.1016/j.tplants.2018.09.007>

How to cite this article: Chomel M, Lavalée JM, Alvarez-Segura N, et al. Drought decreases incorporation of recent plant photosynthate into soil food webs regardless of their trophic complexity. *Glob Change Biol*. 2019;25:3549–3561. <https://doi.org/10.1111/gcb.14754>