



UvA-DARE (Digital Academic Repository)

Supersymmetric Rényi entropy and charged hyperbolic black holes

Hosseini, S.M.; Toldo, C.; Yaakov, I.

DOI

[10.1007/JHEP07\(2020\)131](https://doi.org/10.1007/JHEP07(2020)131)

Publication date

2020

Document Version

Final published version

Published in

Journal of High Energy Physics

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Hosseini, S. M., Toldo, C., & Yaakov, I. (2020). Supersymmetric Rényi entropy and charged hyperbolic black holes. *Journal of High Energy Physics*, 2020(7), [131]. [https://doi.org/10.1007/JHEP07\(2020\)131](https://doi.org/10.1007/JHEP07(2020)131)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Supersymmetric Rényi entropy and charged hyperbolic black holes

Seyed Morteza Hosseini,^a Chiara Toldo^{b,c,d} and Itamar Yaakov^e

^a*Kavli IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan*

^b*Kavli Institute for Theoretical Physics, Kohn Hall, University of California Santa Barbara, CA, 93106, U.S.A.*

^c*Centre de Physique Théorique (CPHT), Ecole Polytechnique, 91128 Palaiseau Cedex, France*

^d*Institut de Physique Théorique, Université Paris Saclay, CEA, CNRS, Orme des Merisiers, 91191 Gif-sur-Yvette Cedex, France*

^e*INFN — Sezione di Milano Bicocca, Dipartimento di Fisica, Edificio U2, Piazza della Scienza 3, I-20126 Milano, MI, Italy*

E-mail: morteza.hosseini@ipmu.jp, chiara.toldo@polytechnique.edu, itamar.yaakov@mib.infn.it

ABSTRACT: The supersymmetric Rényi entropy across a spherical entangling surface in a d -dimensional SCFT with flavor defects is equivalent to a supersymmetric partition function on $\mathbb{H}^{d-1} \times \mathbb{S}^1$, which can be computed exactly using localization. We consider the holographically dual BPS solutions in $(d + 1)$ -dimensional matter coupled supergravity ($d = 3, 5$), which are charged hyperbolically sliced AdS black holes. We compute the renormalized on-shell action and the holographic supersymmetric Rényi entropy and show a perfect match with the field theory side. Our setup allows a direct map between the chemical potentials for the global symmetries of the field theories and those of the gravity solutions. We also discuss a simple case where angular momentum is added.

KEYWORDS: AdS-CFT Correspondence, Black Holes in String Theory, Supersymmetric Gauge Theory

ARXIV EPRINT: [1912.04868](https://arxiv.org/abs/1912.04868)

Contents

1	Introduction	1
2	Field theory	2
2.1	Supersymmetric Rényi entropy	2
2.1.1	Definition of Rényi entropy	3
2.1.2	Definition of supersymmetric Rényi entropy	4
2.1.3	Localization and deformation of SRE	5
2.1.4	The SRE matrix model deformed by defects	5
2.2	Squashed S^3 free energy	6
2.3	Squashed S^5 free energy	8
3	Four-dimensional solutions from the <i>stu</i> model	10
3.1	Warm up: BPS hyperbolic Reissner-Nordström	10
3.2	Hyperbolic black hole solutions of the <i>stu</i> model	12
3.3	Holographic supersymmetric Rényi entropy	13
3.4	Holographic matching	15
4	Six-dimensional hyperbolic solutions	17
4.1	Romans F(4) gauged supergravity coupled to matter	17
4.2	Six-dimensional supersymmetric hyperbolic black holes	18
4.3	Supersymmetric Rényi entropy	19
4.4	Holographic matching	20
5	Concluding remarks	21
A	Explicit construction of the Killing spinor	22
A.1	Four dimensions	22
A.2	Six dimensions	23
B	On-shell action via holographic renormalization	25
C	Computation of the charges	30
D	Rotating charged hyperbolic solutions	31

1 Introduction

The entanglement entropy of the vacuum is an example of a universal observable in quantum field theory, independent of the existence of a particular set of fields, which has many interesting and useful properties. Most prominent among these are its monotonicity properties as a function of the size of the entangling region [1], and the existence of a simple geometric interpretation in the context of holography [2]. We refer the reader to the review [3] for more information.

The Rényi entropy is a one parameter refinement of the entanglement entropy. Besides containing additional information, the Rényi entropy is notable for having a straightforward Euclidean path integral interpretation known as the replica trick [4]. Supersymmetric Rényi entropy (SRE) is a twisted version, in the sense of $(-1)^F$, of Rényi entropy which can be defined for supersymmetric theories in a variety of spacetime dimensions and with varying amounts of supersymmetry [5–8]. Unlike Rényi entropy, SRE can be calculated exactly at arbitrary coupling using the method of supersymmetric localization. It nevertheless shares many of the interesting properties of the untwisted version, including the ability to recover the entanglement entropy as a limit.

In a d -dimensional superconformal field theory (SCFT), the SRE for a $d-2$ -dimensional spherical entangling surface can be computed using the partition function on a d -sphere, branched n times over a maximal $d-2$ -sphere, where the metric has a conical singularity. In holographically dual solutions, gravity becomes dynamical and the issue arises of how to treat such a singularity. By conformally mapping the branched sphere to $\mathbb{H}^{d-1} \times \mathbb{S}^1$, where \mathbb{H} denotes hyperbolic space, the singularity is pushed to infinity. The Rényi entropy is mapped to the thermal entropy in this space, with the new Euclidean time having periodicity $\beta = 2\pi n$. The SRE is likewise mapped to a twisted thermal partition function. The details of the singularity are encoded in the boundary conditions on this space. The gravity duals are hyperbolically sliced solutions, so-called “topological” black holes, whose boundary is indeed of the form $\mathbb{H}^{d-1} \times \mathbb{S}^1$.

The computation of the SRE in d -dimensional models ($d = 2, 3, 4, 5, 6$) with a holographic dual was performed, respectively, in [6, 8–13]. The matching with the gravity computation of the SRE was achieved with supergravity hyperbolic black holes supported by a single gauge field, which corresponds to the graviphoton. Here, we take this one step further by considering supergravity backgrounds with more general couplings, in particular vector multiplets. The corresponding dual field theory computation therefore includes fugacities for the global symmetries of the theory in the thermal picture. Equivalently, co-dimension two flavor vortex defects are included in the d sphere picture. In gravity, we work with four and six-dimensional supergravity solutions, achieving a match with the field theory SRE in $d = 3, 5$ by evaluating the supergravity renormalized on-shell action. We choose to work with d odd because the finite part of the free energy in the field theory is believed to be universal in odd dimensions. For comparison, in the even d case the coefficient of the Weyl anomaly is always universal, while the subleading piece may only be universal in the presence of a sufficient amount of supersymmetry [14]. By working in even-dimensional supergravity, we also avoid subtleties in holographic renormalization

schemes related to the Casimir energy, see e.g. [15–17]. Let us, however, mention that the SRE of supergravity solutions in $d+1 = 5, 7$, coupled to matter were compared to the field theory result, respectively, in [7, 18].

The aim of this paper is twofold. On one hand, we wish to investigate how the SRE is computed holographically in the case where matter couplings are incorporated — in the present case, this consists in considering hyperbolic black hole solutions supported by vector multiplets. On the other hand, our setup allows us to directly map the fugacities appearing in the field theory computation to the black hole chemical potentials. The mapping that we obtain is then rather manifest.¹

The paper is organized as follows. We will first provide results for the supersymmetric Rényi entropy with flavor fugacities for specific models: the ABJM model in $d = 3$, and a $\mathcal{N} = 1$, $\text{USp}(2N)$ gauge theory with N_f fundamental and one anti-symmetric hypermultiplets in $d = 5$. These models have well known gravity dual descriptions. We then focus on the gravity duals to SRE in four and six dimensions, which are hyperbolic black holes. We spell out the solutions, which are new in the $d = 6$ case, and compute their renormalized on-shell action. We show that this matches with the SRE computation. In appendix A, we explicitly construct the Killing spinors for the hyperbolic black holes. Appendix B shows the computation of the renormalized on-shell action via holographic renormalization techniques and appendix C shows that the black hole charges computed from supergravity match those computed in the SCFT. In appendix D, we present a simple example of a rotating hyperbolic black hole which generalizes the static case in section 3.1, and provide the value of its renormalized on-shell action.

2 Field theory

In this section we calculate the free energy of SCFTs on $\mathbb{H}^{d-1} \times \mathbb{S}^1$ that are holographically dual to our hyperbolic BPS black holes. We first introduce the supersymmetric Rényi entropy (SRE) and its deformation by BPS vortex defects. We then describe the relationship of these defects to black hole chemical potentials. Using supersymmetric localization, we construct an appropriate matrix model which captures the exact answer for the free energy. Finally, we use large N techniques to explicitly evaluate the matrix model for field theories dual to the black hole solutions.

2.1 Supersymmetric Rényi entropy

We briefly review the definition of Rényi entropy and its supersymmetric counterpart (SRE). We then show how co-dimension two defect operators alter the localization result

¹For instance, in the case of rotating electric black holes, an elegant prescription to map the black hole chemical potentials to the field theory ones was recently put forward in [19]. This procedure requires taking an extremal limit of a family of supersymmetric, complexified solutions, and the definition of the black hole chemical potentials via appropriate subtraction of the extremal BPS values. In our framework, upon Wick-rotating the BPS black hole solution we are left with a regular geometry with topology $\mathbb{R}^2 \times \mathbb{H}^{d-1}$, where a formal finite temperature can be defined. This allows us to directly map the chemical potentials in gravity into those on the field theory side, with no need for such a subtraction.

for SRE. Finally, we relate such defects to chemical potentials in the partition function on hyperbolic space.

2.1.1 Definition of Rényi entropy

Following the notation in [3], we define entanglement entropy for a vacuum state Ψ by first making a choice of a subregion A of a spatial slice. The complement will be denoted by $B = \bar{A}$. We make the assumption that the Hilbert space of the theory can be likewise locally split as²

$$\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B. \tag{2.1}$$

We then form the reduced density matrix corresponding to A

$$\rho_A \equiv \text{tr}_B |\Psi\rangle \langle \Psi|. \tag{2.2}$$

The entanglement entropy associated to A can be defined as the von Neumann entropy of ρ_A ,

$$S(A) \equiv -\text{Tr} \rho_A \log \rho_A. \tag{2.3}$$

The Rényi entropy is a one parameter refinement of the entanglement entropy defined by

$$S_n(A) \equiv \frac{1}{1-n} \log \text{tr} \rho_A^n, \quad n \in \mathbb{N}. \tag{2.4}$$

It satisfies the relation

$$\lim_{n \rightarrow 1} S_n(A) = S(A), \tag{2.5}$$

where the limit is understood to be taken using an appropriate continuation to non-integer n . We will restrict our attention to the case where A is the $d-1$ ball and the entangling surface is $\partial A = \mathbb{S}^{d-2}$.

The Rényi entropy of a quantum field theory is, in general, divergent. However, for d odd the finite part of the Rényi entropy of a CFT is believed to be a universal observable (see [3] and references within).

The Rényi entropy can alternatively be computed using the replica trick [4]. One considers the path integral on an n -fold cover of the original spacetime branched around the entangling surface ∂A . Denoting the partition function on this space by Z_n , we will *define* the n -th Rényi entropy for a positive integer n by

$$S_n \equiv \frac{1}{1-n} \log \left(\frac{Z_n}{(Z_1)^n} \right). \tag{2.6}$$

This definition is incomplete because the branching means that the spacetime corresponding to Z_n is not smooth but has conical singularities. One could complete the definition by specifying appropriate boundary conditions for all fields at ∂A . We will instead concentrate on the definition of SRE, reviewed in section 2.1, which uses a particular prescription for smoothing out the singularities [5].

²For a critical discussion of the validity of this assumption, see references in footnote 3 of [3]. The subtleties associated with this splitting will not affect our results.

The line element on a branched d sphere is defined as the round sphere metric with a different coordinate range

$$ds^2 = \ell^2 (d\theta^2 + \sin^2(\theta)d\tau^2 + \cos^2(\theta)ds_{\mathbb{S}^{d-2}}^2) , \tag{2.7}$$

$$\theta \in [0, \pi/2], \quad \tau \in [0, 2\pi n) .$$

This metric has a conical singularity along the co-dimension two maximal $d - 2$ sphere at $\theta = 0$.

For n a positive integer, the branched sphere is related by a Weyl transformation to the branched version of \mathbb{R}^d used to define the n -th Rényi entropy [20]. In order to avoid working with a singular space, we can conformally map this space by $\cot(\theta) = \sinh(\chi)$ to $\mathbb{H}^{d-1} \times \mathbb{S}^1$ with line element

$$ds_{\mathbb{H}^{d-1} \times \mathbb{S}^1}^2 = d\tau^2 + d\chi^2 + \sinh(\chi)^2 ds_{\mathbb{S}^{d-2}}^2 , \tag{2.8}$$

$$\chi \in [0, \infty], \quad \tau \in [0, 2\pi n) .$$

The Rényi entropy maps to the thermal entropy in this space with inverse temperature $\beta = 2\pi n$. The singularity at $\theta = 0$ is mapped to $\chi \rightarrow \infty$ [20].

2.1.2 Definition of supersymmetric Rényi entropy

The supersymmetric Rényi entropy (SRE) is a twisted version, in the sense of $(-1)^F$, of Rényi entropy [5–8]. In order to preserve supersymmetry in SRE, one must give nonzero values to additional fields, aside from the metric, in the background supergravity multiplet to which the SCFT is coupled [21–24]. Specifically, one needs to turn on a background R-symmetry gauge field, $A^{(R)}$, which is flat in the bulk of the space and has a delta function like field strength supported on the singularity [5]. For example, in a three-dimensional $\mathcal{N} = 2$ field theory we have [5]³

$$A^{(R)} = -\frac{n-1}{2n} d\tau . \tag{2.9}$$

After the additional Weyl transformation to $\mathbb{H}^{d-1} \times \mathbb{S}^1$, the SRE is related to a twisted, in the sense of $(-1)^F$, version of the thermal partition function which we can call the *hyperbolic index*, in analogy with the superconformal index [25, 26]. A representation of this quantity as a trace over the Hilbert space $\mathcal{H}_{\mathbb{H}^{d-1}}$ of states on \mathbb{H}^{d-1} was given in [27]. Including flavor charges, we can write⁴

$$Z_n^{\text{susy}} = \text{Tr}_{\mathcal{H}_{\mathbb{H}^{d-1}}} e^{-2\pi n (H - i \sum_I \alpha^I Q_I^{\text{flavor}} + i \frac{n-1}{n} Q_R)} , \tag{2.10}$$

where H is the Hamiltonian, Q_R is the R-symmetry charge, Q_I^{flavor} are flavor charges, and the α^I are flavor chemical potentials. The SRE is then defined as⁵

$$S_n^{\text{SRE}} \equiv \frac{1}{1-n} \log \left| \frac{Z_n^{\text{susy}}}{(Z_1^{\text{susy}})^n} \right| . \tag{2.11}$$

³The sign of $A^{(R)}$ chosen here, which is correlated with the choice of Killing spinor preserved by SRE, corresponds to our gravity conventions and is opposite to the one chosen in [5].

⁴As an index, Z_n^{susy} does not change under renormalization group flow, and thus can be computed either in the UV or the IR SCFT. The parameter n is a chemical potential for a combination of charges commuting with the supercharge, similar to those found in [25, 26].

⁵The absolute value, which is absent from the usual definition of Rényi entropy, is used here to avoid some subtleties associated with possible non-universal terms. See [5] for a discussion of the $d = 3$ case.

2.1.3 Localization and deformation of SRE

The partition function defining the SRE can be computed exactly using the method of supersymmetric localization [28, 29]. In the case of SRE in three dimensions, the matrix model one gets from localization coincides with the one used to compute the partition function on the squashed sphere with the squashing parameter related to n in a simple way [5, 30].⁶ This relationship continues to hold for higher dimensions and we consequently make no distinction between the free energy in the two matrix models.

The partition function defining the SRE can be refined by supersymmetric deformations while remaining amenable to localization [5, 31].⁷ Deformations include masses for matter multiplets and Fayet-Iliopoulos (FI) terms for abelian vector multiplets. These deformations break conformal invariance. Additionally, the form of the coupling of the theory to the background supergravity fields, including $A^{(R)}$, depends on a choice of R-symmetry current. If the R-symmetry is abelian, one may choose an arbitrary linear combination of R-symmetry and abelian flavor symmetry currents. In an SCFT, a particular combination, the result of dynamical mixing, is dictated by the superconformal algebra where the R-symmetry transformations appear [32, 33].

Supersymmetric operators can also be added to the SRE. These include Wilson loops and co-dimension two vortex defects [31, 34, 35]. The latter are inserted by demanding that the fields in the path integral have prescribed singularities on the defect worldvolume [36]. If the defect is in a flavor symmetry this is equivalent to introducing background flavor symmetry gauge fields which are flat outside the defect. In fact, the deformation leading from the usual sphere partition function to the SRE is itself such a defect, embedded in the background supergravity multiplet. Due to this, addition of flavor defects to the SRE, oriented along the same sub-manifold, is essentially the same as the R-symmetry mixing effect described above. However, the strength of the defect is now unrelated to the superconformal algebra and represents a deformation of the SRE. In the hyperbolic picture, such a defect is mapped to the holonomy of a flavor symmetry connection along the time direction, i.e. a flavor fugacity. The chemical potentials α for such a fugacity are linearly related to the A_τ^{flavor} flavor gauge fields introduced below, with a proportionality constant which depends on the normalization of the charges.

2.1.4 The SRE matrix model deformed by defects

The matrix model for the round sphere deformed by co-dimension two defects, in dimensions $d = 3, 4, 5$ was derived in [37]. It was shown that a background U(1) flavor symmetry connection A_τ^{flavor} with holonomy $\exp(2\pi i A_\tau^{\text{flavor}})$ induces, after localizing to a matrix model, a mass deformation term

$$m_{\text{defect}} = -i A_\tau^{\text{flavor}}. \tag{2.12}$$

The fact that the mass is imaginary is part of the relationship to R-symmetry mixing. The large N limit of the same matrix models in the presence of R-symmetry mixing or

⁶This is true at the level of the matrix model, not just the final result.

⁷We describe the situation in three dimensions. The situation in five dimensions is analogous.

of mass terms has previously been derived in [38–40]. The mixing parameters are usually called Δ , while masses are denoted by m . Besides being purely imaginary, the mass term induced by the defect also has an origin which is naturally $A_\tau^{\text{flavor}} = 0$. This is true also for the real “physical masses” m . On the other hand, in a theory which has a non-abelian R-symmetry group, the Δ ’s have an origin which is determined by the canonical R-charge, or the canonical dimensions, of matter multiplets. In three dimensions this is $\Delta = 1/2$, while in five dimensions it is $\Delta = 3/2$.

Taking all this into account, and using the relationship between n and the squashing parameter b derived in [37], the defect deformed three-dimensional matrix models are given by those of [38] with the substitution⁸

$$\Delta_{\text{there}} = \frac{1}{2} + \frac{2nA_\tau^{\text{flavor}}}{n+1}, \quad b_{\text{there}} = \frac{1}{\sqrt{n}}. \quad (2.13)$$

For five-dimensional $\mathcal{N} = 1$ theories appearing in [40], we can simply take

$$m_{\text{there}} = -iA_\tau^{\text{flavor}}, \quad \vec{\omega}_{\text{there}} = (1, 1, 1/n). \quad (2.14)$$

We will adopt a democratic convention for the deformation parameters Δ , whereby the physical parameters are augmented by one additional parameter and a constraint is imposed. Interpreting Δ as the result of a flavor defect, we will add a corresponding A^{flavor} . The constraint in terms of A^{flavor} is simply

$$\sum_I A^{\text{flavor},I} = 0. \quad (2.15)$$

2.2 Squashed \mathbb{S}^3 free energy

In this section, we review the squashed \mathbb{S}^3 partition function and its large N limit, as analyzed in [38, 39]. For the purpose of this paper, we consider the ABJM model [41], which is holographically dual to an $\text{AdS}_4 \times \mathbb{S}^7/\mathbb{Z}_k$ background of M-theory. ABJM is a three-dimensional $\mathcal{N} = 6$ supersymmetric Chern-Simons-matter theory with gauge group $U(N)_k \times U(N)_{-k}$ (the subscripts represent the CS levels) with two pairs of bi-fundamental chiral fields A_i and B_i , $i = 1, 2$, in the representation $(\mathbf{N}, \bar{\mathbf{N}})$ and $(\bar{\mathbf{N}}, \mathbf{N})$ of the gauge group, respectively. The chiral fields interact through the quartic superpotential

$$W = \text{Tr} (A_1 B_1 A_2 B_2 - A_1 B_2 A_2 B_1). \quad (2.16)$$

In the $\mathcal{N} = 2$ formulation, the ABJM model has a $U(2) \times U(2)$ action which acts separately on the chiral fields $A_{1,2}$ and $B_{1,2}$. There is a $U(1)^3$ subgroup of the Cartan of this group which preserves the superpotential, a particular linear combination of which is gauged. In addition, there are two topological $U(1)_J$ symmetries. The current for one of these topological symmetries is set to zero by the equations of motion. Due to the appearance of Chern-Simons terms, the action of the other $U(1)_J$ is mixed with the gauge

⁸The setup is symmetric with respect to inversion of b . In order to conform to the notation in [38], we set $b = 1/\sqrt{n}$ instead of $b = \sqrt{n}$ as in [37].

group action. We will work in a gauge in which the fugacity conjugate to the remaining topological symmetry, which could be explicitly added using an FI parameter, is fixed to 1. The remaining global symmetry group, which we will call the flavor group, is given by the $U(1)^3$ compatible with the superpotential acting on the chiral fields. The model admits therefore a three-parameter space of flavor symmetry, or Δ type, deformations.⁹

We introduce the R-charges Δ_I , $I = 1, \dots, 4$, one for each of the four fields $\{A_i, B_i\}$, satisfying

$$\sum_{I=1}^4 \Delta_I = 2. \quad (2.17)$$

The partition function can be written as

$$Z_{\mathbb{S}_b^3} = \int_{-\infty}^{\infty} \left[\prod_{i=1}^N \frac{d\lambda_i}{2\pi} \frac{d\tilde{\lambda}_i}{2\pi} \right] e^{-F_{\mathbb{S}_b^3}(\lambda_i, \tilde{\lambda}_i)}, \quad (2.18)$$

where

$$\begin{aligned} F_{\mathbb{S}_b^3} = & 2 \log N! - \frac{ik}{4\pi b^2} \sum_{i=1}^N (\lambda_i^2 - \tilde{\lambda}_i^2) \\ & - \sum_{i<j}^N \left\{ \log \left[2 \sinh \left(\frac{\lambda_i - \lambda_j}{2} \right) \right] + \log \left[2 \sinh \left(\frac{\lambda_i - \lambda_j}{2b^2} \right) \right] \right\} \\ & - \sum_{i<j}^N \left\{ \log \left[2 \sinh \left(\frac{\tilde{\lambda}_i - \tilde{\lambda}_j}{2} \right) \right] + \log \left[2 \sinh \left(\frac{\tilde{\lambda}_i - \tilde{\lambda}_j}{2b^2} \right) \right] \right\} \\ & - \sum_{i,j=1}^N \sum_{a=1}^2 S_2 \left(\frac{i\Omega}{2} (1 - \Delta_a) - \frac{1}{2\pi b} (\lambda_i - \tilde{\lambda}_j) \middle| b \right) \\ & - \sum_{i,j=1}^N \sum_{b=3}^4 S_2 \left(\frac{i\Omega}{2} (1 - \Delta_b) + \frac{1}{2\pi b} (\lambda_i - \tilde{\lambda}_j) \middle| b \right). \end{aligned} \quad (2.19)$$

Here, $\Omega = b + 1/b$ and $S_2(\lambda|b)$ is the double sine function.

Large N free energy. Consider the following ansatz for the large N saddle point eigenvalue distribution,

$$\lambda_j = N^{1/2} t_j + i v_j, \quad \tilde{\lambda}_j = N^{1/2} t_j + i \tilde{v}_j. \quad (2.20)$$

In the large N limit, we define the continuous functions $t_j = t(j/N)$ and $v_j = v(j/N)$, $\tilde{v}_j = \tilde{v}(j/N)$; and we introduce the density of eigenvalues

$$\rho(t) = \frac{1}{N} \frac{dj}{dt}, \quad s.t. \int dt \rho(t) = 1. \quad (2.21)$$

At large N the sums over N become Riemann integrals, for example,

$$\sum_{j=1}^N \rightarrow N \int dt \rho(t). \quad (2.22)$$

⁹We would like to thank Alberto Zaffaroni for explaining this point.

The large N free energy is then given by [38, 39]

$$\begin{aligned} \frac{F_{\mathbb{S}^3_b}[\rho(t), \delta v(t), \Delta_I | b]}{N^{3/2}} &= \frac{k}{2\pi b^2} \int dt \rho(t) t \delta v(t) - \gamma \left(\int dt \rho(t) - 1 \right) \\ &\quad - \frac{b\Omega^3}{16} \sum_{a=1}^2 (2 - \Delta_a^+) \int dt \rho(t)^2 \left[\left(2 \frac{\delta v(t)}{b\Omega} + \pi \Delta_a^- \right)^2 - \frac{\pi^2}{3} \Delta_a^+ (4 - \Delta_a^+) \right], \end{aligned} \quad (2.23)$$

where we defined $\delta v(t) \equiv v(t) - \bar{v}(t)$, $\Delta_1^\pm \equiv \Delta_1 \pm \Delta_4$, $\Delta_2^\pm \equiv \Delta_2 \pm \Delta_3$, and we added the Lagrange multiplier γ for the normalization of $\rho(t)$. Setting to zero the variation of (2.23) with respect to $\rho(t)$ and $\delta v(t)$ we obtain the following saddle point configuration. We have a central region where

$$\begin{aligned} \rho(t) &= \frac{16b\gamma + 4\Omega kt(\Delta_1\Delta_2 - \Delta_3\Delta_4)}{4\pi^2 b^2 \Omega^3 (\Delta_1 + \Delta_3)(\Delta_2 + \Delta_3)(\Delta_1 + \Delta_4)(\Delta_2 + \Delta_4)}, & -\frac{2b\gamma}{\Omega k \Delta_1} < t < \frac{2b\gamma}{\Omega k \Delta_3}. \\ \delta v(t) &= \frac{2\pi b \Omega^2 kt \sum_{a < b < c} \Delta_a \Delta_b \Delta_c - 4\pi b^2 \Omega \gamma (\Delta_1 \Delta_2 - \Delta_3 \Delta_4)}{8b\gamma + 2\Omega kt(\Delta_1\Delta_2 - \Delta_3\Delta_4)}, \end{aligned} \quad (2.24)$$

When $\delta v = -\pi b \Omega \Delta_2$ on the left the solution reads

$$\rho(t) = \frac{2b\gamma + \Omega kt \Delta_2}{\pi^2 b^2 \Omega^3 (\Delta_1 - \Delta_2)(\Delta_2 + \Delta_3)(\Delta_2 + \Delta_4)}, \quad -\frac{2b\gamma}{\Omega k \Delta_2} < t < -\frac{2b\gamma}{\Omega k \Delta_1}, \quad (2.25)$$

while when $\delta v = \pi b \Omega \Delta_4$ on the right the solution is given by

$$\rho(t) = -\frac{2b\gamma - \Omega kt \Delta_4}{\pi^2 b^2 \Omega^3 (\Delta_1 + \Delta_4)(\Delta_2 + \Delta_4)(\Delta_4 - \Delta_3)}, \quad \frac{2b\gamma}{\Omega k \Delta_3} < t < \frac{2b\gamma}{\Omega k \Delta_4}. \quad (2.26)$$

The normalization of $\rho(t)$ fixes the value of γ as

$$\gamma = \frac{\pi \Omega^2}{\sqrt{2}} \sqrt{k \Delta_1 \Delta_2 \Delta_3 \Delta_4}. \quad (2.27)$$

Plugging the above solution back into (2.23) we obtain the squashed \mathbb{S}^3 free energy¹⁰

$$F_{\mathbb{S}^3_b}(\Delta_I | \Omega) = \frac{2N^{3/2}}{3} \gamma = \frac{\pi N^{3/2} \Omega^2}{3} \sqrt{2k \Delta_1 \Delta_2 \Delta_3 \Delta_4} = \frac{\Omega^2}{4} F_{\mathbb{S}^3}(\Delta_I), \quad (2.28)$$

where $F_{\mathbb{S}^3}$ is the free energy of ABJM on the round \mathbb{S}^3 , i.e. $b = 1$, see [42, section 5]. This is precisely [38, (3.38)].

2.3 Squashed \mathbb{S}^5 free energy

In this section we review the large N limit of the squashed \mathbb{S}^5 free energy of the $\text{USp}(2N)$ gauge theory with N_f hypermultiplets in the fundamental representation and one hypermultiplet in the antisymmetric representation of $\text{USp}(2N)$, as analyzed in [40]. The gauge theories of interest live on the intersection of N D4-branes and N_f D8-branes and orientifold planes in type I' string theory and are holographically dual to a warped $\text{AdS}_6 \times \mathbb{S}^4$ background of massive type IIA supergravity [43] (see also [44–47]).

¹⁰The first equality arises from a virial theorem for the free energy (2.23).

The perturbative partition function can be written as¹¹

$$Z_{\mathbb{S}_\omega^5}^{\text{pert}} = \int_{-\infty}^{\infty} \left[\prod_{i=1}^N \frac{d\lambda_i}{2\pi} \right] e^{-F_{\mathbb{S}_\omega^5}(\lambda_i)}, \quad (2.29)$$

where

$$\begin{aligned} F_{\mathbb{S}_\omega^5} &= N \log 2 + \log N! - N \log S'_3(0|\vec{\omega}) + (N-1) \log S_3 \left(im_a + \frac{\omega_{\text{tot}}}{2} \middle| \vec{\omega} \right) \\ &+ \frac{1}{\omega_1 \omega_2 \omega_3} \frac{4\pi^3}{g_{\text{YM}}^2} \sum_{i=1}^N \lambda_i^2 - \sum_{i>j}^N \log S_3 (i[\pm\lambda_i \pm \lambda_j] | \vec{\omega}) - \sum_{i=1}^N \log S_3 (\pm 2i\lambda_i | \vec{\omega}) \\ &+ \sum_{i>j}^N \log S_3 \left(i[\pm\lambda_i \pm \lambda_j] + im_a + \frac{\omega_{\text{tot}}}{2} \middle| \vec{\omega} \right) + N_f \sum_{i=1}^N \log S_3 \left(\pm i\lambda_i + im_f + \frac{\omega_{\text{tot}}}{2} \middle| \vec{\omega} \right), \end{aligned} \quad (2.30)$$

with $S_3(\lambda|\vec{\omega})$ being the triple sine function. Here, m_a and m_f are the masses for the hypermultiplets in the antisymmetric and fundamental representations of $\text{USp}(2N)$, respectively. We also introduced the notation

$$\omega_{\text{tot}} \equiv \omega_1 + \omega_2 + \omega_3, \quad S_3(\pm z|\vec{\omega}) \equiv S_3(z|\vec{\omega})S_3(-z|\vec{\omega}). \quad (2.31)$$

Large N free energy. We may restrict to $\lambda_i \geq 0$ due to the Weyl reflections of the $\text{USp}(2N)$ group. Consider the following ansatz for the large N saddle point eigenvalue distribution,

$$\lambda_j = N^\alpha t_j, \quad (2.32)$$

where $\alpha \in (0, 1)$ will be determined later. As in the previous section, at large N , we define the continuous function $t_j = t(j/N)$ and we introduce the density of eigenvalues $\rho(t)$, see (2.21). In the large N limit, $\lambda_i = \mathcal{O}(N^{1/2})$ (see (2.32) with $\alpha = 1/2$). Therefore, at large N , the contributions with nontrivial instanton numbers are exponentially suppressed. In the continuum limit, the free energy (2.30) is given by [40]¹²

$$\begin{aligned} F_{\mathbb{S}_\omega^5}[\rho(t), m_a|\vec{\omega}] &= \frac{N^{1+3\alpha}}{\omega_1 \omega_2 \omega_3} \frac{\pi(8-N_f)}{3} \int_0^{t_*} dt \rho(t) |t|^3 - \mu \left(\int_0^{t_*} dt \rho(t) - 1 \right) \\ &- \frac{N^{2+\alpha}}{\omega_1 \omega_2 \omega_3} \frac{\pi(\omega_{\text{tot}}^2 + 4m_a^2)}{8} \int_0^{t_*} dt \rho(t) \int_0^{t_*} dt' \rho(t') [t+t'+|t-t'|], \end{aligned} \quad (2.33)$$

where we added the Lagrange multiplier μ for the normalization of $\rho(t)$. In order to have a consistent saddle point α acquires the value $1/2$, and thus $F_{\mathbb{S}_\omega^5} \propto N^{5/2}$. Setting to zero the variation of (2.33) with respect to $\rho(t)$ we find the following saddle point configuration

$$\begin{aligned} \rho(t) &= \frac{2|t|}{t_*}, \quad t_* = \frac{1}{\sqrt{2}\sqrt{8-N_f}} (\omega_{\text{tot}}^2 + 4m_a^2)^{1/2}, \\ \mu &= -\frac{\pi}{3\sqrt{2}\omega_1 \omega_2 \omega_3} \frac{N^{5/2}}{\sqrt{8-N_f}} (\omega_{\text{tot}}^2 + 4m_a^2)^{3/2}. \end{aligned} \quad (2.34)$$

¹¹We will neglect instanton contributions as they are exponentially suppressed in the large N limit.

¹²Notice, that the free energy at large N does *not* depend on the masses of the N_f fundamental hypermultiplets. As it was shown in [40, (3.22)] their contribution to the large N free energy is of order $\mathcal{O}(N^{3/2})$ and, thus, subleading.

Plugging this back into (2.33) we obtain the squashed \mathbb{S}^5 free energy of the $\text{USp}(2N)$ theory, that reads (cf. [40, (3.38)])¹³

$$F_{\mathbb{S}^5_\omega}(m_a|\vec{\omega}) = \frac{2}{5}\mu = -\frac{\pi\sqrt{2}}{15\omega_1\omega_2\omega_3} \frac{N^{5/2}}{\sqrt{8-N_f}} (\omega_{\text{tot}}^2 + 4m_a^2)^{3/2}. \quad (2.35)$$

Introducing the redundant but *democratic* parameterization

$$\Delta_1 = 1 + \frac{2i}{\omega_{\text{tot}}}m_a, \quad \Delta_2 = 1 - \frac{2i}{\omega_{\text{tot}}}m_a, \quad (2.36)$$

(2.35) can be rewritten as

$$F_{\mathbb{S}^5_\omega}(\Delta_i|\vec{\omega}) = -\frac{\sqrt{2}\pi}{15} \frac{\omega_{\text{tot}}^3}{\omega_1\omega_2\omega_3} \frac{N^{5/2}}{\sqrt{8-N_f}} (\Delta_1\Delta_2)^{3/2}, \quad \Delta_1 + \Delta_2 = 2. \quad (2.37)$$

Finally, setting $\Delta_{1,2} = 1$ and $\omega_{1,2,3} = 1$, we find the round \mathbb{S}^5 free energy [48]

$$F_{\mathbb{S}^5} = -\frac{9\sqrt{2}\pi}{5} \frac{N^{5/2}}{\sqrt{8-N_f}}. \quad (2.38)$$

3 Four-dimensional solutions from the *stu* model

We treat here the four-dimensional gravitational backgrounds used to compute the holographic supersymmetric Rényi entropy. This section is organized as follows: before delving into the more intricate matter coupled solutions, we start by reviewing the simple case of the minimal supergravity BPS hyperbolic Reissner-Nordström and its SRE computation as done in [11, 12]. After this, in 3.2 we first recall the basic features of four-dimensional abelian Fayet-Iliopoulos (FI) gauged supergravity and present the hyperbolic matter coupled black hole solutions which first appeared in [49], leaving the details of the supergravity formalism and the BPS equations to appendix A. In 3.3, we compute the renormalized on-shell action and compare the result with the field theory computation in subsection 3.4, making contact with the minimal case as well. The complete procedure of holographic renormalization is spelled out in appendix B.

3.1 Warm up: BPS hyperbolic Reissner-Nordström

The computation of the SRE for hyperbolic solutions of $\mathcal{N} = 2$ minimal gauged supergravity was treated in [11, 12]. The gravity configurations are solutions to the equations of motion of the bosonic action

$$S = \int d^4x \sqrt{g} \left(R - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{6}{l_{\text{AdS}}} \right), \quad (3.1)$$

and read

$$ds^2 = -\left(\frac{r^2}{l_{\text{AdS}}^2} - 1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \frac{dr^2}{\left(\frac{r^2}{l_{\text{AdS}}^2} - 1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)} + r^2(d\theta^2 + \sinh^2(\theta)d\phi^2), \quad (3.2)$$

¹³The first equality arises from a virial theorem for the free energy (2.33).

with gauge field $A_t = \frac{Q}{r}dt + cdt$. c is a gauge term to be fixed later, in such a way that the gauge field is zero at the horizon r_+ , where g_{tt} vanishes, $g_{tt}(r_+) = 0$. In order for the solution to preserve 1/2 of the supersymmetries, the relation $Q = iM$ should hold. In other words, the charges of the solution should be purely imaginary. As we elaborate later on, this is not a problem because our aim is to study an analytically continued solution in Euclidean signature, obtained by $t \rightarrow -i\tau$, where the metric nevertheless remains real. With a slight abuse of terminology, consistent with the literature, we will continue referring to these solutions as “topological” or hyperbolic black holes. We set for simplicity $l_{\text{AdS}} = 1$.

First of all, imposing the BPS relation $M = -iQ$ and the fact that $g_{tt}(r_+) = 0$ we have that

$$Q = ir_+(1 \pm r_+). \quad (3.3)$$

The Wick rotated solution is characterized by a temperature T , found as the inverse periodicity of the τ coordinate, once we impose that the metric caps off smoothly at r_+ . Indeed, for $r \rightarrow r_+$ the metric, upon changing coordinates to $R = \sqrt{\frac{2(r-r_+)}{2r_+-1}}$, approaches

$$ds^2 = dR^2 + R^2 d\tau^2 (2r_+ - 1) + r_+^2 (d\theta^2 + \sinh^2(\theta) d\phi^2). \quad (3.4)$$

Therefore, the periodicity of the τ coordinate should be $\beta \equiv \Delta\tau = \frac{2\pi}{2r_+-1}$. The temperature¹⁴ is the inverse of this period:

$$T = \frac{2r_+ - 1}{2\pi}. \quad (3.5)$$

In order for the gauge field not to be singular at the horizon

$$A(r_+) = \frac{Q}{r_+} dt + c dt = 0, \quad (3.6)$$

we set $c = -\frac{Q}{r_+}$. We define the chemical potential ϕ as the asymptotic value of the gauge field, therefore $\phi \equiv \lim_{r \rightarrow \infty} A_t = c$.

To find the SRE, one identifies T with T_0/n , where T_0 is the temperature of the neutral black hole and n is the replica parameter. In this way,

$$T = \frac{1}{2\pi n}. \quad (3.7)$$

Combining (3.5) and (3.7), we can extract the value of r_+ as a function of the replica parameter n :

$$r_+ = \frac{n \mp 1}{2n}. \quad (3.8)$$

We choose the lower branch since, for $n = 1$, r_+ should go to unity. Similar reasoning makes us choose the lower sign in (3.3). The expression for the free energy found in [11, 12] reads

$$I = \frac{\text{Vol}(\mathbb{H}^2)\beta}{8\pi G_4} \left(-r_+^3 + iQ - \frac{Q^2}{r_+} \right), \quad (3.9)$$

¹⁴Once again this we denote this as “temperature of the black hole” but indeed we stress that its meaning comes from the Euclidean solution.

which, upon using (3.3), (3.8) becomes

$$I = \frac{\text{Vol}(\mathbb{H}^2)\beta}{8\pi G_4} \frac{(n+1)^2}{n^2} = \frac{\pi}{8G_4} \frac{(n+1)^2}{n}. \quad (3.10)$$

This matches the branched sphere partition function on the field theory side [11, 12], upon setting $\Delta_I = 1/2$, $I = 1, \dots, 4$, in (2.28) and using the standard AdS₄/CFT₃ relation $\frac{1}{G_4} = \frac{2\sqrt{2}}{3} N^{3/2}$ and the regularized volume $\text{Vol}(\mathbb{H}^2) = -2\pi$ [11].

Finally, we notice that the chemical potential takes the form

$$\phi = -\frac{Q}{r_+} = -i(1 - r_+) = -i\frac{n-1}{2n}, \quad (3.11)$$

matching the value of the R-symmetry background field (2.9). We record this expression as it will be useful later on in the computation of the SRE in the matter coupled case.

3.2 Hyperbolic black hole solutions of the *stu* model

The AdS₄ black holes with hyperbolic horizon we are after are solutions to abelian FI gauged supergravity in four spacetime dimensions. U(1) FI gauged supergravity arises as a truncation to the Cartan subalgebra, U(1)⁴, of $\mathcal{N} = 8$ gauged supergravity. The model thus obtained, called the *stu* model, corresponds to the prepotential

$$F(X) = -2i\sqrt{X^0 X^1 X^2 X^3}, \quad (3.12)$$

in the standard notation of $\mathcal{N} = 2$ supergravity. We will deal with a purely electric solution that has a hyperbolic horizon, supported by purely real scalars. In the BPS limit, the solution correspond to a 1/2 BPS black hole, preserving 4 out of the original 8 supercharges.

Spherical black holes of this model were constructed in [50, 51], and later elaborated upon in [52]. The hyperbolic solution, along with its uplift to eleven dimensions, first appeared in [49]. It is a static black hole characterized by the following metric

$$ds^2 = -\frac{U(r)}{4} dt^2 + \frac{dr^2}{U(r)} + h^2(r)(d\theta^2 + \sinh^2(\theta)d\phi^2), \quad (3.13)$$

with

$$U(r) = \frac{1}{\sqrt{\mathcal{H}}} f(r), \quad f(r) = -1 - \frac{\mu}{r} + 4g^2 r^2 \mathcal{H}, \quad h^2(r) = \sqrt{\mathcal{H}} r^2, \quad (3.14)$$

and

$$\mathcal{H} = H_1 H_2 H_3 H_4, \quad H_I = 1 + \frac{b_I}{r}, \quad I = 1, \dots, 4. \quad (3.15)$$

We set $g = 1$ from now on, and notice that we have rescaled time to match the asymptotic geometry (2.8). The non-vanishing components of the vector fields supporting the configurations are

$$A^I = \frac{1}{2} \left(1 - \frac{1}{H_I} \right) \frac{q_I}{b_I} dt + c^I dt, \quad (3.16)$$

where we have included four constant parameters c^I (to be determined later) which are required so that the gauge fields are non-singular at the horizon.

The equations of motion are satisfied if the parameters satisfy the following relation:

$$b_I = \mu \sin^2(\zeta_I), \quad q_I = \mu \sin(\zeta_I) \cos(\zeta_I). \quad (3.17)$$

Uppercase indices I, J run from 1 to 4, while lowercase ones i, j run from 1 to 3. The magnetic charges are set to zero, hence this is a purely electric configuration. The scalar fields z^i are *real* and parameterized by the holomorphic sections X^i , $z^i = X^i/X^0$. They assume the form [50]

$$z^1 = \frac{H_1 H_2}{H_3 H_4}, \quad z^2 = \frac{H_1 H_3}{H_2 H_4}, \quad z^3 = \frac{H_1 H_4}{H_2 H_3}. \quad (3.18)$$

The uplift of the solution to eleven-dimensional supergravity was performed in [49], where the solution was interpreted as the decoupling limit of spinning M2-branes. The BPS branch, which provides the solutions of interest here, is obtained by setting $\mu = 0$ and by taking

$$q_I = i b_I. \quad (3.19)$$

This configuration solves the BPS equations, as shown in appendix A.1. Notice that the electric charge assumes a purely imaginary value, as it did in the minimal case studied in [11, 12]. This is not a problem, as our aim is to study an analytically continued solution preserving supersymmetry. For this purpose, it is legitimate to take some parameters to be genuinely complex, since the Killing spinor equation, being analytic in the supergravity fields, will still admit a solution in the complexified background. Nevertheless, the Euclideanized metric in this case will remain purely real. It would be desirable to find a suitable solution directly in Euclidean supergravity coupled to matter multiplets, however in the following we will content ourselves with (a Wick-rotated version of) the Lorentzian solutions at hand.

The hyperbolic Reissner-Nordström solution discussed in the previous subsection is recovered from our setup upon taking the scalars to be constant

$$H_1 = H_2 = H_3 = H_4 = H, \quad z^i = 1, \quad i = 1, 2, 3, \quad (3.20)$$

taking all the gauge fields equal, and redefining the stu fields A^I (see [49, (3.15)]) as $A^I = A/2$. By doing so, the number of independent electric charges reduces to one, that of the graviphoton A .

3.3 Holographic supersymmetric Rényi entropy

From the stu black hole at our disposal, we can compute the temperature (see footnote 14)

$$T = \frac{1}{4\pi} \left. \frac{dU}{dr} \right|_{r_+}, \quad (3.21)$$

which turns out to be

$$T = \frac{(r_+^3(b_3 + b_4 + 2r_+) - b_1(b_2 b_3(2b_4 + r_+) + b_2 b_4 r_+ + b_3 b_4 r_+ - r_+^3) + b_2(r_+^3 - b_3 b_4 r_+))}{2\pi r_+ \sqrt{b_1 + r_+} \sqrt{b_2 + r_+} \sqrt{b_3 + r_+} \sqrt{b_4 + r_+}}. \quad (3.22)$$

Here, r_+ is the location of the horizon, obtained by requiring $U(r_+) = 0$. We leave the quantity r_+ implicit for the moment: trying to solve for r_+ from the vanishing of the warp factor yields a quartic equation whose explicit expression is quite cumbersome to manipulate.

Consider the uncharged black hole $q_1 = q_2 = q_3 = q_4 = 0$. In this case, the requirement $U(r_+) = 0$ gives $4r_+^2 - 1 = 0$, hence r_+ takes the simple form

$$r_+ = \frac{1}{2}. \tag{3.23}$$

Denoting by T_0 the temperature of the uncharged black hole, we have

$$T_0 = \frac{1}{2\pi}, \tag{3.24}$$

which will be useful later when defining the supersymmetric Rényi entropy.

In order for the gauge field to be non-singular at the horizon, we require $A^I(r_+) = 0$. Given the expression (3.16), this leads to

$$c^I = -\frac{i}{2} \left(1 - \frac{1}{H_I(r_+)} \right), \quad I = 1, \dots, 4. \tag{3.25}$$

The chemical potentials ϕ_I are defined as the asymptotic values of the gauge fields. They assume the form (we do not distinguish here between upper and lower indices on the chemical potentials)

$$\phi_I = c^I = -\frac{i}{2} \frac{b_I}{b_I + r_+}, \quad I = 1, \dots, 4. \tag{3.26}$$

By inserting (3.26) into (3.22), we can express the temperature as a function of the chemical potentials in the following way:

$$T = \frac{-i(\phi_1 + \phi_2 + \phi_3 + \phi_4) + 1}{\pi(\sqrt{1 - 2i\phi_1}\sqrt{1 - 2i\phi_2}\sqrt{1 - 2i\phi_3}\sqrt{1 - 2i\phi_4})} r_+, \tag{3.27}$$

where we have once again left r_+ implicit. We also point out that the quantities ϕ_I are *imaginary*, therefore T is *real*, as it should be. At this point, we can define

$$T = \frac{T_0}{n} = \frac{1}{2\pi n}. \tag{3.28}$$

Solving this equation for r_+ we obtain

$$r_+ = \frac{1}{2n} \frac{\sqrt{1 - 2i\phi_1}\sqrt{1 - 2i\phi_2}\sqrt{1 - 2i\phi_3}\sqrt{1 - 2i\phi_4}}{1 - i(\phi_1 + \phi_2 + \phi_3 + \phi_4)}. \tag{3.29}$$

Additionally, we know that the quantity r_+ must satisfy the relation $U(r_+) = 0$. Inserting the definitions (3.26) into $U(r_+) = 0$ yields the condition

$$1 + n^2(\phi_1 + \phi_2 + \phi_3 + \phi_4 + i)^2 = 0, \tag{3.30}$$

which is solved by

$$\phi_1 + \phi_2 + \phi_3 + \phi_4 = \frac{i(1 \pm n)}{n}. \tag{3.31}$$

We choose the lower sign since for $n = 1$ we should have zero chemical potential. As we will see in a moment, the choice of the upper branch translates in the dual field theory to a constraint on the value of the R-symmetry background field. To recapitulate, at this point we have obtained the expression (3.29) for r_+ in terms of the chemical potentials and the Rényi parameter n , supplemented by the constraint (3.31).

The renormalized on-shell action I is computed by adapting the procedure of [53] to the case of hyperbolic horizons. The computation, reported in appendix B, is tedious and not particularly illuminating. In the end, the thermodynamical potential reads

$$I = \frac{\beta \text{Vol}(\mathbb{H}^2)}{8\pi c G_4} \left(-\frac{\mu}{2} + r_+ \right), \tag{3.32}$$

where $\text{Vol}(\mathbb{H}^2)$ is the (regularized) volume of \mathbb{H}^2 , and $\beta = 1/T$ is the period of the Euclidean time direction. For the BPS case $\mu = 0$, we have

$$I = \frac{\beta \text{Vol}(\mathbb{H}^2) r_+}{8\pi c G_4} = \frac{2\pi \text{Vol}(\mathbb{H}^2)}{8\pi c G_4} \left(\frac{i\sqrt{1-2i\phi_1}\sqrt{1-2i\phi_2}\sqrt{1-2i\phi_3}\sqrt{1-2i\phi_4}}{2(i+\phi_1+\phi_2+\phi_3+\phi_4)} \right). \tag{3.33}$$

This expression is useful when comparing with the field theory result Z_n (2.28). The free energy of the black hole is given by

$$I = -\log Z(\phi_I, T). \tag{3.34}$$

The state variables are computed according to

$$E = \left(\frac{\partial I}{\partial \beta} \right)_\phi - \frac{\phi_I}{\beta} \left(\frac{\partial I}{\partial \phi_I} \right)_\beta, \quad S_{\text{BH}} = \beta \left(\frac{\partial I}{\partial \beta} \right)_\phi - I, \quad Q_I = -\frac{1}{\beta} \left(\frac{\partial I}{\partial \phi_I} \right)_\beta. \tag{3.35}$$

The renormalized on-shell action, (3.33), is computed in the grand canonical ensemble. In this ensemble, the Gibbs potential W is given by (see appendix B)

$$W = \frac{I}{\beta} = E - T S_{\text{BH}} - \phi^I Q_I + \Lambda \left(\phi_1 + \phi_2 + \phi_3 + \phi_4 - i \frac{(1-n)}{n} \right), \tag{3.36}$$

where Q_I are the electric charges of the black hole, and we inserted the Lagrange multiplier Λ which enforces the constraint (3.31) among the chemical potentials.

3.4 Holographic matching

In this section, we perform the holographic matching. The asymptotic value of the four-dimensional bulk gauge fields is related to the dual field theory flavor symmetry connection, defined in section 2.1.4, as

$$A_{\text{bulk}}^I(r \rightarrow \infty) = \phi_I dt = (A^{\text{flavor}, I}(\mathbb{S}_n^3) + A_{\text{bulk}}^{(R)}) d\tau, \tag{3.37}$$

where we have used $t = -i\tau$. To preserve supersymmetry, the background R-symmetry gauge field must have the form (2.9). The background R-symmetry gauge field is identified

with the chemical potential related to the R-symmetry gauge field in supergravity, which is the diagonal combination¹⁵

$$A_{\text{bulk}}^{(R)}(r \rightarrow \infty) = \frac{1}{4}(\phi_1 + \phi_2 + \phi_3 + \phi_4) dt = i \frac{1-n}{4n} dt = \frac{1-n}{4n} d\tau, \quad (3.38)$$

that appears in the supercovariant derivative of the spinor parameter in the susy variations (A.1). Notice that $A_{\text{bulk}}^{(R)} = \frac{1}{2}A^{(R)}$. As a simple consistency check, (3.38) is precisely the relation (3.31) obtained previously.

We are now ready to make contact with the field theory. The bulk fields correspond to the holonomies, shifted by the amount $(1-n)/(4n)$ due to the R-symmetry connection. In other words, we use the mapping (2.13) between the holonomies $A^{\text{flavor},I}$ and the parameters Δ_I , supplemented by the shift due to the R-symmetry:

$$A^I = A^{\text{flavor},I} + A_{\text{bulk}}^{(R)} = \left(\Delta_I - \frac{1}{2}\right) \left(\frac{n+1}{2n}\right) + \frac{1-n}{4n} = \left(\frac{(1+n)\Delta_I}{2n} - \frac{1}{2}\right). \quad (3.39)$$

Thus, we have

$$\phi_I = i \left(\frac{(1+n)\Delta_I}{2n} - \frac{1}{2}\right), \quad I = 1, \dots, 4. \quad (3.40)$$

Taking the sum of the l.h.s. and the r.h.s. we obtain the constraint

$$\frac{n-1}{n} = 2 - \frac{n+1}{2n} \sum_I \Delta_I \quad \Rightarrow \quad \sum_I \Delta_I = 2, \quad (3.41)$$

which reproduces the usual constraint on the parameters Δ_I . We use the standard relation

$$\frac{l_{\text{AdS}}^2}{G_4} = \frac{2\sqrt{2}}{3} N^{3/2}, \quad (3.42)$$

where we have taken into account $l_{\text{AdS}}^2 = 1/4$ from (3.14). Inserting (3.40) into (3.33), with $\mathbf{c} = 2$, the expression of the free energy becomes

$$I = -\frac{\sqrt{2}\pi N^{3/2} (n+1)^2}{3n} \sqrt{\Delta_1 \Delta_2 \Delta_3 \Delta_4} = -\log Z_{\mathbb{S}_n^3}, \quad \sum_{I=1}^4 \Delta_I = 2, \quad (3.43)$$

exactly matching the field theory computation (2.28) upon identifying $b \equiv 1/\sqrt{n}$, see (2.13). Note that we have defined the regularized volume as $\text{Vol}(\mathbb{H}^2) = -2\pi$ as in [11]. One easily sees that at the conformal point, $\Delta_I = 1/2$, which corresponds to the minimal supergravity case, the on-shell action reduces as expected to the one found in [11, 12].

We are now going to compute the supersymmetric Rényi entropy. First, notice that the partition function on the field theory side, see (2.28), satisfies

$$\log Z_{\mathbb{S}_n^3} = \frac{(n+1)^2}{4n} \log Z_{\mathbb{S}^3}. \quad (3.44)$$

¹⁵Note that the factor of 1/2 between (3.38) and (2.9) is due to the fact that the gauge fields in the *stu* model are defined with a factor of 1/2 with respect to the graviphoton in minimal supergravity [49].

The supersymmetric Rényi entropy is defined as (2.11)

$$S_n^{\text{SRE}} = \frac{n \log Z_{\mathbb{S}^3} - \log Z_{\mathbb{S}_n^3}}{n - 1}. \quad (3.45)$$

Therefore, we have

$$S_n = \frac{3n + 1}{4n} S_1, \quad S_1 = \log Z_{\mathbb{S}^3}, \quad (3.46)$$

as expected.

4 Six-dimensional hyperbolic solutions

We introduce here the six-dimensional hyperbolic solutions necessary for the holographic computation of the supersymmetric Rényi entropy. We first give some details regarding six-dimensional Romans $F(4)$ gauged supergravity coupled to one vector multiplet. We then present the hyperbolic black hole solutions coupled to matter, which have not previously appeared in the literature. In 4.3, we compute the holographic Rényi entropy, using the result of appendix B, and show the matching with the field theory computation in section 2.3.

4.1 Romans $F(4)$ gauged supergravity coupled to matter

In what follows, we will consider the six-dimensional $F(4)$ gauged supergravity coupled to one vector multiplet. Relevant references for this theory are [54, 55]. While the massive type IIA supergravity origin of this theory as a truncation of the supersymmetric warped $\text{AdS}_6 \times \mathbb{S}^4$ solution has not been established, there is evidence for it based on previous holographic matchings, see for instance [56, 57]. Taking the pragmatic approach of these latter papers, we work out supersymmetric solutions and proceed with the comparison of our result with its field theory counterpart. The five-dimensional SCFT dual to the warped $\text{AdS}_6 \times \mathbb{S}^4$ background is the one described in section 2.3. Solutions relevant for the supersymmetric Rényi entropy computation in the minimal theory (no vector multiplets) [58] were studied in [6, 13]. The non-minimal case is characterized by the presence of an additional flavor symmetry.

The bosonic fields of the six-dimensional Romans supergravity theory [58] consist of the metric $g_{\mu\nu}$, a scalar field X , a two-form potential $B_{\mu\nu}$, a one-form potential A , and an $\text{SU}(2)$ gauge field A^j with $j = 1, 2, 3$. In addition, there are fermionic fields comprising a pair of gravitini ψ_μ^A , $A = 1, 2$ and one spin 1/2 fermion χ^A . The vector multiplets consist of one gauge field A_μ , four scalar fields ϕ_α , with $\alpha = 0, 1, 2, 3$, and one gaugino λ_A . The scalar fields parameterize the coset space $\frac{\text{SO}(4,1)}{\text{SO}(4)}$. For additional details on the model, we refer the reader to [56, 57].

In finding the solution, we may take the Romans supergravity solution as example. In this solution, only one of the components of the $\text{SU}(2)$ gauge field, which we take to be A^3 [13], is nonzero. This gauge field is purely electric, meaning that the only nonzero component of the field strength is F_{rt} . This allows us to set the two-form potential $B_{\mu\nu}$

to zero, as there is no source for it.¹⁶ In our setup with an additional vector multiplet, we will still require the B field to vanish. Moreover, as in [57], we require the scalar fields in the vector multiplet ϕ_α to be neutral under A^3 . This restricts the nonzero components to ϕ_0 and ϕ_3 . We are further able to find a solution with only ϕ_3 turned on, namely $\phi_0 = 0$. Thus, we are left with the bosonic content: the metric, two gauge fields, the dilaton X , and the scalar field ϕ_0 .

4.2 Six-dimensional supersymmetric hyperbolic black holes

For the non-minimal case, we adapt the solutions of [60, section 3.2] to the \mathbb{H}^4 horizon topology. The solution is a static black hole characterized by the following metric

$$ds^2 = -U(r)dt^2 + \frac{dr^2}{V(r)} + h(r)ds_{\mathbb{H}^4}^2, \quad (4.1)$$

with $ds_{\mathbb{H}^4}^2$ the area element of four-dimensional hyperbolic space

$$ds_{\mathbb{H}^4}^2 = d\chi^2 + \sinh(\chi)^2 (d\theta^2 + \sin^2(\theta)d\psi^2 + \sin^2(\theta)\sin^2(\chi)d\phi^2), \quad (4.2)$$

and

$$U(r) = \frac{9}{2} \frac{f(r)}{\mathcal{H}^{3/4}}, \quad V(r) = \frac{f(r)}{\mathcal{H}^{1/4}} \quad h(r) = \mathcal{H}^{1/4} r^2, \quad (4.3)$$

with¹⁷

$$f(r) = -1 - \frac{\mu}{r^3} + \frac{2}{9} r^2 \mathcal{H} \quad \mathcal{H} = H_1 H_2, \quad H_I = 1 + \frac{b_I}{r^3}. \quad (4.4)$$

Here, $I = 1, 2$. The vector fields supporting the configuration read

$$A_t^I = \frac{3}{2} \left(1 - \frac{1}{H_I} \right) \frac{q_I}{b_I} - c^I dt, \quad I = 1, 2, \quad (4.5)$$

with parameters

$$b_I = \mu \sin^2(\xi_I), \quad q_I = \mu \sin(\xi_I) \cos(\xi_I), \quad (4.6)$$

and the scalars, in the notation of [60] are given by

$$X_1 = H_1^{-5/8} H_2^{3/8}, \quad X_2 = H_1^{3/8} H_2^{-5/8}. \quad (4.7)$$

The configuration with spherical slicing first appeared in [60], and the solution presented here is its generalization to hyperbolic slicing. However, the origin of the original configuration as a solution of a supergravity theory was unclear. It is easy to verify that the configuration is a solution to the equations of motion of F(4) gauged supergravity coupled to one vector multiplet, which are reported in [62]. One first truncates the theory to the $U(1) \times U(1)$ sector, as was done in [63], obtaining the Lagrangian [62, (3.2)]. One can then see that the field φ_1 can be consistently set to zero. Moreover, since all the field strengths

¹⁶This is in contrast to the six-dimensional solutions of [57, 59] of the form $\text{AdS}_2 \times \Sigma_{g_1} \times \Sigma_{g_2}$, which realizes the partial topological twist on $\Sigma_{g_1} \times \Sigma_{g_2}$. In that case, there is magnetic flux on Σ_{g_1} and Σ_{g_2} . This creates a source for the $H_{\mu\nu}$ field, which needs to be canceled by a nonzero value of B , in order to have a solution with $H = 0$.

¹⁷As in [13], we have conveniently rescaled the time direction by a factor of $3/\sqrt{2}$ with respect to [61].

are electric, there is no source term for the field $B_{\mu\nu}$, hence the latter can be set to zero as well. The remaining fields in our solutions can be mapped to those in [57, 62] via¹⁸

$$\begin{aligned} F_1 &= dA_1 = F_3 - F_{i1}, & F_2 &= dA_2 = F_3 + F_{i1}, \\ X_1 &= e^{\sigma-\phi_3}, & X_2 &= e^{\sigma+\phi_3}. \end{aligned} \tag{4.8}$$

With this mapping, and once we impose the truncations described above, one can show that the equations of motion are solved. The gauging parameters g, m are set to $g = 3m$ and $m = 1/(3\sqrt{2})$, justifying the factor $2/9$ in the warp factor $f(r)$ in (4.3).

The BPS branch is obtained, as usual, by setting $\mu = 0$ and $q_I = ib_I$. The solution is 1/2 BPS, and its Killing spinor is explicitly constructed in A.2. These solutions, once a Wick rotation to Euclidean spacetime is performed and setting $b_1 = b_2$, reduce to those considered in [6, 13].

4.3 Supersymmetric Rényi entropy

As in the previous case, we start the procedure by computing the period of the Euclidean time circle, namely the temperature of the hyperbolically sliced black hole. Given the expression for the warp factor (4.3), we have

$$T = -\frac{(4b_1b_2 + b_1r_+^3 + b_2r_+^3 - 2r_+^6)}{6\sqrt{2}\pi r_+^2 \sqrt{b_1 + r_+^3} \sqrt{b_2 + r_+^3}}. \tag{4.9}$$

Once we impose that the gauge field vanishes at the black hole horizon, we introduce the chemical potentials ϕ_I , $I = 1, 2$, as the asymptotic value of the gauge fields (4.5). We obtain

$$\phi_I = -\frac{3}{2} \frac{q_I}{b_I + r_+^3} = -\frac{3}{2} \frac{ib_I}{b_I + r_+^3}, \quad I = 1, 2, \tag{4.10}$$

where in the second equality we have used the BPS relation $q_I = ib_I$. The temperature can then be rewritten as

$$T = \frac{1}{\sqrt{2}\pi} \frac{1 - i(\phi_1 + \phi_2)}{\sqrt{3 - 2i\phi_1}\sqrt{3 - 2i\phi_2}} r_+. \tag{4.11}$$

By equating $T = T_0/n = 1/(2\pi n)$, we obtain an expression for r_+ in terms of the chemical potentials and the Rényi parameter n :

$$r_+ = \frac{\sqrt{3 - 2i\phi_1}\sqrt{3 - 2i\phi_2}}{\sqrt{2}n(1 - i(\phi_1 + \phi_2))}, \tag{4.12}$$

taking into account once more that these quantities are related via

$$\phi_1 + \phi_2 = \frac{i(1 \pm n)}{n}. \tag{4.13}$$

As explained in the previous section, we choose the lower sign so that the configuration reduces to a neutral black hole for $n = 1$.

¹⁸The field we call ϕ_3 and F_{i1} coincides respectively with ϕ_2 and F_6 of [62].

The renormalized on-shell action can be computed easily (see appendix B) by imposing supersymmetry. Using $\mathbf{c} = \sqrt{2}/3$, we obtain

$$I = \frac{\beta \text{Vol}(\mathbb{H}^4)}{8\pi\mathbf{c}G_6} \left(-r_+^3 - \frac{\mu}{2}\right) = -\frac{3n}{4\sqrt{2}G_6} \text{Vol}(\mathbb{H}^4)r_+^3. \quad (4.14)$$

This is consistent with the result of [13], which is valid in the absence of vector multiplets. (4.14) combined with the previous expression, (4.12) for r_+ yields

$$I = \frac{\pi^2 n}{\sqrt{2}G_6} \left(\frac{\sqrt{3 - 2i\phi_1}\sqrt{3 - 2i\phi_2}}{\sqrt{2n(1 - i(\phi_1 + \phi_2))}} \right)^3, \quad (4.15)$$

supplemented by the constraint (4.13) between the chemical potentials. We have also used the normalized volume $\text{Vol}(\mathbb{H}^4) = 4\pi^2/3$ [13].

4.4 Holographic matching

We recall the expression that relates the asymptotic value of the bulk gauge field to the corresponding dual quantities:

$$A_{\text{bulk}}^I(r \rightarrow \infty) = \phi_I dt = (A^I(\mathbb{S}_n^5) + A_{\text{bulk}}^{(R)})d\tau. \quad (4.16)$$

Recall that, on the field theory side, the R-symmetry background gauge field has the expression (2.9). The corresponding chemical potential in the supergravity notation reads

$$A_{\text{bulk}}^{(R)} = \frac{\phi_1 + \phi_2}{2} dt = i \frac{1-n}{2n} dt = \frac{1-n}{2n} d\tau. \quad (4.17)$$

We are ready now to make contact with the field theory chemical potentials. Indeed, the bulk fields correspond to (2.14), which are related to Δ_I via (2.36), shifted by the amount $(1-n)/(2n)$ due to the R-symmetry connection, resulting in

$$A^I = (\Delta_I - 1) \left(\frac{2n+1}{2n} \right) + \frac{1-n}{2n} = \frac{3}{2} \left(\frac{(1+2n)\Delta_I}{3n} - 1 \right). \quad (4.18)$$

Therefore, we have

$$\phi_I = i \frac{3}{2} \left(\frac{\Delta_I(2n+1)}{3n} - 1 \right), \quad I = 1, 2. \quad (4.19)$$

Notice that taking the sum over the index I and using (4.13) we get the relation $\Delta_1 + \Delta_2 = 2$. Taking into account (4.19), noting that $l_{\text{AdS}}^2 = 9/2$, and using the relation [48]

$$\frac{l_{\text{AdS}}^4}{G_6} = \frac{27\sqrt{2}}{\sqrt{8-N_f}} \frac{N^{5/2}}{5\pi}, \quad (4.20)$$

the gravitational on-shell action in (4.15) yields exactly

$$I = \frac{\sqrt{2}\pi N^{5/2}}{15\sqrt{8-N_f}} \frac{(2n+1)^3}{n^2} (\Delta_1 \Delta_2)^{3/2}, \quad \Delta_1 + \Delta_2 = 2. \quad (4.21)$$

This perfectly agrees with the prediction from the field theory (2.37), once we set $\vec{\omega} = (1, 1, 1/n)$. In the absence of flavor symmetry (or masses), we obtain the result of the minimal case. Indeed, imposing $\Delta_1 = \Delta_2 = 1$ we retrieve the result of [6, 13], which reads

$$I = \frac{\sqrt{2}\pi (2n+1)^3 N^{5/2}}{15n^2 \sqrt{8-N_f}} = -\log Z_{\mathbb{S}_n^5}. \quad (4.22)$$

One can easily work out the value of S_n^{SRE} as

$$S_n^{\text{SRE}} = \frac{n \log Z_{\mathbb{S}^5} - \log Z_{\mathbb{S}_n^5}}{n-1} = \frac{19n^2 + 7n + 1}{27n^2} S_1, \quad S_1 = \log Z_{\mathbb{S}^5}. \quad (4.23)$$

5 Concluding remarks

Following the work on magnetically charged AdS₄ black holes in [64], intense efforts have been put into the holographic computation of entropy for BPS black holes with compact horizons, using localization (see [65, 66] and references within). Some of the computations involve a rather subtle treatment of the matrix integrals which compute the relevant SCFT partition function. For instance, progress has been made on the longstanding problem of computing the entropy of rotating BPS black holes in AdS₅ from the superconformal index of $\mathcal{N} = 4$ SYM using such a treatment [67]. Our computation is somewhat similar, the black holes in question having no magnetic flux, but does not involve the same subtleties. This may be due to the observation, made in [19], that the Killing spinors relevant to the computation in the bulk, and hence in the SCFT, should be anti-periodic in the Euclidean time direction. While this can be arranged for partition functions like the one used to compute the superconformal index [68], it arises naturally in the context of the SRE, i.e. the hyperbolic index, when viewed as a Weyl transformation of the branched sphere. This fact still awaits a satisfactory physical explanation.

Regarding possible future directions, it would be interesting to incorporate magnetic charges in the black hole background, and compare the resulting free energy with the corresponding field theory computation generalized by magnetic fluxes. Moreover, one could compute the subleading N corrections to the Supersymmetric Rényi and compare with the supergravity computation, along the lines of [69]. Finally, it would be interesting to investigate in our setup the expansion of the SRE around $n = 1$. In [5, 22, 70] it was found that the first correction to the entanglement entropy is proportional to the coefficient of the stress tensor vacuum two-point function, and it would be interesting to find the interpretation of this statement in the supergravity picture. We hope to come back to these points in the future.

Acknowledgments

We would like to thank Laura Andrianopoli, Davide Cassani, Martin Fluder, Márk Mezei, Ioannis Papadimitriou, Julian Sonner for discussions, Kiril Hristov, Tatzuma Nishioka and Alberto Zaffaroni for carefully reading a first draft of the manuscript. The work of SMH was supported by World Premier International Research Center Initiative (WPI Initiative),

MEXT, Japan. CT acknowledges support from the NSF Grant PHY-1125915 and the Agence Nationale de la Recherche (ANR) under the grant Black-dS-String (ANR-16-CE31-0004) and would like to thank the Simons Center for Geometry and Physics, Stony Brook University, Kavli IPMU and Universita' di Parma for hospitality during some steps of this paper. The work of IY was financially supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 754496 — FELLINI.

A Explicit construction of the Killing spinor

A.1 Four dimensions

To show the resolution of the Killing spinor equations (KSE) of four-dimensional abelian FI gauged supergravity in presence of vector multiplets we follow essentially the conventions of [51], where the Killing spinor for configurations with spherical spatial section was worked out (see also [50]). The modification to configurations with hyperbolic horizon is an easy task that we perform in what follows.

The supersymmetry transformation of the gravitini and gaugini in terms of complex spinors read

$$\begin{aligned}\delta\psi_\mu &= \nabla_\mu\varepsilon + \frac{i}{4}T_{\rho\sigma}^-\gamma^\rho\gamma^\sigma\gamma_\mu\varepsilon - \frac{g}{2}\xi_\Lambda L^\Lambda\gamma_\mu\varepsilon, \\ \delta_\varepsilon\lambda^i &= i\partial_\mu z^i\gamma^\mu\varepsilon + iG_{\mu\nu}^-\gamma^{\mu\nu}\varepsilon + gg^{i\bar{j}}\bar{f}_{\bar{j}}^\Lambda\xi_\Lambda\varepsilon.\end{aligned}\tag{A.1}$$

The supercovariant derivative of the gravitino appearing in the supersymmetry variations is given by

$$\nabla_\mu\varepsilon = \left(\partial_\mu - \frac{1}{4}\omega_\mu^{ab}\gamma_{ab}\right)\varepsilon + \frac{1}{4}(\mathcal{K}_i\partial_\mu z^i - \mathcal{K}_{\bar{i}}\partial_\mu\bar{z}^{\bar{i}})\varepsilon + ig\xi_\Lambda A_\mu^\Lambda\varepsilon,\tag{A.2}$$

and we have defined

$$T_{\mu\nu}^- = 2i\text{Im}\mathcal{N}_{\Lambda\Sigma}L^I F_{\mu\nu}^{J-}, \quad G_{\mu\nu}^i = -g^{i\bar{j}}\bar{f}_{\bar{j}}^\Lambda\text{Im}\mathcal{N}_{\Lambda\Sigma}F_{\mu\nu}^{\Sigma-},\tag{A.3}$$

where L^I are the upper part of the covariantly holomorphic section \mathcal{V}

$$\mathcal{V} = \begin{pmatrix} L^\Lambda \\ M_\Lambda \end{pmatrix} \equiv e^{\mathcal{K}/2}\Omega = e^{\mathcal{K}/2} \begin{pmatrix} X^\Lambda \\ F_\Lambda \end{pmatrix},\tag{A.4}$$

\mathcal{K} is the Kähler potential, and f_i^Λ are defined as

$$\begin{pmatrix} f_i^\Lambda \\ h_{\Sigma,i} \end{pmatrix} \equiv \nabla_i\mathcal{V} = \left(\partial_i + \frac{1}{2}\partial_i\mathcal{K}\right)\mathcal{V}.\tag{A.5}$$

Further definitions can be found for instance in [71]. Finally $[\gamma_a, \gamma_b]$ denotes the anti-symmetrized product with unit weight, i.e. $[\gamma_a, \gamma_b] = \frac{1}{2}(\gamma_a\gamma_b - \gamma_b\gamma_a)$. We set the Fayet-Iliopoulos parameters $\xi_\Lambda = 1$ for $\Lambda = 0, \dots, 3$. For the four-dimensional solution described in section 3.2 (see (3.13) and (3.16)), we choose the following vierbeins

$$\begin{aligned}e_t^0 &= \mathcal{H}(r)^{-1/4}\sqrt{f(r)}, & e_r^1 &= \frac{\mathcal{H}(r)^{1/4}}{\sqrt{f(r)}}, \\ e_\theta^2 &= r\mathcal{H}(r)^{1/4}, & e_\phi^4 &= r\mathcal{H}(r)^{1/4}\sinh(\theta),\end{aligned}\tag{A.6}$$

and the non-vanishing components of the spin connection are then

$$\omega_t^{01} = U'(r), \quad \omega_\theta^{12} = h'(r), \quad \omega_\phi^{13} = h'(r) \sinh(\theta), \quad \omega_\phi^{23} = \cosh(\theta). \quad (\text{A.7})$$

Assuming a Killing spinor that fulfills the following relation

$$\varepsilon = (a i \gamma_0 + b \gamma_1) \varepsilon, \quad (\text{A.8})$$

with

$$a = \frac{i}{\sqrt{f(r)}}, \quad b = -\frac{2gr}{\sqrt{f(r)}} \mathcal{H}^{1/2}, \quad (\text{A.9})$$

the supersymmetry equations (A.1) simplify considerably, and one obtains the following explicit solution for the Killing spinor

$$\varepsilon = \frac{1}{2\sqrt{gr}} e^{\frac{t}{2n}} \mathcal{H}^{-1/8} e^{-\frac{1}{2}\gamma_{012}\theta} e^{-\frac{1}{2}\gamma_{23}\phi} \left(\sqrt{f(r)} - i - i\gamma_1 \sqrt{f(r)} + i \right) (1 - \gamma_0) \varepsilon_0, \quad (\text{A.10})$$

where ε_0 is an arbitrary spinor in four dimensions. Notice that in the limit of constant scalars, namely $b_1 = b_2 = b_3 = b_4 = Q$ we recover the Killing spinor of [72].

A.2 Six dimensions

In this section we explicitly construct the Killing spinor from the BPS equations of six-dimensional F(4) Romans gauged supergravity coupled to one vector multiplet. We mostly follow the conventions of [57], briefly recapping only the quantities relevant in our case, and referring to that paper for the details we omitted here for brevity. The supersymmetry variations of the fermions are

$$\begin{aligned} \delta\psi_{A\mu} &= \nabla_\mu \varepsilon_A - \frac{1}{2} g \sigma_{AC}^x A_{x\mu} \varepsilon^C + \frac{1}{16} e^{-\sigma} [\tilde{T}_{[AB]\nu\lambda} \gamma_7 - T_{(AB)\nu\lambda}] (\gamma_\mu^{\nu\lambda} - 6\delta_\mu^\nu \gamma^\lambda) \varepsilon^B + S_{AC} \gamma^\mu \varepsilon^C \\ &\quad + \frac{i}{32} e^{2\sigma} H_{\nu\lambda\rho} \gamma_7 (\gamma_\mu^{\nu\lambda\rho} - 3\delta_\mu^\nu \gamma^{\lambda\rho}) \varepsilon_A, \\ \delta\lambda_A^I &= iP_{x\mu}^I \sigma_{AC}^x \gamma^\mu \varepsilon^C - iP_{0\nu}^I \varepsilon_{AC} \gamma^7 \gamma^\nu \varepsilon^C + \frac{i}{2} e^{-\sigma} T_{\nu\rho}^I \gamma^{\nu\rho} \varepsilon_A + M_{AB}^I \varepsilon^B, \\ \delta\chi_A &= \frac{i}{2} \gamma^\nu \partial_\nu \sigma \varepsilon_A + \frac{i}{16} e^{-\sigma} [\tilde{T}_{[AB]\nu\lambda} \gamma_7 - T_{(AB)\nu\lambda}] \gamma^{\nu\lambda} \varepsilon^B + \frac{1}{32} e^{2\sigma} H_{\nu\lambda\rho} \gamma_7 \gamma^{\nu\lambda\rho} \varepsilon_A + N_{AB} \varepsilon^B, \end{aligned} \quad (\text{A.11})$$

where the capital Greek indices are raised and lowered with the $\text{SO}(4, n_V)$ invariant metric and the indices A, B, \dots with the antisymmetric tensor ε_{AB} . The objects appearing in the susy equations are defined as

$$\tilde{T}_{[AB]\nu\lambda} = \varepsilon_{AB} L_{0\Sigma}^{-1} \hat{F}_{\nu\lambda}^\Sigma, \quad T_{(AB)\nu\lambda} = \sigma_{AB}^x L_{x\Sigma}^{-1} F_{\nu\lambda}^\Sigma, \quad T_{I\nu\lambda} = L_{I\Sigma}^{-1} \hat{F}_{\nu\lambda}^\Sigma, \quad (\text{A.12})$$

and the matrices N_{AB} , S_{AB} , M_{AB}^I , along with a convenient parameterization of the scalar coset L_Λ^Σ are defined in [57], to which we refer for all missing definitions. In our case they boil down to

$$\begin{aligned} N_{AB} &= \frac{1}{4} (g \cosh(\phi_3) e^\sigma - 3m e^{-3\sigma}) \varepsilon_{AB}, \\ S_{AB} &= \frac{i}{4} (g \cosh(\phi_3) e^\sigma + m e^{-3\sigma}) \varepsilon_{AB}, \\ M_{AB} &= -2g \sinh(\phi_3) e^\sigma \sigma_{AB}^3. \end{aligned} \quad (\text{A.13})$$

As in [13], also in our case the only component of the SU(2) gauge field is the one in the $i = 3$ direction. With reference to (4.3) we choose the following vielbeins

$$\begin{aligned}
 e_r^0 &= \frac{\mathcal{H}^{1/8}}{f(r)^{1/2}}, & e_t^1 &= \frac{3}{\sqrt{2}} \frac{f(r)^{1/2}}{\mathcal{H}^{3/8}}, & e_\chi^2 &= r\mathcal{H}^{1/8}, & e_\theta^3 &= r\mathcal{H}^{1/8} \sinh(\chi), \\
 e_\psi^4 &= r\mathcal{H}^{1/8} \sinh(\chi) \sin(\theta), & e_\phi^5 &= r\mathcal{H}^{1/8} \sinh(\chi) \sin(\theta) \sin(\chi),
 \end{aligned}
 \tag{A.14}$$

The non-vanishing components of the spin connection read

$$\begin{aligned}
 w_t^{01} &= \frac{U'(r)\sqrt{V(r)}}{2\sqrt{U(r)}}, & w_\chi^{02} &= -\frac{h'(r)\sqrt{V(r)}}{2\sqrt{h(r)}}, \\
 w_\theta^{03} &= -\frac{\sinh(\chi)h'(r)\sqrt{V(r)}}{2\sqrt{h(r)}}, & w_\theta^{23} &= -\cosh(\chi), \\
 w_\psi^{04} &= -\frac{\sinh(\chi)h'(r)\sqrt{V(r)}\sin\theta}{2\sqrt{h(r)}}, & w_\psi^{24} &= -\cosh(\chi)\sin(\theta), \\
 w_\psi^{34} &= -\cos(\theta), & w_\phi^{05} &= -\frac{\sinh(\chi)h'(r)\sqrt{V(r)}\sin(\theta)\sin(\psi)}{2\sqrt{h(r)}}, \\
 w_\phi^{25} &= -\cosh(\chi)\sin(\theta)\sin(\psi), & w_\phi^{35} &= -\cos(\theta)\sin(\psi), \\
 w_\phi^{45} &= -\cos(\psi).
 \end{aligned}
 \tag{A.15}$$

We are going to consider first the variation of the dilatino χ_A . Given our truncation, we can see that imposing the relation

$$(\delta_A^B + ix(r)\gamma^0\sigma_{AC}^x\epsilon^{CB} + y(r)\gamma^1\delta_A^B)\epsilon_B = 0,
 \tag{A.16}$$

with

$$\begin{aligned}
 x(r) &= -\frac{3ir^2}{\sqrt{2b_1(b_2+r^3)+2b_2r^3+2r^6-9r^4}}, \\
 y(r) &= -\frac{\sqrt{2}\sqrt{b_1+r^3}\sqrt{b_2+r^3}}{\sqrt{2b_1(b_2+r^3)+r^3(2b_2+2r^3-9r)}},
 \end{aligned}
 \tag{A.17}$$

where $x(r)^2 + y(r)^2 = 1$, the gravitino equation reduces to

$$\begin{aligned}
 \delta\psi_{A,t} &= \left(\partial_t - \frac{1}{2}(1 - i(\phi_1 + \phi_2))m_A^B\right)\epsilon_B = \left(\partial_t - \frac{1}{2n}m_A^B\right)\epsilon_B, \\
 \delta\psi_{A,r} &= (\partial_r + f_1(r)m_A^B\gamma^0 + f_2(r)\delta_A^B)\epsilon_B, \\
 \delta\psi_{A,\chi} &= \left(\partial_\chi - \frac{1}{2}\gamma_{012}m_A^B\right)\epsilon_B, \\
 \delta\psi_{A,\theta} &= \left(\partial_\theta - \frac{1}{2}\sinh(\chi)\gamma_{013}m_A^B - \frac{1}{2}\cosh(\chi)\gamma_{23}\right)\epsilon_B, \\
 \delta\psi_{A,\psi} &= \left(\partial_\psi - \frac{1}{2}\sinh(\chi)\sin(\theta)\gamma_{014}m_A^B - \frac{1}{2}\cosh(\chi)\sin(\theta)\gamma_{24} - \frac{1}{2}\cos\theta\gamma_{34}\right)\epsilon_B, \\
 \delta\psi_{A,\phi} &= \left(\partial_\phi - \frac{1}{2}\sin(\theta)\sinh(\chi)\sin(\psi)\gamma_{015}m_A^B + \frac{1}{2}\sin(\theta)\cosh(\chi)\sin(\psi)\gamma_{25} \right. \\
 &\quad \left. - \frac{1}{2}\cos(\theta)\sin(\psi)\gamma_{35} - \frac{1}{2}\cos(\psi)\gamma_{45}\right)\epsilon_B.
 \end{aligned}
 \tag{A.18}$$

Here, we defined the functions $f_1(r)$ and $f_2(r)$ as

$$\begin{aligned}
 f_1(r) &= \frac{9(b_1(2b_2 + r^3) + b_2r^3)}{16r(b_1 + r^3)(b_2 + r^3)}, \\
 f_2(r) &= \frac{(r^3(b_1 + b_2) + 4b_1b_2 - 2r^6)}{2\sqrt{2r}\sqrt{b_1 + r^3}\sqrt{b_2 + r^3}\sqrt{2r^3(b_1 + b_2) + 2b_1b_2 + 2r^6 - 9r^4}}.
 \end{aligned}
 \tag{A.19}$$

and we defined $m_A{}^B = \sigma_{AC}^x \epsilon^{CB}$. The $t, \chi, \theta, \psi, \phi$ equation can be solved immediately by the following [73]:

$$\varepsilon_A(r, t, \chi, \theta, \psi, \phi) = e^{\frac{t}{2n}m_A{}^B} e^{\frac{\chi}{2}\gamma_{012}m_A{}^B} e^{\frac{\theta}{2}\gamma_{23}} e^{\frac{\psi}{2}\gamma_{34}} e^{\frac{\phi}{2}\gamma_{45}} \varepsilon_A(r),
 \tag{A.20}$$

and the radial component of (A.18), together with the relation (A.16) can be solved by standard methods of [72], resulting in

$$\varepsilon_A(r) = (u(r) + v(r)\gamma^1)(\delta_{AB} - i\bar{\Gamma}_{AB})\varepsilon_{0,B}.
 \tag{A.21}$$

Here $\varepsilon_{0,B}$ is a doublet of constant spinors, $\bar{\Gamma}_{AB} = \gamma_0\sigma_{AC}^3\epsilon^{CB}$ and

$$\begin{aligned}
 u(r) &= \sqrt{\frac{1+x(r)}{y(r)}} e^{w(r)} = \sqrt{\frac{\sqrt{2b_1(b_2 + r^3) + r^3(2b_2 + 2r^3 - 9r)} + 3ir^2}{\sqrt{2}\sqrt{b_1 + r^3}\sqrt{b_2 + r^3}}} e^{w(r)}, \\
 v(r) &= -\sqrt{\frac{1-x(r)}{y(r)}} e^{w(r)} = -\sqrt{\frac{\sqrt{2b_1(b_2 + r^3) + r^3(2b_2 + 2r^3 - 9r)} - 3ir^2}{\sqrt{2}\sqrt{b_1 + r^3}\sqrt{b_2 + r^3}}} e^{w(r)},
 \end{aligned}
 \tag{A.22}$$

with

$$w(r) = \int^r f_1(r') dr' = \frac{9}{16} \left(-\frac{1}{3} \log(b_1 + r^3) - \frac{1}{3} \log(b_2 + r^3) + 2 \log(r) \right),
 \tag{A.23}$$

hence

$$e^{w(r)} = \frac{r^{9/8}}{(b_1 + r^3)^{3/16} (b_2 + r^3)^{3/16}}.
 \tag{A.24}$$

The total Killing spinor is then given by combining (A.20) with the radial dependent part in (A.21). Notice that the second bracket of (A.21) projects out half of the supersymmetries, which signals the fact that the solution indeed is 1/2 BPS. It is easy to check that this expression also solves the gaugino equation $\delta\lambda_A^I = 0$ in (A.11).

B On-shell action via holographic renormalization

In this section we compute the renormalized on-shell action in the grand-canonical ensemble for the solutions we described in the main sections. In [53] the on-shell action for the corresponding $\mathcal{N} = 2$ four-dimensional gauged supergravity and Romans $F(4)$ spherical solutions was computed. Generalizing the computation to hyperbolic horizons of different topology requires a minimal modification of their procedure, which we explain in this appendix, following their notation closely. Other relevant references for holographic renormalization in this context are for instance [17, 74, 75] and we will make use of them when deriving the counterterms. Notice that here d denotes the dimension of the boundary.

For both setups the action can be cast in the following form (see [60, (5.1)]):

$$S = -\frac{1}{16\pi G_{d+1}} \int_M d^{d+1}x \sqrt{-g} \left(R - \frac{1}{2} G_{ij} \partial_\mu z^i \partial^\mu z^j - \frac{1}{4} M_{IJ} F_{\mu\nu}^I F^{\mu\nu,J} - V(z^i) \right) + \frac{1}{8\pi G_{d+1}} \int_{\partial M} d^d x \sqrt{-h} \Theta, \quad (\text{B.1})$$

where M is a $(d+1)$ -dimensional spacetime with metric $g_{\mu\nu}$, boundary ∂M with induced metric $h_{\mu\nu}$. In this case Θ is the trace of the extrinsic curvature $\Theta_{\mu\nu}$ of the boundary $\Theta_{\mu\nu} = -\frac{1}{2}(\nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu)$, where ξ^μ is the outward-pointing normal to ∂M .

We can massage the bulk term of the action (B.1) by making use of the trace of the Einstein's equation, to obtain

$$I_{\text{bulk}} = -\frac{1}{16\pi G_{d+1}} \int_M d^{d+1}x \sqrt{-g} \left[-\frac{1}{2(d-1)} M_{IJ} F_{\mu\nu}^I F^{\mu\nu,J} + \frac{2}{d-1} V(z^i) \right]. \quad (\text{B.2})$$

This latter expression can be rewritten as [53]

$$I_{\text{bulk}} = -\frac{1}{8\pi G_{d+1}} \int_M d^{d+1}x \sqrt{-g} R_\phi^\phi. \quad (\text{B.3})$$

We use an ansatz for the metric of the form

$$ds^2 = -\frac{\mathcal{H}^{-(d-2)/(d-1)} f(r)}{c^2} dt^2 + \frac{\mathcal{H}^{1/(d-1)}}{f(r)} dr^2 + \mathcal{H}^{1/(d-1)} r^2 ds_{\mathbb{H}^{d-1}}^2, \quad (\text{B.4})$$

with

$$f(r) = -1 - \frac{\mu}{r^{d-2}} + \tilde{g} r^2 \mathcal{H}(r)^2, \quad \mathcal{H} = \prod_I H_I, \quad (\text{B.5})$$

which encompasses both the four-dimensional configurations of section 3.2 and those of section 4.2 for a suitable choice of \mathbf{c} and \tilde{g} . A direct computation of the term in (B.3), once we define $B(r) = \frac{1}{2(d-1)} \log \mathcal{H}(r)$, gives

$$\sqrt{-g} R_\phi^\phi = -\frac{1}{c} \frac{d}{dr} \left(B'(r) r^{d-1} f(r) + r^{d-2} (f(r) + 1) \right). \quad (\text{B.6})$$

This term differs from the spherical case treated in [53] by the sign of the last addendum. The bulk term therefore yields

$$I_{\text{bulk}} = \frac{\text{Vol}(\mathbb{H}^{d-1})\beta}{16\pi \mathbf{c} G_{d+1}} \left(B'(r_{\text{inf}}) r_{\text{inf}}^{d-1} f(r_{\text{inf}}) + r_{\text{inf}}^{d-2} (f(r_{\text{inf}}) + 1) - r_+^{d-2} \right), \quad (\text{B.7})$$

where we used that $f(r_+) = 0$. As for the Gibbons-Hawking term, the normal outward pointing is given by $n^r = \sqrt{f(r)} \mathcal{H}^{-1/(2(d-1))} = \sqrt{f(r)} e^{-B(r)}$. The extrinsic curvature reads

$$\Theta = -\frac{e^{-B(r)} (2f(r) (rB'(r) + d - 1) + r f'(r))}{2r \sqrt{f(r)}}, \quad (\text{B.8})$$

which yields a Gibbons-Hawking term of the form

$$I_{\text{GH}} = -\frac{\text{Vol}(\mathbb{H}^{d-1})\beta}{8\pi \mathbf{c} G_{d+1}} r_{\text{inf}}^{d-2} \left[f(r) (rB'(r) + d - 1) + \frac{1}{2} r f'(r) \right]_{r=r_{\text{inf}}}. \quad (\text{B.9})$$

Here we used

$$\sqrt{-h} = \frac{e^{B(r)}}{\mathbf{c}} r^{d-1} \sqrt{f(r)} v_{d-1}, \quad (\text{B.10})$$

where $v_2 = \sinh(\theta)$ and $v_4 = \sinh^3(\chi) \sin^2(\theta) \sin(\psi)$. This leads to

$$I_{\text{bulk}} + I_{\text{GH}} = \frac{\text{Vol}(\mathbb{H}^{d-1})\beta}{8\pi\mathbf{c}G_{d+1}} \left(-(d-2)r_{\text{inf}}^{d-2} f(r_{\text{inf}}) - \frac{1}{2}r_{\text{inf}}^{d-1} f'(r_{\text{inf}}) + r_{\text{inf}}^{d-2} - r_+^{d-2} \right). \quad (\text{B.11})$$

We will now spell out the relevant counterterms for the different cases, specializing to the different $d+1 = D = 4$ and $D = 6$ cases, dealing first with the former.

Four dimensions. The holographic renormalization procedure in $D = 4$ follows from [75], where the counterterms for $\mathcal{N} = 2$ U(1)-gauged supergravity coupled to three vector multiplets were derived. In particular our solution is purely electric hence the counterterms boil down to

$$I_{\text{ct}} = \frac{1}{8\pi G_4} \int d^3x \sqrt{-h} \left(\mathcal{W}(z^i) + \frac{1}{2\tilde{g}} \mathcal{R}_3 + \dots \right), \quad (\text{B.12})$$

where the ellipsis denotes the terms which are subleading once the cutoff is send to infinity. \mathcal{R} is the Ricci scalar of the boundary, and \mathcal{W} is a function of the scalar fields called superpotential. We have the following expression for the Ricci scalar of the boundary \mathcal{R}_3 :

$$\mathcal{R}_3 = -\frac{2}{r^2} e^{-2B(r)}, \quad (\text{B.13})$$

and the superpotential that drives the flow is given by [53]

$$\mathcal{W} = \frac{\tilde{g}}{2} \sum_{I=0}^3 X^I, \quad (\text{B.14})$$

which coincides with that used in [75–77] and amounts to imposing Neumann boundary conditions on the scalar fields, a procedure that is compatible with supersymmetry [78]. Adding this to the action (B.11) we finally find

$$I \equiv I_{\text{bulk}} + I_{\text{GH}} + I_{\text{ct}} = -\frac{\beta \text{Vol}(\mathbb{H}^2)}{8\pi\mathbf{c}G_4} \left(r_+ + \frac{\mu}{2} \right), \quad (\text{B.15})$$

which indeed reduces to $I = -\frac{\beta \text{Vol}(\mathbb{H}^2)}{8\pi\mathbf{c}G_4} r_+$ in the BPS limit.

Six dimensions. In $D = 6$ we have the following counterterms [53]:

$$I_{\text{ct}} = \frac{1}{8\pi G_d} \int d^3x \sqrt{-h} \left(\mathcal{W}(z^i) + \frac{1}{6g} \mathcal{R}_5 + \frac{1}{18g^3} \left(\mathcal{R}_{5,ab} \mathcal{R}_5^{ab} - \frac{5}{16} \mathcal{R}_5^2 \right) \right), \quad (\text{B.16})$$

where \mathcal{R}_5 and $\mathcal{R}_{5,ab}$ are respectively the Ricci scalar and the Ricci tensor of the five-dimensional boundary metric.¹⁹ Notice that the terms of higher power in the curvature

¹⁹A full treatment of the supersymmetric boundary counterterms for matter coupled $D = 6$ supergravity to our knowledge is still unknown (see [17] for a treatment for $D = 5$). Nevertheless the scalar and vector falloff is very rapid at infinity, so that there is no contribution from the matter fields for our configurations. See also the discussion later in the text.

this time contribute to the free energy once the cutoff is removed. We give here the form for the Ricci scalar:

$$\mathcal{R}_5 = -\frac{12}{r^2}e^{-2B(r)}. \tag{B.17}$$

In our case one could for instance choose as counterterm the superpotential \mathcal{W} appearing in the susy variations, $S_{AB} = \mathcal{W}\epsilon_{AB}$, which shows the same falloff behaviour and reduces to that of [53] for $X^1 = X^2$. For the six-dimensional configurations taken into consideration, however, the asymptotic falloff of the scalars is very rapid. The expansion of the superpotential \mathcal{W} contains terms which are at least quadratic in the fields (see for instance [79]) therefore it turns out that the scalars do not contribute to the boundary counterterm,²⁰ and indeed a term $\mathcal{W} = 4\tilde{g}$ suffices to renormalize the on-shell action. Putting together expressions (B.11) and (B.16), we get

$$I = I_{\text{bulk}} + I_{\text{GH}} + I_{\text{ct}} = -\frac{\beta \text{Vol}(\mathbb{H}^4)}{8\pi\mathbf{c}G_6} \left(r_+^3 + \frac{\mu}{2} \right), \tag{B.18}$$

which indeed reduces to $I = -\frac{\beta \text{Vol}(\mathbb{H}^4)}{8\pi\mathbf{c}G_6} r_+^3$ in the BPS limit.

Thermodynamics relation and conserved charges in d dimensions. In this section we prove the formula

$$W = \frac{I}{\beta} = E - TS_{\text{BH}} - \phi^I Q_I, \tag{B.19}$$

again generalizing the computation of [53] to the hyperbolic case, following closely their notation. We start from (B.2) and we rewrite it with the help of the R_{tt} component of the Einstein's equations, assuming that all matter fields are independent of time, as it is the case for the solutions considered in this paper. We obtain

$$R_{tt}^t = \frac{1}{2}M_{IJ}F_{tr}^I F^{tr,J} - \frac{1}{4(d-1)}M_{IJ}F_{\mu\nu}^I F^{\mu\nu,J} + \frac{1}{d-1}V, \tag{B.20}$$

hence (B.2) becomes

$$I_{\text{bulk}} = -\frac{1}{8\pi G_{d+1}} \int d^{d+1}x \sqrt{-g} \left(R_{tt}^t - \frac{1}{2}M_{IJ}F_{tr}^I F^{J,tr} \right). \tag{B.21}$$

We have verified that for the metric of the form (B.4) the following holds

$$R_{tt}^t = \frac{1}{\mathbf{c}\sqrt{-g}} \frac{d}{dr} \left(\sqrt{-h}\Theta_t^t \right). \tag{B.22}$$

Moreover, we have the Maxwell's equation $\partial_r(\sqrt{-g}M_{IJ}F^{J,rt}) = 0$. We define the following conserved charges \mathbf{q}_I as

$$\mathbf{q}_I = \frac{\sqrt{-g}}{v_{d-1}} M_{IJ}F^{J,rt}. \tag{B.23}$$

²⁰The scalar behaviour at infinity is $z^i \sim \text{const} + \mathcal{O}(r^{-3})$, while $\sqrt{-h} \sim r^5$. Indeed, also for the known cases [13, 53] the counterterm contribution is just a constant independent of the scalars falloff, $\mathcal{W} = 4\tilde{g}$.

Plugging these expressions into (B.21) we arrive at the following formula for I_{bulk} :

$$\begin{aligned}
 I_{\text{bulk}} &= -\frac{1}{8\pi G_{d+1}} \int d^d x \int_{r_+}^{r_{\text{inf}}} dr \frac{d}{dr} \left(\frac{1}{2} A_t^I \mathbf{q}_I + \frac{\sqrt{-h}}{\mathbf{c}} \Theta_t^t \right) \\
 &= -\frac{\beta \text{Vol}(\mathbb{H}^{d-1})}{8\pi G_{d+1}} \left(\frac{1}{2} A_t^I \mathbf{q}_I + \frac{\sqrt{-h}}{v_{d-1} \mathbf{c}} \Theta_t^t \right) \Big|_{r_+}^{r_{\text{inf}}}.
 \end{aligned}
 \tag{B.24}$$

To regularize the action we need to add the Gibbons-Hawking term, therefore, the full regularized action reads

$$I_{\text{reg}} = \beta W = \frac{\beta \text{Vol}(\mathbb{H}^{d-1})}{8\pi G_{d+1}} \left(-\frac{1}{2} \phi^I \mathbf{q}_I + \frac{\sqrt{-h}}{v_{d-1} \mathbf{c}} (\Theta - \Theta_t^t) \Big|_{r_{\text{inf}}} + \frac{\sqrt{-h}}{v_{d-1} \mathbf{c}} \Theta_t^t \Big|_{r_+} \right).
 \tag{B.25}$$

The first term gives directly the product of the chemical potentials, defined as

$$\phi^I = A^t(r_{\text{inf}}) - A^t(r_+),
 \tag{B.26}$$

and the electric charges, once we define the charges as $Q_I = \frac{\text{Vol}(\mathbb{H}^{d-1})}{16\pi G_{d+1}} \mathbf{q}_I$. We see that the second term is related to the ADM mass of the system, while the third one is related to the product of the temperature T and the Bekenstein-Hawking entropy S_{BH} . Let us focus on the latter. Given the definitions

$$T = \frac{1}{4\pi \mathbf{c} \sqrt{\mathcal{H}(r_+)}} \frac{df}{dr} \Big|_{r_+}, \quad S_{\text{BH}} = \frac{A}{4G_{d+1}} = \frac{\text{Vol}(\mathbb{H}^{d-1})}{8\pi G_{d+1}} (2\pi \sqrt{\mathcal{H}(r_+) r_+^{d-1}}),
 \tag{B.27}$$

we obtain

$$TS_{\text{BH}} = -\frac{\text{Vol}(\mathbb{H}^{d-1})}{8\pi \mathbf{c} G_{d+1}} \frac{\sqrt{-h}}{v_{d-1}} \Theta_t^t \Big|_{r_+},
 \tag{B.28}$$

which holds for a metric of the form (B.4).

The energy is extracted from the renormalized boundary stress energy tensor $T^{ab} = \frac{2}{\sqrt{-h}} \frac{\delta I}{\delta h_{ab}}$ in this way:

$$E = \frac{1}{8\pi G_{d+1}} \int_{\Sigma} \sqrt{\sigma} u_a T^{ab} K_b,
 \tag{B.29}$$

where K_a is the Killing vector field associated with an isometry of the boundary induced metric (in this case, time translations). Σ is the spacelike section of the boundary, σ_{ab} is the induced metric on Σ , and $u^a = \sqrt{-h^{tt}}(1, 0, 0)$ is the unit normal vector to Σ . We will first compute the regulated energy E_{reg} , discussing the counterterms later. The regularized energy reads

$$E_{\text{reg}} = \frac{\text{Vol}(\mathbb{H}^{d-1})}{8\pi G_{d+1}} \frac{\sqrt{-h}}{v_{d-1} \mathbf{c}} (-\Theta_t^t + \Theta).
 \tag{B.30}$$

Plugging all these relations into (B.25) we get

$$W_{\text{reg}} = E_{\text{reg}} - TS_{\text{BH}} - \phi^I Q_I.
 \tag{B.31}$$

This is the relation valid for the regularized quantities. The renormalized ones are obtained by adding the counterterms spelled out in the previous subsections. The counterterms

contribute only to the renormalization of the mass, giving E_{ren} . Hence, the thermodynamics relation (B.19) holds, as expected.

For the records, we report here the explicit values of the energy $E_{\text{ren}} = E$ (black hole mass), entropy S_{BH} and charges Q^I for the solutions considered in the main text. For the four-dimensional solutions in section 3.2 we have

$$\begin{aligned} E &= \frac{\text{Vol}(\mathbb{H}^2)}{16\pi G_4} \left(\mu - \frac{1}{2}(b_1 + b_2 + b_3 + b_4) \right), \\ S_{\text{BH}} &= \frac{1}{4} \sqrt{H_1(r_+)H_2(r_+)H_3(r_+)H_4(r_+)} r_+^2, \\ Q_I &= ib_I \frac{\text{Vol}(\mathbb{H}^2)}{16\pi G_4}, \quad I = 1, \dots, 4. \end{aligned} \tag{B.32}$$

For the six-dimensional solutions of section 4.2 we find

$$\begin{aligned} E &= \frac{3\text{Vol}(\mathbb{H}^4)}{8\pi\sqrt{2}G_6} \left(\mu - \frac{3}{4}(b_1 + b_2) \right), \\ S_{\text{BH}} &= \frac{1}{4} \sqrt{H_1(r_+)H_2(r_+)} r_+^4, \\ Q_I &= i \frac{3}{\sqrt{2}} b_I \frac{\text{Vol}(\mathbb{H}^4)}{16\pi G_4}, \quad I = 1, 2. \end{aligned} \tag{B.33}$$

C Computation of the charges

We now demonstrate that the black hole charges computed from supergravity match those computed in the SCFT. We do so only for the ABJM model.

The trace representation in (2.10) contains three independent flavor charges which correspond to some choice of basis for chemical potentials, represented by flavor gauge fields, satisfying the constraint (2.15). In order to compare with the bulk charges, it is useful to implement the constraint using a Lagrange multiplier charge Λ

$$Z_n^{\text{susy}} = \text{Tr}_{\mathbb{H}^{d-1}} e^{-2\pi n(H - i\sum_I \alpha^I Q_I + i\frac{n-1}{n} Q_R - i\Lambda \sum_I \alpha^I)}. \tag{C.1}$$

From this expression, we can calculate the following

$$\partial_n (-\log Z_n^{\text{susy}}) = 2\pi H - i \sum_I \alpha^I Q_I + i Q_R - i\Lambda \sum_I \alpha^I, \tag{C.2}$$

and

$$\frac{1}{2\pi n} \partial_{\alpha^I} (-\log Z_n^{\text{susy}}) = -i(Q_I + \Lambda). \tag{C.3}$$

In order to compare with the bulk, we first recast (C.1) in terms of the bulk quantities I, β , and ϕ^I :

$$I = -\log \left[\text{Tr}_{\mathbb{H}^{d-1}} e^{-\beta \left(E - \sum_I \phi^I Q_I + i\frac{2\pi-\beta}{4\beta} \sum_I Q_I + i\frac{\beta-2\pi}{\beta} Q_R - \Lambda \sum_I \phi^I + i\frac{2\pi-\beta}{\beta} \Lambda \right)} \right]. \tag{C.4}$$

As a check on this expression, the constraint charge indeed now imposes

$$\sum_{I=1}^4 \phi^I = i \frac{2\pi - \beta}{\beta} = i \frac{1 - n}{n}, \tag{C.5}$$

that is (3.31).

In terms of bulk variables, we can now calculate

$$-\frac{1}{\beta}\partial_{\phi^I}I\Big|_{\beta} = Q_I + \Lambda, \quad (\text{C.6})$$

which, from the definition (3.35), implies,

$$\mathcal{Q}_I = Q_I + \Lambda. \quad (\text{C.7})$$

We also find that

$$\partial_{\beta}I\Big|_{\phi^I} = E + \sum_I \phi^I \mathcal{Q}_I - i\Lambda - \frac{i}{4} \sum_I Q_I + iQ_R, \quad (\text{C.8})$$

yielding

$$E = \partial_{\beta}I\Big|_{\phi^I} - \frac{1}{\beta} \sum_I \phi^I \partial_{\phi^I}I\Big|_{\beta} + i \left(-Q_R + \frac{1}{4} \sum_I Q_I \right) + i\Lambda, \quad (\text{C.9})$$

which is compatible with (3.35) only if we set

$$Q_R = \Lambda + \frac{1}{4} \sum_I Q_I = \frac{1}{4} \sum_I \mathcal{Q}_I. \quad (\text{C.10})$$

We expect the subleading terms of the bulk gauge fields to capture the vacuum expectation value of the charges, i.e.

$$Q_I = \mathcal{N}b_I, \quad (\text{C.11})$$

where \mathcal{N} is a normalization constant. We cannot extract all the \mathcal{Q}_I , because we do not know Q_R from the field theory. However, we do expect the following equations for one less variable

$$\mathcal{Q}_I - \frac{1}{4} \sum_J \mathcal{Q}_J = i\mathcal{N} \left(b_I - \frac{1}{4} \sum_J b_J \right). \quad (\text{C.12})$$

Recall that $\mathcal{Q}_I = -\frac{1}{\beta}\partial_{\phi^I}I\Big|_{\beta}$, see (3.35). One may now check, using the expressions (3.26) and (3.33) for b_I and I as a function of the ϕ^I , that this indeed holds, with

$$\mathcal{N} = i \frac{\text{Vol}(\mathbb{H}^2)}{8\pi\mathbf{c}G_4}, \quad (\text{C.13})$$

after imposing (3.31). Therefore setting $\mathbf{c} = 2$ the value of the charges coincide with the supergravity ones (B.32).

D Rotating charged hyperbolic solutions

In this last section we take into consideration supersymmetric rotating black holes with hyperbolic event horizon that generalize the solutions of section 3.1.²¹ We compute their

²¹Hyperbolic rotating black holes with nontrivial scalar fields exist as well [80], along with analogous magnetic configurations realizing the topological twist [81, 82] but we do not consider them here and we focus instead on the simple minimal gauged supergravity (“universal” truncation) solution.

on-shell action and we show that it assumes a simple form, once the BPS constraints are enforced. We will make contact with the limiting procedure of [19], which allows to approach an extremal BPS limit in the complexified solution.

The Kerr-Newman hyperbolic solution with purely electric charge reads:

$$ds^2 = -\frac{\Delta_r}{\Xi^2 \rho^2} (dt + a \sinh^2(\theta) d\phi)^2 + \frac{\rho^2}{\Delta_r} dr^2 + \frac{\rho^2}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sinh^2(\theta)}{\Xi^2 \rho^2} (adt - (r^2 + a^2) d\phi)^2, \quad (\text{D.1})$$

with

$$\begin{aligned} \Delta_r &= (r^2 + a^2) \left(\frac{r^2}{l^2} - 1 \right) - 2mr + Q^2, \\ \rho^2 &= r^2 + a^2 \cosh^2(\theta), \quad \Delta_\theta = 1 + \frac{a^2}{l^2} \cosh^2(\theta), \quad \Xi = 1 + \frac{a^2}{l^2}, \end{aligned} \quad (\text{D.2})$$

and the gauge field

$$A = -\frac{Qr}{\Xi \rho^2} (dt + a \sinh^2(\theta) d\phi). \quad (\text{D.3})$$

The on-shell action satisfies the thermodynamics relation (we set $G_4 = 1$)

$$I = \beta(M - TS_{\text{BH}} - \phi_e Q_e - \Omega J), \quad (\text{D.4})$$

where

$$\begin{aligned} M &= \frac{m}{(1 + \frac{a^2}{l^2})}, & S_{\text{BH}} &= 4\pi \frac{r_+ + a^2}{(1 + \frac{a^2}{l^2})}, & J &= \frac{am}{(1 + \frac{a^2}{l^2})}, \\ \phi_e &= \frac{Qr_+}{r_+^2 + a^2}, & Q_e &= \frac{Q}{(1 + \frac{a^2}{l^2})}, & \Omega &= \frac{a(l^2 - r_+^2)}{l^2(a^2 + r_+^2)}, \end{aligned} \quad (\text{D.5})$$

and

$$\beta = \frac{4\pi(r_+^2 + a^2)}{r_+ \left(-1 + \frac{a^2}{l^2} + \frac{3r_+^2}{l^2} - \frac{(Q^2 - a^2)}{r_+^2} \right)}. \quad (\text{D.6})$$

One can see that the boundary of spacetime takes the form

$$ds^2 = -\frac{dt^2}{\Xi^2} + \frac{l^2 d\theta^2}{\Delta_\theta} + \frac{l^2}{\Xi} \sinh^2(\theta) d\phi^2, \quad (\text{D.7})$$

that can be cast in

$$ds^2 = \frac{\Delta_\theta}{\Xi^2} (-d\tau^2 + l^2 (d\Theta^2 + \sinh^2(\Theta) d\Phi^2)), \quad (\text{D.8})$$

via the change of coordinates [83]

$$\tau = \frac{t}{\Xi}, \quad \cosh(\Theta) = \cosh(\theta) \sqrt{\frac{\Xi}{\Delta_\theta}}, \quad \Phi = \phi - \frac{at}{l^2 \Xi}. \quad (\text{D.9})$$

The metric (D.8) describes a space which is conformal to (part of) $\mathbb{R} \times \mathbb{H}^2$.

The BPS condition, which can be read off from [84] is given by

$$m^4 + 2 \left(1 - \frac{a^2}{l^2}\right) m^2 Q^2 + \left(1 + \frac{a^2}{l^2}\right)^2 Q^4 = 0, \quad (\text{D.10})$$

which has no solution for real Q , as expected from the previous sections. However, it has solutions for *imaginary* Q and a , which, as stated before, makes sense if we have in mind to work with a Euclidean solution (obtained by Wick rotating $t \rightarrow -i\tau$), for which the gauge field and metric will then be real.

We define $a \rightarrow ij$ and $Q_e \rightarrow iq_e$ with j and q_e real, and we set $l = 1$ for simplicity. We use the BPS condition (D.10) written in function of the latter parameters, to read off the value of the mass

$$m = (j + 1)q_e, \quad (\text{D.11})$$

where we chose the positive branch for regularity. We plug this relation into Δ_r in (D.2) to express the charge q_e as a function of the outer radius r_+ , using the fact that $\Delta_r(r_+) = 0$:

$$q_e = -(j \pm r_+)(r_+ \pm 1). \quad (\text{D.12})$$

Given these relations, the on-shell action (D.4) assumes the simple form

$$I = \frac{\pi(r_+ \pm j)^2}{(j - 1)(j + 1 \pm 2r_+)}. \quad (\text{D.13})$$

In terms of the chemical potentials and β , the on-shell action reads

$$I = \pm i \frac{\beta(\phi_e + i)^2}{2(\Omega + i)}. \quad (\text{D.14})$$

Notice that the chemical potentials Ω and ϕ_e satisfy the relation

$$2i\phi_e - i\Omega - 1 = \pm 2\pi T, \quad (\text{D.15})$$

where $T = 1/\beta$. At this point one can then introduce the replica parameter by imposing $T = 1/(2\pi n)$, and write the on-shell action in terms of two out of the three parameters in (D.15), achieving a generalization of the SRE. A field theory computation, starting for instance from the results in [85], is still unknown. Notice that (D.15) is the generalization of eq. (3.31) to the presence of chemical potential for angular momentum. Defining the shifted potentials, whose meaning will be clear in a moment,

$$\varphi \equiv \beta(\phi_e - \phi_*), \quad \omega \equiv \beta(\Omega - \Omega_*), \quad (\text{D.16})$$

where $\Omega_* = -i$, $\phi_* = -i$, we are able to write (D.13) as

$$I = i \frac{\varphi^2}{2\omega}. \quad (\text{D.17})$$

The variables φ and ω satisfy $2\varphi - \omega = \mp 2\pi i$, like in [86]. This redefinition is similar in spirit to that performed in [19, 77], where Ω_* and ϕ_* are the values of the chemical potentials

computed on the extremal BPS solution. In our case indeed these values give $T = 0$, however the corresponding horizon radius r_* is imaginary, $r_* = \pm i\sqrt{j}$. While it is hard to make sense of this as a proper “extremal BPS” limit, the similarity that arises with [19, 77] is suggestive (r_* in these latter papers is real and corresponds to a well-defined extremal BPS black hole). As a final remark, the form of the on-shell action (D.17) is compatible with the more general form

$$I = i \frac{\sqrt{\varphi_1 \varphi_2 \varphi_3 \varphi_4}}{2\omega}, \quad \sum_{I=1}^4 \frac{\varphi_i}{2} - \omega = -2\pi i, \quad (\text{D.18})$$

expected from the study of [86] carried out for black holes with a spherical horizon. Indeed, (D.18) reduces to (D.17) if we set all φ_I equal, as is the case for minimal gauged supergravity.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] H. Casini and M. Huerta, *On the RG running of the entanglement entropy of a circle*, *Phys. Rev. D* **85** (2012) 125016 [[arXiv:1202.5650](https://arxiv.org/abs/1202.5650)] [[INSPIRE](#)].
- [2] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](https://arxiv.org/abs/hep-th/0603001)] [[INSPIRE](#)].
- [3] T. Nishioka, *Entanglement entropy: holography and renormalization group*, *Rev. Mod. Phys.* **90** (2018) 035007 [[arXiv:1801.10352](https://arxiv.org/abs/1801.10352)] [[INSPIRE](#)].
- [4] P. Calabrese and J.L. Cardy, *Entanglement entropy and quantum field theory*, *J. Stat. Mech.* **0406** (2004) P06002 [[hep-th/0405152](https://arxiv.org/abs/hep-th/0405152)] [[INSPIRE](#)].
- [5] T. Nishioka and I. Yaakov, *Supersymmetric Rényi Entropy*, *JHEP* **10** (2013) 155 [[arXiv:1306.2958](https://arxiv.org/abs/1306.2958)] [[INSPIRE](#)].
- [6] N. Hama, T. Nishioka and T. Ugajin, *Supersymmetric Rényi entropy in five dimensions*, *JHEP* **12** (2014) 048 [[arXiv:1410.2206](https://arxiv.org/abs/1410.2206)] [[INSPIRE](#)].
- [7] X. Huang and Y. Zhou, *$\mathcal{N} = 4$ Super-Yang-Mills on conic space as hologram of STU topological black hole*, *JHEP* **02** (2015) 068 [[arXiv:1408.3393](https://arxiv.org/abs/1408.3393)] [[INSPIRE](#)].
- [8] M. Crossley, E. Dyer and J. Sonner, *Super-Rényi entropy & Wilson loops for $\mathcal{N} = 4$ SYM and their gravity duals*, *JHEP* **12** (2014) 001 [[arXiv:1409.0542](https://arxiv.org/abs/1409.0542)] [[INSPIRE](#)].
- [9] A. Giveon and D. Kutasov, *Supersymmetric Rényi entropy in CFT_2 and AdS_3* , *JHEP* **01** (2016) 042 [[arXiv:1510.08872](https://arxiv.org/abs/1510.08872)] [[INSPIRE](#)].
- [10] H. Mori, *Supersymmetric Rényi entropy in two dimensions*, *JHEP* **03** (2016) 058 [[arXiv:1512.02829](https://arxiv.org/abs/1512.02829)] [[INSPIRE](#)].
- [11] T. Nishioka, *The Gravity Dual of Supersymmetric Rényi Entropy*, *JHEP* **07** (2014) 061 [[arXiv:1401.6764](https://arxiv.org/abs/1401.6764)] [[INSPIRE](#)].
- [12] X. Huang, S.-J. Rey and Y. Zhou, *Three-dimensional SCFT on conic space as hologram of charged topological black hole*, *JHEP* **03** (2014) 127 [[arXiv:1401.5421](https://arxiv.org/abs/1401.5421)] [[INSPIRE](#)].

- [13] L.F. Alday, P. Richmond and J. Sparks, *The holographic supersymmetric Rényi entropy in five dimensions*, *JHEP* **02** (2015) 102 [[arXiv:1410.0899](#)] [[INSPIRE](#)].
- [14] E. Gerchkovitz, J. Gomis and Z. Komargodski, *Sphere Partition Functions and the Zamolodchikov Metric*, *JHEP* **11** (2014) 001 [[arXiv:1405.7271](#)] [[INSPIRE](#)].
- [15] P. Benetti Genolini, D. Cassani, D. Martelli and J. Sparks, *The holographic supersymmetric Casimir energy*, *Phys. Rev. D* **95** (2017) 021902 [[arXiv:1606.02724](#)] [[INSPIRE](#)].
- [16] I. Papadimitriou, *Supercurrent anomalies in 4d SCFTs*, *JHEP* **07** (2017) 038 [[arXiv:1703.04299](#)] [[INSPIRE](#)].
- [17] O.S. An, *Anomaly-corrected supersymmetry algebra and supersymmetric holographic renormalization*, *JHEP* **12** (2017) 107 [[arXiv:1703.09607](#)] [[INSPIRE](#)].
- [18] S. Yankielowicz and Y. Zhou, *Supersymmetric Rényi entropy and Anomalies in 6d (1,0) SCFTs*, *JHEP* **04** (2017) 128 [[arXiv:1702.03518](#)] [[INSPIRE](#)].
- [19] A. Cabo-Bizet, D. Cassani, D. Martelli and S. Murthy, *Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS₅ black holes*, *JHEP* **10** (2019) 062 [[arXiv:1810.11442](#)] [[INSPIRE](#)].
- [20] H. Casini, M. Huerta and R.C. Myers, *Towards a derivation of holographic entanglement entropy*, *JHEP* **05** (2011) 036 [[arXiv:1102.0440](#)] [[INSPIRE](#)].
- [21] G. Festuccia and N. Seiberg, *Rigid Supersymmetric Theories in Curved Superspace*, *JHEP* **06** (2011) 114 [[arXiv:1105.0689](#)] [[INSPIRE](#)].
- [22] C. Closset, T.T. Dumitrescu, G. Festuccia and Z. Komargodski, *Supersymmetric Field Theories on Three-Manifolds*, *JHEP* **05** (2013) 017 [[arXiv:1212.3388](#)] [[INSPIRE](#)].
- [23] T.T. Dumitrescu and G. Festuccia, *Exploring Curved Superspace (II)*, *JHEP* **01** (2013) 072 [[arXiv:1209.5408](#)] [[INSPIRE](#)].
- [24] T.T. Dumitrescu, G. Festuccia and N. Seiberg, *Exploring Curved Superspace*, *JHEP* **08** (2012) 141 [[arXiv:1205.1115](#)] [[INSPIRE](#)].
- [25] J. Kinney, J.M. Maldacena, S. Minwalla and S. Raju, *An index for 4 dimensional super conformal theories*, *Commun. Math. Phys.* **275** (2007) 209 [[hep-th/0510251](#)] [[INSPIRE](#)].
- [26] C. Romelsberger, *Counting chiral primaries in $N = 1$, $d = 4$ superconformal field theories*, *Nucl. Phys. B* **747** (2006) 329 [[hep-th/0510060](#)] [[INSPIRE](#)].
- [27] Y. Zhou, *Information Theoretic Inequalities as Bounds in Superconformal Field Theory*, [arXiv:1607.05401](#) [[INSPIRE](#)].
- [28] E. Witten, *Topological Quantum Field Theory*, *Commun. Math. Phys.* **117** (1988) 353 [[INSPIRE](#)].
- [29] V. Pestun, *Localization of gauge theory on a four-sphere and supersymmetric Wilson loops*, *Commun. Math. Phys.* **313** (2012) 71 [[arXiv:0712.2824](#)] [[INSPIRE](#)].
- [30] N. Hama, K. Hosomichi and S. Lee, *SUSY Gauge Theories on Squashed Three-Spheres*, *JHEP* **05** (2011) 014 [[arXiv:1102.4716](#)] [[INSPIRE](#)].
- [31] A. Kapustin, B. Willett and I. Yaakov, *Exact Results for Wilson Loops in Superconformal Chern-Simons Theories with Matter*, *JHEP* **03** (2010) 089 [[arXiv:0909.4559](#)] [[INSPIRE](#)].
- [32] D.L. Jafferis, *The Exact Superconformal R-Symmetry Extremizes Z*, *JHEP* **05** (2012) 159 [[arXiv:1012.3210](#)] [[INSPIRE](#)].

- [33] C. Closset, T.T. Dumitrescu, G. Festuccia, Z. Komargodski and N. Seiberg, *Contact Terms, Unitarity and F-Maximization in Three-Dimensional Superconformal Theories*, *JHEP* **10** (2012) 053 [[arXiv:1205.4142](#)] [[INSPIRE](#)].
- [34] A. Kapustin, B. Willett and I. Yaakov, *Exact results for supersymmetric abelian vortex loops in 2+1 dimensions*, *JHEP* **06** (2013) 099 [[arXiv:1211.2861](#)] [[INSPIRE](#)].
- [35] N. Drukker, T. Okuda and F. Passerini, *Exact results for vortex loop operators in 3d supersymmetric theories*, *JHEP* **07** (2014) 137 [[arXiv:1211.3409](#)] [[INSPIRE](#)].
- [36] S. Gukov, *Surface Operators*, [arXiv:1412.7127](#) [[INSPIRE](#)].
- [37] T. Nishioka and I. Yaakov, *Supersymmetric Rényi entropy and defect operators*, *JHEP* **11** (2017) 071 [[arXiv:1612.02894](#)] [[INSPIRE](#)].
- [38] D. Martelli, A. Passias and J. Sparks, *The gravity dual of supersymmetric gauge theories on a squashed three-sphere*, *Nucl. Phys. B* **864** (2012) 840 [[arXiv:1110.6400](#)] [[INSPIRE](#)].
- [39] Y. Imamura and D. Yokoyama, *$N = 2$ supersymmetric theories on squashed three-sphere*, *Phys. Rev. D* **85** (2012) 025015 [[arXiv:1109.4734](#)] [[INSPIRE](#)].
- [40] C.-M. Chang, M. Fluder, Y.-H. Lin and Y. Wang, *Romans Supergravity from Five-Dimensional Holograms*, *JHEP* **05** (2018) 039 [[arXiv:1712.10313](#)] [[INSPIRE](#)].
- [41] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, *$N = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091 [[arXiv:0806.1218](#)] [[INSPIRE](#)].
- [42] D.L. Jafferis, I.R. Klebanov, S.S. Pufu and B.R. Safdi, *Towards the F-Theorem: $N = 2$ Field Theories on the Three-Sphere*, *JHEP* **06** (2011) 102 [[arXiv:1103.1181](#)] [[INSPIRE](#)].
- [43] K.A. Intriligator, D.R. Morrison and N. Seiberg, *Five-dimensional supersymmetric gauge theories and degenerations of Calabi-Yau spaces*, *Nucl. Phys. B* **497** (1997) 56 [[hep-th/9702198](#)] [[INSPIRE](#)].
- [44] A. Brandhuber and Y. Oz, *The D-4 – D-8 brane system and five-dimensional fixed points*, *Phys. Lett. B* **460** (1999) 307 [[hep-th/9905148](#)] [[INSPIRE](#)].
- [45] O. Bergman and D. Rodriguez-Gomez, *5d quivers and their AdS_6 duals*, *JHEP* **07** (2012) 171 [[arXiv:1206.3503](#)] [[INSPIRE](#)].
- [46] D.R. Morrison and N. Seiberg, *Extremal transitions and five-dimensional supersymmetric field theories*, *Nucl. Phys. B* **483** (1997) 229 [[hep-th/9609070](#)] [[INSPIRE](#)].
- [47] N. Seiberg, *Five-dimensional SUSY field theories, nontrivial fixed points and string dynamics*, *Phys. Lett. B* **388** (1996) 753 [[hep-th/9608111](#)] [[INSPIRE](#)].
- [48] D.L. Jafferis and S.S. Pufu, *Exact results for five-dimensional superconformal field theories with gravity duals*, *JHEP* **05** (2014) 032 [[arXiv:1207.4359](#)] [[INSPIRE](#)].
- [49] M. Cvetič et al., *Embedding AdS black holes in ten-dimensions and eleven-dimensions*, *Nucl. Phys. B* **558** (1999) 96 [[hep-th/9903214](#)] [[INSPIRE](#)].
- [50] M.J. Duff and J.T. Liu, *Anti-de Sitter black holes in gauged $N = 8$ supergravity*, *Nucl. Phys. B* **554** (1999) 237 [[hep-th/9901149](#)] [[INSPIRE](#)].
- [51] W.A. Sabra, *Anti-de Sitter BPS black holes in $N = 2$ gauged supergravity*, *Phys. Lett. B* **458** (1999) 36 [[hep-th/9903143](#)] [[INSPIRE](#)].

- [52] C. Toldo and S. Vandoren, *Static nonextremal AdS₄ black hole solutions*, *JHEP* **09** (2012) 048 [[arXiv:1207.3014](#)] [[INSPIRE](#)].
- [53] A. Batrachenko, J.T. Liu, R. McNees, W.A. Sabra and W.Y. Wen, *Black hole mass and Hamilton-Jacobi counterterms*, *JHEP* **05** (2005) 034 [[hep-th/0408205](#)] [[INSPIRE](#)].
- [54] L. Andrianopoli, R. D'Auria and S. Vaula, *Matter coupled F(4) gauged supergravity Lagrangian*, *JHEP* **05** (2001) 065 [[hep-th/0104155](#)] [[INSPIRE](#)].
- [55] R. D'Auria, S. Ferrara and S. Vaula, *F(4) supergravity and 5-D superconformal field theories*, *Class. Quant. Grav.* **18** (2001) 3181 [[hep-th/0008209](#)] [[INSPIRE](#)].
- [56] M. Gutperle, J. Kaidi and H. Raj, *Janus solutions in six-dimensional gauged supergravity*, *JHEP* **12** (2017) 018 [[arXiv:1709.09204](#)] [[INSPIRE](#)].
- [57] S.M. Hosseini, K. Hristov, A. Passias and A. Zaffaroni, *6D attractors and black hole microstates*, *JHEP* **12** (2018) 001 [[arXiv:1809.10685](#)] [[INSPIRE](#)].
- [58] L.J. Romans, *The F(4) Gauged Supergravity in Six-dimensions*, *Nucl. Phys. B* **269** (1986) 691 [[INSPIRE](#)].
- [59] M. Suh, *Supersymmetric AdS₆ black holes from F(4) gauged supergravity*, *JHEP* **01** (2019) 035 [[arXiv:1809.03517](#)] [[INSPIRE](#)].
- [60] D.D.K. Chow, *Single-rotation two-charge black holes in gauged supergravity*, [arXiv:1108.5139](#) [[INSPIRE](#)].
- [61] M. Cvetič, H. Lü and C.N. Pope, *Gauged six-dimensional supergravity from massive type IIA*, *Phys. Rev. Lett.* **83** (1999) 5226 [[hep-th/9906221](#)] [[INSPIRE](#)].
- [62] M. Suh, *Supersymmetric AdS₆ black holes from matter coupled F(4) gauged supergravity*, *JHEP* **02** (2019) 108 [[arXiv:1810.00675](#)] [[INSPIRE](#)].
- [63] P. Karndumri, *Twisted compactification of N = 2 5D SCFTs to three and two dimensions from F(4) gauged supergravity*, *JHEP* **09** (2015) 034 [[arXiv:1507.01515](#)] [[INSPIRE](#)].
- [64] F. Benini, K. Hristov and A. Zaffaroni, *Black hole microstates in AdS₄ from supersymmetric localization*, *JHEP* **05** (2016) 054 [[arXiv:1511.04085](#)] [[INSPIRE](#)].
- [65] A. Zaffaroni, *Lectures on AdS Black Holes, Holography and Localization*, [arXiv:1902.07176](#) [[INSPIRE](#)].
- [66] S.M. Hosseini, *Black hole microstates and supersymmetric localization*, Ph.D. Thesis, Milan Bicocca University, Milan Italy 2018, [arXiv:1803.01863](#) [[INSPIRE](#)].
- [67] F. Benini and P. Milan, *Black holes in 4d N = 4 Super-Yang-Mills*, *Phys. Rev. X* **10** (2020) 021037 [[arXiv:1812.09613](#)] [[INSPIRE](#)].
- [68] C. Closset, H. Kim and B. Willett, *Supersymmetric partition functions and the three-dimensional A-twist*, *JHEP* **03** (2017) 074 [[arXiv:1701.03171](#)] [[INSPIRE](#)].
- [69] J. Nian and X. Zhang, *Entanglement Entropy of ABJM Theory and Entropy of Topological Black Hole*, *JHEP* **07** (2017) 096 [[arXiv:1705.01896](#)] [[INSPIRE](#)].
- [70] E. Perlmutter, *A universal feature of CFT Rényi entropy*, *JHEP* **03** (2014) 117 [[arXiv:1308.1083](#)] [[INSPIRE](#)].
- [71] L. Andrianopoli et al., *N = 2 supergravity and N = 2 superYang-Mills theory on general scalar manifolds: Symplectic covariance, gaugings and the momentum map*, *J. Geom. Phys.* **23** (1997) 111 [[hep-th/9605032](#)] [[INSPIRE](#)].

- [72] L.J. Romans, *Supersymmetric, cold and lukewarm black holes in cosmological Einstein-Maxwell theory*, *Nucl. Phys. B* **383** (1992) 395 [[hep-th/9203018](#)] [[INSPIRE](#)].
- [73] H. Lü, C.N. Pope and J. Rahmfeld, *A construction of Killing spinors on S^n* , *J. Math. Phys.* **40** (1999) 4518 [[hep-th/9805151](#)] [[INSPIRE](#)].
- [74] I. Papadimitriou, *Holographic Renormalization of general dilaton-axion gravity*, *JHEP* **08** (2011) 119 [[arXiv:1106.4826](#)] [[INSPIRE](#)].
- [75] A. Cabo-Bizet, U. Kol, L.A. Pando Zayas, I. Papadimitriou and V. Rathee, *Entropy functional and the holographic attractor mechanism*, *JHEP* **05** (2018) 155 [[arXiv:1712.01849](#)] [[INSPIRE](#)].
- [76] A. Gneccchi and C. Toldo, *First order flow for non-extremal AdS black holes and mass from holographic renormalization*, *JHEP* **10** (2014) 075 [[arXiv:1406.0666](#)] [[INSPIRE](#)].
- [77] D. Cassani and L. Papini, *The BPS limit of rotating AdS black hole thermodynamics*, *JHEP* **09** (2019) 079 [[arXiv:1906.10148](#)] [[INSPIRE](#)].
- [78] D.Z. Freedman and S.S. Pufu, *The holography of F-maximization*, *JHEP* **03** (2014) 135 [[arXiv:1302.7310](#)] [[INSPIRE](#)].
- [79] I. Papadimitriou, *Non-Supersymmetric Membrane Flows from Fake Supergravity and Multi-Trace Deformations*, *JHEP* **02** (2007) 008 [[hep-th/0606038](#)] [[INSPIRE](#)].
- [80] K. Hristov, S. Katmadas and C. Toldo, *Matter-coupled supersymmetric Kerr-Newman-AdS₄ black holes*, *Phys. Rev. D* **100** (2019) 066016 [[arXiv:1907.05192](#)] [[INSPIRE](#)].
- [81] D. Klemm, *Rotating BPS black holes in matter-coupled AdS₄ supergravity*, *JHEP* **07** (2011) 019 [[arXiv:1103.4699](#)] [[INSPIRE](#)].
- [82] K. Hristov, S. Katmadas and C. Toldo, *Rotating attractors and BPS black holes in AdS₄*, *JHEP* **01** (2019) 199 [[arXiv:1811.00292](#)] [[INSPIRE](#)].
- [83] D. Klemm and A. Maiorana, *Fluid dynamics on ultrastatic spacetimes and dual black holes*, *JHEP* **07** (2014) 122 [[arXiv:1404.0176](#)] [[INSPIRE](#)].
- [84] M.M. Caldarelli and D. Klemm, *Supersymmetry of Anti-de Sitter black holes*, *Nucl. Phys. B* **545** (1999) 434 [[hep-th/9808097](#)] [[INSPIRE](#)].
- [85] K. Murata, T. Nishioka, N. Tanahashi and H. Yumisaki, *Phase Transitions of Charged Kerr-AdS Black Holes from Large- N Gauge Theories*, *Prog. Theor. Phys.* **120** (2008) 473 [[arXiv:0806.2314](#)] [[INSPIRE](#)].
- [86] S. Choi, C. Hwang, S. Kim and J. Nahmgoong, *Entropy Functions of BPS Black Holes in AdS₄ and AdS₆*, *J. Korean Phys. Soc.* **76** (2020) 101 [[arXiv:1811.02158](#)] [[INSPIRE](#)].