

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Freeze tolerance of the yeast *Torulaspota delbrueckii*: cellular and biochemical basis

Cecília Alves-Araújo^a, Maria Judite Almeida^a, Maria João Sousa^{a*}, Cecília Leão^b

^a Center of Biology/Department of Biology, University of Minho, *Campus de Gualtar*, 4710-057 Braga, Portugal. ^b Life and Health Sciences Research Institute, School of Health Sciences, University of Minho, *Campus de Gualtar*, 4710-057 Braga, Portugal.

Running title: High Freeze-Tolerance of *Torulaspota delbrueckii*

Key words: *Torulaspota delbrueckii*, freeze-tolerance, membrane-integrity, baker's yeast

* Author for correspondence. Phone: 351 253604310; Fax: 351 253678980

E-mail address: mjsousa@bio.uminho.pt

Present address: Department of Biology, University of Minho, *Campus de Gualtar*, 4710-057 Braga, Portugal

1 **Abstract**

2 The freeze-stress responses to prolonged storage at - 20 °C in *Torulaspota delbrueckii* PYCC5323
3 were investigated. In this yeast no loss of cell viability was observed for at least 120 days during
4 freezing at - 20°C, whereas a loss of 80% was observed in a commercial baker's yeast after 15 days.
5 In the former strain, freeze resistance was dependent on an adaptation process. The primary cell
6 target of freeze stress was the plasma membrane, preservation of it's integrity being related with a
7 lower increase of lipid peroxidation and with higher resistance to H₂O₂, but not with intracellular
8 trehalose concentration.

9

10

1 **1. Introduction**

2 Frozen-dough technology is well established in the baking industry, making it easier for bakers to
3 supply oven-fresh bakery products to consumers and improving labour conditions. However, storage
4 of frozen bread-dough may lead to the loss of baker's yeast cell viability as well as of its baking
5 capacity, and consequently to economic losses. Thus, bread-making industry keeps a high demand
6 for yeast strains with improved freeze resistance [1,2]. Most research on this field has focused on
7 strains of *Saccharomyces cerevisiae*, which is the species currently used as baker's yeast. In this
8 species, tolerance to freezing has generally been correlated with the intracellular trehalose
9 concentration, but no direct correlation has been found above a threshold value [3-6]. Prior to the
10 frozen storage, once the yeast cells are mixed with flour the fermentation takes place and a rapid loss
11 of stress resistance occurs [7]. This has also been associated with the degradation of intracellular
12 trehalose [8]. However, it has been shown that retention of high trehalose levels in fermenting cells
13 does not prevent the loss of fermentation capacity during freezing, and that other factors - not yet
14 identified - are required for the maintenance of freeze stress resistance [9, 10]. Accumulation of other
15 solutes such as amino acids and glycerol, and expression of aquaporins were also reported to
16 increase freezing resistance [11-15]. In addition, oxidative damage has been considered to be a factor
17 underlying freeze-thaw damage, since an oxidative burst has been predicted to occur during thawing
18 [16]. In agreement with this, Park et al. found that respiratory ability and functional mitochondria are
19 necessary to confer full resistance to freeze-thaw stress [17]. It has also been shown that freeze
20 tolerance is correlated with tolerance to H₂O₂, and free radicals were detected in *S. cerevisiae* after
21 the freeze-thaw process [5, 18].

22 The strain PYCC5323 of *Torulaspota delbrueckii*, isolated from traditional corn and rye bread
23 dough from the North of Portugal, besides presenting dough-raising capacity, growth rates and
24 biomass yields similar to commercial baker's yeast, displays high freeze and osmotic tolerance [19-
25 21]. Therefore this yeast emerges as a powerful candidate for the bread making industry, and the
26 elucidation of such a peculiar behaviour reveals to be of great interest.

27 In this work the freeze-stress responses to prolonged storage at - 20 °C in *Torulaspota delbrueckii*
28 PYCC5323 were investigated. The results obtained were compared to the ones of a commercial
29 baker's strain of *S. cerevisiae*. The following cellular and biochemical parameters were analysed: cell

1 viability, plasma membrane integrity, oxidative damages, intracellular trehalose content and
2 trehalase(s) activity.

3

4 **2. Materials and methods**

5 *2.1. Microorganisms, growth and freezing conditions*

6 The strains used were *Torulasporea delbrueckii* PYCC5323, isolated from homemade corn and rye
7 bread dough, and *Saccharomyces cerevisiae* PYCC5325, isolated from commercial compressed
8 baker's yeast - both supplied by the Portuguese Yeast Culture Collection, Faculdade de Ciências e
9 Tecnologia, Universidade Nova de Lisboa, Caparica, Portugal. Stock cultures were maintained on
10 glucose-yeast extract-peptone-agar (2% (wt/vol) glucose, 1% (wt/vol) peptone, 0.5% (wt/vol) yeast
11 extract, 2% (wt/vol) agar), at 4 °C. Yeast strains were grown on yeast extract-peptone-sucrose (YPS)
12 medium containing 2% (wt/vol) sucrose, 4% (wt/vol) peptone, 2% (wt/vol) yeast extract, 0.2% (wt/vol)
13 KH_2PO_4 , and 0.1% (wt/vol) $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, at 30 °C. Peptone (ref. 0118-17) and yeast extract (ref.
14 0127-17) were from Difco (Becton Dickinson, Sparks, MD, USA) and sucrose from Merck (E. Merck,
15 Darmstadt, Germany).

16 For freezing assays, cells were harvested at initial stationary phase (24 hours of culture, 2.4-
17 2.7×10^8 cells ml^{-1} for *T. delbrueckii* and 1.1-1.5 $\times 10^8$ cells ml^{-1} for *S. cerevisiae*), washed twice with
18 deionised water, and suspended in a quarter of the initial volume in sterile water to an A_{640} of 12-20.
19 Aliquots (5 ml) of cells were transferred into 15-ml polycarbonate tubes, centrifuged and the pellet
20 (400-500 mg of wet weight) suspended in 500 μl of the storing medium: LF medium or as described in
21 results. LF medium is a liquid medium formulated to simulate the fermenting ability of yeast in bread
22 dough, with the following composition: 1% (wt/vol) glucose, 1% (wt/vol) sucrose, 3% (wt/vol) maltose,
23 0.25% (wt/vol) $(\text{NH}_4)_2\text{SO}_4$, 0.5% (wt/vol) urea, 1.6% (wt/vol) KH_2PO_4 , 0.5% (wt/vol) $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$,
24 0.06% (wt/vol) MgSO_4 , 22.5 ppm nicotinic acid, 5.0 ppm pantothenic acid, 2.5 ppm thiamine, 1.25 ppm
25 pyridoxine, 1.0 ppm riboflavin and 0.5 ppm folic acid [22]. The samples were then frozen at -20 °C for
26 different time periods (cooling rate approximately 3 °C min^{-1}) [5], and then thawed at 30 °C for 2 min.
27 For fast freezing, 15-ml polycarbonate tubes containing the cell suspensions were directly immersed in
28 liquid nitrogen (cooling rate approximately 200 °C min^{-1}) [5]. For pre-fermentation treatments, cells
29 were subjected to a fermentation period before freezing as follows: the pellets in the 15-ml
30 polycarbonate tubes were suspended in LF medium to a final $\text{OD}_{640 \text{ nm}}$ of 0.3-0.5, and incubated for

1 120 min, at 30 °C. After this, the suspension was centrifuged and the pellet was suspended in LF
2 medium and frozen as described above.

3 4 *2.2. Extraction and assay of trehalose*

5 Cells were harvested by centrifugation, washed twice with cold deionised water and sampled for
6 dry weight contents. Trehalose was extracted from cold cell pellets with 5% (wt/vol) trichloroacetic acid
7 (Merck, Darmstadt, Germany) for 45 min with occasional shaking. Cells were then centrifuged at 735
8 g, for 10 min. Extraction was repeated once more, and supernatants from the two extractions were
9 combined and used for the determination of trehalose by high-performance liquid chromatography.
10 The apparatus used was a Gilson chromatograph (132-RI Detector) equipped with a carbohydrate H⁺
11 column (SS-100, H⁺, Hypersil) which was maintained at 30 °C. A solution of H₂SO₄ (0.0025 M) was
12 used as the mobile phase at a flow rate of 0.45 ml min⁻¹. The relative values (%) of intracellular
13 trehalose concentration after different periods of freezing were calculated by dividing the values of
14 intracellular trehalose concentration obtained for the frozen samples, by those obtained for the
15 unfrozen samples.

16 17 *2.3. Trehalase activity and protein assay*

18 Pellets containing 75-100 mg (wet wt) of washed cells were suspended in 1 ml ice-cold 50 μM
19 MES (4-morpholineethanesulfonic acid) (Boehringer Mannheim, GmbH-Germany)/KOH buffer, pH 7,
20 containing 50 μM CaCl₂. Cells were broken by vortexing with 500 μl of glass beads (0.5 mm diameter),
21 for four periods of 1 min, with 1 min intervals on ice between them. The crude enzyme extract was
22 centrifuged for 3 min at 13 200 g, at 4 °C. The supernatant was dialysed overnight at 4 °C against 10
23 mM Mes/KOH buffer, pH 7, containing 50 μM CaCl₂. Trehalase was assayed as described previously
24 [23]. The glucose liberated was determined by glucose oxidase/peroxidase method (Glucose GOD –
25 Perid, Boehringer Mannheim, GmbH-Germany). Protein determination was carried out according to
26 Lowry et al. [24]. Specific activity of trehalase was expressed as units (U, nmol glucose released per
27 min) per mg protein.

28 29 *2.4. Measurement of cell viability and membrane integrity*

30 The viability of yeast cells was determined by counting CFU. For this assay, yeast cell
31 suspensions were washed twice with deionised water and, after convenient dilution, spread on YPDA

1 medium plates. The plates were incubated for 48 hours at 30 °C before counting. The relative values
2 (%) of viable cells after different periods of freezing were calculated by dividing the values of CFU
3 counts obtained for the frozen samples, by those obtained for the unfrozen samples.

4 Membrane integrity was analysed by flow cytometry using the membrane exclusion dye,
5 propidium iodide (PI). In these assays, cells with preserved membrane integrity are not permeated by
6 propidium iodide (PI⁻ cells) while those that lost their membrane integrity do incorporate the
7 fluorochrome (PI⁺ cells) [25]. Cell suspensions (about 10⁷ cells ml⁻¹) were incubated for 10 min in the
8 dark with a 20 µg/ml PI solution (ratio of 2:1, respectively) and injected on a Partec PAS III flow
9 cytometer equipped with an argon-ion laser emitting a 15 mW beam at 488 nm . From each sample
10 2x10⁴ cells were analysed. Control suspensions of membrane-disrupted cells were prepared by boiling
11 cell suspensions. The relative values (%) of PI⁻ cells after different periods of freezing were calculated
12 by dividing the values of PI⁻ cells obtained for the frozen samples, by those obtained for the unfrozen
13 samples.

14

15

16 *2.5. Oxidative stress evaluation*

17 For evaluating the effect of pre-treatment with the radical scavenger N-tert-butyl- α -phenylnitron
18 (PBN) (Aldrich Chem. Co. Milwaukee, WI 53201), cells were grown for 24 hours to initial stationary
19 phase in YPS medium at 30 °C. PBN was added to the culture at the final concentrations of 0.5, 5.0
20 and 15 mM, and cells were cultured for 30 min before being harvested.. PBN was removed by
21 washing cells twice with deionised water and cells were frozen in LF medium as described above. In
22 assays where a pre-fermentation was performed, PBN was added to the fermenting cell suspensions
23 30 min before the end of the fermentation period.

24 For treatment with oxidising agents, cells were grown to mid-exponential (OD_{640 nm} 0.5-1.0) and
25 initial stationary phase (24 hours of culture), harvested and washed twice with deionised water.
26 Subsequently, these cells were suspended in water in order to achieve a concentration of OD_{640nm} 0.6-
27 0.7, and 0.1 ml of this suspension was spread on solid YPDA plates (50 mm diameter). A paper disc
28 (6 mm diameter - BBL, 231039, Becton Dickinson, Sparks, MD, USA) containing 10 µl of one oxidising
29 agent, was laid on each inoculated plate. The oxidants used were: menadione at concentrations of
30 0.05, 0.10, 1.0 mM; diamide at concentrations of 0.3, 3.0 M; hydrogen peroxide at 1.1 and 11 M, all

1 purchased from Sigma Chemical Co (St. Louis, MO). The plates were incubated at 26 °C during two
2 days, after which the diameters of the inhibition halos around the paper disks were measured. The
3 value obtained was the average of two perpendicular diameters, excluding the disc diameter [26].
4

5 *2.6. Thiobarbituric acid (TBA) reaction for lipid peroxide analysis*

6 For TBA-reactive substances (TBARS) quantification, pellets containing 350-400 mg (wet weight)
7 of cells were washed with ice-cold deionised water and suspended in 0.75 ml ice-cold sodium
8 phosphate (E. Merck, Darmstadt, Germany) buffer, pH 7.2. Cells were broken by vortexing with 500 µl
9 of glass beads (0.5 mm diameter) for six periods of 1 min with 1 min intervals on ice between them.
10 TBARS were then determined according to Buege and Aust [27]. The TBARS concentration was
11 expressed as µmol malondialdehyde per mg protein.
12

13 *2.7. Reproducibility of the results*

14 All the experiments were repeated at least three times, and the data reported are mean values
15 and SD. When statistical analyses were performed, the significance was tested by analysis of
16 variance (Anova, Microsoft Excel 2000).
17

18 **3. Results**

19 *3.1. Cell viability and membrane integrity along freeze storage*

20 To characterise the freeze resistance of *T. delbrueckii* PYCC5323, we first studied cell viability,
21 assessed as colony-forming units (CFU), during freezing at -20 °C for up to 120 days as described in
22 materials and methods (Fig. 1A). For a comparative analysis, a commercial baker's yeast, *S.*
23 *cerevisiae* PYCC5325, was used as a reference strain (Fig. 1B). In *T. delbrueckii* no loss of cell
24 viability was observed for the entire storage period, whereas a loss of 80% was obtained in *S.*
25 *cerevisiae* after 15 days. In both species, viability loss was faster when cells were stored in freezing
26 medium without sugars, even when glycerol was added to keep the osmotic pressure. These results
27 are consistent with previous reports showing that the presence of the disaccharide trehalose in the
28 extracellular medium has a protective effect in cell viability during freezing [28].

29 To elucidate whether the loss of membrane integrity along freezing was directly conditioning cell
30 viability, membrane damage was monitored by flow cytometry using propidium iodide (PI) cell staining.

1 The decrease of cells with preserved plasma membrane (PI⁻ cells) was much less pronounced in *T.*
2 *delbrueckii* (Fig. 2 A) than in *S. cerevisiae*. As shown in Fig. 2B, there was a direct correlation between
3 the percentage of PI⁻ cells and the percentage of CFU counts for both *T. delbrueckii* and *S. cerevisiae*
4 yeast strains. This was evident either in cells subjected to a short fermentation period (120 min) or not
5 subjected to fermentation before freezing. The results show that, independently of the physiological
6 state of the cells, membrane integrity is directly conditioning cell viability, expressed as CFU counts,
7 and therefore the plasma membrane seems to be one of the first freezing targets.

8

9 3.2. Cell oxidative stress responses during the freeze-thaw process

10 The results described above indicate that *T. delbrueckii* displays a higher freeze resistance when
11 compared with *S. cerevisiae*, which is mainly due to plasma membrane integrity. To examine whether
12 this capacity to preserve plasma membrane integrity was correlated with oxidative stress resistance,
13 we directly assessed oxidative damage in the membranes of both *T. delbrueckii* and *S. cerevisiae*
14 during frozen storage by measuring TBA-reactive substances (TBARS). The TBARS test quantifies
15 lipid peroxides in the thiobarbituric acid derivatized form. As shown in Table 1, for both yeast strains,
16 frozen cells presented higher amounts of products of lipid peroxidation than control cells (cells before
17 freezing). However, the percentage increase in TBARS levels was significantly enhanced ($P < 0.05$)
18 much earlier in *S. cerevisiae* (approximately 61% for the fifth day), which was associated to its rapid
19 decrease in cell membrane integrity. As shown in the previous section, the presence of glycerol did
20 not protect cells of *T. delbrueckii* during freezing. Therefore we tested *T. delbrueckii* under these
21 conditions to evaluate the increase in the percentage of TBARS. The results showed that the values of
22 percentage increase of TBARS in *T. delbrueckii* frozen in a medium with glycerol, for 60 and 84 days,
23 were similar to those observed for *S. cerevisiae* (respectively 64% and 83%). Moreover, samples of *T.*
24 *delbrueckii* frozen under conditions where no loss of cell viability was observed, presented much lower
25 percentage increase in TBARS production (Fig. 1, Table 1).

26 The role of oxidative stress during freezing, was examined by using the oxygen radical scavenger
27 PBN. In *S. cerevisiae* PYCC5325, pre-incubation of cells with PBN (1 or 5 mM) resulted in higher
28 levels of membrane integrity preservation for the first days of frozen storage, but no protective effect
29 by PBN was observed in cells frozen after a period of pre-fermentation (results not shown). This result
30 might reflect the different physiological status of the cells and it is conceivable that oxidative stress is

1 being carried over by other factors [17]. Next, the oxidative stress responses of both strains to
2 hydrogen peroxide (H₂O₂, toxicity mainly due to hydroxyl radicals), menadione (a superoxide-
3 generating agent) and diamide (thiol-oxidizing drug) were evaluated. The results obtained in a disk
4 diffusion assay are shown in Fig. 3. *T. delbrueckii* was more sensitive to menadione ($P < 0.001$) and
5 diamide ($P < 0.05$) while more tolerant to H₂O₂ ($P < 0.01$) when compared with *S. cerevisiae*.
6 Therefore, like it was previously reported for *S. cerevisiae*, in *T. delbrueckii* freeze resistance is also
7 correlated with H₂O₂ resistance [5].

8

9 3.3. Intracellular trehalose content during the storage period

10 The results described above pointed to the possibility that oxidative stress was not the only
11 condition influencing the cell freeze resistance, mainly in cells frozen after a short period of pre-
12 fermentation. Intracellular trehalose accumulation has been described to protect cells from oxygen
13 radicals and also to be involved in the stabilisation of the plasma membrane structure during freezing
14 [29-31]. Hence, it was also investigated whether trehalose was involved on the higher capacity of *T.*
15 *delbrueckii* to maintain plasma membrane integrity compared to *S. cerevisiae*. Immediately before
16 freezing, the values of the intracellular trehalose content were high and similar in both species: 109.34
17 ± 8.72 and 112.79 ± 20.97 mg g⁻¹ dry wt for *S. cerevisiae* and *T. delbrueckii*, respectively. During the
18 entire freezing storage period, these values were kept high and constant in *T. delbrueckii* but
19 decreased quickly in *S. cerevisiae*. For both species the relative values of the intracellular trehalose
20 content during freezing followed closely the percentage of PI⁻ cells (Fig. 4). In *S. cerevisiae*, the
21 decrease in the intracellular trehalose content was accompanied by an increase of the extracellular
22 trehalose content and of the medium OD_{260nm} (used as a measure of leakage of cell contents)
23 (Fig.4B). At the end of the assay, the amount of trehalose found in the media was about the same as
24 the one lost from the cells, indicating that total trehalose amount (intracellular + extracellular) remained
25 unchanged. Thus, cell leakage appears to be responsible for the decrease in trehalose content of the
26 cells, which is consistent with the observed loss of plasma membrane integrity. In addition, a decrease
27 in the intracellular trehalose content was observed in cells that were subjected to a short fermentation
28 period before freezing (cells more sensitive to freeze stress), which was similar for both strains (from
29 109.34 ± 8.72 to 9.79 ± 0.93 mg g⁻¹ dry wt for *S. cerevisiae* and from 112.79 ± 20.97 to 17.16 ± 6.24
30 mg g⁻¹ dry wt for *T. delbrueckii*, during a 120 min fermentation period).

1 These results seemed to indicate that a similar activation pattern of trehalase(s) by glucose was
2 present in both yeast strains. To address this point, trehalase(s) activities were monitored in cell free
3 extracts prepared from cells in the absence of glucose (stationary phase cells) as well as after a
4 glucose pulse. A similar behaviour was presented by both strains with a two to three-fold increase in
5 activity almost immediately after glucose addition (from 15.0 ± 0.4 to 32.1 ± 1.3 mU/mg protein for *S.*
6 *cerevisiae* and from 11.1 ± 0.2 to 31.0 ± 0.7 mU/mg protein for *T. delbrueckii*, respectively before and
7 after glucose addition). The results are in agreement with the mobilisation of trehalose observed
8 during the pre-fermentation period.

9 The results described above are consistent with the conclusion that the higher freeze resistance
10 displayed by *T. delbrueckii* can not be attributed to higher intracellular trehalose contents.

12 3.4. Cell adaptation during the freeze period

13 We evaluated the ability of *T. delbrueckii* and *S. cerevisiae* to adapt to freezing by inhibiting
14 protein synthesis with cycloheximide. The results obtained when cells were frozen at -20 °C, showed
15 that for *T. delbrueckii*, contrasting with *S. cerevisiae*, the presence of cycloheximide in the freezing
16 medium increased the loss of cell viability throughout frozen storage (Fig. 1). To assess if
17 cycloheximide could have a toxic effect in *T. delbrueckii* cells, being responsible for the observed
18 decrease in cell viability, a cell suspension of this yeast was incubated in LF medium with
19 cycloheximide, and the number of CFU counts was determined without previous freezing the
20 suspension. No differences were found between the number of CFU counts estimated for this cell
21 suspension and for the control without cycloheximide. These results indicated that at the concentration
22 tested, cycloheximide was not having a toxic effect. In addition, when cells of *T. delbrueckii* were
23 frozen at a much faster rate, in liquid nitrogen (-196 °C), and subsequent storage at -80 °C, loss of cell
24 viability rapidly increased, CFU counts dropping approximately 50% after one day of frozen storage.
25 Together these results suggest that cell viability of *T. delbrueckii* is dependent on *de novo* protein
26 synthesis, and therefore that yeast cells can adapt in a slow freezing process, most probably during
27 the initial cooling period. In accordance with these findings, the capacity of *T. delbrueckii* cells to adapt
28 might be essential for the yeast high freeze tolerance phenotype.

1 4. Discussion

2 Our findings clearly indicate that the strain *T. delbrueckii* PYCC5323 exhibits a high freeze
3 tolerance, when compared with the baker's yeast *S. cerevisiae*, which reinforces previous reports
4 claiming for its useful exploitation in baking industry. Evidence was presented that the primary cell
5 target of freezing stress is the plasma membrane and that the capacity to preserve membrane integrity
6 displayed by *T. delbrueckii* PYCC5323 is correlated with higher resistance to lipid oxidative damage.
7 Hydroxyl radicals appear to be the agents responsible for cell membrane damage in freeze stress in
8 this yeast. This is the opposite of a previous report on *S. cerevisiae* where superoxide radicals were
9 considered as the agents responsible for cell damage in these stress conditions [18]. To ascertain the
10 key role of these radicals during freezing we are developing a catalase null mutant in *T. delbrueckii*
11 PYCC5323. Loss of cell viability seems to be correlated with the percentage increase in TBARS levels
12 and not with their absolute values, suggesting that a higher content in unsaturated fatty acids could
13 allow the cell to cope with higher absolute levels of lipid oxidation without compromising membrane
14 integrity. This hypothesis would also agree with previous work, showing that freeze-tolerant yeast
15 strains have larger amounts of unsaturated fatty acids when compared with sensitive strains [32].
16 Previous studies have pointed to a lack of capacity in *S. cerevisiae* to adapt to cold stress [17],
17 although evidence for cold-induced expression changes associated with improved cryoresistance has
18 also been provided more recently [33, 34]. According to our results, in *T. delbrueckii*, contrasting with
19 *S. cerevisiae*, the surviving capacity (evaluated by cell viability) is dependent on *de novo* protein
20 synthesis. An adaptation process during slow freezing appears to be determinant for the yeast high
21 freeze tolerance phenotype. In the light of these observations, the yeast response to freeze stress
22 seems to be strongly dependent on the yeast strain, culture and freeze conditions. However, further
23 studies will be necessary to clarify the molecular basis of cell adaptation to cold/freeze. In the case of
24 the strain *T. delbrueckii* PYCC5323, the observed behaviour is particularly relevant in view to its
25 utilisation for frozen dough production, and implies that the dough should be frozen at a slow, rather
26 than at a fast rate.

27 In addition and noteworthy from a methodological point of view, the results regarding the
28 correlation observed along freezing between the loss of membrane integrity and cell proliferative
29 capacity, validate the application of flow cytometry and the use of the fluorochrome PI as a measure of
30 viability of cells subjected to freeze stress. Therefore, and contrary to other stress conditions [25], the

1 assessment of PI⁻ cells by flow cytometry, as a method to determine cell viability either in *T.*
2 *delbrueckii* or *S. cerevisiae*, can replace the more laborious and time consuming determination of CFU
3 counts.

4

5 **Acknowledgements**

6 We are grateful to Professor Victor Costa for laboratory support and for critical reading of the
7 manuscript. Cecília Alves-Araújo was supported by a PhD grant (PRAXIS XXI/BD/21543/99) from
8 Fundação para a Ciência e a Tecnologia, Portugal.

9

10 **References**

11 [1] Randez-Gil, F., Sanz, P. and Prieto, J.A. (1999) Engineering baker's yeast: room for improvement.

12 TIBTECH 17, 237-244.

13 [2] Randez-Gil, F., Aguilera, J., Codón, A., Rincón, A. M., Estruch, F. and Prieto, J. A. (2003)

14 Baker's yeast: challenges and future prospects. In: Functional Genetics of Industrial Yeasts, De
15 Winde JH (ed). Springer-Verlag, Heidelberg; 57-97.

16 [3] Attfield, P.V., Raman, A. and Northcott, C. (1992) Construction of *Saccharomyces cerevisiae*

17 strains that accumulate relatively low concentrations of trehalose, and their application in testing
18 the contribution of the disaccharide to stress tolerance. FEMS Microbiol. Lett. 94, 271-276.

19 [4] Yokoigawa, K., Murakami, Y. and Kawai, H. (1995) Trehalase activity and trehalose content in a

20 freeze-tolerant yeast, *Torulasporea delbrueckii*, and its freeze-sensitive mutant. Biosci. Biotechnol.
21 Biochem. 59(11), 2143-2145.

22 [5] Lewis, J.G., Learmonth, R.P., Attfield, P.V. and Watson, K. (1997) Stress co-tolerance and

23 trehalose content in baking strains of *Saccharomyces cerevisiae*. J. Ind. Microbiol. Biotechnol. 18,
24 30-36.

25 [6] Sano, F., Asakawa, N., Inoue, Y. and Sakurai, M. (1999) A dual role for intracellular trehalose in

26 the resistance of yeast cells to water stress. Cryobiology 39, 80-87.

27 [7] Hino, A., Tacano, H. and Tanaka, Y. (1987) New freeze-tolerant yeast for frozen dough

28 preparations. Cereal Chem. 64(4), 269-275.

- 1 [8] Shima, J., Hino, A., Yamada-Iyo, C., Suzuki, Y., Nakajima, R., Watanabe, H., Mori, K. and Takano,
2 H. (1999) Stress tolerance in doughs of *Saccharomyces cerevisiae* trehalase mutants derived
3 from commercial baker's yeast. *Appl. Environ. Microbiol.* 65, 2841-2846.
- 4 [9] Van Dijck, P., Colavizza, D., Smet, P. and Thevelein, J. M. (1995) Differential importance of
5 trehalose in stress in fermenting and nonfermenting *Saccharomyces cerevisiae* cells. *Appl.*
6 *Environ. Microbiol.* 61(1), 109-115.
- 7 [10] Versele, M., Thevelein, J.M. and Van Dijck, P. (2004) The high general stress resistance of the
8 *Saccharomyces cerevisiae* fil1 adenylate cyclase mutant (Cyr1Lys1682) is only partially
9 dependent on trehalose, Hsp104 and overexpression of Msn2/4-regulated genes. *Yeast* 21(1), 75-
10 86.
- 11 [11] Shima, J., Sakata-Tsuda, Y., Suzuki, Y., Nakajima, R., Watanabe, H., Kawamoto, S. and Takano,
12 H. (2003) Disruption of the *CAR1* gene encoding arginase enhances freeze tolerance of the
13 commercial baker's yeast *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 69(1), 715-718.
- 14 [12] Terao, Y., Nakamori, S. and Takagi, H. (2003) Gene dosage effect of L-proline biosynthetic
15 enzymes on L-proline accumulation and freeze tolerance in *Saccharomyces cerevisiae*. *Appl.*
16 *Environ. Microbiol.* 69(11), 6527-6532.
- 17 [13] Izawa, S., Sato, M., Yokoigawa, K. and Inoue, Y. (2004) Intracellular glycerol influences
18 resistance to freeze stress in *Saccharomyces cerevisiae*: analysis of a quadruple mutant in
19 glycerol dehydrogenase genes and glycerol-enriched cells. *Appl. Microbiol. Biotechnol.* May 4
20 [Epub ahead of print]
- 21 [14] Tanghe, A., Van Dijck, P., Dumortier, F., Teunissen, A., Hohmann, S. and Thevelein, J.M. (2002)
22 Aquaporin expression correlates with freeze tolerance in baker's yeast, and overexpression
23 improves freeze tolerance in industrial strains. *Appl. Environ. Microbiol.* 68(12), 5981-5989.
- 24 [15] Tanghe, A., Van Dijck, P., Colavizza, D. and Thevelein, J.M. (2004) Aquaporin-mediated
25 improvement of freeze tolerance of *Saccharomyces cerevisiae* is restrict to rapid freezing
26 conditions. *Appl. Environ. Microbiol.* 70(6), 3377-3382.
- 27 [16] Hermes-Lima, M. and Storey, K.B. (1993) Antioxidant defenses in the tolerance of freezing and
28 anoxia by garter snakes. *Am. J. Physiol.* 265 (3 Pt 2), R646-652.

- 1 [17] Park, J.-I., Grant, C.M., Atfield, P.V. and Dawes, I.W. (1997) The freeze-thaw stress response of
2 the yeast *Saccharomyces cerevisiae* is growth phase specific and is controlled by nutritional state
3 via the RAS-cyclic AMP signal transduction pathway. *Appl. Environ. Microbiol.* 63, 3818-3824.
- 4 [18] Park, J.-I., Grant, C.M., Davies, M.J. and Dawes, I.W. (1998) The cytoplasmic Cu, Zn superoxide
5 dismutase of *Saccharomyces cerevisiae* is required for resistance to freeze-thaw stress.
6 Generation of free radicals during freezing and thawing. *J. Biol. Chem.* 273(36), 22921-22928.
- 7 [19] Almeida, M.J. and Pais, C.S. (1996a) Characterization of yeast population from traditional corn
8 and rye bread doughs. *Lett. Appl. Microbiol.* 23, 154-158.
- 9 [20] Almeida, M.J. and Pais, C.S. (1996b) Leavening ability and freeze tolerance of yeasts isolated
10 from traditional corn and rye bread doughs. *Appl. Environ. Microbiol.* 62, 4401-4401.
- 11 [21] Hernandez-Lopez, M.J., Prieto, J.A. and Randez-Gil, F. (2003) Osmotolerance and leavening
12 ability in sweet and frozen sweet dough. Comparative analysis between *Torulaspota delbrueckii*
13 and *Saccharomyces cerevisiae* baker's yeast strains. *Antonie Van Leeuwenhoek.* 84(2), 125-134.
- 14 [22] Hino, A., Mihara, K., Nakashima, K. and Takano, H. (1990) Trehalose levels and survival ratio of
15 freeze-tolerant versus freeze-sensitive yeasts. *Appl. Environ. Microbiol.* 56(5), 1386-1391.
- 16 [23] Durnez, P., Pernambuco, M.B., Oris, E., Arguelles, J.C., Mergelsberg, H. and Thevelein, J.M.
17 (1994) Activation of trehalase during growth induction by nitrogen sources in the yeast
18 *Saccharomyces cerevisiae* depends on the free catalytic subunits of cAMP-dependent protein
19 kinase, but not on functional Ras proteins. *Yeast* 10(8), 1049-1064.
- 20 [24] Lowry, O.H., Rosenbrough, N.J., Farr, A.L. and Randall, R.J. (1951) Protein measurement with
21 the Folin phenol reagent. *J. Biol. Chem.* 193, 265-275.
- 22 [25] Prudêncio, C., Sansonetty, F. and Côte-Real, M. (1998) Flow cytometric assessment of cell
23 structural and functional changes induced by acetic acid in the yeasts *Zygosaccharomyces bailii*
24 and *Saccharomyces cerevisiae*. *Cytometry* 31(4), 307-313.
- 25 [26] Buege, J.A. and Aust, S.D. (1978) Microsomal lipid peroxidation. *Methods Enzymol.* 52, 302-310.
- 26 [27] Krems, B., Charizanis, C. and Entian, K.D. (1995) Mutants of *Saccharomyces cerevisiae* are
27 involved in oxidative stress resistance. *Curr. Genet.* 27, 427-434.
- 28 [28] Diniz-Mendes, L., Bernardes, E., Araujo, P.S., Panek, A.D. and Paschoalin, V.M.F. (1999)
29 Preservation of frozen yeast cells by trehalose. *Biotechnol. Bioeng.* 65(5), 572-578.

- 1 [29] Benaroudj, N., Lee, D. H. and Goldberg, A. L. (2001) Trehalose accumulation during cellular
2 stress protects cells and cellular proteins from damage by oxygen radicals. *J. Biol. Chem.* 276,
3 24261–24267.
- 4 [30] Crowe, J.H., Crowe, L.M. and Chapman, D. (1984) Preservation of membranes in anhydrobiotic
5 organisms: the role of trehalose. *Science* 223: 701-703.
- 6 [31] Crowe, J.H., Carpenter, J.F. and Crowe, L.M. (1998) The role of vitrification in anhydrobiosis.
7 *Ann. Rev. Physiol.* 60, 73-103.
- 8 [32] Murakami, Y., Yokoigawa, K. and Kawai, H. (1995) Lipid composition of a freeze-tolerant yeast,
9 *Torulaspota delbrueckii*, and its freeze-sensitive mutant. *Appl. Microbiol. Biotechnol.* 44, 167-171.
- 10 [33] Sahara, T., Goda T. and Ohgiya, S. (2002) Comprehensive expression analysis of time-
11 dependent genetic responses in yeast cells to low temperature. *J. Biol. Chem.* 277(51), 50015-
12 50021.
- 13 [34] Rodriguez-Vargas, S., Estruch, F. and Randez-Gil, F. (2002) Gene expression analysis of cold
14 and freeze stress in baker's yeast. *Appl. Environ. Microbiol.* 68(6), 3024-3030.
- 15
16

Fig. 1. Relative values (%) of colony forming units (CFU) in stationary growth phase cells of *T. delbrueckii* (A), and *S. cerevisiae* (B) during frozen storage at $-20\text{ }^{\circ}\text{C}$, for 120 and 30 days, respectively. Cells were grown in YPS medium during 24 hours at $30\text{ }^{\circ}\text{C}$, harvested, washed with water and suspended in different storage media: (\square) LF medium; (\bullet) LF medium + 0.01% cycloheximide; (\circ) LF medium without sugars + 0.16 M glycerol; (Δ) Sugars solution (1% glucose + 1% sucrose + 3% maltose).

Fig. 2. Relative values (%) of negative propidium iodide cells (PI⁻ cells) of *T. delbrueckii* (close symbols) and *S. cerevisiae* (open symbols) during storage at $-20\text{ }^{\circ}\text{C}$ (A) and correlation between relative values (%) of PI⁻ cells and colony forming units (CFU) (B). Cells grown in YPS medium during 24 hours at $30\text{ }^{\circ}\text{C}$ (initial stationary growth phase), harvested, washed with water and suspended in LF medium were frozen directly (\blacktriangle , \triangle) or after a pre-fermentation period of 120 min (\bullet , \circ), as described in material methods. In Fig. 2A, for some results error bars are within the data point labels.

Fig. 3. Effect of oxidising agents on the radial growth of *T. delbrueckii* (black bars) and *S. cerevisiae* (open bars). Cells were grown in YPS medium at $30\text{ }^{\circ}\text{C}$, to mid-exponential growth phase (B) or to initial stationary growth phase (A), and 0.1 ml of a suspension ($\text{OD}_{640\text{ nm}}$ 0.6-0.7) of these cells was spread on YPDA plates. A paper disc, 6 mm diameter, containing 10 μl of one oxidising agent was laid on each inoculated plate. The plates were incubated during two days, after which the diameters of the inhibition halos were measured. The values shown are those obtained for the highest concentration of oxidant tested.

Fig. 4. Cells of *T. delbrueckii* (A) and *S. cerevisiae* (B) were grown in YPS medium during 24 hours at $30\text{ }^{\circ}\text{C}$, harvested, washed with water, suspended LF media and frozen at $-20\text{ }^{\circ}\text{C}$. During freezing samples were taken and assessed for: (\circ) relative values (%) of negative propidium iodide cells (PI⁻ cells); (\blacktriangle) relative values (%) of intracellular trehalose concentration (T_{in}); (\square) extracellular trehalose concentration (T_{out} , g/l); (\bullet) medium optical density at 260 nm, used as a measure of leakage of cell contents ($\text{OD}_{260\text{ nm}}$) The relative values (%) were estimated as described in material and methods.