

Synthesis of formyl-thienylpyrroles: versatile building blocks for NLO materials

M. Manuela M. Raposo,^{a*} Ana M. R. C. Sousa,^a A. Maurício C. Fonseca^a and G. Kirsch^b

^a Centro de Química, Universidade do Minho, Campus de Gualtar
4710-057 Braga, Portugal

^b Laboratoire d'Ingénierie Moléculaire et Biochimie Pharmacologique,
UFR SciFA, Université de Metz, 1, Boulevard Arago, Metz Technopôle,
57078 Metz Cedex 3, France

Abstract - Several formyl-substituted 1-alkyl(aryl)-2-(2'-thienyl)pyrroles **3-7** were synthesized by functionalization of the pyrrole or thiophene ring of thienylpyrroles **2** using different methods: Vilsmeier formylation or metalation followed by reaction with DMF.

Keywords: formyl-substituted thienylpyrroles, reactivity studies, Vilsmeier formylation, $m\text{-}[\text{C}_6\text{H}_4]_2\text{N}$, UV-Visible spectroscopy.

1. Introduction

Formylation is a key process in organic synthesis, with the resulting aldehyde function being a “crossroads” intermediate. Not surprisingly, a large number of methods have been developed for this reaction. Reagents for electrophilic formylation are mostly of the form Y-CH=X^+ . Thus the reactions attributed to Vilsmeier (ClCH=NR_2^+), Rieche (*e.g.* $\text{MeOCHCl}_2 \rightarrow \text{MeO=CHCl}^+$), Gatterman ($\text{Zn}[\text{CN}]_2/\text{HCl} \rightarrow \text{HC=NH}_2^{2+}$), Gatterman-Koch ($\text{CO/HCl/Lewis acid} \rightarrow \text{HC=O}^+$) and even Duff ($\text{CH}_2=\text{NH}_2^+$ - followed by dehydrogenation of initially formed RCH_2NH_2) all fit this pattern.¹

Organolithiums are beyond any doubt the most useful metalated heterocycles. Usually they are prepared by direct deprotonation of acidic hydrogens using strong bases or, particularly useful in the case of the less acidic sites in aromatic rings, by halogen exchange between a halogenated heterocycle and an organolithium compound or

*Corresponding author. Phone: +351 253 604381; Fax: +351 253 678983; e-mail: mfox@quimica.uminho.pt

lithium metal. Another frequent alternative is the so-called *ortho*-lithiation or “directed *ortho*-metalation” (DoM) which is the metalation of an aromatic ring adjacent to a heteroatom-containing functional group by providing the lithium base with a point of coordination, thus increasing reactivity close to the coordination site. The lithiated species generated by all these methods are able to react with all kinds of electrophiles.²⁻³

Vilsmeier formylation and metalation followed by quenching with DMF constitute the most significant routes for the preparation of formyl-substituted pyrroles and thiophenes.⁴⁻⁵

The formyl- derivatives obtained can further react to afford more complex molecules which have made of formyl-thiophenes and formyl-pyrroles some of the most important molecules to be used as building blocks in biological active compounds, supramolecular chemistry and molecular electronics.⁶⁻²⁵

During the last years we have been concerned with the studies of several organic and organometallic compounds such as oligothiophenes, thienylpyrroles, thienylphthalazines, thienyl and bithienyl- Mo complexes bearing donor and electron acceptor units motivated by their potential applications in optical and electronic devices.²⁵⁻³⁴ Recently, we have investigated donor-acceptor substituted thienylpyrroles. Due to their solvatochromic, electrochemical and non-linear optical properties, donor-acceptor thienylpyrrole derivatives could be used for the manufacture of semi-conductor materials or materials with strong non-linear optical (NLO) properties.²⁸⁻³¹ Conjugated 1-(alkyl)aryl-2-(2'-thienyl)pyrroles, as strong π -electron donor moieties functionalized with the formyl- group on the thiophene or on the pyrrole moiety, could be used as precursors in the preparation of functional π -conjugated systems with several applications (e.g. NLO).

As part of our continuing interest in non-linear optical material, in this paper we describe the synthesis of formyl-thienylpyrroles, **3-7** prepared from 1-propyl-2-(2'-thienyl)pyrrole **2a** and 1-aryl-2-(2'-thienyl)pyrroles **2b-f**.³⁵

2. Results and discussion

2.1. Synthesis

2.1.1. Synthesis of 1-*n*-propyl-2-(2'-thienyl)pyrrole **2a**

In order to compare the reactivity of 1-alkyl- and 1-aryl-2-(2'-thienyl)pyrroles through the Vilsmeier reaction and through the metalation followed by quenching with DMF, 1-propyl-2-(2'-thienyl)pyrrole **2a** was synthesized from *N*-propyl-4-(2'-thienyl)-4-oxobutanamide **1a** using a combination of the Friedel-Crafts and the Lawesson reactions.³⁵⁻³⁶ Direct amidation of 4-oxo-(2'-thienyl)butanoic acid³⁶ with propylamine was carried out through DCC-BtOH mediated reaction. Amide **1a** was obtained as a colourless solid in good yield (80%).

Recently we have demonstrated that the reaction of secondary aryl-(2'-thienyl)-4-oxobutanamides with Lawesson's reagent (LR) can yield thienylpyrroles and/or bithiophenes in different ratios depending on the substituent(s) of the precursor arylamines.³⁵ In the case of the secondary *n*-propyl amide **1a**, treatment with an equimolar amount of LR gave only the thienylpyrrole **2a** in 47% yield.

2.1.2. Vilsmeier formylation

Electrophilic substitution reactions of thienylpyrroles were found to be very selective. According to earlier reports, the pyrrole nitrogen atom has a greater ability to delocalize the positive charge of σ -complexes than the sulfur atom in thiophene; pyrrole is

therefore considerably more reactive towards electrophilic substitution than thiophene. Even when both α -positions of the pyrrole ring are occupied, electrophilic substitution will preferentially occur in the β -position of the pyrrole ring rather than the α -position of the thiophene ring.^{4,37-40} The reactivity of these systems has been demonstrated with the use of electrophilic reactions producing derivatives with the electrophile substituted primarily on the pyrrole ring.^{28-31,39-46}

To our knowledge, there is only one previous study describing the formylation of thienylpyrroles and this study was performed through the Vilsmeier-Haack reaction on the simple 2-(2'-thienyl)pyrrole and 2-(3'-thienyl)pyrrole. Bouka *et al* obtained exclusively derivatives formylated on the α -position of the pyrrole ring.⁴⁰

Therefore, we decided to study the reactivity of different thienylpyrroles bearing *N*-alkyl or *N*-aryl groups on the pyrrole ring through the Vilsmeier-Haack reaction and metalation followed by reaction with DMF.

Accordingly, Vilsmeier formylation of thienylpyrroles **2a-c** proceeded selectively in the pyrrole ring to form the corresponding formyl-substituted thienylpyrroles **3-4**.

In our studies of Vilsmeier-Haack formylation of 1-propyl-2-(2'-thienyl)pyrrole **2a**, the 5-position of the pyrrole ring was found to be much more reactive than the 3-position. The Vilsmeier-Haack formylation of **2a**, with DMF/POCl₃ at 60 °C for 2 h produced a mixture of 5-formyl- **3a** (63%), and 3-formyl- derivative **4a**, in lower yield (5%) (Table 1, entries 1-2). Under the same experimental conditions 1-aryl-2-(2'-thienyl)pyrroles **2b-c** behave quite differently giving a mixture of the 5- and 3-formyl- derivatives in similar yields. Formylation of **2b** gave a mixture of **3b** (19%) and **4b** (22%), (Table 1, entries 3-4) and formylation of **2c** gave a mixture of **3c** (12%) and **4c** (10%). In comparison to alkylpyrrole **2a** the formyl derivatives of 1-aryl-2-(2'-thienyl)pyrroles **2b-c** were obtained in lower yields.

<Scheme 1>

<Table 1>

In order to interpret the results obtained we consider several factors: (i) an appreciably larger nucleophilicity of the pyrrole ring compared to that of thiophene; (ii) a decrease in the electron density in the pyrrole ring due to the competitive conjugation of the *N*-aryl group with the unshared electron pair on nitrogen and also due to its negative inductive effect; (iii) possible steric influence due to the *n*-propyl group for attack at the α -position of the pyrrole ring.⁴⁴

3-Substituted pyrroles are the most difficult to synthesize since most electrophilic aromatic substitution reactions and lithiation reactions of *N*-substituted pyrroles occur at the 2-position and so functionalization at the 3-position of pyrrole is a challenging goal in synthetic research. Bulky substituents on the nitrogen atom promote 3-substitution. This observation led to new approaches to the synthesis of 3-substituted pyrroles, as these are normally found only as by products in reactions leading predominantly to 2-substitution.⁴⁷⁻⁴⁹ As 3-formyl *N*-arylpyrroles are key synthetic intermediates to highly biologically active compounds the preparation of new derivatives, even in fair yields, still remains an attractive goal.^{10-11,44,46}

The structures of formyl-substituted pyrrole derivatives **3-4** were unambiguously confirmed by their analytical and spectral data. In the ¹H NMR spectrum of 5-formyl-substituted pyrrole derivatives **3a-c** two signals at about 6.40-6.64 and 6.96-7.17 ppm were detected. Both signals appear as doublets with coupling constants of 4.2-4.5 Hz indicating the presence of two adjacent protons (3-H and 4-H) at the corresponding pyrrole moiety. In the ¹H NMR spectrum of 3-formyl-substituted pyrrole derivatives **4a-c** two signals at about 6.47-6.79 and 6.78-7.19 ppm were detected. Both signals appear as doublets with coupling constants of 3.0-3.3 Hz. These signals were attributed

to the 5-H and 4-H in the pyrrole moiety. The position of the formyl group in pyrrole derivatives **4a-c** was also confirmed through the analysis of 2D NOESY spectra. Thus, the 2D NOESY spectra of compounds **4a-c** shows a cross peak of the pyrrole ring 4-H with the formyl group and also a cross peak of the formyl group with the thiophene ring 3'-H. On the contrary, the 2D NOESY spectrum of compound **4a** shows a cross peak of the NCH₂ protons of the *n*-propyl group with the 5-H proton of the pyrrole ring but there is no cross peak of the pyrrole ring 5-H with the formyl group. This is taken as evidence for the attachment of the formyl group at ring position 3. In all the ¹H NMR spectra of formyl-substituted pyrrole derivatives **3a-c** and **4a-c** three signals at about 6.90-7.20 (multiplet), 6.72-7.20 (double doublet) and 7.19-7.63 (double doublet) were detected. These signals were attributed respectively, to the 4', 3' and 5'-H protons in the thiophene moiety.

2.1.3 Metalation followed by reaction with DMF

As (5'-formyl-2'-thienyl)pyrroles could not be synthesized solely by the Vilsmeier-Haack reaction, we tried to prepare these compounds by lithiation followed by treatment with DMF.

The electron-rich five member aromatic *N*-substituted pyrrole, furan and thiophene are lithiated at C-2 by direct deprotonation with a lithium-containing base. Several authors have reported the α -lithiation of *N*-arylpyrroles using different experimental conditions: *n*-BuLi-TMEDA chelate, *n*-BuLi-*t*BuOK (LiCKOR) superbases, *ter*-BuLi-secondary amides, Na/dry ether/0 °C /*ter*-BuLi, *ter*-BuLi/- 78 °C/THF, *n*-BuLi/THF/- 78 °C, *n*-BuLi-TMEDA/THF/- 75 °C.⁴⁹⁻⁵⁴

Recently, novel methods for site-selective lithiation in α or benzylic positions of 1-(methoxyphenyl), 1-(chlorophenyl), 1-(bromophenyl), 1-(trifluoromethylphenyl) and 1-(methylphenyl)pyrroles have also been reported.⁵⁵⁻⁵⁹

To our knowledge, the formylation of thienylpyrroles through lithiation followed by quenching with DMF has not been previously reported, and success would open the way to a new range of formyl-functionalized thienylpyrroles.

The metalation of thienylpyrroles **2** was carried out with *n*-BuLi in dry ether at 0 °C for 1 h. Subsequently, the organolithium derivatives were converted to the corresponding formyl compounds, by addition of DMF followed by refluxing the mixture for 1h (Scheme 2, Table 2). In order to compare the reactivity of pyrroles **2** under the experimental conditions described above the reaction time studied for all derivatives (for the metalation and for the reaction with DMF) was 1h. As a consequence of the limited reaction time, in some cases, unreacted starting materials remained in the reaction mixtures.

Through this method thienylpyrroles **2a-b** and **2e-f** were selectively lithiated at the α -position of the thiophene ring giving formyl-derivatives **5a-b** and **5e-f**.

The methoxy group is known as a moderately strong *ortho* directing substituent with electron withdrawal and electron donor properties.^{3,55,60} At the same time, for compounds **2c-d** the 4-methoxy and the 2,4-dimethoxy group(s) have an α -directing effect in the aromatic ring. Consequently, the formylation of the aromatic ring was also observed for thienylpyrroles **2c-d** (Table 2, entries 4, 6 and 7, compounds **6c-d**, **7d**) due to the *ortho* directing effect of the methoxy groups (Scheme 2) giving a mixture of several formylated derivatives with 2-(5'-formyl-2'-thienyl)pyrroles **5c-d** being the major products.

For compound **2d** we studied also the effect of the reaction time for the metalation step and for the reaction with DMF. Metalation of thienylpyrrole **2d** for 2h followed by 2 h of reflux with DMF gave a mixture of 4 compounds: thienylpyrrole **2d** (18%), 1-(2'',4''-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole **5d** (34%), 1-(3''-formyl-

2'',4''-dimethoxyphenyl)-2-(2'-thienyl)pyrrole **7d** (14%) and 1-(3''-formyl-2'',4''-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole **6d** (18%) (Table 2, entries 5-7). The experiment showed that, instead of improving the yield of compound **5d** we obtained the diformylated compound **6d** and a higher yield for the formyl-aryl derivative **7d** as a result of the increase of the reaction time.

Compounds **5e** and **5f** were obtained but in lower yields. It should be recorded that we could not isolate any benzylic formyl product from these reactions.

¹H NMR spectra of (5'-formyl-2'-thienyl)pyrrole derivatives **5a-f** showed two signals at about 6.61-7.12 and 7.49-7.71 ppm. Both signals appear as doublets with coupling constants of 3.9-4.2 Hz indicating the presence of two adjacent protons (3'-H and 4'-H) at the corresponding thiophene moiety. In all the ¹H NMR spectra of formyl-substituted thiophene derivatives **5a-f** three signals at about 6.20-6.42 (multiplet) 6.53-6.77 (double doublet), and 6.82-6.97 (multiplet) were detected. These signals were attributed respectively, to the 4, 3 and 5-H protons in the pyrrole moiety.

<Scheme 2>

<Table 2>

2.2. UV-visible study of formyl-substituted thienylpyrroles

The electronic spectra of formyl-thienylpyrrole derivatives were recorded in ethanol (Tables 1-2).

All the formyl-substituted thienylpyrroles **3-7** synthesized exhibit intense absorptions in the UV-visible range. The position of these absorptions is influenced by the structure of the compounds, for example by the substituent on the nitrogen atom of the pyrrole ring

and by the position of substitution of the formyl group on the thiophene, pyrrole or on the aromatic ring(s).

Communication between the electron donating and accepting termini can be evaluated by comparing the λ_{\max} values. Dramatic differences in energy occur upon formyl-substitution of thienylpyrroles **2**. For example, thienylpyrrole **2d** ($\lambda_{\max} = 286.5$ nm) is shifted 98 nm upon formyl substitution (formyl- derivative **5d**, $\lambda_{\max} = 384.5$ nm) (Table 2, entry 5). This effect has been attributed to the stabilization of LUMO by the electron-withdrawing groups.⁶¹ The influence of the substituent on the nitrogen atom of the pyrrole ring is demonstrated by comparison of the absorption maxima of compounds **5a** and **5d** as the longest wavelength transition is shifted from 374.0 nm in 1-(*n*-propyl)-2-(5'-formyl-2'-thienyl)pyrrole **5a** (Table 2, entry 1) to 384.5 nm for 1-(2',4'-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole **5d** (Table 1, entry 5). The influence of the position of the formyl group on the pyrrole ring on λ_{\max} of absorption for formyl derivatives **3** and **4** is noteworthy. The difference in λ_{\max} values ($\Delta\lambda_{\max}$) between compounds (**3a-b** and **4a-b**) is in the range of 15-28 nm (Table 1, entries 1-4). As expected, the presence of the formyl group on the 5-position of the pyrrole ring (**3a-3b**), relative to the same acceptor group in the 3-position (**4a-4b**), results in a bathochromic shift in the λ_{\max} of absorption for **3a-3b** due to more extensive electron delocalization.

3. Conclusions

Starting from the readily available thienylpyrroles **2**, commercial reagents and simple and convenient procedures were used to synthesize new formyl-, thienylpyrroles in fair to good yields, *via* two methods: Vilsmeier formylation and lithiation followed by reaction with DMF.

Vilsmeier-Haack formylation of **2** was made at the 3- and 5-positions on the pyrrole ring to give compounds **3-4**. These results are in accordance with the greater nucleophilicity of the pyrrole ring *versus* the thiophene ring as has been shown earlier in the case of the tricyanovinyl reaction and azo coupling reaction of 1-(alkyl)aryl-2-(2'-thienyl)pyrroles **2**.²⁸⁻²⁹

The lithiation of thienylpyrroles **2a-b** and **2e-f**, followed by quenching with DMF, occurred selectively on the α - position of the thiophene ring giving formyl derivatives **5a-b** and **5e-f**. For compounds **2c-d** with methoxy group(s) on the aryl ring was obtained a mixture of formyl-derivatives **5c-d**, **6c-d** and **7d** with the formyl group on the thiophene ring and/or on the aryl ring. The major compounds formed were (5'-formyl-2'-thienyl)pyrroles **5c-d**.

The formyl- derivatives **3-7** studied exhibit an absorption band in the UV-visible range influenced by the structure of the compounds: the type of substituent on the nitrogen atom of the pyrrole ring and also by the position of the formyl group on the thiophene or on the pyrrole ring.

The conjugated formyl derivatives of 1-alkyl(aryl)-2-(2'-thienyl)pyrroles will be used in the future, as precursors in the preparation of compounds with a stronger electronwithdrawing group for potential applications in NLO.^{22,25}

4. Experimental

4.1. General

¹H NMR spectra were obtained on a Varian Unity Plus Spectrometer at 300 MHz and ¹³C NMR spectra were determined on a Varian Unity Plus Spectrometer at 75.4 MHz using the solvent peak as internal reference. The solvents are indicated in parenthesis before the chemical shift values (δ relative to TMS). Mps were determined on a Gallenkamp apparatus and are uncorrected. Infrared spectra were recorded on a Perkin

Elmer 1600 FTIR spectrophotometer. UV-visible absorption spectra were obtained using a Shimadzu UV/2501PC spectrophotometer. EI mass spectra EI (70 eV) and HRMS were run on a Unicam GC-MS 120. Elemental analysis was carried out on a Leco CHNS-932. Column chromatography was performed on Merck silica gel 60 (Art 9385). Light petroleum refers to solvent boiling in the range 40-60 °C.

The synthesis of 1-aryl-2-(2'-thienyl)pyrroles **2b-g** has been described elsewhere.³⁵

4.2. Synthesis of 1-propyl-2-(2'-thienyl)pyrrole **2a**

(i) *Synthesis of N-propyl-4-(2'-thienyl)-4-oxobutanamide **1a***. Amide **1a** was obtained using the experimental method described in refs. 35-36, by reacting 4-oxo-(2-thienyl)butanoic acid (5.4 mmol) in CH₂Cl₂ with 1,3-dicyclohexylcarbodiimide (7.1 mmol) and BtOH (7.1 mmol) and adding propylamine (5.4 mmol) at rt.

N-Propyl-4-(2'-thienyl)-4-oxobutanamide **1a**. Colourless solid (80%). Mp 96.2-97.6 °C (EtOH). IR (liquid film) ν 3311 (NH), 3029, 1661 (C=O), 1645 (C=O), 1552, 1523, 1420, 1397, 1248, 1179, 1063, 983, 954, 910, 851, 717 cm⁻¹. ¹H NMR (CDCl₃) δ 0.92 (t, 3H, *J*=7.2 Hz, CH₃), 1.52 (m, 2H, CH₂CH₃), 2.61 (t, 2H, *J*=7.8 Hz, CONHCH₂), 5.78 (br s, 1H, NH), 7.13-7.15 (m, 1H, 4'-H) 7.65 (dd, 1H, *J*=5.1 and 1.2 Hz, 5'-H), 7.78 (dd, 1H, *J*=3.7 and 1.2 Hz, 3'-H). ¹³C NMR (CDCl₃) δ 11.30, 22.76, 30.26, 34.62, 41.27, 128.16, 132.23, 133.73, 143.59, 171.71, 192.09. Anal. Calcd for C₁₁H₁₅NO₂S: C, 58.63; H, 6.72; N, 6.22; S, 14.23. Found: C, 58.63; H, 6.71; N, 6.23; S, 13.98.

(ii) *Reaction of amide **1a** with Lawesson's Reagent*. Thienylpyrrole **2a** was obtained using the experimental method described in refs. 35-36, by heating the amide **1a** (2.3 mmol) in toluene (12 ml) with the Lawesson reagent (2.3 mmol) at reflux during 30 min..

1-Propyl-2-(2'-thienyl)pyrrole **2a**. Orange oil (47%). UV (EtOH): λ_{max} nm (ϵ /M⁻¹ cm⁻¹) 291.0 (1803), 225.5 (2054). IR (liquid film) ν 3102, 2964, 2932, 2874, 1508, 1470, 1430, 1383, 1345, 1299, 1234, 1201, 1108, 1070, 941, 896, 844, 783, 711, 613

cm⁻¹. ¹H NMR (CDCl₃) δ 0.92 (t, 3H, *J*=7.2 Hz, (CH₂)₂CH₃), 1.65-1.68 (m, 2H, CH₂CH₂CH₃), 3.97 (t, 2H, *J*=7.2 Hz, NCH₂), 6.17-6.22 (m, 1H, 4-H), 6.31 (dd, 1H, *J*=3.6 and 1.8 Hz, 3-H), 6.76-6.88 (m, 1H, 5-H) 7.01 (dd, 1H, *J*=3.6 and 1.2 Hz, 3'-H), 7.06-7.09 (m, 1H, 4'-H), 7.29 (dd, 1H, *J*=5.1 and 1.2 Hz, 5'-H). ¹³C NMR (CDCl₃) δ 11.17, 24.74, 48.99, 107.71, 110.16, 122.68, 124.71, 125.29, 126.28, 127.16, 134.97. MS (EI) *m/z* (%): 191 (M⁺, 100), 162 (50), 149 (34), 130 (7), 121 (13), 111 (40), 104 (15). HRMS: *m/z* (EI) for: C₁₁H₁₃NS; calcd 191.0768; found: 191.0763.

4.3. General procedure for the synthesis of formyl derivatives 3-4 of 1-(alkyl)aryl-2-(2'-thienyl)pyrroles 2a-c through Vilsmeier formylation

POCl₃ (0.46 mmol) was added to DMF (0.46 mmol) at 0 °C and the mixture was stirred for 15 min. at 0 °C. After this time pyrroles **2** (0.39 mmol) dissolved in DMF (1 ml) were added dropwise with stirring. The reaction mixture was then heated 2 h at 60 °C. The solution was then poured slowly into 5 ml saturated sodium acetate aqueous solution and stirred 30 min.. The organic layer was diluted with ether, washed with saturated NaHCO₃ aqueous solution, and dried with anhydrous MgSO₄. Evaporation of the organic extract under reduced pressure gave a mixture of 5-formyl- **3** and 3-formylpyrroles **4** which were purified by "flash" chromatography on silica with increasing amounts of ether in light petroleum as eluent.

Vilsmeier formylation of **2a** gave a mixture of 5-formyl-1-propyl-2-(2'-thienyl)pyrrole **3a** and 3-formyl-1-propyl-2-(2'-thienyl)pyrrole **4a**. The first component eluted was 5-formyl-1-propyl-2-(2'-thienyl)pyrrole **3a** as a green oil (63%). UV (EtOH): λ_{max} nm (ε/M⁻¹ cm⁻¹), 321.5 (24650). IR (liquid film): ν 2964, 1658 (C=O), 1509, 1473, 1428, 1396, 1314, 1294, 1251, 1225, 1198, 1154, 1042, 847, 776, 702 cm⁻¹. ¹H NMR (CDCl₃) δ 0.90 (t, 3H, *J*=7.5Hz, (CH₂)₂CH₃), 1.70-1.82 (m, 2H, CH₂CH₂CH₃), 4.41 (t, 2H, *J*=7.8 Hz, NCH₂), 6.40 (d, 1H, *J*=4.2 Hz, 3-H), 6.96 (d, 1H, *J*=4.2 Hz, 4-H), 7.12-7.16 (m, 1H, 4'-H), 7.18 (dd, 1H, *J*=3.6 and 1.2 Hz, 3'-H), 7.44 (dd, 1H, *J*=5.1 and 1.2 Hz, 5'-H), 9.54 (s, 1H, CHO). ¹³C NMR (CDCl₃) δ 10.80, 24.68, 47.50, 111.89, 124.73 (two overlapped signals), 127.00, 127.42, 127.59, 132.17, 132.51, 136.36. MS (EI) *m/z* (%): 219 (M⁺, 100), 218 (21), 204 (16), 202 (74), 190 (28), 177 (80), 176 (77), 162 (18), 148 (16), 121 (30), 104 (7). HRMS: (EI) *m/z* (%) for C₁₂H₁₃NOS; calcd

219.0718; found 219.0720. The second component eluted was *3-formyl-1-propyl-2-(2'-thienyl)pyrrole* **4a** as a brown oil (5%). UV (EtOH): λ_{\max} nm ($\epsilon/M^{-1} \text{ cm}^{-1}$), 293.5 (5440), 248.0 (10388), 211.0 (8678). IR (liquid film): ν 1662 (C=O), 1485, 1446, 1382, 1257, 849, 765, 706 cm^{-1} . ^1H NMR (CDCl_3) δ 0.87 (t, 3H, $J=7.5$ Hz, $(\text{CH}_2)_2\text{CH}_3$), 1.65-1.82 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 3.87 (t, 2H, $J=7.2$ Hz, NCH_2), 6.47 (d, 1H, $J=3.3$ Hz, 5-H), 6.78 (d, 1H, $J=3.3$ Hz, 4-H), 7.13-7.20 (m, 2H, 3' and 4'-H), 7.54 (dd, 1H, $J=5.1$ and 1.2 Hz, 5'-H), 9.54 (s, 1H, CHO). MS (EI) m/z (%): 219 (M^+ , 100), 218 (19), 190 (12), 174 (11), 162 (24), 149 (10), 130 (4), 121 (6), 104 (7), 89 (5). HRMS: (EI) m/z (%) for $\text{C}_{12}\text{H}_{13}\text{NOS}$; calcd 219.0718; found 219.0708.

Vilsmeier formylation of **2b** gave a mixture of *5-formyl-1-phenyl-2-(2'-thienyl)pyrrole* **3b** and *3-formyl-1-phenyl-2-(2'-thienyl)pyrrole* **4b**. The first component eluted was *5-formyl-1-phenyl-2-(2'-thienyl)pyrrole* **3b** as a pale yellow solid (19%). Mp: 72.6-73.9 °C. UV (EtOH): λ_{\max} nm ($\epsilon/M^{-1} \text{ cm}^{-1}$), 340.5 (20100), 253.0 (8630). IR (liquid film): ν 1660 (CHO), 1596, 1511, 1497, 1469, 1433, 1406, 1361, 1318, 1046, 847, 775, 695, 547 cm^{-1} . ^1H NMR (CDCl_3) δ 6.64 (d, 1H, $J=4.2$ Hz, 3-H), 6.72 (dd, 1H, $J=3.9$ and 1.2 Hz, 3'-H), 6.86-6.90 (m, 1H, 4'-H), 7.17 (d, 1H, $J=4.2$ Hz, 4-H), 7.19 (dd, 1H, $J=5.1$ and 1.2 Hz, 5'-H), 7.33-7.39 (m, 2H, 2xAr-H), 7.49-7.56 (m, 3H, 3xAr-H) 9.39 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 109.72, 110.96, 120.51, 126.43, 126.62, 127.21, 128.87, 129.34, 129.46, 132.66, 136.45, 137.15, 178.76. MS (EI) m/z (%): 253 (M^+ , 100), 225 (12), 175 (11), 147 (10), 121 (20), 77 (12). HRMS: (EI) m/z (%) for $\text{C}_{15}\text{H}_{11}\text{NOS}$; calcd 253.0561; found 253.0558. The second component eluted was *3-formyl-1-phenyl-2-(2'-thienyl)pyrrole* **4b** as pale yellow solid (22%). Mp: 106.9-107.8 °C. UV (EtOH): λ_{\max} nm ($\epsilon/M^{-1} \text{ cm}^{-1}$) 325.0 (7440) IR (liquid film): ν 1661 (CHO), 1596, 1496, 1473, 1449, 1413, 1241, 848, 762, 695 cm^{-1} . ^1H NMR (Acetone- d_6) δ 6.79 (d, 1H, $J=3.0$ Hz, 5-H), 7.10-7.13 (m, 1H, 4'-H), 7.15-7.19 (m, 2H, 4-H and 3'-H), 7.30-7.36 (m, 2H, 2xAr-H), 7.43-7.49 (m, 3H, 3xAr-H), 7.63 (dd, 1H, $J=5.1$ and 1.2 Hz, 5'-H), 9.82 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 108.14, 110.95, 125.44, 126.27, 127.09, 128.20, 128.48, 129.18, 130.61, 132.66, 134.92, 138.60, 186.76. MS (EI) m/z (%): 253 (M^+ , 100), 224 (19), 209 (14), 180 (5), 121 (11), 77 (22). HRMS: (EI) m/z (%) for $\text{C}_{15}\text{H}_{11}\text{NOS}$; calcd 253.0561; found 253.0576.

Vilsmeier formylation of **2c** gave a mixture of 5-formyl-1-(4'-methoxyphenyl)-2-(2'-thienyl)pyrrole **3c** and 3-formyl-1-(4'-methoxyphenyl)-2-(2'-thienyl)pyrrole **4c**. The first component eluted was 5-formyl-1-phenyl-2-(2'-thienyl)pyrrole **3c** as a yellow solid (12%). Mp: 110.5-111.9 °C. UV (EtOH): λ_{max} nm ($\epsilon/M^{-1} \text{ cm}^{-1}$), 343.0 (21559), 257.5 (8165), 226.5 (14668). IR (liquid film): ν 1655 (CHO), 1512, 1495, 1395, 1337, 1319, 1248, 1230, 1193, 1162, 1103, 1052, 832, 805, 779, 765, 709 cm^{-1} . ^1H NMR (CDCl_3) δ 3.89 (s, 3H, OCH_3), 6.63 (d, 1H, $J=4.5$ Hz, 3-H), 6.80 (dd, 1H, $J=3.8$ and 1.2 Hz, 3'-H), 6.88-6.92 (m, 1H, 4'-H), 7.00 (d, 2H, $J=9.0$ Hz, 2xAr-H), 7.15 (d, 1H, $J=4.5$ Hz, 4-H), 7.19 (dd, 1H, $J=5.1$ and 1.2 Hz, 5'-H), 7.26 (d, 2H, $J=9.0$ Hz, 2xAr-H), 9.38 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 55.49, 110.58, 114.45, 120.38, 126.41, 126.51, 127.16, 129.54, 129.88, 132.71, 134.76, 136.69, 160.17, 178.88. MS (EI) m/z (%): 283 (M^+ , 100), 254 (19), 240 (30), 175 (48), 147 (18), 121 (16), 108 (6). HRMS: (EI) m/z (%) for $\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$; calcd 283.0667; found 283.0664. Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$: C, 67.83; H, 4.69; N, 4.94; S, 11.33. Found: C, 67.58; H, 4.96; N, 4.91; S, 10.82. The second component eluted was 3-formyl-1-(4'-methoxyphenyl)-2-(2'-thienyl)pyrrole **4c** as pale yellow solid (10%). Mp: 99.7-101.0 °C. IR (liquid film): ν 1661 (CHO), 1512, 1449, 1300, 1242, 1028, 835, 763 cm^{-1} . ^1H NMR (Acetone- d_6) δ 3.86 (s, 3H, OCH_3), 6.76 (d, 1H, $J=3.0$ Hz, 5-H), 7.00 (d, 2H, $J=9.0$ Hz, 2xAr-H), 7.08 (dd, 1H, $J=3.0$ and 1.0 Hz, 4-H), 7.09-7.13 (m, 1H, 4'-H), 7.16 (dd, 1H, $J=3.4$ and 1.2 Hz, 3'-H), 7.25 (d, 2H, $J=9.0$ Hz, 2xAr-H), 7.61 (dd, 1H, $J=5.1$ and 1.2 Hz, 5'-H), 9.81 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 55.46, 107.85, 114.24, 125.43, 125.68, 127.03, 127.55, 128.40, 129.54, 129.90, 130.51, 131.47, 135.23, 186.71. MS (EI) m/z (%): 283 (M^+ , 100), 254 (13), 240 (26), 210 (5), 175 (11), 121 (8). HRMS: (EI) m/z (%) for $\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$; calcd 283.0667; found 283.0666. Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$: C, 67.83; H, 4.59; N, 4.94; S, 11.33. Found: C, 67.64; H, 4.98; N, 4.91; S, 11.03.

4.4. General procedure for the synthesis of formyl- derivatives of 1-(alkyl)aryl-2-(2'-thienyl)pyrroles **2a-f**, via metalation with *n*-BuLi followed by reaction with DMF

A 2.5 M solution of *n*-BuLi in hexanes (0.44 ml, 1.1 mmol) was dropped under Ar at 0° C to a stirred solution of thienylpyrroles **2** (1.0 mmol) in anhydrous ether. The reaction mixture was then stirred 1 h at 0° C and was allowed to stand 15 min. at room

temperature. DMF (0.05 ml, 1.0 mmol) dissolved in anhydrous ether (2 ml) was added dropwise at r.t. The mixture was heated at reflux for 1 h. The mixture was poured into water (20 ml) and extracted with (3 x 50 ml) of ethyl acetate. The combined organic extracts were washed with H₂O (100 ml), dried with MgSO₄ and the solvent was evaporated under reduced pressure to give the crude 1-(alkyl)aryl-2-(5'-formyl-2'-thienyl)pyrroles **5** or a mixture of formyl- derivatives **5**, **6** and **7**, which were purified by "flash" chromatography on silica with increasing amounts of ether in light petroleum as eluent.

Metalation of thienylpyrrole **2a** followed by a reaction with DMF gave a mixture of **2a** and *1-propyl-2-(5'-formyl-2'-thienyl)pyrrole 5a*. The first compound eluted was thienylpyrrole **2a** (19%). The second compound eluted was *1-propyl-2-(5'-formyl-2'-thienyl)pyrrole 5a* as an orange oil (68%). UV (EtOH): λ_{max} nm ($\epsilon/\text{M}^{-1} \text{cm}^{-1}$) 374.0 (9474). IR (liquid film): ν 1659 (C=O), 1554, 1513, 1475, 1436, 1381, 1283, 1229, 1061, 941, 808, 724, 668, 611, 506 cm^{-1} . ¹H NMR (CDCl₃) δ 0.93 (t, 3H, $J=7.5\text{Hz}$, (CH₂)₂CH₃), 1.77-1.90 (m, 2H, CH₂CH₂CH₃), 4.08 (t, 2H, $J=7.5 \text{ Hz}$, NCH₂), 6.20-6.22 (m, 1H, 4-H), 6.53 (dd, 1H, $J=3.9$ and 1.8 Hz , 3-H), 6.82-6.86 (m, 1H, 5-H), 7.12 (d, 1H, $J=3.9 \text{ Hz}$, 3'-H), 7.71 (d, 1H, $J=3.9 \text{ Hz}$, 4'-H), 9.87 (s, 1H, CHO). ¹³C NMR (CDCl₃) δ 11.06, 24.61, 49.64, 108.70, 112.38, 112.44, 125.56, 125.70, 137.22, 140.94, 145.32, 182.54. MS (EI) m/z (%): 219 (M⁺, 100), 218 (9), 191 (19), 177 (26), 176 (23), 162 (17), 148 (7), 104 (7), 78 (2). HRMS: (EI) m/z (%) for C₁₂H₁₃NOS; calcd 219.0718; found 219.0718.

Metalation of thienylpyrrole **2b** followed by a reaction with DMF gave a mixture of **2b** and *1-phenyl-2-(5'-formyl-2'-thienyl)pyrrole 5b*. The first compound eluted was thienylpyrrole **2b** (9%). The second compound eluted was *1-phenyl-2-(5'-formyl-2'-thienyl)pyrrole 5b* as a dark orange oil (78%). UV (EtOH): λ_{max} nm ($\epsilon/\text{M}^{-1} \text{cm}^{-1}$) 374.0 (19180), 260.0 (7000). IR (liquid film): ν 1659 (CHO), 1597, 1498, 1461, 1438, 1230, 1062, 916, 802, 724, 696, 667 cm^{-1} . ¹H NMR (CDCl₃) δ 6.36-6.38 (m, 1H, 4-H), 6.61 (d, 1H, $J=4.2 \text{ Hz}$, 3'-H), 6.71 (dd, 1H, $J=3.6$ and 1.5 Hz , 3-H), 6.94-6.97 (m, 1H, 5-H), 7.28-7.33 (m, 2H, 2xAr-H), 7.42-7.47 (m, 3H, 3xAr-H), 7.49 (d, 1H, $J=4.2 \text{ Hz}$, 4'-H), 9.75 (s, 1H, CHO). ¹³C NMR (CDCl₃) δ 109.99, 113.05, 124.42, 126.88, 126.94, 128.42, 129.40 (two overlapped signals), 136.88, 139.38, 140.73, 145.25, 182.49. Anal.

Calcd for C₁₅H₁₁NOS: C, 71.13; H, 4.35; N, 5.53; S, 12.67. Found: C, 70.98; H, 4.54; N, 5.58; S, 12.69.

Metalation of thienylpyrrole **2c** followed by a reaction with DMF gave a mixture of *1-(4''-methoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole* **5c** and *1-(3''-formyl-4''-methoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole* **6c**. The first compound eluted was *1-(4''-methoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole* **5c** as a green solid (63%). Mp: 96.6-97.4 °C. UV (EtOH): λ_{max} nm (ε/M⁻¹ cm⁻¹) 379.0 (18613), 270.0 inf. (6000), 230.5 (16603), 203.5 (24294). IR (liquid film): ν 1659 (CHO), 1609, 1539, 1515, 1462, 1444, 1412, 1299, 1250, 1231, 1182, 1169, 1158, 1106, 1061, 917, 836, 797, 755, 724, 666, 619 cm⁻¹. ¹H NMR (CDCl₃) δ 3.87 (s, 3H, OCH₃), δ 6.32-6.36 (m, 1H, 4-H), 6.67 (d, 1H, *J*=4.2 Hz, 3'-H), 6.70 (dd, 1H, *J*=3.6 and 1.8 Hz, 3-H), 6.88-6.91 (m, 1H, 5-H), 6.95 (d, 2H, *J*=9.0 Hz, 2xAr-H), 7.22 (d, 2H, *J*=9.0 Hz, 2xAr-H), 7.49 (d, 1H, *J*=4.2 Hz, 4'-H), 9.74 (s, 1H, CHO). ¹³C NMR (CDCl₃) δ 55.45, 109.63, 112.33, 114.41, 124.02, 127.13, 127.19, 128.21, 132.04, 136.98, 140.43, 145.32, 159.52, 182.46. MS (EI) *m/z* (%): 283 (M⁺, 100), 268 (54), 240 (6), 121 (8), 103 (5). HRMS: (EI) *m/z* (%) for C₁₆H₁₃NO₂S; calcd 283.0667; found 283.0665. Anal. Calcd for C₁₆H₁₃NO₂S: C, 67.83; H, 4.59; N, 4.94; S, 11.33. Found: C, 67.78; H, 4.82; N, 5.02; S, 11.25. The second compound eluted was *1-(3''-formyl-4''-methoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole* **6c** as a orange oil (5%). UV (EtOH): λ_{max} nm (ε/M⁻¹ cm⁻¹) 374.0 (10605), 270.0 inf. (5000). IR (liquid film): ν 1684 (C=O), 1659 (C=O), 1611, 1499, 1462, 1394, 1273, 1234, 1122, 1019, 869, 817, 755, 726, 666 cm⁻¹. ¹H NMR (Acetone-d₆) δ 4.11 (s, 3H, OCH₃), 6.37-6.40 (m, 1H, 4-H), 6.80 (dd, 1H, *J*=3.8 and 1.2 Hz, 3-H), 6.91 (d, 1H, *J*=3.9 Hz, 3'-H), 7.09-7.11 (m, 1H, 5-H), 7.41 (d, 2H, *J*=8.7 Hz, 5''-H), 7.65 (dd, 1H, *J*=8.7 and 2.4 Hz, 6''-H), 7.69 (d, 1H, *J*=2.4 Hz, 2''-H), 7.75 (d, 1H, *J*=3.9 Hz, 4'-H), 9.81 (s, 1H, CHO), 10.48 (s, 1H, CHO). ¹³C NMR (CDCl₃) δ 56.06, 110.18, 112.56, 112.92, 124.49, 125.07, 126.56, 127.02, 127.07, 132.36, 134.26, 137.01, 140.92, 144.81, 161.43, 182.53, 188.68. MS (EI) *m/z* (%): 311 (M⁺, 100), 285 (5), 268 (30), 235 (6). HRMS: (EI) *m/z* (%) for C₁₇H₁₃NO₃S; calcd 311.0616; found 311.0625.

Metalation of thienylpyrrole **2d** followed by a reaction with DMF gave a mixture of thienylpyrrole **2d**, *1-(2'',4''-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole* **5d** and

1-(3''-formyl-2'',4''-dimethoxyphenyl)-2-(2'-thienyl)pyrrole 7d. The first compound eluted was pyrrole **2d** as a green solid (40%). The second component eluted was *1-(3''-formyl-2'',4''-dimethoxyphenyl)-2-(2'-thienyl)pyrrole 7d* as a dark green oil (8%). IR (liquid film): ν 3108, 1693 (C=O), 1659, 1581, 1492, 1442, 1397, 1334, 1288, 1232, 1186, 1164, 1121, 1096, 1013, 948, 842, 814, 716 cm^{-1} . ^1H NMR (CDCl_3) δ 3.48 (s, 3H, OCH_3), 3.96 (s, 3H, OCH_3), 6.33-6.38 (m, 1H, 4-H), 6.54 (dd, 1H, $J=4.2$ and 1.8 Hz, 3-H), 6.70 (dd, 1H, $J=3.6$ and 1.2 Hz, 3'-H), 6.75 (d, 1H, $J=9.0$ Hz, 6''-H or 5''-H), 6.80-6.83 (m, 1H, 5-H), 6.86-6.90 (m, 1H, 4'-H), 7.08 (dd, 1H, $J=5.1$ and 1.2 Hz, 5'-H), 7.42 (d, 1H, $J=9.0$ Hz, 6''-H or 5''-H), 10.45 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 56.28, 61.44, 106.44, 109.59, 109.72, 119.05, 123.71, 123.88, 125.04, 125.72, 127.07, 128.54, 134.70, 135.92, 158.82, 161.18, 189.14. MS (EI) m/z (%): 313 (M^+ , 100), 298 (17), 284 (13), 270 (9), 255 (5), 121 (9). HRMS: (EI) m/z (%) for $\text{C}_{17}\text{H}_{15}\text{NO}_3\text{S}$; calcd 313.0772; found 313.0785. The third compound eluted was *1-(2'',4''-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5d* as a dark green oil (48 %). UV (EtOH): λ_{max} nm ($\epsilon/\text{M}^{-1} \text{cm}^{-1}$) 384.5 (18158), 277.5 (6264), 233.0 (1312), 207.0 (30609). IR (liquid film): ν 2962, 2838, 1652 (CHO), 1613, 1589, 1541, 1516, 1462, 1443, 1416, 1383, 1306, 1286, 1260, 1232, 1210, 1161, 1135, 1092, 1062, 1046, 1030, 928, 911, 803, 728 cm^{-1} . ^1H NMR (CDCl_3) δ 3.66 (s, 3H, OCH_3), 3.89 (s, 3H, OCH_3), 6.34-6.38 (m, 1H, 4-H), 6.52-6.58 (m, 2H, 3''-H and 5''-H), 6.74 (dd, 1H, $J=3.9$ and 1.8 Hz, 3-H), 6.80-6.84 (m, 2H, 3'-H and 5-H), 7.17 (d, 1H, $J=9.3$ Hz, 6''-H), 7.51 (d, 1H, $J=4.2$ Hz, 4'-H), 9.73 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 55.54, 55.69, 99.72, 104.41, 109.56, 111.33, 121.03, 123.11, 127.34, 128.00, 129.78, 137.08, 140.00, 145.78, 156.40, 161.39, 182.51. MS (EI) m/z (%): 313 (M^+ , 100), 270 (8), 255 (8). HRMS: (EI) m/z (%) for $\text{C}_{17}\text{H}_{15}\text{NO}_3\text{S}$; calcd 313.0773; found 313.0770. Anal. Calcd for $\text{C}_{17}\text{H}_{15}\text{NO}_3\text{S}$: C, 65.16; H, 4.79; N, 4.47; S, 10.24. Found: C, 65.14; H, 5.07; N, 4.58; S, 10.04.

Metalation of thienylpyrrole **2d** during 2 h followed by 2 h of reflux with DMF gave a mixture of 4 compounds: thienylpyrrole **2d** (18%), *1-(2'',4''-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5d* (34%), *1-(3''-formyl-2'',4''-dimethoxyphenyl)-2-(2'-thienyl)pyrrole 7d* (14%) and *1-(3''-formyl-2'',4''-dimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole 6d* as a dark orange oil (18%). UV (EtOH): λ_{max} nm ($\epsilon/\text{M}^{-1} \text{cm}^{-1}$) 377.0 (16040), 260.5 (14720). IR (liquid film): ν 1690 (C=O), 1657 (C=O), 1492, 1461,

1288, 1231, 1094, 1019, 812, 608 cm^{-1} . ^1H NMR (CDCl_3) δ 3.48 (s, 3H, OCH_3), 4.00 (s, 3H, OCH_3), 6.39-6.42 (m, 1H, 4-H), 6.77 (dd, 1H, $J=3.6$ and 1.5 Hz, 3-H), 6.82 (d, 1H, $J=9.0$ Hz, 5''-H), 6.85-6.90 (m, 2H, 3'-H and 5-H), 7.46 (d, 1H, $J=9.0$ Hz, 6''-H), 7.54 (d, 1H, $J=4.2$ Hz, 4'-H), 9.75 (s, 1H, CHO), 10.45 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 56.35, 61.79, 106.96, 110.46, 112.23, 119.37, 123.62, 125.26, 127.47, 127.87, 135.43, 137.30, 140.62, 144.74, 158.44, 161.99, 182.49, 188.79. MS (EI) m/z (%): 341 (M^+ , 100), 326 (7), 312 (13), 298 (7), 83 (9). HRMS: (EI) m/z (%) for $\text{C}_{18}\text{H}_{15}\text{NO}_4\text{S}$; calcd 341.0722; found 341.0710.

1-(3'',4'',5''-Trimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5e. Metalation of thienylpyrrole **2e** followed by a reaction with DMF gave a mixture of thienylpyrrole **2e** and *1-(3'',4'',5''-Trimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5e*. The first component eluted thienylpyrrole **2e** as a pale green solid (40%). The second component eluted was *1-(3'',4'',5''-Trimethoxyphenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5e* as a yellow oil (12%). Mp: 113.5-114.7 $^\circ\text{C}$. UV (EtOH): λ_{max} nm ($\epsilon/\text{M}^{-1} \text{cm}^{-1}$) 377.0 (11860), 265.0 inf. (6820). IR (liquid film): ν 2935, 1659 (CHO), 1596, 1507, 1463, 1417, 1262, 1230, 1127, 1066, 1033, 1004, 840, 808, 727, 668 cm^{-1} . ^1H NMR (CDCl_3) δ 3.79 (s, 6H, $2\times\text{OCH}_3$), 3.92 (s, 3H, OCH_3), 6.34-6.37 (m, 1H, 4-H), 6.52 (br s, 2H, 2'' and 6''-H), 6.66 (d, 1H, $J=4.2$ Hz, 3'-H), 6.71 (dd, 1H, $J=3.8$ and 1.5 Hz, 3-H), 6.89-6.93 (m, 1H, 5-H), 7.52 (d, 1H, $J=3.9$ Hz, 4'-H), 9.77 (s, 1H, CHO). ^{13}C NMR (CDCl_3) δ 55.45, 109.63, 112.33, 114.41, 124.02, 127.13, 127.19, 128.21, 132.04, 136.98, 140.43, 145.32, 159.52, 182.46. MS (EI) m/z (%): 343 (M^+ , 100), 328 (50), 300 (6). HRMS: (EI) m/z (%) for $\text{C}_{18}\text{H}_{17}\text{NO}_4\text{S}$; calcd 343.0878; found 343.0896.

1-(4''-Fluorophenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5f. Metalation of thienylpyrrole **2f** followed by a reaction with DMF gave a mixture of thienylpyrrole **2f** and *1-(4''-Fluorophenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5f*. The first component eluted thienylpyrrole **2f** as a pale yellow solid (15%). The second component eluted was the *1-(4''-Fluorophenyl)-2-(5'-formyl-2'-thienyl)pyrrole 5f* as a brown solid (25%). Mp: 112-114 $^\circ\text{C}$. UV (EtOH): λ_{max} nm ($\epsilon/\text{M}^{-1} \text{cm}^{-1}$) 373.5 (15191), 257.6 (6386), 226.0 (8830), 204.0 (17041). IR (liquid film): ν 1659 (CHO), 1511, 1462, 1439, 1412, 1349, 1221, 1189, 1154, 1094, 1061, 1039, 916, 842 cm^{-1} . ^1H NMR (CDCl_3) δ 6.30-6.40 (m, 1H, 4-H), 6.63 (d, 1H, $J=4.2$ Hz, 3'-H), 6.65-6.75 (m, 1H, 3-H), 6.85-6.90 (m, 1H, 5-H), 7.13

(t_{ap} , 2H, $J=8.4$ Hz, 3'' and 5''-H), 7.25-7.35 (m, 2H, 2'' and 6''-H), 7.49 (d, 1H, $J=4.2$ Hz, 4'-H), 9.75 (s, 1H, CHO). MS (EI) m/z (%): 271 (M^+ , 100), 243 (3), 242 (8), 209 (5), 198 (6), 185 (5), 150 (9), 133 (6), 121 (6), 95 (5), 75 (4). HRMS: (EI) m/z (%) for $C_{15}H_{10}FNOS$; calcd 271.0467; found 271.0465.

Acknowledgements

Thanks are due to Foundation for Science and Technology (Portugal) for financial support through IBQF (UM) and through POCTI, FEDER (ref. POCTI/QUI/37816/2001).

References and notes

1. March, J., *Advanced Organic Chemistry*; 4th edn; Wiley-Interscience: New York, 1992, 542 and references cited therein.
2. Schlosser, M. *Angew. Chem. Int. Ed.* **2005**, *44*, 376 and references cited therein.
3. Chinchilla, R.; Nájera, C.; Yus, M. *Chem. Rev.* **2004**, *104*, 2667.
4. Jackson, A. H., Pyrroles In *The Chemistry of Heterocyclic Compounds*; Vol. 48, Part 1; Jones, R. A., Ed.; Wiley: New York, 1990, 295 and references cited therein.
5. Bird, C. W., Ed.; *Comprehensive Heterocyclic Chemistry II*; Vol. 2, Pergamon: Oxford, 1996, 1.
6. Castiñeiras, A.; Carballo, R.; Pérez, T. *Polyhedron* **2001**, *20*, 441.
7. Alonso, R.; Bermejo, E.; Castiñeiras, A.; Carballo, R.; Pérez, T. *J. Mol. Struct.* **2002**, *606*, 155.
8. Rodríguez-Argüelles, M.; López-Silva, E. C.; Sanmartín, J.; Pelagatti, P.; Zani, F., *J. Inorg. Biochem.* **2005**, *99*(11), 2231.
9. Won, D.-H.; Lee, C.-H. *Tetrahedron Lett.* **2003**, *44*(35), 6695.
10. Beecher, J. E.; Tirrell, D. A. *Tetrahedron Lett.* **1998**, *39*(23), 3927.
11. Haldar, P.; Guin, J.; Ray, J. K. *Tetrahedron Lett.* **2005**, *46*(7), 1071.
12. Pichon-Santander, C.; Scott, A. I. *Tetrahedron Lett.* **2000**, *41*(16), 2825.
13. Rodríguez, J. G.; Lafuente, A.; Rubio, L. *Tetrahedron Lett.* **2004**, *45*(29), 5685.

14. Toba, M.; Takeoka, Y.; Rikukawa, M.; Sanui, K. *Synthetic Metals*. **2005**, *152*(1-3), 197.
15. Fan, Q.-L.; Zhang, G.-W.; Lu, X.-M.; Chen, Y.; Huang, Y.-Q.; Zhou, Y.; Chan, H. S. O.; Lai, Y.-H.; Xu, G.-Q.; Huang, W. *Polymer* **2005**, *46*(24), 11165.
16. Purkarthofer, T.; Gruber, K.; Fechter, M. H.; Griengl, H. *Tetrahedron* **2005**, *61*(32), 7661.
17. Takekuma, S.-I.; Takahashi, K.; Sakaguchi, A.; Shibata, I.; Sasaki, M.; Minematsu, T.; Takekuma, H. *Tetrahedron* **2005**, *61*(32), 10349.
18. Hareau, G. P.-J.; Neya, S.; Funasaki, N.; Taniguchi, I. *Tetrahedron Lett.* **2002**, *43*(17), 3109.
19. Simionescu, C. I.; Grigoras, M.; Cianca, I.; Olaru, N. *Eur. Polym. J.* **1998**, *34*(7), 891.
20. Mignani G.; Leising F.; Meyrueix R.; Samson M., *Tetrahedron Lett.* **1990**, *31* (33), 4743.
21. Effenberger, F.; Wurthner, F.; Steybe F. *J. Org. Chem.* **1995**, *60*, 2082.
22. Eckert, K.; Schroder, A.; Hartmann, H. *Eur. J. Org. Chem.* **2000**, 1327 and references cited therein.
23. Turbiez, M., P.; Frère, P.; Roncali, J. *Tetrahedron* **2005**, *61*, 3045.
24. Mason, C. R.; Skabara, P. J.; Cupertino, D.; Schofield, J.; Meghdadi, F.; Ebner, B.; Sariciftci, N. S. *J. Mater. Chem.* **2005**, *15*, 1446.
25. Raposo, M. M. M.; Kirsch, G. *Tetrahedron* **2003**, *59*(26), 4891.
26. Raposo, M. M. M.; Fonseca, A. M. C.; Kirsch, G. *Tetrahedron* **2004**, *60*(18), 4071.
27. Batista, R. M. F.; Costa, S. P. G.; Raposo, M. M. M. *Tetrahedron Lett.* **2004**, *45*(13), 2825.
28. Raposo, M. M. M.; Sousa, A. M. R. C.; Fonseca, A. M. C.; Kirsch, G. *Tetrahedron*, **2005**, *61*(34), 8249-8256.
29. Raposo, M. M. M.; Sousa, A. M. R. C.; Kirsch, G.; Ferreira, F.; Belsey, M.; Matos Gomes; Fonseca, A. M. C. *Tetrahedron*, **2005**, *61*(50), 11991-11998.

30. Raposo, M. M. M.; Fonseca, A. M. C.; Sousa, A. M. R. C.; Kirsch, G. *Mater. Sci. Forum* **2006**, 514-516, 98-102.
31. Raposo, M. M. M.; Fonseca, A. M. C.; Sousa, A. M. R. C.; Kirsch, G. *Mater. Sci. Forum* **2006**, 514-516, 103-107.
32. Raposo, M. M. M.; Sampaio, A. M. B. A.; Kirsch, G. *J. Heterocyclic Chem.* **2005**, 42(7), 1245.
33. Costa, F., Silva; C. J. R.; Raposo, M. M. M.; Fonseca, A. M.; Neves, I. C.; Carvalho, A. P.; Pires, J. *Microporous Mesoporous Mater.* **2004**, 72, 111.
34. Fonseca, A. M. C.; Raposo, M. M. M.; Sousa, A. M. R. C.; Kirsch, G.; Beley, M. *Eur. J. Inorg. Chem.* **2005**, 21, 4361.
35. Raposo, M. M. M.; Sampaio, A. M. B. A.; Kirsch, G. *Synthesis* **2005**, 2, 199.
36. Raposo, M. M. M.; Kirsch G. *Heterocycles* **2001**, 55(8), 1487.
37. Belen'kii, L. I. *Kim. Geterotsikl. Soed.* **1980**, 12, 1587.
38. Belen'kii, L. I. *Heterocycles* **1994**, 37(3), 2029.
39. Boukou-Poba, J.-P.; Farnier, M.; Guillard, R. *Tetrahedron Lett.* **1979**, 20(19), 1717.
40. Boukou-Poba, J.-P.; Farnier, M.; Guillard, R. *Can. J. Chem.* **1981**, 59, 2962.
41. Trofimov, B. A.; Korostova, S. E.; Mikhaleva, A. I.; Nesterenko, R.N.; Sigalov, M. V.; Voronov, V. K.; Polovnikova, R. I. *Zh. Org. Kim.* **1982**, 18, 894.
42. Korostova, S. E.; Nesterenko, R. N.; Mikhaleva, A. I.; Shevchenko, S. G.; Kalabin, G. A.; Polovnikova, R. I. *Kim Geterotsikl. Soedin.* **1990**, 273.
43. Korostova, S. E.; Nesterenko, R. N.; Mikhaleva, A. I.; Shevchenko, S. G.; Kalabin, G. A.; Polovnikova, R. I. *Kim. Geterotsikl. Soed.* **1991**, 337.
44. Korostova, S. E.; Mikhaleva, A. I.; Trofimov, B. A. *Russ. Chem. Rev. (Engl. Transl.)* **1999**, 68(6), 459.
45. Trofimov, B. A.; Sobenina, L. N.; Demenev, A. P.; Mikhaleva, A. I.; Ushakov, I. A.; Tarasova, O. A.; Smirnov, V. I.; Petrushenko, K. B.; Vokin, A. I.; Murzina, N. M.; Myachina, G. F. *Arkivok* **2001**, ix, 37.
46. Niziursky-Mann; R. E.; Cava, M. P. *Heterocycles* **1992**, 34(10), 2003.
47. Majchrzak, M. W. and Simchen, G. *Tetrahedron* **1986**, 42, 1299.

48. Muchowski, J. M.; Naef, R. *Helv. Chim. Acta* **1984**, *67*, 1168 and references cited therein.
49. Kozikowski, A. P.; Cheng, X.-M. *J. Org. Chem.* **1984**, *49*, 3239.
50. Stefan, K.-P.; Schuhmann, W.; Parlar, H.; Korte, Friedhelm *Chem. Ber.* **1989**, *122*, 169.
51. Lomas, J. S.; Vaissermann, J. *J. Chem. Soc Perkin Trans. 2* **1998**, 1777.
52. Robertson, J.; Kuhnert, N.; Zhao, Y. *Heterocycles* **2000**, *53*(11), 2415.
53. Selvi, S.; Pu, S.-C.; Cheng, Y.-M.; Fang, J.-M.; Chou, P.-T. *J. Org. Chem.* **2004**, *69*, 6674.
54. Martineau, D.; Gros, P.; Fort, Y. *J. Org. Chem.* **2004**, *69*, 7914.
55. Faigl, F.; Fogassy, K.; Thurner, A.; Toke, László, T. *Tetrahedron* **1997**, *53*(13), 4883.
56. Faigl, F.; Fogassy, K.; Szántó, Z.; Lopata, A.; Toke, L. *Tetrahedron* **1998**, *54*(17), 4367.
57. Faigl, F.; Fogassy, K.; Szucs, E.; Kovács, K.; Keserú, Gy.-M.; Harmat, V.; Böcskei, Zs.; Toke, L. *Tetrahedron*, **1999** *55*(25), 7881.
58. Fogassy, K.; Kovács, K.; Keserú, G. M.; Toke, L.; Faigl, F.; *J. Chem. Soc. Perkin Trans. 1* **2001**, 1039.
59. Faigl, F.; Thurner, A.; Vass, B.; Toke, L. *J. Chem. Research (S)* **2003**, 132.
60. Snieckus, V. *Chem. Rev.* **1990**, *90* (6), 879 and references cited therein.
61. Casado, J.; Pappenfus, T. M.; Miller, L. L.; Mann, K. R.; Orti, E.; Viruela, P. M.; Pou-Amerigo, R.; Hernandez, V.; Navarrete, J. T. L. *J. Am. Chem. Soc.* **2003**, *125*, 2524.

Captions

Scheme 1. Synthesis of formyl-thienylpyrroles **3-4** from thienylpyrroles **2** by Vilsmeier-Haack reaction with DMF.

Scheme 2. Synthesis of formyl-thienylpyrroles **5-7** from thienylpyrroles **2** by m followed by reaction with DMF.

Table 1. Yields, ^1H NMR, IR and UV-vis data of formyl-thienylpyrroles **3-4**.

^a For the *CHO* proton of formyl-thienylpyrroles **3-4** (300 MHz, CDCl_3)

^b For the *CHO* proton of formyl-thienylpyrroles **3-4** (300 MHz, Acetone-d_6).

^c All the UV/vis spectra were run in ethanol.

Table 2. Yields, ^1H NMR, IR and UV-vis data of formyl-thienylpyrroles **5-7**.

^a Yields for compounds **5d-7d** obtained for 2 h of lithiation followed by 2 h of reaction with DMF.

^b For the *CHO* proton of formyl-thienylpyrroles **5-7** (300 MHz, CDCl_3).

^c For the *CHO* proton of formyl-thienylpyrroles **5-7** (300 MHz, acetone-d_6).

^d All the UV/vis spectra were run in ethanol.

Tables


Table 1

Entry	Pyrrole	Formyl- pyrrole	Yield (%)	δ_{H} (ppm) ^a	IR ν_{CHO} [cm ⁻¹]	UV/vis λ_{max} [nm] ^c (ϵ)
1	2a	3a	63	9.54	1658	321.5 (24,650)
2	2a	4a	5	9.54	1662	293.5 (5440)
3	2b	3b	19	9.39	1660	340.5 (20,100)
4	2b	4b	22	9.82 ^b	1661	325.0 (7440)
5	2c	3c	12	9.38	1655	343.0 (21,559)
6	2c	4c	10	9.81 ^b	1661	-----


Table 2

Entry	Pyrrole	UV/vis. λ_{\max} [nm] ^d (ϵ)	Formyl- pyrrole	Yield (%)	δ_{H} (ppm) ^b	IR ν_{CHO} [cm ⁻¹]	UV/vis. λ_{\max} [nm] ^d (ϵ)
1	2a	291.0 (1800)	5a	68	9.87	1659	374.0 (9474)
2	2b	294.5 (9208)	5b	78	9.75	1659	374.0 (19,180)
3	2c	290.0 (11,410)	5c	63	9.74	1659	379.0 (18,613)
4	2c	-----	6c	5	9.81 ^c , 10.48 ^c	1659, 1684	374.0 (10,605)
5	2d	286.5 (10,093)	5d	48, 34 ^a	9.73	1652	384.5 (18,158)
6	2d	-----	6d	18 ^a	9.75, 10.45	1657, 1690	377.0 (16,040)
7	2d	-----	7d	8, 14 ^a	10.45	1693	-----
8	2e	281.5 (8477)	5e	12	9.77	1659	377.0 (11,860)
9	2f	289.5 (7939)	5f	25	9.75	1659	373.5 (15,191)

Scheme 1



Scheme 2



Graphical Abstract

Synthesis of formyl-thienylpyrroles: versatile building blocks for NLO materials

M. Manuela M. Raposo^{a*}, Ana M. R. C. Sousa^a, A. Maurício C. Fonseca^a and G. Kirsch^b

^aCentro de Química, Universidade do Minho, Campus de Gualtar 4710-057 Braga, Portugal

^bLaboratoire d'Ingénierie Moléculaire et Biochimie Pharmacologique, UFR SciFA, Université de Metz, 1, Boulevard Arago, Metz Technopôle, 57078 Metz Cedex 3, France

