

Interaction of a potential antitumoral benzothieno[3,2-*b*]pyrrole with lipid membranes and salmon sperm DNA

Ana S. Abreu,^{a,b} M. Solange D. Carvalho,^a Elisabete M.S. Castanheira,^b
Maria-João R.P. Queiroz,^a Paula M.T. Ferreira^a

a) Centre of Chemistry, b) Centre of Physics, Univ. of Minho, Campus de Gualtar,
4710-057 Braga, Portugal. e-mail: anabreu@quimica.uminho.pt

In this work, the interaction of a potential antitumoral benzothieno[3,2-*b*]pyrrole (**BTP**) with lipid membranes and natural salmon sperm DNA was studied by fluorescence techniques. Studies of the influence of **BTP** on the growth of human tumor cell lines showed that this compound highly inhibit the growth of NCI-H460 (non-small cell lung cancer) cells with a $GI_{50} = 3.9 \mu\text{M}$. The interaction of the fluorescent **BTP** with ds-DNA allowed the determination of a binding constant of $K_i = (2.9 \pm 0.3) \times 10^6 \text{ M}^{-1}$ and a binding site size of $n = 2.0 \pm 0.7$, pointing to a high affinity of this compound to the macromolecule. Fluorescence quenching experiments using I point to an intercalative mode of binding.

Fluorescence studies of **BTP** incorporated in lipid aggregates of DPPC, DOPE and Egg-PC (Fig. 1) indicate that this compound is located mainly near the hydrophobic lipid tails and is able to distinguish between the rigid gel phase and fluid phases.





Fig. 1 - Fluorescence spectra of **BTP** in lipid membranes of DOPE, Egg-PC and DPPC. Inset: Fluorescence spectra of **BTP** in cyclohexane and ethanol at 55 °C.

Acknowledgements: This work was funded by FCT and FEDER through CQ-UM and CFUM, project POCI/QUI/59407/2004 and post-Doc grant (SFRH/BPD/24548/2005) of A.S.A.