



UvA-DARE (Digital Academic Repository)

Games and cardinalities in inquisitive first-order logic

Grilletti, G.; Ciardelli, I.

DOI

[10.1017/S1755020321000198](https://doi.org/10.1017/S1755020321000198)

Publication date

2023

Document Version

Final published version

Published in

Review of Symbolic Logic

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Grilletti, G., & Ciardelli, I. (2023). Games and cardinalities in inquisitive first-order logic. *Review of Symbolic Logic*, 16(1), 241-267. <https://doi.org/10.1017/S1755020321000198>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

GAMES AND CARDINALITIES IN INQUISITIVE FIRST-ORDER LOGIC

GIANLUCA GRILLETTI

Institute for Logic, Language and Computation
and

IVANO CIARDELLI

Munich Center for Mathematical Philosophy

Abstract. Inquisitive first-order logic, InqBQ , is a system which extends classical first-order logic with formulas expressing questions. From a mathematical point of view, formulas in this logic express properties of sets of relational structures. This paper makes two contributions to the study of this logic. First, we describe an Ehrenfeucht–Fraïssé game for InqBQ and show that it characterizes the distinguishing power of the logic. Second, we use the game to study cardinality quantifiers in the inquisitive setting. That is, we study what statements and questions can be expressed in InqBQ about the number of individuals satisfying a given predicate. As special cases, we show that several variants of the question *how many individuals satisfy $\alpha(x)$* are not expressible in InqBQ , both in the general case and in restriction to finite models.

§1. Introduction. According to the traditional view, the semantics of a logical system specifies truth-conditions for the sentences in the language. This focus on truth restricts the scope of logic to a special kind of sentences, namely, statements, whose semantics can be adequately characterized in terms of truth-conditions. In recent years, a more general view of semantics has been developed, which goes under the name of *inquisitive* semantics (see [4] for a language-oriented introduction and [2] for a logic-oriented one). In this approach, the meaning of a sentence is laid out not by specifying when the sentence is true relative to a state of affairs, but rather by specifying when it is supported by a given state of information. This view allows us to interpret in a uniform way both statements and questions: for instance, the statement *it rains* will be supported by an information state s if the information available in s implies that it rains, while the question *whether it rains* will be supported by s if the information available in s determines whether or not it rains.

In its first-order version, referred to as InqBQ , inquisitive logic can be seen as a conservative extension of classical first-order logic with formulas expressing questions. Thus, in addition to standard first-order formulas like Pa and $\forall x.Px$, we also have formulas like $?Pa$ (“*does a have property P?*”), $\exists x.Px$ (“*what is an instance of an*

Received: July 7, 2020.

2020 *Mathematics Subject Classification*: 03B60, 03B65, 91A44.

Key words and phrases: inquisitive logic, Ehrenfeucht–Fraïssé games, quantification, cardinality quantifiers, generalized quantifiers.

© The Author(s), 2021. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

individual having property P?”) and $\forall x. ?Px$ (“*which individuals have property P?*”). A model for this logic is based on a set W of possible worlds, each representing a possible state of affairs, corresponding to a standard first-order structure. An information state is modeled as a subset $s \subseteq W$. The idea, which goes back to the work of Hintikka [12], is that a set of worlds s stands for a body of information that is compatible with the actual world being one of the worlds $w \in s$ and incompatible with it being one of the worlds $w \notin s$. The semantics of the language takes the form of a support relation holding between information states in a model and sentences of the language.

From a mathematical point of view, a sentence of InqBQ expresses a property of a set s of first-order structures. The crucial difference between statements and questions is that statements express *local* properties of information states—which boil down to requirements on the individual worlds $w \in s$ —while questions express *global* requirements, having to do with the way the worlds in s are related to each other. Thus, for instance, the formula $?Pa$ requires that the truth-value of Pa be the same in all worlds in s ; the formula $\exists x.Px$ requires that there be an individual that has property P uniformly in all worlds in s ; and the formula $\forall x. ?Px$ requires that the extension of property P be the same across s . Global properties can also take the form of *dependencies*: thus, e.g., $?Pa \rightarrow ?Qa$ requires that the truth-value of Qa be functionally determined by the truth-value of Pa in s , while $\forall x. ?Px \rightarrow \forall x. ?Qx$ requires that the extension of property Q be functionally determined by the extension of property P in s . Thus, inquisitive first-order logic provides a language that can be used to talk about both local and global features of an information state.

In contrast to inquisitive propositional logic, which has been thoroughly investigated (see, among others, [1, 3, 5, 10, 19, 20, 22]), inquisitive *first-order* logic has received comparatively little attention [2, 11]. In particular, a detailed investigation of the expressive power of the logic has so far been missing. This paper makes a first, important step in this direction.

In the classical setting, a powerful tool to study the expressiveness of first-order logic is given by Ehrenfeucht–Fraïssé games (also known as EF games or back-and-forth games), introduced in 1967 by Ehrenfeucht [7], developing model-theoretic results presented by Fraïssé [9]. These games provide a particularly perspicuous way of understanding what differences between models can be detected by means of first-order formulas of a certain quantifier rank. Reasoning about winning strategies in this game, one can prove that two first-order structures are elementarily equivalent, or one can find a formula telling them apart.

One of the main merits of EF games is that they allow for relatively easy proofs that certain properties of first-order structures are not first-order expressible. A classical application of this kind is the characterization of the cardinality quantifiers definable in classical first-order logic. This characterization says that the only cardinality quantifiers definable in classical first-order logic are those which, for some natural number m , are insensitive to the difference between any cardinals larger than m . This characterization yields a range of interesting undefinability results: for instance, it implies that the quantifiers *an even number of individuals* and *infinitely many individuals* are not first-order definable.

The basic idea of EF games has proven to be very flexible and adaptable to a wide range of logical settings, including fragments of first-order logic with finitely many variables [14]; extensions of first-order logic with generalized quantifiers [15]; monadic second order logic [8]; modal logic [24]; and intuitionistic logic [18, 25]. In each case,

the game provides an insightful characterization of the distinctions that can and cannot be made by means of formulas in the logic.

In this paper, we make two contributions to the study of inquisitive first-order logic. First, we introduce an EF-style game for InqBQ and show that this game provides a characterization of the expressive power of the logic. Second, we introduce a notion of *inquisitive cardinality quantifier*, and use the game to study which of these quantifiers are definable in InqBQ.

The notion of inquisitive cardinality quantifier is a natural generalization of the standard notion of cardinality quantifier: besides standard quantifiers like *infinitely many*, which combine with a property to form a statement, we will now also have quantifiers like *how many*, which combine with a property to form a question. Using the EF-game, we will be able to characterize exactly the range of cardinality quantifiers expressible in InqBQ. The characterization is similar to the one for classical first-order logic: the definable cardinality quantifiers are those that, for some finite threshold m , are incapable of distinguishing between cardinalities larger than m .

The result implies that many natural kinds of questions about cardinalities are not expressible in InqBQ. The prime example is the question *how many individuals have property P*, which is supported in a state s if the extension of P has the same cardinality in all the worlds in s . We show that this question is not expressible in InqBQ, even in restriction to finite models. This means that a logical treatment of *how many* questions in inquisitive logic requires a proper extension of InqBQ. Other examples of cardinality questions which we show not to be expressible in InqBQ are: whether the number of P is finite or infinite; whether it is even or odd; whether it is countable or uncountable.

From a meta-theoretical point of view, our characterization result is especially interesting in light of the fact that it is still an open question how the expressive power of InqBQ compares to those of first- and second-order logics. It is not known, e.g., whether InqBQ is compact and whether an entailment-preserving translation to first-order logic exists. Our result indicates that, at least with respect to the expression of cardinality properties, InqBQ is much more similar to first-order logic than to second-order logic, where quantifiers like ‘*infinitely many*’ and ‘*an even number of*’ can be expressed.

The paper is structured as follows: in Section 2 we provide some technical background on the logic InqBQ. In Section 3 we describe the game and show that it characterizes the distinguishing power of the logic. In Section 4 we characterize the cardinality quantifiers definable in InqBQ. In Section 5 we summarize our findings and mention some directions for future work.

§2. Inquisitive first-order logic. In this section we provide a basic introduction to inquisitive first-order logic. For a more comprehensive introduction, the reader is referred to [2].

Syntax. Let Σ be a predicate logic signature. For simplicity, we first restrict to the case in which Σ is a *relational* signature, i.e., contains no function symbols. The extension to an arbitrary signature, which involves some subtleties familiar from the classical case [13], is discussed in Section 3.4. The set \mathcal{L} of formulas of InqBQ over Σ is defined as follows, where $R \in \Sigma$ is an n -ary relation symbol:

$$\varphi ::= R(x_1, \dots, x_n) \mid (x_1 = x_2) \mid \perp \mid \varphi \wedge \varphi \mid \varphi \rightarrow \varphi \mid \forall x.\varphi \mid \varphi \vee \varphi \mid \exists x.\varphi.$$

We will take negation to be a defined operator:

$$\neg\varphi := \varphi \rightarrow \perp.$$

Formulas without occurrences of \vee and \exists are referred to as *classical* formulas and can be identified with standard FOL-formulas, that is, first-order logic formulas. That is, the set \mathcal{L}_c of classical formulas is given by:

$$\alpha ::= R(x_1, \dots, x_n) \mid (x_1 = x_2) \mid \perp \mid \alpha \wedge \alpha \mid \alpha \rightarrow \alpha \mid \forall x.\alpha.$$

As usual, classical formulas may be viewed as formalizing statements, such as, for instance, ‘every object has property P ’. In the following, the variables α, β, γ will range over classical formulas, while φ, ψ, χ will range over arbitrary formulas. If α and β are classical formulas, then we can define: ¹

- $\alpha \vee \beta := \neg(\neg\alpha \wedge \neg\beta)$
- $\exists x.\alpha := \neg\forall x.\neg\alpha$

The connective \vee and the quantifier \exists , referred to respectively as *inquisitive disjunction* and *inquisitive existential quantifier*, allow us to form questions. For instance, if α is a classical formula then the formula $\alpha \vee \neg\alpha$ represents the question *whether* α . We abbreviate this formula as $?\alpha$:

- $?\alpha := \alpha \vee \neg\alpha$

The formula $\exists x.\alpha(x)$ represents the question *what is an object satisfying* $\alpha(x)$; and the formula $\forall x.?\alpha(x)$ represents the question *which objects satisfy* $\alpha(x)$.

Semantics. A model for InqBQ consists of: a set W of worlds, representing possible states of affairs; a set D of individuals, the objects that the first-order variables range over; and an interpretation function I , which determines at each world the extension of all relation symbols, including identity.

DEFINITION 2.1 (Models). *A model for the signature Σ is a tuple $M = \langle W, D, I \rangle$ where W and D are sets and I is a function mapping each world $w \in W$ and each n -ary relation symbol $R \in \Sigma \cup \{=\}$ to a corresponding n -ary relation $I_w(R) \subseteq D^n$ —the extension of R at w . The interpretation of identity is subject to the following condition:*

$$I_w(=) \text{ is a congruence, i.e., an equivalence relation } \sim_w \text{ such that, if } R \in \Sigma \text{ and } d_i \sim_w d'_i \text{ for } i \leq n, \text{ then } \langle d_1, \dots, d_n \rangle \in I_w(R) \iff \langle d'_1, \dots, d'_n \rangle \in I_w(R).$$

As discussed in the introduction, in inquisitive logic the semantics of the language specifies when a formula is supported at an information state $s \subseteq W$, rather than when a formula is true at a possible world $w \in W$. As usual, to handle open formulas and quantification, the support relation is defined relative to an assignment, which is a function from variables to the set D of individuals; if g is an assignment and $d \in D$, then $g[x \mapsto d]$ is the assignment which maps x to d and behaves like g on all other variables.

¹ We could in principle define \vee and \exists for arbitrary formulas; however, these operators are only natural and useful when applied to classical formulas, on which they yield the standard disjunction and existential quantifier of classical logic.

DEFINITION 2.2 (Support). *Let $M = \langle W, D, I \rangle$ be a model and let $s \subseteq W$.*

$$\begin{aligned}
 M, s \models_g R(x_1, \dots, x_n) &\iff \forall w \in s : \langle g(x_1), \dots, g(x_n) \rangle \in I_w(R) \\
 M, s \models_g x_1 = x_2 &\iff \forall w \in s : g(x_1) \sim_w g(x_2) \\
 M, s \models_g \perp &\iff s = \emptyset \\
 M, s \models_g \varphi \wedge \psi &\iff M, s \models_g \varphi \text{ and } M, s \models_g \psi \\
 M, s \models_g \varphi \vee \psi &\iff M, s \models_g \varphi \text{ or } M, s \models_g \psi \\
 M, s \models_g \varphi \rightarrow \psi &\iff \forall t \subseteq s : M, t \models_g \varphi \text{ implies } M, t \models_g \psi \\
 M, s \models_g \forall x. \varphi &\iff M, s \models_{g[x_i \mapsto d]} \varphi \text{ for all } d \in D \\
 M, s \models_g \exists x. \varphi &\iff M, s \models_{g[x_i \mapsto d]} \varphi \text{ for some } d \in D
 \end{aligned}$$

As usual, if $\varphi(x_1, \dots, x_n)$ is a formula whose free variables are among x_1, \dots, x_n , then the value of g on variables other than x_1, \dots, x_n is irrelevant. If $d_1, \dots, d_n \in D$, we can therefore write $M, s \models \varphi(d_1, \dots, d_n)$ to mean that $M, s \models_g \varphi$ holds with respect to an assignment g that maps x_i to d_i . In particular, if φ is a sentence we can drop reference to g altogether. Moreover, we write $M \models \varphi$ as a shorthand for $M, W \models \varphi$ and we say that M supports φ .

It is easy to verify that the support relation has the following two basic features:

- Persistency: if $M, s \models_g \varphi$ and $t \subseteq s$ then $M, t \models_g \varphi$;
- Empty state property: if $M, \emptyset \models_g \varphi$ for all φ .

Recovering classical logic. In restriction to classical formulas, the above definition of support gives a non-standard semantics for classical first-order logic. To see why, let us associate to each world $w \in M$ a corresponding relational structure \mathcal{M}_w , having as its domain the quotient D/\sim_w and with the interpretation of relation symbols induced by $I_w(R)$. Then we have the following connection.

PROPOSITION 2.3. *For all classical formulas $\alpha \in \mathcal{L}_c$, all models M , assignments g , and information states s :*

$$M, s \models_g \alpha \iff \forall w \in s : \mathcal{M}_w \models_{\bar{g}} \alpha \text{ holds in first-order logic}$$

where \bar{g} is the assignment mapping x to the \sim_w -equivalence class of $g(x)$.

Thus, as far as the standard fragment of the language is concerned, the relation of support is essentially a recursive definition of global truth with respect to a set of structures sharing the same domain. Notice that the standard definition of truth can be recovered as a special case of support by taking s to be a singleton. We will also write $M, w \models_g \alpha$ as an abbreviation for $M, \{w\} \models_g \alpha$.

Questions. As we just saw, evaluating a classical formula on an information state s amounts to evaluating it at each world in s and determining whether it is satisfied at each world. The same is not true for formulas that contain the operators \forall and \exists ; typically, such formulas allow us to express global requirements on a state, which cannot be reduced to requirements on the single worlds in the state. We will illustrate this point by means of some examples. First take a classical sentence α and consider the formula $?\alpha := \alpha \vee \neg\alpha$. We have:

$$M, s \models ?\alpha \iff (\forall w \in s : M, w \models \alpha) \text{ or } (\forall w \in s : M, w \models \neg\alpha).$$

Thus, in order for s to support $?\alpha$, all the worlds in s must agree on the truth-value of α . In other words, $?\alpha$ is supported at s only if the information available in s determines

whether or not α is true. Thus, $?\alpha$ can be taken as a formal representation of the question “*is α true?*”

Next take $\alpha(x)$ to be a classical formula having only the variable x free, and consider the formula $\exists x.\alpha(x)$. We have:

$$M, s \models \exists x.\alpha(x) \iff \exists d \in D \text{ such that } \forall w \in s : M, w \models \alpha(d).$$

Thus, in order for s to support $\exists x.\alpha(x)$ there must be an individual d which satisfies $\alpha(x)$ at all worlds in s . In other words, $\exists x.\alpha(x)$ is supported at s if the information available in s implies for some specific individual that it satisfies $\alpha(x)$ —i.e., gives us a specific witness for $\alpha(x)$. Thus, $\exists x.\alpha(x)$ can be taken as a formal representation of the question “*what is an object satisfying $\alpha(x)$?*”

Finally, let again $\alpha(x)$ be a classical formula having only x free, and let us denote by $I_w(\alpha)$ the set of objects which satisfy $\alpha(x)$ at w , i.e., $I_w(\alpha) := \{d \in D \mid M, w \models \alpha(d)\}$. Consider the formula $\forall x.? \alpha(x) := \forall x.(\alpha(x) \vee \neg \alpha(x))$. We have:

$$M, s \models \forall x.? \alpha(x) \iff \forall w, w' \in s : I_w(\alpha) = I_{w'}(\alpha).$$

Thus, in order for s to support $\forall x.? \alpha(x)$, all the worlds in s must agree on which objects satisfy $\alpha(x)$. In other words, $\forall x.? \alpha(x)$ is supported at s if the information available in s determines exactly which individuals satisfy $\alpha(x)$. Thus, $\forall x.? \alpha(x)$ can be taken as a formal representation of the question “*which objects satisfy $\alpha(x)$?*”

Identity and cardinalities. An aspect of InqBQ which is worth commenting on is the interpretation of identity. In InqBQ, the interpretation of identity may differ at different worlds. This allows us to deal with uncertainty about the identity relation: e.g., one may have information about two individuals, a and b (say, one knows Pa and Qb) and yet be uncertain whether a and b are distinct individuals, or the same. This also allows for uncertainty about how many individuals there are. Indeed, although a model is based on a fixed set D of epistemic individuals—objects to which information can be attributed—the domain of actual individuals at a world w is given by the equivalence classes modulo \sim_w ; the number of actual individuals that exist at w is the number of such equivalence classes, i.e., the cardinality of the quotient $D_w := D/\sim_w$. Similarly, if $\alpha(x)$ is a classical formula having at most x free, then the number of individuals that satisfy $\alpha(x)$ at w is given by the cardinality of the set $\alpha_w := I_w(\alpha)/\sim_w$.

Notice that, as a special case, we could take \sim_w to be the actual relation of identity on D at each world. A model in which identity is treated in this way is called an id-model.

This section is only intended as a summary of the key definitions and features of InqBQ and as a quick illustration of how questions can be captured by formulas in this logic. With these basic notions in place, let us now turn to the first novel contribution of the paper: an Ehrenfeucht–Fraïssé game for InqBQ.

§3. An Ehrenfeucht–Fraïssé game for InqBQ. The EF game for InqBQ is played by two players, S (Spoiler) and D (Duplicator), using two inquisitive models M_0, M_1 as a board. As in the classical case, the game proceeds in turns: at each turn, S picks an object from one of the two models and D must respond by picking a corresponding object from the other model. At the end of the game, a winner is decided by comparing the atomic formulae supported by the sub-structures built during the game.

However, there are two crucial differences with the classical EF game. First, the objects that are picked during the game are not just individuals $d \in D_i$, but also

information states $s \subseteq W_i$. This is because the logical repertoire of InqBQ contains not only the operators \forall and $\bar{\exists}$, which quantify over individuals, but also the operator \rightarrow , which quantifies over information states. Second, the roles of the two models in the game are not symmetric. This is connected to the absence of a classical negation in the language of InqBQ; unlike in classical logic, it could be that a model M_0 supports all the formulas supported by a model M_1 , but not vice versa. This directionality is reflected by the game.

3.1. The game. A position in an EF game for InqBQ is a tuple

$$\langle M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1 \rangle,$$

where:

- $M_0 = \langle W_0, D_0, I_0 \rangle$ and $M_1 = \langle W_1, D_1, I_1 \rangle$ are models for InqBQ;
- s_0 and s_1 are information states in the models M_0 and M_1 respectively; and
- \bar{a}_0 and \bar{a}_1 are tuples of equal length of elements from D_0 and D_1 respectively.

If not otherwise specified, a game between the models M_0 and M_1 starts from position $\langle M_0, W_0, \varepsilon; M_1, W_1, \varepsilon \rangle$, where ε indicates the empty tuple.

Starting a round from a position $\langle M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1 \rangle$, S can choose between the following possible moves:²

- $\bar{\exists}$ -move: S picks an element $b_0 \in D_0$; D responds with an element $b_1 \in D_1$; the game continues from the position $\langle M_0, s_0, \bar{a}_0 b_0; M_1, s_1, \bar{a}_1 b_1 \rangle$;
- \forall -move: S picks an element $b_1 \in D_1$; D responds with an element $b_0 \in D_0$; the game continues from the position $\langle M_0, s_0, \bar{a}_0 b_0; M_1, s_1, \bar{a}_1 b_1 \rangle$; and
- \rightarrow -move: S picks a sub-state $t_1 \subseteq s_1$; D responds with a sub-state $t_0 \subseteq s_0$; S picks $i \in \{1, 0\}$. The game continues from $\langle M_i, t_i, \bar{a}_i; M_{1-i}, t_{1-i}, \bar{a}_{1-i} \rangle$.

Notice the asymmetry between the roles of the two models: by performing an \rightarrow -move, S can pick an information state from M_1 , but not a state in M_0 .

With respect to termination condition, we consider different versions of the game. In the bounded version of the game, a pair of numbers $\langle i, q \rangle \in \mathbb{N}^2$ is fixed in advance. This number constrains the development of the game: in total, S can play only i implication moves and only q quantifier moves (i.e., $\bar{\exists}$ -move or a \forall -move). When there are no more moves available, the game ends. If $\langle M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1 \rangle$ is the final position, the game is won by Player D if the following condition is satisfied, and by player S otherwise:

- Winning condition for D: for all atomic formulas $\alpha(x_1, \dots, x_n)$ where n is the size of the tuples \bar{a}_0 and \bar{a}_1 , we have:

$$M_0, s_0 \models \alpha(\bar{a}_0) \implies M_1, s_1 \models \alpha(\bar{a}_1). \tag{1}$$

In the unbounded version of the game, no restriction is placed at the outset on the number of moves to be performed. Instead, player S has the option to declare the game over at the beginning of each round: in this case, the winner is determined as in the bounded version of the game. If the game never stops, then D is the winner.³

² In the following, the notation $\bar{a}b$ indicates the sequence obtained by adding the element b at the end of the sequence \bar{a} .

³ Here we consider games in which the rounds of play are indexed by natural numbers. To define games of transfinite length, one would have to specify how to determine the game position corresponding to a limit ordinal. We leave this for future work.

M_0	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr> <td style="border: none;"></td> <td style="border: none;">w_0</td> <td style="border: none;">w_1</td> </tr> <tr> <td style="border: none;">d_1</td> <td style="text-align: center;">•</td> <td style="text-align: center;">×</td> </tr> <tr> <td style="border: none;">d_2</td> <td style="text-align: center;">×</td> <td style="text-align: center;">•</td> </tr> </table>		w_0	w_1	d_1	•	×	d_2	×	•
	w_0	w_1								
d_1	•	×								
d_2	×	•								

M_1	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr> <td style="border: none;"></td> <td style="border: none;">v_0</td> <td style="border: none;">v_1</td> </tr> <tr> <td style="border: none;">e_1</td> <td style="text-align: center;">•</td> <td style="text-align: center;">•</td> </tr> <tr> <td style="border: none;">e_2</td> <td style="text-align: center;">×</td> <td style="text-align: center;">•</td> </tr> </table>		v_0	v_1	e_1	•	•	e_2	×	•
	v_0	v_1								
e_1	•	•								
e_2	×	•								

	M	N
S	$a_1 := d_1$	
D		$b_1 := e_2$
S	$s_1 := \{w_0\}$	$t_1 := \{v_1\}$
D		$i := 1$
S		$b_2 := e_1$
D	$a_2 := d_2$	

Fig. 1. On the left, two inquisitive models in the signature $\Sigma = \{P\}$. The top row represents the set of worlds of the models (e.g., $\{w_0, w_1\}$ for the model M_0) and the left column represents the domain (e.g., $\{d_1, d_2\}$ for M_0); $I(P)$ is encoded by the entries of the table: a • indicates that P holds, while a × indicates that P does not hold (e.g., $d_1 \in I_{w_0}(P)$ and $d_2 \notin I_{w_0}(P)$). On the right, a run of the Ehrenfeucht-Fraïssé game.

EXAMPLE 3.1. Take the signature $\Sigma = \{P\}$, where P is a unary predicate symbol. Given the models M_0 and M_1 in Figure 1, in the table we show a run of the bounded game with $\langle i, q \rangle = \langle 1, 2 \rangle$ between M_0 and M_1 . At the end of the run, the position is $\langle M_1, \{v_1\}; \langle e_2, e_1 \rangle; M_0, \{w_0\}, \langle d_1, d_2 \rangle \rangle$. The winner is Spoiler, since

$$M_1, \underbrace{\{v_1\}}_{t_1} \models P(\underbrace{e_1}_{b_1}) \not\models M_0, \underbrace{\{w_0\}}_{s_1} \models P(\underbrace{d_2}_{a_1}).$$

As usual, a winning strategy for a player is a strategy which guarantees victory to them, no matter what the opponent plays. If D has a winning strategy in the EF game of length $\langle i, q \rangle$ starting from position $\langle M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1 \rangle$ we write:

$$(M_0, s_0, \bar{a}_0) \preceq_{i,q} (M_1, s_1, \bar{a}_1).$$

We write $\approx_{i,q}$ for the relation $\preceq_{i,q} \cap \succeq_{i,q}$. Notice that the game with bounds i, q is finite (since the number of turns is bounded by $i + q$), zero-sum (as can be seen from the winning condition) and has perfect information. Therefore, if $(M_0, s_0, \bar{a}_0) \preceq_{i,q} (M_1, s_1, \bar{a}_1)$ does not hold, then it follows from the Gale-Stewart Theorem that Spoiler has a winning strategy in the EF game of length $\langle i, q \rangle$ starting from the position $\langle M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1 \rangle$.

We write $M_0 \preceq_{i,q} M_1$ as a shorthand for $(M_0, W_0, \varepsilon) \preceq_{i,q} (M_1, W_1, \varepsilon)$.

The following two propositions follow easily from the definition of the game.

PROPOSITION 3.2. If $(M_0, s_0, \bar{a}_0) \preceq_{i,q} (M_1, s_1, \bar{a}_1)$ then for all $i' \leq i$ and $q' \leq q$ it holds $(M_0, s_0, \bar{a}_0) \preceq_{i',q'} (M_1, s_1, \bar{a}_1)$.

Proof. We prove the result by contraposition. Suppose that Spoiler has a winning strategy in the game $EF_{i',q'}(M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1)$ for some $i' \leq i$ and $q' \leq q$. This means that, no matter which choices Duplicator makes during a run, Spoiler can perform i' implication moves and q' quantifier move and force the game to end up in a position

$$\begin{aligned} & \langle M_0, t_0, \bar{a}_0 \bar{b}_0; M_1, t_1, \bar{a}_1 \bar{b}_1 \rangle && \text{(case 1)} \\ \text{or} & \langle M_1, t_1, \bar{a}_0 \bar{b}_0; M_1, t_1, \bar{a}_1 \bar{b}_1 \rangle && \text{(case 2)} \end{aligned}$$

for which there exists an atomic formula $\alpha(\bar{x}, \bar{y})$ such that

$$\begin{aligned} & M_0, t_0 \models \alpha(\bar{a}_0, \bar{b}_0) \text{ and } M_1, t_1 \not\models \alpha(\bar{a}_1, \bar{b}_1) && \text{in case 1} \\ \text{or} & M_1, t_1 \models \alpha(\bar{a}_1, \bar{b}_1) \text{ and } M_0, t_0 \not\models \alpha(\bar{a}_0, \bar{b}_0) && \text{in case 2.} \end{aligned}$$

We will show that, from this point on, Spoiler can still win the game after performing $i - i'$ additional implication moves and $q - q'$ additional quantifier moves: This amounts to a winning strategy in the game $\text{EF}_{i,q}(M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1)$. We will focus here on case 1, since the other case is completely analogous.

- Firstly, Spoiler performs $i - i'$ implication moves, always picking t_1 as the information state and choosing $i := 0$ (that is, maintaining the order of the models). Since Duplicator can only choose substates of t_0 , the position of the game after these moves will be of the form:

$$\langle M_0, t'_0, \bar{a}_0 \bar{b}_0; M_1, t_1, \bar{a}_1 \bar{b}_1 \rangle$$

with $t'_0 \subseteq t_0$.

- Secondly, Spoiler performs $q - q'$ quantification moves in an arbitrary way; no matter what Duplicator responds, we end up in a position of the form:

$$\langle M_0, t'_0, \bar{a}_0 \bar{b}_0 \bar{c}_0; M_1, t_1, \bar{a}_1 \bar{b}_1 \bar{c}_1 \rangle$$

for \bar{c}_0 and \bar{c}_1 of length $q - q'$.

This is indeed a winning position for Spoiler, since

$$\begin{cases} M_0, t'_0 \models \alpha(\bar{a}_0, \bar{b}_0) & \text{(by persistency, since } t'_0 \subseteq t_0) \\ M_1, t_1 \not\models \alpha(\bar{a}_1, \bar{b}_1). \end{cases} \quad \square$$

PROPOSITION 3.3. *Suppose $\langle i, q \rangle \neq \langle 0, 0 \rangle$. $(M_0, s_0, \bar{a}_0) \preceq_{i,q} (M_1, s_1, \bar{a}_1)$ iff the following three conditions are satisfied:*

- If $i > 0$, then $\forall t_1 \subseteq s_1 \exists t_0 \subseteq s_0 : (M_0, t_0, \bar{a}_0) \approx_{i-1,q} (M_1, t_1, \bar{a}_1)$.
- If $q > 0$, then $\forall b_0 \in D_0 \exists b_1 \in D_1 : (M_0, s_0, \bar{a}_0 b_0) \preceq_{i,q-1} (M_1, s_1, \bar{a}_1 b_1)$.
- If $q > 0$, then $\forall b_1 \in D_1 \exists b_0 \in D_0 : (M_0, s_0, \bar{a}_0 b_0) \preceq_{i,q-1} (M_1, s_1, \bar{a}_1 b_1)$.

Proof. The three conditions amount precisely to the fact that, for every move available to Spoiler, there is a corresponding move for Duplicator that leads to a sub-game in which Duplicator has a winning strategy. This is precisely what is needed for Duplicator to have a winning strategy in the original game. \square

3.2. IQ degree and types. We define the *implication degree* (Ideg) and *quantification degree* (Qdeg) of a formula by the following inductive clauses, where p stands for an atomic formula:

$$\begin{array}{ll} \text{Ideg}(p) & = 0 & \text{Qdeg}(p) & = 0 \\ \text{Ideg}(\perp) & = 0 & \text{Qdeg}(\perp) & = 0 \\ \text{Ideg}(\varphi_1 \wedge \varphi_2) & = \max(\text{Ideg}(\varphi_1), \text{Ideg}(\varphi_2)) & \text{Qdeg}(\varphi_1 \wedge \varphi_1) & = \max(\text{Qdeg}(\varphi_1), \text{Qdeg}(\varphi_2)) \\ \text{Ideg}(\varphi_1 \vee \varphi_2) & = \max(\text{Ideg}(\varphi_1), \text{Ideg}(\varphi_2)) & \text{Qdeg}(\varphi_1 \vee \varphi_1) & = \max(\text{Qdeg}(\varphi_1), \text{Qdeg}(\varphi_2)) \\ \text{Ideg}(\varphi_1 \rightarrow \varphi_2) & = \max(\text{Ideg}(\varphi_1), \text{Ideg}(\varphi_2)) + 1 & \text{Qdeg}(\varphi_1 \rightarrow \varphi_1) & = \max(\text{Qdeg}(\varphi_1), \text{Qdeg}(\varphi_2)) \\ \text{Ideg}(\forall x.\varphi) & = \text{Ideg}(\varphi) & \text{Qdeg}(\forall x.\varphi) & = \text{Qdeg}(\varphi) + 1 \\ \text{Ideg}(\exists x.\varphi) & = \text{Ideg}(\varphi) & \text{Qdeg}(\exists x.\varphi) & = \text{Qdeg}(\varphi) + 1 \end{array}$$

The combined degree of a formula is defined as $\text{IQdeg}(\varphi) = \langle \text{Ideg}(\varphi), \text{Qdeg}(\varphi) \rangle$. We define a partial order \leq on such degrees by setting:

$$\langle a, b \rangle \leq \langle a', b' \rangle \iff a \leq a' \text{ and } b \leq b'.$$

We denote by $\mathcal{L}_{i,q}^l$ the set of formulas φ such that $\text{IQdeg}(\varphi) \leq \langle i, q \rangle$ and the set of free variables in φ is included in $\{x_1, \dots, x_l\}$. We can then define the key notion of $\langle i, q \rangle$ -type.

DEFINITION 3.4 $\langle i, q \rangle$ -types. *Let M be a model, s an information state, and \bar{a} a tuple of elements in M of length l . The $\langle i, q \rangle$ -type of $\langle M, s, \bar{a} \rangle$ is the set*

$$\text{tp}_{i,q}(M, s, \bar{a}) := \left\{ \varphi \in \mathcal{L}_{i,q}^l \mid M, s \models \varphi(\bar{a}) \right\}.$$

We also define the following notation:

$$(M_0, s_0, \bar{a}_0) \sqsubseteq_{i,q} (M_1, s_1, \bar{a}_1) \stackrel{\text{def}}{\iff} \text{tp}_{i,q}(M_0, s_0, \bar{a}_0) \subseteq \text{tp}_{i,q}(M_1, s_1, \bar{a}_1)$$

$$(M_0, s_0, \bar{a}_0) \equiv_{i,q} (M_1, s_1, \bar{a}_1) \stackrel{\text{def}}{\iff} \text{tp}_{i,q}(M_0, s_0, \bar{a}_0) = \text{tp}_{i,q}(M_1, s_1, \bar{a}_1)$$

EXAMPLE 3.5. *Consider the models M_0, M_1 in Figure 1. Since $P(x) \in \mathcal{L}_{0,0}^1$ and $M_1, \{v_0, v_1\} \models P(e_1)$, we have $(M_1, \{v_0, v_1\}, \langle e_1 \rangle) \not\sqsubseteq_{0,0} (M_0, \{w_0, w_1\}, \langle d_1 \rangle)$; while $M_0, \{w_0, w_1\} \not\models P(d_1)$.*

Notice that, if the signature is finite, there are only a finite number of non-equivalent formulas of combined degree at most $\langle i, q \rangle$, and consequently only a finite number of $\langle i, q \rangle$ -types. This can be shown inductively as follows:

- The quotient $\mathcal{L}_{0,0}^l / \equiv$ is a distributive lattice under the operations \wedge and \vee . Moreover, since we are working with a finite relational signature, $\mathcal{L}_{0,0}^l$ contains only finitely many atomic formulas, and the equivalence classes of these formulas generate the whole lattice. Since every finitely generated distributive lattice is finite, $\mathcal{L}_{0,0}^l / \equiv$ is finite, which means that $\mathcal{L}_{0,0}^l$ contains only finitely many formulas up to logical equivalence.
- Formulas in $\mathcal{L}_{i,q}^l$ are equivalent to Boolean combinations of formulas in $A \cup B$, for $A = \left\{ \varphi \rightarrow \psi \mid \varphi, \psi \in \mathcal{L}_{i-1,q}^l \right\}$ and $B = \left\{ \exists x. \varphi, \forall x. \varphi \mid \varphi \in \mathcal{L}_{i,q-1}^{l+1} \right\}$ —where we impose by definition $\mathcal{L}_{i,q}^l = \emptyset$ if $i < 0$ or $q < 0$. By induction hypothesis, A and B contain only finitely many non-equivalent formulas.

3.3. The EF theorem. What follows is the first main result of the paper: the relations $\preceq_{i,q}$ and $\sqsubseteq_{i,q}$ coincide.

THEOREM 3.6. *Suppose the signature Σ is finite. Then*

$$(M_0, s_0, \bar{a}_0) \preceq_{i,q} (M_1, s_1, \bar{a}_1) \iff (M_0, s_0, \bar{a}_0) \sqsubseteq_{i,q} (M_1, s_1, \bar{a}_1).$$

Proof. We will prove this by well-founded induction on $\langle i, q \rangle$. For the basic case, $\langle i, q \rangle = \langle 0, 0 \rangle$, we just have to verify that, if Condition (1) holds for all atomic formulas, then it holds for all formulas $\varphi \in \mathcal{L}_{0,0}^l$. This is straightforward. Next, suppose $\langle i, q \rangle > \langle 0, 0 \rangle$ and suppose the claim holds for all $\langle i', q' \rangle < \langle i, q \rangle$. For the left-to-right direction, proceed by contraposition. Suppose that for some $\varphi \in \mathcal{L}_{i,q}^l$ the following conditions

hold:

$$M_0, s_0 \models \varphi(\bar{a}_0) \quad M_1, s_1 \not\models \varphi(\bar{a}_1).$$

We proceed by induction on the structure of φ ; some cases are easy to consider:

- If φ is an atom, it follows $(M_0, s_0, \bar{a}_0) \not\leq_{0,0} (M_1, s_1, \bar{a}_1)$; so, by Proposition 3.2 also $(M_0, s_0, \bar{a}_0) \not\leq_{i,q} (M_1, s_1, \bar{a}_1)$. Thus, in this case the conclusion follows.
- If φ is a conjunction $\psi \wedge \chi$ then we have:

$$\begin{cases} M_0, s_0 \models \psi(\bar{a}_0) \wedge \chi(\bar{a}_0) \implies M_0, s_0 \models \psi(\bar{a}_0) \text{ and } M_0, s_0 \models \chi(\bar{a}_0) \\ M_1, s_1 \not\models \psi(\bar{a}_1) \wedge \chi(\bar{a}_1) \implies M_1, s_1 \not\models \psi(\bar{a}_1) \text{ or } M_1, s_1 \not\models \chi(\bar{a}_1). \end{cases}$$

So, either ψ or χ is a less complex witness of $(M_0, s_0, \bar{a}_0) \not\leq_{i,q} (M_1, s_1, \bar{a}_1)$.

- If φ is a disjunction $\psi \vee \chi$, we can reach a conclusion analogous to the one we reached for conjunction.

The remaining cases are those in which φ is of the form $\psi \rightarrow \chi$, $\forall x.\psi$ or $\exists x.\psi$ (cases $\Rightarrow^1, \Rightarrow^2, \Rightarrow^3$ respectively). Let us consider the three cases separately.

Case \Rightarrow^1 : φ is an implication $\psi \rightarrow \chi$. In this case we have:

$$\begin{aligned} M_1, s_1 \not\models \psi(\bar{a}_1) \rightarrow \chi(\bar{a}_1) &\implies (\exists t_1 \subseteq s_1) [M_1, t_1 \models \psi(\bar{a}_1) \text{ and } M_1, t_1 \not\models \chi(\bar{a}_1)], \\ M_0, s_0 \models \psi(\bar{a}_0) \rightarrow \chi(\bar{a}_0) &\implies (\nexists t_0 \subseteq s_0) [M_0, t_0 \models \psi(\bar{a}_0) \text{ and } M_0, t_0 \not\models \chi(\bar{a}_0)]. \end{aligned}$$

Thus there exists a state $t_1 \subseteq s_1$ with a different $\langle i-1, q \rangle$ -type than every $t_0 \subseteq s_0$ —either because it supports ψ or because it does not support χ . So by induction hypothesis, if S performs a \rightarrow -move and chooses t_1 , for every choice t_0 of D we have $(M_0, t_0, \bar{a}_0) \not\leq_{i-1,q} (M_1, t_1, \bar{a}_1)$. It follows by Proposition 3.3 that $(M_0, s_0, \bar{a}_0) \not\leq_{i,q} (M_1, s_1, \bar{a}_1)$ as wanted.

Case \Rightarrow^2 : φ is a universal $\forall x.\psi$. In this case we have:

$$\begin{aligned} M_1, s_1 \not\models \forall x.\psi(\bar{a}_1, x) &\implies (\exists b_1 \in D_1) M_1, s_1 \not\models \psi(\bar{a}_1, b_1), \\ M_0, s_0 \models \forall x.\psi(\bar{a}_0, x) &\implies (\forall b_0 \in D_0) M_0, s_0 \models \psi(\bar{a}_0, b_0). \end{aligned}$$

Thus if S performs a \forall -move and chooses b_1 , for every choice b_0 of D, by induction hypothesis we have

$$(M_0, s_0, \bar{a}_0 b_0) \not\leq_{i,q-1} (M_1, s_1, \bar{a}_1 b_1).$$

It follows by Proposition 3.3 that $(M_0, s_0, \bar{a}_0) \not\leq_{i,q} (M_1, s_1, \bar{a}_1)$ as wanted.

Case \Rightarrow^3 : φ is an inquisitive existential $\exists x.\psi$. This case is similar to the previous one: S can perform an \exists -move and pick an element b_0 in D_0 with no counterpart in D_1 , and by Proposition 3.3 we get the result.

This completes the proof of the left-to-right direction of the inductive step. Now consider the converse direction. Again, we proceed by contraposition. Suppose that S has a winning strategy in the EF game of length $\langle i, q \rangle$ starting from $(M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1)$. We consider again three cases, depending on the first move of the winning strategy (cases $\Leftarrow^1, \Leftarrow^2, \Leftarrow^3$ respectively).

Case \Leftarrow^1 : the first move is a \rightarrow -move. Suppose S starts by choosing $t_1 \subseteq s_1$. As this is a winning strategy for S, for every choice $t_0 \subseteq s_0$ of D we have

$$(M_0, t_0, \bar{a}_0) \not\leq_{i-1,q} (M_1, t_1, \bar{a}_1) \quad \text{or} \quad (M_1, t_1, \bar{a}_1) \not\leq_{i-1,q} (M_0, t_0, \bar{a}_0).$$

By inductive hypothesis, this translates to

$$\exists \psi_{t_0} \in \text{tp}(t_0) \setminus \text{tp}(t_1) \quad \text{or} \quad \exists \theta_{t_0} \in \text{tp}(t_1) \setminus \text{tp}(t_0),$$

where $\text{tp}(t_0) := \text{tp}_{i-1,q}(M_0, t_0, \bar{a}_0)$ and $\text{tp}(t_1) := \text{tp}_{i-1,q}(M_1, t_1, \bar{a}_1)$.

Given this, there exist two families $\{\psi_{t_0} \mid t_0 \subseteq s_0\}$ and $\{\theta_{t_0} \mid t_0 \subseteq s_0\}$ such that:

$$\begin{cases} \psi_{t_0} \in \text{tp}(t_0) \setminus \text{tp}(t) & \text{if } \text{tp}(t_0) \setminus \text{tp}(t) \neq \emptyset \\ \psi_{t_0} := \perp & \text{otherwise} \end{cases} \quad \begin{cases} \theta_{t_0} \in \text{tp}(t) \setminus \text{tp}(t_0) & \text{if } \text{tp}(t) \setminus \text{tp}(t_0) \neq \emptyset \\ \theta_{t_0} := \top & \text{otherwise.} \end{cases}$$

Moreover, we can suppose the two families to be finite, as there are only a finite number of formulas of degree $\langle i - 1, q \rangle$ up to logical equivalence (see Section 3.2). Define now the formula φ as follows:

$$\varphi := \bigwedge_{t_0 \subseteq s_0} \theta_{t_0} \rightarrow \bigvee_{t_0 \subseteq s_0} \psi_{t_0}.$$

We have: (i) $\text{IQdeg}(\varphi) \leq \langle i, q \rangle$, (ii) $\varphi \notin \text{tp}_{i,q}(M_0, s_0, \bar{a}_0)$ (since by construction φ is falsified at $t_1 \subseteq s_1$) and (iii) $\varphi \in \text{tp}_{i,q}(M_1, s_1, \bar{a}_1)$ (since by construction φ holds at every state $t_0 \subseteq s_0$). Thus we have $(M_0, t_0, \bar{a}_0) \not\sqsubseteq_{i-1,q} (M_1, t_1, \bar{a}_1)$, as we wanted.

Case \Leftarrow^2 : the first move is a \forall -move. Suppose S starts by choosing $b_1 \in D_1$. As this is a winning strategy for S , for every choice $b_0 \in D_0$ of D we have

$$(M_0, s_0, \bar{a}_0 b_0) \not\sqsubseteq_{i,q-1} (M_1, s_1, \bar{a}_1 b_1).$$

By induction hypothesis, the above translates to

$$\exists \psi_{b_0} \in \text{tp}(b_0) \setminus \text{tp}(b_1),$$

where $\text{tp}(b_0) := \text{tp}_{i,q-1}(M_0, s_0, \bar{a}_0 b_0)$ and $\text{tp}(b_1) := \text{tp}_{i,q-1}(M_1, s_1, \bar{a}_1 b_1)$.

Now the formula

$$\varphi := \forall x. \bigvee_{b_0 \in D_0} \psi_{b_0}$$

has IQ-degree at most $\langle i, q \rangle$, and by construction we have $\varphi \in \text{tp}_{i,q}(M_0, s_0, \bar{a}_0)$ and $\varphi \notin \text{tp}_{i,q}(M_1, s_1, \bar{a}_1)$. Thus, we have $(M_0, t_0, \bar{a}_0) \not\sqsubseteq_{i-1,q} (M_1, t_1, \bar{a}_1)$.

Case \Leftarrow^3 : the first move is a \exists -move. Reasoning as in the previous case, we find that there exists a $b_0 \in D_0$ —the element chosen by S —such that for every $b_1 \in D_1$

$$\exists \theta_{b_1} \in \text{tp}(t_0) \setminus \text{tp}(t_1).$$

In particular, it follows that the formula

$$\varphi := \exists x. \bigwedge_{b_1 \in D_1} \psi_{b_1}$$

is a formula of complexity at most $\langle i, q \rangle$ such that $\varphi \in \text{tp}_{i,q}(M_0, s_0, \bar{a}_0)$ and $\varphi \notin \text{tp}_{i,q}(M_1, s_1, \bar{a}_1)$. Again, it follows that $(M_0, t_0, \bar{a}_0) \not\sqsubseteq_{i-1,q} (M_1, t_1, \bar{a}_1)$. \square

As a corollary, we also get a game-theoretic characterization of the distinguishing power of formulas in the $\langle i, q \rangle$ -fragment of InqBQ .

COROLLARY 3.7. *For a finite signature Σ , we have:*

$$(M_0, s_0, \bar{a}_0) \approx_{i,q} (M_1, s_1, \bar{a}_1) \iff (M_0, s_0, \bar{a}_0) \equiv_{i,q} (M_1, s_1, \bar{a}_1).$$

3.4. Extending the result to function symbols. The results we just obtained assume that the signature Σ is relational. However, it is not hard to extend them to the case in which Σ contains function symbols (including nullary function symbols, i.e., constant symbols). In InqBQ, function symbols are interpreted rigidly: if $f \in \Sigma$ is an n -ary function symbol, then the interpretation function I of a model M must assign to all worlds w in the model the same function $I_w(f) : D^n \rightarrow D$.⁴

As in the case of classical logic [13] Section 3.3, the presence of function symbols requires some care in formulating the EF game. The reason is that allowing atomic formulas to contain arbitrary occurrences of function symbols allows us to generate with a finite number of choices in the game an infinite sub-structure of the model—which spoils the crucial locality feature of the game. Technically, a simple way to circumvent the problem this is to follow [13] Section 3.3 and work with formulas which are *unnested*.

DEFINITION 3.8 (Unnested formula). *An unnested atomic formula is a formula of one of the following forms:*

$$x = y \qquad c = y \qquad f(\bar{x}) = \bar{y} \qquad R(\bar{x}).$$

An unnested formula is a formula that contains only unnested atoms.

Examples of *nested formulas*—i.e., non-unnested formulas—are $f(x) = g(y)$, $R(f(x))$ and $f(c) = x$.

We can now make the following amendments to the definition above: (i) the winning conditions for the game are determined by looking at whether Equation (1) is satisfied for all *unnested* atomic formulas, and (ii) the $\langle i, q \rangle$ -types are re-defined as sets of *unnested* formulas of degree at most $\langle i, q \rangle$. Other than that, the statement of the result and the proof are the same as above.

Using identity we can turn an arbitrary formula into an equivalent unnested one (e.g., replacing $P(f(x))$ with $\forall y.((y = f(x)) \rightarrow Py)$) so the restriction to unnested formula is not a limitation to the generality of the game-theoretic characterization; rather, it can be seen as an indirect way of assigning formulas containing function symbols with the appropriate $\langle i, q \rangle$ -degree—making explicit a quantification which is implicit in the presence of a function symbol.

3.5. A symmetric version of the game. As noticed before, a difference between the Ehrenfeucht–Fraïssé game for classical logic and the game introduced in Section 3 is that the latter is asymmetric—the two models under consideration do not play the same role. This allows us to study the relation $\sqsubseteq_{i,q}$ in addition to the relation $\equiv_{i,q}$. This contrast with the situation in classical first-order logic, where the relations \sqsubseteq_q and \equiv_q coincide due to the semantics of negation. A natural question is whether we can define a symmetric version of the game which directly characterizes the relation $\equiv_{i,q}$. In what follows we will consider a naïve modification of the game to obtain a symmetric version and study the induced equivalence relation between models. The symmetric way is just

⁴ In the general case, non-rigid function symbols are also allowed; however, such symbols can be dispensed with as usual in favor of relation symbols constrained by suitable axioms. See Section 4.3.5 of [2] for the details.

like the game introduced above, except that we replace the \rightarrow -move and the winning condition as follows.

- \rightarrow -move. S picks a sub-state $t_i \subseteq s_i$ for $i = 0$ or 1 . D responds with a sub-state $t_{1-i} \subseteq s_{1-i}$. The game continues from $\langle M_0, t_0, \bar{a}_0; M_1, t_1, \bar{a}_1 \rangle$.
- Winning condition for D: for all atomic formulas $\alpha(x_1, \dots, x_n)$ where n is the size of the tuples \bar{a}_0 and \bar{a}_1 , we have:

$$M_0, s_0 \models \alpha(\bar{a}_0) \iff M_1, s_1 \models \alpha(\bar{a}_1).$$

Notice that, compared to the original version, the implication has become a biconditional.

We will indicate with $M_0, s_0, \bar{a}_0 \approx_{i,q}^s M_1, s_1, \bar{a}_1$ the existence of a winning strategy for Duplicator in the symmetric game from position $\langle M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1 \rangle$ with bounds $\langle i, q \rangle$; we indicate by $M_0, s_0, \bar{a}_0 \approx^s M_1, s_1, \bar{a}_1$ the existence of a winning strategy for D in the unbounded version of the symmetric game. We will also use notational conventions analogous to the ones introduced for the original game. Notice that, as anticipated, the roles of the two models in the game are interchangeable:

LEMMA 3.9. $\approx_{i,q}^s$ and \approx^s are symmetric relations.

Comparing this version with the original one, we clearly made Spoiler’s life much easier: now he can perform \rightarrow -moves without any restrictions on the model from which he can choose the state; and the winning condition for Duplicator is more restrictive than the original one. So the following result should not come as a surprise:

LEMMA 3.10. If $M_0, s_0, \bar{a}_0 \approx_{i,q}^s M_1, s_1, \bar{a}_1$, then $M_0, s_0, \bar{a}_0 \preceq_{i,q} M_1, s_1, \bar{a}_1$.

Proof. The idea of the proof is simple: given a winning strategy for Duplicator in the game $EF_{i,q}^s(M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1)$, this is also a winning strategy in the game $EF_{i,q}(M_0, s_0, \bar{a}_0; M_1, s_1, \bar{a}_1)$. The details are left to the reader. \square

As an immediate corollary we obtain the following result.

COROLLARY 3.11. Suppose the signature Σ is finite. Then

$$(M_0, s_0, \bar{a}_0) \approx_{i,q}^s (M_1, s_1, \bar{a}_1) \implies (M_0, s_0, \bar{a}_0) \equiv_{i,q} (M_1, s_1, \bar{a}_1).$$

However, the converse of Corollary 3.11 does not hold in general.


PROPOSITION 3.12.

$$(M_0, s_0, \bar{a}_0) \equiv_{i,q} (M_1, s_1, \bar{a}_1) \not\Rightarrow (M_0, s_0, \bar{a}_0) \approx_{i,q}^s (M_1, s_1, \bar{a}_1).$$

Proof. Consider the models M and N in Figure 2a. We have:

- $M \preceq_{0,1} N$: the winning strategy for Duplicator is described in Table 2b;
- $M \succeq_{0,1} N$: the winning strategy for Duplicator is described in Table 2c;
- $M \not\approx_{0,1}^s N$: if Spoiler picks the element d_2 , every move of Duplicator leads to Spoiler’s victory. \square

Thus, unlike the original version, the symmetric version of the game does not provide us with an exact characterization of the relation $\equiv_{i,q}$ determined by the logic.



(A) The two models considered in Example 3.5.

S plays ...	D responds ...
d_1	e_1
d_2	e_1
d_3	e_3
e_1	d_1
e_2	d_3
e_3	d_3

(B) Winning strategy for D in the game $EF_{0,1}(M, N)$.

S plays ...	D responds ...
d_1	e_1
d_2	e_3
d_3	e_3
e_1	d_1
e_2	d_1
e_3	d_3

(C) Winning strategy for D in the game $EF_{0,1}(N, M)$.

Fig. 2.

Nevertheless, the symmetric game is still useful: if we can show that Duplicator has a winning strategy in the symmetric game with bounds $\langle i, q \rangle$ between two models, this suffices to show that these models are indistinguishable by formulae of degree $\langle i, q \rangle$. This is convenient, since strategies are often easier to describe in the symmetric game than in the original game, since we do not have to keep track of how the role of the models gets swapped in the course of the game. Indeed, we use this strategy in the proof of Theorem 4.5.

§4. Characterizing the cardinality quantifiers definable in InqBQ. In this section we will use the EF-game for InqBQ to study in detail what InqBQ can express about the number of individuals satisfying a predicate P . The sentences we are concerned with include not only statements about the number of individuals satisfying P , like those in (1), but also questions about the number of individuals satisfying P , like those in (2).

- (1)
 - a. There is no P .
 - b. There are at least three P .
 - c. The number of P is even.
 - d. There are infinitely many P .
- (2)
 - a. Are there any P ?
 - b. How many P are there?
 - c. Is the number of P even, or odd?
 - d. Are there infinitely many P ?

Which among the statements in (1) and the questions in (2) can be expressed in InqBQ? Instead of pursuing a direct answer to this question, we will tackle the problem from a more general perspective. We will see that, in an inquisitive setting, all these sentences instantiate the form $Qx.Px$, where Q is a quantifier which is sensitive only to the cardinality of its argument. Thus—interestingly—in the inquisitive setting, not only *no* and *at least three*, but also *how many* can be viewed as generalized quantifiers. We can then ask which cardinality quantifiers are expressible in InqBQ. In this section, we will establish a simple answer to this question. From this answer, a verdict about the definability of the examples above, as well as many other similar examples, can be easily reached.

We will first look at cardinality quantifiers in the setting of standard first-order logic, FOL, and recall the characterization of cardinality quantifiers expressible in FOL; we will then present a generalization of the notion of a cardinality quantifier to InqBQ, which encompasses also inquisitive quantifiers like *how many*; finally, we will use the Ehrenfeucht–Fraïssé game introduced in Section 3 to provide a characterization of the cardinality quantifiers expressible in InqBQ and use this characterization to show that, just like many interesting statements about cardinalities are not expressible in FOL, so many interesting questions about cardinalities are not expressible in InqBQ.

4.1. Cardinality quantifiers in classical first-order logic. In classical logic, a formula $\alpha(x)$, with at most the variable x free, determines, relative to a model \mathcal{M} , a corresponding set of individuals:

$$\alpha_{\mathcal{M}} := \{d \in D \mid \mathcal{M} \models \alpha(d)\}.$$

Let K be a class of cardinals. This is an operator that can be added to classical first-order logic by stipulating that if $\alpha(x)$ is a classical formula with at most x free, then $Q_K x.\alpha(x)$ is a formula, with the following semantics ($\#$ denotes the cardinality of a set):⁵

$$\mathcal{M} \models Q_K x.\alpha(x) \iff \#\alpha_{\mathcal{M}} \in K.$$

By a cardinality quantifier we mean a quantifier which is of the form Q_K for some class of cardinals K . Notice that the existential quantifier \exists is a cardinality quantifier, since $\exists = Q_{\text{Card} \setminus \{0\}}$, for Card the class of all cardinals. By contrast, the universal quantifier \forall is not a cardinality quantifier, since the condition $\mathcal{M} \models \forall x.P(x)$, namely, $P_{\mathcal{M}} = D$, cannot be formulated solely in terms of $\#P_{\mathcal{M}}$.⁶

Let $\chi_K[P]$ be a FOL-formula (thus, not containing Q_K). We say that $\chi_K[P]$ defines Q_K if $Q_K x.Px \equiv \chi_K[P]$. It is not hard to see that if this is the case, then for every

⁵ One can, more generally, allow the formation of the formula $Q_K x.\alpha$ for any formula α , even when α contains free variables besides x . Extending the semantic clause to this case is straightforward: we just have to relativize the clause to an assignment function g . However, we restrict to the case in which $Q_K x.\alpha$ is a sentence, since this does not lead to a loss of generality for our purposes, and it is convenient not to have assignments around all the time.

⁶ In this paper, we focus on cardinality quantifiers of type (1), which operate on a single unary predicate. More generally, one could consider cardinality quantifiers of type $\langle n_1, \dots, n_k \rangle$, which operate on k predicates of arities n_1, \dots, n_k respectively. It seems quite possible that the characterization result given here can be extended to this general setting. However, we leave this extension for future work.

formula $\alpha(x)$ we have $Q_K x. \alpha(x) \equiv \chi_K[\alpha]$. We say that the quantifier Q_K is definable in FOL if there is a FOL-formula which defines it.

The statements in (1) can all be seen as having the form $Qx.Px$, where Q is a cardinality quantifier. Indeed, we have the following characterizations, where $[3, \dots)$ is the class of cardinals ≥ 3 ; Even is the set of even natural numbers; Inf is the class of infinite cardinals.

- (3) a. $\mathcal{M} \models (1.a) \iff P_{\mathcal{M}} = \emptyset \iff \#P_{\mathcal{M}} \in \{0\}$
- b. $\mathcal{M} \models (1.b) \iff \#P_{\mathcal{M}} \geq 3 \iff \#P_{\mathcal{M}} \in [3, \dots)$
- c. $\mathcal{M} \models (1.c) \iff \#P_{\mathcal{M}} \text{ is even} \iff \#P_{\mathcal{M}} \in \text{Even}$
- d. $\mathcal{M} \models (1.d) \iff \#P_{\mathcal{M}} \text{ is infinite} \iff \#P_{\mathcal{M}} \in \text{Inf}$.

What cardinality quantifiers are definable in classical first-order logic? That is, for what classes K of cardinals is the quantifier Q_K definable? The answer is given by the following theorem, which is an easy application of EF-games for FOL (and seems, to the best of our knowledge, to be folklore).

THEOREM 4.1. *Let K be a class of cardinals. The quantifier Q_K is definable in first-order logic if and only if there exists a natural number n such that K contains either all or none of the cardinals $\kappa \geq n$.*

Consider again the statements in (1), repeated below for convenience with the corresponding classes of cardinals given on the right. It follows immediately from the characterization that the first two statements are expressible in classical first-order logic, while the third and fourth are not.

- (4) a. There is no P . $K = \{0\}$
- b. There are at least three P . $K = [3, \dots)$
- c. The number of P is even. $K = \text{Even}$
- d. There are infinitely many P . $K = \text{Inf}$

4.2. Cardinality quantifiers in InqBQ. Let us now turn to the inquisitive case. A model M for inquisitive first-order logic represents a variety of states of affairs, one for each possible world w . At each world w , the state of affairs is represented by the first-order structure \mathcal{M}_w , having as its domain the set $D_w := D/\sim_w$. Let $\alpha(x)$ be a classical formula with at most the variable x free. Relative to each world w , $\alpha(x)$ determines an extension α_w , which is a set of individuals from D_w :

$$\alpha_w := \{d \in D_w \mid \mathcal{M}_w \models \alpha(d)\}.$$

Therefore, relative to an information state s , the formula $\alpha(x)$ determines a corresponding set of cardinals, $\{\#\alpha_w \mid w \in s\}$. We refer to this set of cardinals as the *cardinality trace* of $\alpha(x)$ in s .

DEFINITION 4.2 (Cardinality trace). *Let M be a model, s an information state, and $\alpha(x)$ a classical formula where at most the variable x occurs free. The cardinality trace of $\alpha(x)$ in s is the set of cardinals:*

$$\text{tr}_s(\alpha) = \{\#\alpha_w \mid w \in s\}.$$

A cardinal κ is in $\text{tr}_s(\alpha)$ if, according to the information available in s , κ might be the number of elements satisfying $\alpha(x)$; that is, if it might be the case that the extension of $\alpha(x)$ has cardinality κ . Thus, $\text{tr}_s(\alpha)$ captures exactly the information available in s about the number of individuals satisfying $\alpha(x)$.

Now let \mathbb{K} be a class of sets of cardinals. We associate with \mathbb{K} a corresponding quantifier $Q_{\mathbb{K}}$. We can add this quantifier to InqBQ by stipulating that if $\alpha(x)$ is a classical formula with at most x free, then $Q_{\mathbb{K}x}.\alpha(x)$ is a formula, interpreted by the following clause⁷:

$$M, s \models Q_{\mathbb{K}x}.\alpha(x) \iff \text{tr}_s(\alpha) \in \mathbb{K}.$$

A cardinality quantifier is a quantifier which is of the form $Q_{\mathbb{K}}$, where \mathbb{K} is a class of sets of cardinals.

Let $\chi_{\mathbb{K}}[P]$ be an InqBQ -formula (thus, without cardinality quantifiers). We say that $\chi_{\mathbb{K}}[P]$ defines the quantifier $Q_{\mathbb{K}}$ if $Q_{\mathbb{K}x}.Px \equiv \chi_{\mathbb{K}}[P]$. Again, it is not hard to see that if this holds, then for every classical formula $\alpha(x)$ we have $Q_{\mathbb{K}x}.\alpha(x) \equiv \chi_{\mathbb{K}}[\alpha]$. We say that $Q_{\mathbb{K}}$ is definable in InqBQ if there is an InqBQ -formula that defines it.

In order to make the notion of a cardinality quantifier more concrete, let us see how the statements in (1) and the questions in (2) can be seen as instantiating the form $Qx.Px$ where Q is a cardinality quantifier in the sense of inquisitive logic.

Consider first the statements in (1). In general, in inquisitive semantics a statement α is supported by a state s iff the information available in s implies that α is true. This means that α is true at all worlds $w \in s$. Keeping this in mind, we can see that the statements in (1) have the following semantics:

- (5) a. $M, s \models (1.a) \iff \forall w \in s : P_w = \emptyset \iff \text{tr}_s(P) \subseteq \{0\}$
- b. $M, s \models (1.b) \iff \forall w \in s : \#P_w \geq 3 \iff \text{tr}_s(P) \subseteq [3, \dots)$
- c. $M, s \models (1.c) \iff \forall w \in s : \#P_w \text{ is even} \iff \text{tr}_s(P) \subseteq \text{Even}$
- d. $M, s \models (1.d) \iff \forall w \in s : \#P_w \text{ is infinite} \iff \text{tr}_s(P) \subseteq \text{Inf}$

Let us now check that all these statements correspond to statements of the form $Qx.Px$ for Q a cardinality quantifier. For this, we introduce a useful notation.

DEFINITION 4.3 (Downward closure of a class). *Let K be a class. We denote by K^\downarrow the class consisting of all sets X such that $X \subseteq K$.*

Thus, if K is a set, then $K^\downarrow = \wp(K)$. However, if K is a proper class, then K^\downarrow will not be a set either; moreover, K^\downarrow will not contain K , since K is not a set.

Now consider the cardinality quantifiers Q_1 – Q_4 determined by the following classes:

$$\mathbb{K}_1 = \{0\}^\downarrow \quad \mathbb{K}_2 = [3, \dots)^\downarrow \quad \mathbb{K}_3 = \text{Even}^\downarrow \quad \mathbb{K}_4 = \text{Inf}^\downarrow.$$

We have:

$$\begin{aligned} M, s \models Q_1x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_1 \iff \text{tr}_s(P) \subseteq \{0\} \iff M, s \models (1.a) \\ M, s \models Q_2x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_2 \iff \text{tr}_s(P) \subseteq [3, \dots) \iff M, s \models (1.b) \\ M, s \models Q_3x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_3 \iff \text{tr}_s(P) \subseteq \text{Even} \iff M, s \models (1.c) \\ M, s \models Q_4x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_4 \iff \text{tr}_s(P) \subseteq \text{Inf} \iff M, s \models (1.d) \end{aligned}$$

Next, consider the questions in (2). Start with (2.a), the question whether there are any P . This question is settled in an information state s in case the information in

⁷ The reason for restricting the application of Q_K to classical formulas is that $Q_Kx.\alpha(x)$ only looks at the semantics of α with respect to worlds. Non-classical formulas only become significant when interpreted relative to information states; relative to single worlds, the operators \forall and \exists collapse on their classical counterparts \forall and \exists . Therefore, while extending our quantifiers to operate on non-classical formulas is not problematic, it is also not interesting.

s implies that there are no P , or it implies that there are some P . The former is the case if the extension of P is empty in all worlds $w \in s$. The latter is the case if the extension of P is non-empty in all worlds $w \in s$. Thus, the semantics of (2.a) is as follows.

$$\begin{aligned} M, s \models (2.a) &\iff (\forall w \in W : P_w = \emptyset) \text{ or } (\forall w \in W : P_w \neq \emptyset) \\ &\iff (\forall w \in W : \#P_w = 0) \text{ or } (\forall w \in W : \#P_w \geq 1) \\ &\iff \text{tr}_s(P) = \{0\} \text{ or } \text{tr}_s(P) \subseteq [1, \dots]. \end{aligned}$$

Second, consider the question (2.b), how many individuals are P . This question is settled in an information state s if the information available in s determines exactly how many individuals are P . This is the case if there is a cardinal κ such that at every world $w \in s$, the extension P_w contains κ elements.⁸

$$\begin{aligned} M, s \models (2.b) &\iff \exists \kappa \forall w \in W : \#P_w = \kappa \\ &\iff \text{tr}_s(P) \text{ contains at most one element} \\ &\iff \text{tr}_s(P) \subseteq \{\kappa\} \text{ for some cardinal } \kappa. \end{aligned}$$

Next, consider (2.c), the question whether the number of P is even or odd. This is settled in an information state s in case the information available in s implies that the number of P is even, or that the number of P is odd.⁹ The former holds if the extension of P is even at every world in s . The latter holds if the extension of P is odd at every world in s .

$$\begin{aligned} M, s \models (2.c) &\iff (\forall w \in W : \#P_w \text{ is even}) \text{ or } (\forall w \in W : \#P_w \text{ is odd}) \\ &\iff \text{tr}_s(P) \subseteq \text{Even} \text{ or } \text{tr}_s(P) \subseteq \text{Odd}. \end{aligned}$$

Finally, consider (2.d), the question whether there are infinitely many P . This is settled in an information state s in case the information available in s implies that there are infinitely many P , or it implies that there aren't infinitely many P . The former is the case if the extension of P is infinite at every world $w \in s$, while the latter is the case if the extension of P is finite at every world $w \in s$.

$$\begin{aligned} M, s \models (2.d) &\iff (\forall w \in s : \#P_w \text{ is finite}) \text{ or } (\forall w \in s : \#P_w \text{ is infinite}) \\ &\iff \text{tr}_s(P) \subseteq \text{Fin} \text{ or } \text{tr}_s(P) \subseteq \text{Inf}. \end{aligned}$$

Now consider four cardinality quantifiers, Q_5 – Q_8 , determined by the following classes:

- (6) a. $\mathbb{K}_5 = \{0\}^\downarrow \cup [1, \dots]^\downarrow$
- b. $\mathbb{K}_6 = \bigcup \{ \{\kappa\}^\downarrow \mid \kappa \text{ a cardinal} \}$
- c. $\mathbb{K}_7 = \text{Even}^\downarrow \cup \text{Odd}^\downarrow$
- d. $\mathbb{K}_8 = \text{Fin}^\downarrow \cup \text{Inf}^\downarrow$

⁸ An equivalent way of formulating the same condition is to say that (2.b) is settled in s iff the number of P is the same at all the worlds in s : $M, s \models (2.b) \iff \forall w, w' \in s : \#P_w = \#P_{w'}$.

⁹ Notice that the question presupposes that the number of P is either even or odd. Since all and only the finite cardinals are even or odd, the question presupposes that the number of P is finite. About the way presuppositions of questions are interpreted in inquisitive logic, see Section 1.3 of [2].

Then we have:

$$\begin{aligned}
 M, s \models Q_5x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_5 &\iff \text{tr}_s(P) \subseteq \{0\} \text{ or } \text{tr}_s(P) \subseteq [1, \dots] \\
 &&\iff M, s \models (2.a) \\
 M, s \models Q_6x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_6 &\iff \text{tr}_s(P) \subseteq \{\kappa\} \text{ for some } \kappa \\
 &&\iff M, s \models (2.b) \\
 M, s \models Q_7x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_7 &\iff \text{tr}_s(P) \subseteq \text{Even} \text{ or } \text{tr}_s(P) \subseteq \text{Odd} \\
 &&\iff M, s \models (2.c) \\
 M, s \models Q_8x.Px &\iff \text{tr}_s(P) \in \mathbb{K}_8 &\iff \text{tr}_s(P) \subseteq \text{Fin} \text{ or } \text{tr}_s(P) \subseteq \text{Inf} \\
 &&\iff M, s \models (2.d)
 \end{aligned}$$

So, in the inquisitive setting, a new range of “inquisitive” cardinality quantifiers come into play, which combine with a property to yield questions like those exemplified in (2). In addition to standard cardinality quantifiers like ‘no’, ‘at least three’, ‘infinitely many’, we also have new, question-forming cardinality quantifiers like ‘how many’ and ‘whether finitely or infinitely many’.

4.3. Characterization. What cardinality quantifiers can be expressed in InqBQ? Given that, in the inquisitive setting, cardinality quantifiers are in one-to-one correspondence with classes of sets of cardinals, this question can be made precise as follows.

QUESTION 4.4. *For which classes of sets of cardinals \mathbb{K} is the quantifier $Q_{\mathbb{K}}$ definable in InqBQ?*

The next theorem provides an answer to this question. In essence, what the theorem says is that the cardinality quantifiers definable in InqBQ are all and only the inquisitive disjunctions of cardinality quantifiers definable in classical first-order logic.¹⁰ Before stating the Theorem, let us fix some useful notations. For any natural number n , we let:

- $[0, n] := \{m \in \text{Card} \mid m \leq n\}$
- $[n, \dots] := \{\kappa \in \text{Card} \mid \kappa \geq n\}$

Moreover, we introduce an equivalence relation $=_n$ that disregards differences between cardinals larger than n . More precisely, if κ and κ' are two cardinals:

$$\kappa =_n \kappa' \iff \kappa = \kappa' \text{ or } \kappa, \kappa' > n.$$

If A and B are sets of cardinals, we write $A =_n B$ if A and B are the same set, modulo identifying all cardinals larger than n :

$$\begin{aligned}
 A =_n B &\iff \forall \kappa \in A \exists \kappa' \in B \text{ such that } \kappa =_n \kappa' \text{ and} \\
 &\quad \forall \kappa' \in B \exists \kappa \in A \text{ such that } \kappa =_n \kappa'.
 \end{aligned}$$

¹⁰ While we have not specified a general notion of inquisitive generalized quantifier here, a natural notion should allow as an instance the quantifier Q_0 whose semantics is given by: $M, s \models Q_0x.Px \iff \forall w, w' \in s : I_w(P) = I_{w'}(P)$. Informally, $Q_0x.Px$ expresses the question “which elements are P ?” Now this quantifier is definable in InqBQ by the formula $\forall x.?Px$, which is clearly not equivalent to an inquisitive disjunction of classical formulas. This shows that the result we show here is really specific for cardinality quantifiers.

Moreover, we say that a class of sets of cardinals \mathbb{K} is:

- $=_n$ -invariant, if whenever $B \in \mathbb{K}$ and $A =_n B$ we have $A \in \mathbb{K}$ and
- downward-closed, if whenever $B \in \mathbb{K}$ and $A \subseteq B$ we have $A \in \mathbb{K}$.

We can now state our second main result.

THEOREM 4.5 (Characterization of cardinality quantifiers definable in InqBQ). *Let \mathbb{K} be a class of sets of cardinals. The following are equivalent:*

1. *The cardinality quantifier $Q_{\mathbb{K}}$ is definable in InqBQ.*
2. *$\mathbb{K} = K_1^{\downarrow} \cup \dots \cup K_n^{\downarrow}$ where for each $K_i \subseteq \text{Card}$ there exists a natural number m such that K_i contains either all or none of the cardinals $\kappa \geq m$.*
3. *\mathbb{K} is downward closed and $=_m$ -invariant for some natural number m .*

Proof. We show that $2 \Rightarrow 1 \Rightarrow 3 \Rightarrow 2$.

[2 \Rightarrow 1] Suppose $\mathbb{K} = K_1^{\downarrow} \cup \dots \cup K_n^{\downarrow}$ where for each K_i there exists a natural number m such that K_i contains either all or none of the cardinals $\kappa \geq m$. By Theorem 4.1, for each K_i we have a classical formula χ_i such that, in classical first-order logic:

$$\mathcal{M} \models \chi_i \iff \#P_{\mathcal{M}} \in K_i.$$

These formulas are also formulas of InqBQ, and it follows from Proposition 2.3 that we have:

$$\begin{aligned} M, s \models \chi_i &\iff \forall w \in s : M_w \models \chi_i \\ &\iff \forall w \in s : \#P_{M_w} \in K_i \\ &\iff \text{tr}_s(P) \subseteq K_i \\ &\iff \text{tr}_s(P) \in K_i^{\downarrow}. \end{aligned}$$

Now consider the inquisitive disjunction $\chi_1 \vee \dots \vee \chi_n$. We have:

$$\begin{aligned} M, s \models \chi_1 \vee \dots \vee \chi_n &\iff M, s \models \chi_1 \text{ or } \dots \text{ or } M, s \models \chi_n \\ &\iff \text{tr}_s(P) \in K_1^{\downarrow} \text{ or } \dots \text{ or } \text{tr}_s(P) \in K_n^{\downarrow} \\ &\iff \text{tr}_s(P) \in K_1^{\downarrow} \cup \dots \cup K_n^{\downarrow} \\ &\iff \text{tr}_s(P) \in \mathbb{K} \\ &\iff M, s \models Q_{\mathbb{K}}x.Px. \end{aligned}$$

This shows that the InqBQ formula $\chi_1 \vee \dots \vee \chi_n$ defines the quantifier $Q_{\mathbb{K}}$.

[1 \Rightarrow 3] Next, consider the implication from 1 to 3. Suppose $Q_{\mathbb{K}}$ is definable in InqBQ by a formula $\varphi_{\mathbb{K}}$. We need to show that \mathbb{K} is downward closed and $=_m$ invariant for some natural number m .

We firstly show that \mathbb{K} is downward closed. Suppose $A \subseteq B \in \mathbb{K}$. This means that there exists a model M and an information state s such that $M, s \models \varphi_{\mathbb{K}}$ and $\text{tr}_s(P) = B$. Consider now the state $t := \{w \in s \mid \#P_w \in A\} \subseteq s$. By definition we have $\text{tr}_t(P) = A$; and by persistency $M, t \models \varphi_{\mathbb{K}}$. Thus $A \in \mathbb{K}$, as wanted.

Next, we show that \mathbb{K} is closed under $=_m$ for some m . We want to show that the condition above holds for $m = q$, where q is the quantifier degree of the defining formula $\varphi_{\mathbb{K}}$. So, suppose $A \in \mathbb{K}$ and $A =_q B$. If we find two information models M, N such that $\text{tr}_{WM}(P) = A$, $\text{tr}_{WN}(P) = B$ and $M \approx_{i,q} N$ then we are done, since in this case:

$$A \in \mathbb{K} \iff M \models \varphi_{\mathbb{K}} \iff N \models \varphi_{\mathbb{K}} \iff B \in \mathbb{K}.$$

Consider enumerations of the sets A and B :¹¹ $A := \{\kappa_\alpha \mid \alpha < \lambda\}$ and $B := \{\kappa'_\alpha \mid \alpha < \lambda\}$ which both start with the same initial sequence $\langle \kappa_1, \dots, \kappa_l \rangle = \langle \kappa'_1, \dots, \kappa'_l \rangle$ enumerating $A \cap [0, q] = B \cap [0, q]$. Let M, N be the models defined by the following clauses:

$$\begin{array}{ll}
 W^M := \{w_\alpha \mid \alpha < \lambda\} & W^N := \{w_\alpha \mid \alpha < \lambda\} = W^M \\
 D^M := \left\{ d_\beta^\alpha \mid \alpha < \lambda \ \& \ \beta < \kappa_\alpha \right\} & D^N := \left\{ e_\beta^\alpha \mid \alpha < \lambda \ \& \ \beta < \kappa'_\alpha \right\} \\
 I_{w_\gamma}^M(P) \left(d_\beta^\alpha \right) \iff \alpha = \gamma & I_{w_\gamma}^N(P) \left(e_\beta^\alpha \right) \iff \alpha = \gamma \\
 M \text{ is an id-model} & N \text{ is an id-model.}
 \end{array}$$

An example of these models is given in Figure 3. Notice that $\#I_{w_\alpha}^M(P) = \kappa_\alpha$ and $\#I_{w_\alpha}^N(P) = \kappa'_\alpha$. In particular, it follows that $\text{tr}_{W^M}(P) = A$ and $\text{tr}_{W^N}(P) = B$. So if we show that $M \approx_{i,q} N$ then we are done. In order to show this, we present here a winning strategy for Duplicator in the symmetric version of the EF-game between M and N (cf. Section 3.5):

- If Spoiler plays an implication move and chooses an information state s from either of the models, then Duplicator responds by choosing the same state s from the other model (this is possible since $W^M = W^N$).
- If Spoiler plays a quantifier move and chooses an element d_β^α from the model M , we consider two separate cases:
 - If $d_\beta^\alpha = a_i$ for some i , that is, it has already been picked during the run—by either Spoiler or Duplicator—then Duplicator responds by choosing b_i .
 - If d_β^α has not been previously picked, then Duplicator chooses an element e_γ^α (notice that the elements have the same superscript and possibly different subscripts) which has not been previously picked during the run. The fact that duplicator can find such an element is guaranteed by $A =_q B$: this means that either $\kappa_\alpha = \kappa'_\alpha$, or else $\kappa_\alpha, \kappa'_\alpha > q$. In the former case the number of elements d_β^α and e_γ^α is exactly the same; in the latter case the number of elements e_γ^α is larger than the number of quantifier moves in the game.
- If Spoiler plays a quantifier move and chooses an element e_β^α from the model N , then Duplicator applies the same strategy as in the previous case, swapping the roles of the models M and N .

Notice that with this strategy Duplicator ensures that at the end of the run the final position:

1. has the same state s for both models;
2. $a_i = a_j$ if and only if $b_i = b_j$;
3. corresponding elements a_i, b_i in the two models have the same superscripts, that is, a_i and b_i are of the form d_β^α and e_β^α respectively.

¹¹ In the enumerations, we allow for repetitions of the same elements with different indices. This allows us to use the same cardinal λ as the set of indices for both sets A and B .

<i>M</i>			
	w_0	w_1	w_2
d_0^0	•	×	×
d_1^0	•	×	×
d_0^1	×	•	×
d_1^1	×	•	×
d_2^1	×	•	×
d_0^2	×	×	•
d_1^2	×	×	•
d_2^2	×	×	•
d_3^2	×	×	•
d_4^2	×	×	•

<i>N</i>			
	w_0	w_1	w_2
e_0^0	•	×	×
e_1^0	•	×	×
e_0^1	×	•	×
e_1^1	×	•	×
e_2^1	×	•	×
e_3^1	×	•	×
e_0^2	×	×	•
e_1^2	×	×	•
e_2^2	×	×	•
e_3^2	×	×	•

Fig. 3. Suppose $q = 2$, and consider the sets $A = \{2, 3, 5\}$ and $B = \{2, 4\}$. Notice that $A =_2 B$. We enumerate these sets as $\langle 2, 3, 5 \rangle$ and $\langle 2, 4, 4 \rangle$. The figure shows the models M and N derived from this enumeration. These models are indistinguishable in the EF-game with only 2 quantifier moves, regardless of the number of implication moves.

This is indeed a winning strategy, since:

$$M, s \models P(d_\beta^\alpha) \iff s \subseteq \{\alpha\} \iff N, s \models P(e_\gamma^\alpha)$$

$$M, s \models a_i = a_j \iff N, s \models b_i = b_j.$$

[3 \Rightarrow 2] Suppose \mathbb{K} is downward closed and $=_m$ -invariant for some number m . Let A_1, \dots, A_n be the subsets of $[0, m + 1]$ which are contained in \mathbb{K} . Now define:

$$K_i = \begin{cases} A_i & \text{if } m + 1 \notin A_i \\ A_i \cup [m + 1, \dots) & \text{if } m + 1 \in A_i. \end{cases}$$

We claim that $\mathbb{K} = K_1^\downarrow \cup \dots \cup K_n^\downarrow$. Start with the right-to-left inclusion. Let $B \in K_1^\downarrow \cup \dots \cup K_n^\downarrow$. This means that $B \subseteq K_i$ for some i . Now we distinguish two cases.

- Case 1: $K_i = A_i$. Then $A_i \in \mathbb{K}$ by definition, and since \mathbb{K} is downward closed, also $B \in \mathbb{K}$.
- Case 2: $K_i = A_i \cup [m + 1, \dots)$. We claim that in this case, $A_i =_m A_i \cup B$: if so, since $A_i \in \mathbb{K}$ and \mathbb{K} is $=_m$ -invariant, we have $A_i \cup B \in \mathbb{K}$, which in turn by downward closure yields $B \in \mathbb{K}$. To see that $A_i =_m A_i \cup B$, the only non-trivial step is to show that for all $\kappa \in B$ there exists some $\kappa' \in A_i$ such that $\kappa' =_m \kappa$. So, take $\kappa \in B$: if $\kappa \leq m$ then $\kappa \in A_i$ (since $B \subseteq K_i = A_i \cup [m + 1, \dots)$), so we can take $\kappa' = \kappa$; if on the other hand $\kappa > m$, then $\kappa =_m m + 1 \in A_i$.

Either way, we conclude $B \in \mathbb{K}$, which gives the right-to-left inclusion.

For the converse inclusion, suppose $B \in \mathbb{K}$. Again, we distinguish two cases.

- Case 1: $B \subseteq [0, m]$. In this case, $B = A_i$ for some $i \leq n$, and thus $B \in K_i^\downarrow$.
- Case 2: $B \ni \kappa$ for some $\kappa > m$. In this case, $B =_m B \cup \{m + 1\}$, since $\kappa =_m m + 1$. Since $B \in \mathbb{K}$ and \mathbb{K} is $=_m$ -invariant, also $B \cup \{m + 1\} \in \mathbb{K}$. Now take $(B \cup \{m + 1\}) \cap [0, m + 1]$: by downward closure, this set is in \mathbb{K} , and since it is a subset of $[0, m + 1]$, it coincides with A_i for some $i \leq n$. Notice that $m + 1 \in A_i$, and thus, $K_i = A_i \cup [m + 1, \dots)$. Therefore, $B \subseteq K_i$, which implies $B \in K_i^\downarrow$.

In either case, we conclude that $B \in K_i$ for some $i \leq n$, which gives the left-to-right inclusion. □

Theorem 4.5 allows us to tell immediately which among the questions in (2) are expressible in InqBQ: (2.a), the question whether there is any P , is expressible, since it has the form $Q_{\mathbb{K}}x.Px$ for the class $\mathbb{K}_5 = \{0\}^\downarrow \cup [1, \dots)^\downarrow$, where both $\{0\}$ and $[1, \dots)$ are definable in classical first-order logic. Indeed, the defining formula is simply $\exists x.Px$, which abbreviates $\exists x.Px \vee \neg \exists x.Px$.

The remaining questions, (2.b), (2.c), and (2.d) are not expressible, since they have the form $Q_{\mathbb{K}}x.Px$ for the following classes \mathbb{K} :

$$\mathbb{K}_6 = \bigcup \{ \{\kappa\}^\downarrow \mid \kappa \text{ a cardinal} \} \quad \mathbb{K}_7 = \text{Even}^\downarrow \cup \text{Odd}^\downarrow \quad \mathbb{K}_8 = \text{Fin}^\downarrow \cup \text{Inf}^\downarrow$$

Clearly, these classes are not the form $K_1^\downarrow \cup \dots \cup K_n^\downarrow$ for K_1, \dots, K_n definable in classical first-order logic. In a similar way, we can see that none of the following questions about the cardinality of P is expressible in InqBQ.

- (7)
- a. How many P are there, modulo k ? (for $k \geq 2$)
 - b. Is the number of P even, odd, or infinite?
 - c. Is the number of P a prime number, or a composite one?
 - d. Are there uncountably many P ?

It is worth pausing to remark that, while InqBQ can express the question “*what objects are P?*” (by means of the formula $\forall x.?Px$, see Section 2), it cannot express the corresponding cardinality question “*how many object are P?*” From the perspective of logical modeling of questions, this means that analyzing *how many* questions—an important class of questions—requires a proper extension of the logic InqBQ. Developing and investigating such an extension is an interesting prospect for future work.

Since the proof of Theorem 4.5 is quite flexible, the characterization result can be seen to hold also when we restrict to certain salient classes of models. For instance, since the proof uses only id-models, we obtain the following Corollary.

COROLLARY 4.6. *The characterization given in Theorem 4.5 holds also when we restrict to the class of id-models.*

Moreover, it is an easy exercise to adapt the proof to show the following result, concerning the class of finite models and the class of finite id-models.

COROLLARY 4.7. *Let \mathbb{K} be a set of sets of finite cardinals. The following are equivalent:*

1. *The cardinality quantifier $Q_{\mathbb{K}}$ is definable in InqBQ with respect to the class of finite models (resp. finite id-models). That is, there is a formula $\chi_{\mathbb{K}}$ of InqBQ such that $Q_{\mathbb{K}}x.Px$ is equivalent to $\chi_{\mathbb{K}}[P]$ in restriction to finite models (resp. finite id-models).*

2. $\mathbb{K} = K_1^\downarrow \cup \dots \cup K_n^\downarrow$ for some sets $K_1, \dots, K_n \subseteq \mathbb{N}$, where for each K_i there exists $m \in \mathbb{N}$ such that K_i contains all or none of the numbers $k \geq m$.

§5. Conclusion. EF games often provide an insightful perspective on a logic and a useful characterization of its expressive power. In this paper we have described an EF-game for inquisitive first-order logic, InqBQ, showing that it characterizes exactly the distinguishing power of the logic, and we used the game to study the expressive power of the logic with respect to certain cardinality properties.

In comparison to its classical counterpart, the game presents two novelties. Firstly, the roles of the two models on which the game is played are not symmetric: certain moves have to be performed mandatorily in one of the models. This feature reflects the fact that InqBQ lacks a classical negation and that the theory of a model—unlike in the classical case—can be properly included in that of another. Secondly, the objects that are picked in the course of the game are not just individuals $d \in D$, but also information states, i.e., subsets $s \subseteq W$ of the universe of possible worlds. This feature reflects the fact that InqBQ contains not only the quantifiers \forall, \exists over individuals, but also the implication \rightarrow , which allows for a restricted kind of quantification over information states.

Moreover, we introduced the notion of a cardinality quantifier in InqBQ, that is, a quantifier which is sensitive only to the number of individuals which satisfy a given property. As we illustrated, the inquisitive setting provides a more general perspective on this notion: besides quantifiers like *infinitely many*, which combine with a property to form a statement, we now also have quantifiers like *how many*, which combine with a property to form a question.

Using the EF-game, we were able to characterize exactly the range of cardinality quantifiers expressible in InqBQ. The characterization is similar to the one for classical first-order logic: the definable quantifiers are those that, for some natural number n , do not make any distinctions between cardinals larger than n . As we anticipated in the introduction, this is particularly interesting as it sheds some light on the expressive power of InqBQ, which is still poorly understood. At present, we do not know exactly how InqBQ relates to standard first- and second-order predicate logic: we do not know, e.g., whether InqBQ is compact and whether an entailment-preserving translation to first-order logic exists. Our characterization result shows that, at least with respect to the expression of cardinality properties, InqBQ is very similar to first-order logic, sharing the same kind of expressive limitations, and very different from second-order logic.

We also saw that the characterization result yields a number of interesting examples of questions which are not expressible in InqBQ. Crucially, this includes the question *how many individuals satisfy P* , both in the general case and in restriction to finite models. This means that in order to capture *how many* questions in predicate logic, a proper extension of InqBQ is needed. Other interesting examples of questions which were proved not to be expressible are *whether the number of P is even or odd* (also in restriction to finite models) and *whether the number of P is finite or infinite*.

The work presented in this paper can be taken further in several directions. Firstly, in the context of classical logic, several variants of the EF game have been studied. For example, [23] presents a *dynamic* EF game, corresponding to a more fine-grained classification of classical structures. In the inquisitive case, an analogous refinement

could lead to interesting insights into the structure of inquisitive models. Ehrenfeucht–Fraïssé games can also be used to compare different extensions of a fixed logic, as shown in [15]. In this regard, the results presented in Section 4 already yield some interesting corollaries. For example, adding to InqBQ the quantifier *how many* (the operator Q_5 in Section 4.2) yields a logic which is strictly more expressive than InqBQ. More generally, the techniques introduced in this paper are likely to provide a useful tool for a systematic study of quantifiers in inquisitive logic.

Second, as pointed out in Footnote 6, in this paper we only studied the simplest cardinality quantifiers, namely, those that operate on a single unary predicate. In further work, it would be natural to look at how our characterization result extends to the general case of cardinality quantifiers operating on several predicates, possibly of arities different from 1. This is not just a technical exercise: there are interesting cardinality questions involving multiple predicates, such as “*are there more P or more Q?*”.

Finally, one major goal for future work is to look beyond *cardinality* quantifiers and study *generalized quantifiers* in the inquisitive setting. The classical theory of generalized quantifiers is well-established [16, 17] and an important topic across logic, linguistics, and cognitive science. As illustrated in this paper and discussed in more detail in [6], the inquisitive perspective leads to a more general perspective on quantifiers. Among other things, this perspective allows us to bring interrogative words like *who*, *which*, and *how many* within the purview of generalized quantifier theory (on this enterprise, see also [21]). How does the classical theory of generalized quantifiers scale up to this more general setting? What novelties arise? The Ehrenfeucht–Fraïssé game presented in this paper will likely prove to be a fundamental tool in answering these questions.

Acknowledgment Financial support by the European Research Council under the EU’s Horizon 2020 research and innovation program (grant No. 680220) is gratefully acknowledged.

BIBLIOGRAPHY

- [1] Ciardelli, I. (2009). Inquisitive Semantics and Intermediate Logics. M.Sc. Thesis, University of Amsterdam.
- [2] ———. (2016). Questions in Logic. Ph.D. Thesis, Institute for Logic, Language and Computation, University of Amsterdam.
- [3] ———. (2018). Questions as information types. *Synthese*, **195**, 321–365.
- [4] Ciardelli, I., Groenendijk, J., & Roelofsen, F. (2019). *Inquisitive Semantics*. Oxford: Oxford University Press.
- [5] Ciardelli, I., & Roelofsen, F. (2011). Inquisitive logic. *Journal of Philosophical Logic*, **40**(1), 55–94.
- [6] ———. (2018). An inquisitive perspective on modals and quantifiers. *Annual Review of Linguistics*, **4**, 129–149.
- [7] Ehrenfeucht, A. (1967). An application of games to the completeness problem for formalized theories. *The Journal of Symbolic Logic*, **32**(2), 281–282.
- [8] Fagin, R. (1975). Monadic generalized spectra. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, **21**, 89–96.
- [9] Fraïssé, R. (1954). Sur quelques classifications des systèmes de relations. *Publications Scientifiques de l’Université D’Alger*, **1**(1), 35–182.

- [10] Frittella, S., Greco, G., Palmigiano, A., & Yang, F. (2016). A multi-type calculus for inquisitive logic. In *International Workshop on Logic, Language, Information, and Computation*. Heidelberg: Springer, pp. 215–233.
- [11] Grilletti, G. (2019). Disjunction and existence properties in inquisitive first-order logic. *Studia Logica*, **107**(6), pp. 1199–1234.
- [12] Hintikka, J. (1962). *Knowledge and Belief: An Introduction to the Logic of the Two Notions*. Ithaca, NY: Cornell University Press.
- [13] Hodges, W. (1997). *A Shorter Model Theory*. New York: Cambridge University Press.
- [14] Immerman, N. (1982). Upper and lower bounds for first order expressibility. *Journal of Computer and System Sciences*, **25**(1), 76–98.
- [15] Kolaitis, P., & Väänänen, J. (1995). Generalized quantifiers and pebble games on finite structures. *Annals of Pure and Applied Logic*, **74**(1), 23–75.
- [16] Lindström, P. (1966). First order predicate logic with generalized quantifiers. *Theoria*, **32**(3), 186–195.
- [17] Mostowski, A. (1957). On a generalization of quantifiers. *Foundations of Mathematics*, **44**, 12–36.
- [18] Połacik, T. (2008). Back and forth between first-order Kripke models. *Logic Journal of the IGPL*, **16**(4), 335–355.
- [19] Punčochář, V. (2015). Weak negation in inquisitive semantics. *Journal of Logic, Language, and Information*, **24**(3), 323–355.
- [20] ———. (2016). A generalization of inquisitive semantics. *Journal of Philosophical Logic*, **45**(4), 399–428.
- [21] Rexach, J. G. (1997). *Questions and Generalized Quantifiers*. Netherlands, Dordrecht: Springer, pp. 409–452.
- [22] Roelofsens, F. (2013). Algebraic foundations for the semantic treatment of inquisitive content. *Synthese*, **190**(1), 79–102.
- [23] Väänänen, J. (2011). *Models and Games* (first edition). New York: Cambridge University Press.
- [24] van Benthem, J. (1976). Modal Correspondence Theory. Dissertation, Universiteit van Amsterdam, Instituut voor Logica en Grondslagenonderzoek van Exacte Wetenschappen, pp. 1–148.
- [25] Visser, A. (2001). Submodels of Kripke models. *Archive for Mathematical Logic*, **40**(4), 277–295.

INSTITUTE FOR LOGIC, LANGUAGE AND COMPUTATION, UVA
SCIENCE PARK 107, 1098 XG AMSTERDAM, THE NETHERLANDS
E-mail: grilletti.gianluca@gmail.com

MUNICH CENTER FOR MATHEMATICAL PHILOSOPHY, LMU
GESCHWISTER-SCHOLL-PLATZ 1, 80539 MUNICH, GERMANY
E-mail: ivano.ciardelli@lmu.de