ON THE ENUMERATION OF FINITE L-ALGEBRAS

C. DIETZEL, P. MENCHÓN, L. VENDRAMIN

ABSTRACT. We use Constraint Satisfaction Methods to construct and enumerate finite L-algebras up to isomorphism. These objects were recently introduced by Rump and appear in Garside theory, algebraic logic, and the study of the combinatorial Yang–Baxter equation. There are 377,322,225 isomorphism classes of L-algebras of size eight. The database constructed suggest the existence of bijections between certain classes of L-algebras and well-known combinatorial objects. We prove that Bell numbers enumerate isomorphism classes of finite linear L-algebras. We also prove that finite regular L-algebras are in bijective correspondence with infinite-dimensional Young diagrams.

1. Introduction

In the study of set-theoretical solutions of the Yang–Baxter equation, the *cycloid* equation,

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z),$$

plays a fundamental role, see for example [3,11,13]. The formula first occurred in algebraic logic [2,7,19] and also appears in the theory of Garside groups [4] and in connection with von Neumann algebras and orthomodular lattices [16,17]. To combine similarities between these different objects, Rump introduced L-algebras in [12]. It turns out that L-algebras appear in connection with other topics in algebraic logic [20], in the combinatorial study of the Yang–Baxter equation [12] and in number theory [15].

An L-algebra is a set X together with a binary operation $(x, y) \mapsto x \cdot y$ on X that satisfies the cycloid equation and such that there exists a distinguished element $e \in X$ with some particular properties (see Definition 2.1 for details). The distinguished element e is crucial to define a partial order on X:

$$(1.1) x \leqslant y \Longleftrightarrow x \cdot y = e.$$

If P is a partially ordered set (*poset*, for short), we call an L-algebra X with the same underlying set as P an L-algebra on P, when (1.1) holds for all $x, y \in P$. In this case, P has a greatest element that coincides with the logical unit of X.

L-algebras have everything to become fundamental objects in algebraic logic. On the one hand, when the binary operation of X is interpreted as implication and equality as logical equivalence between propositions, L-algebras provide a significant extension of classical logic. For instance, the cycloid equation holds in classical and intuitionistic logic, and also in logics based on a continuous t-norm. On the other hand, L-algebras have a rich algebraic structure that provides new tools to study logical structures. For example, every L-algebra X admits a structure group G(X), and this group can be used to give a conceptual and short proof of Mundici's embedding theorem [8].

One of our main results here is an explicit construction of isomorphism classes of L-algebras of small size. For small size, the approach is somewhat inspired by [1] and the cycloid equation.

²⁰¹⁰ Mathematics Subject Classification. Primary:03G25; Secondary: 06D20.

Theorem 1.1. Up to isomorphism, there are 924,071 L-algebras of $size \leq 7$.

Since Hilbert algebras [5] are particular cases of L-algebras, we also obtain the construction of small Hilbert algebras up to isomorphism.

Theorem 1.2. Up to isomorphism, there are 1,085,642 Hilbert algebras of size ≤ 10 .

The method developed to prove Theorem 1.1 cannot be applied for L-algebras of size eight. This happens because the search space concerning L-algebras of size eight is "too large". Additionally, checking lexicographical symmetry breaking for the whole symmetric group Sym_{n-1} is already computationally expensive for n=8. For these reasons, the enumeration problem of L-algebras of size eight requires new ideas

Theorem 1.3. Up to isomorphism, there are 377,322,225 L-algebras of size eight.

A detailed account regarding the number of isomorphism classes of L-algebras and Hilbert algebras of size n appears in Tables 1.1 and 1.2.

Table 1.1. The number of isomorphism classes of (discrete) L-algebras of size ≤ 8 .

n	1	2	3	4	5	6	7	8
L(n)	1	1	5	44	632	15582	907,806	377,322,225
discrete	1	1	3	15	101	1241	151,846	149,390,095

As a by-product, we produce a database of finite L-algebras and Hilbert algebras. The proofs of Theorems 1.1, 1.2 and 1.3 use constraint programming techniques. Constraint programming is a paradigm for solving combinatorial problems. The idea is to search for variables that satisfy a certain number of constraints.

The main tools used are the constraint modeling assistant Savile Row [9] and [6]. Savile Row is a flexible modelling assistant for constraint programming; it provides a high-level language fitting perfectly with our particular combinatorial problem.

The databases produced are available at https://github.com/vendramin/L, with DOI name 10.5281/zenodo/6630229.

The calculations of Theorems 1.1, 1.2 and 1.3 were performed in an Intel(R) Core(TM) i7-8700 CPU @ 3.20GHz, with 32GB RAM.

By inspection of the database, we find connections involving certain finite families of *L*-algebras and well-known combinatorial objects. Two families of *L*-algebras are particular relevant for this work. We prove that finite *linear L*-algebras (Definition 4.1) are enumerated by Bell numbers (Theorem 4.5). We also prove that the elements of another important family of finite *L*-algebras, that of *regular L*-algebras (Definition 5.3), are in bijective correspondence with infinite-dimensional Young diagrams (Theorem 5.9).

2. Preliminaries

We refer to [12] for the basic theory of L-algebras.

Table 1.2. The numbers of isomorphism classes of Hilbert algebras of size ≤ 10 .

n	1	2	3	4	5	6	7	8	9	10
Hilbert	1	1	2	6	21	95	550	4036	37602	1,043,328

Table 1.3. The construction of isomorphism classes of L-algebras of size eight.

	discrete	non-discrete		
Poset	trivial	diamond	other	
Nr. cases	73	156	2043	
Nr. solutions	149,390,095	138,219,543	89,712,587	
Run-time	4 days	5 hours	11 hours	
Data (compressed)	208MB	212MB	153MB	

Definition 2.1. An *L*-algebra is a set *X* together with a function $X \times X \to X$, $(x,y) \mapsto x \cdot y$, such that

- (1) there exists an element $e \in X$ such that $e \cdot x = x$ and $x \cdot e = x \cdot x = e$ for all $x \in X$,
- (2) $(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z)$ holds for all $x, y, z \in X$, and
- (3) if $x \cdot y = y \cdot x = e$, then x = y.

The element e is unique and is called the *logical unit* of X.

Definition 2.2. A *Hilbert algebra* is an *L*-algebra *X* such that

$$(2.1) x \cdot (y \cdot z) = (x \cdot y) \cdot (x \cdot z)$$

holds for all $x, y, z \in A$.

Definition 2.3. If X and Y are L-algebras, a homomorphism of L-algebras is a map $f: X \to Y$ such that $f(x \cdot y) = f(x) \cdot f(y)$ for all $x, y \in X$.

The derived notions of isomorphism and automorphism are clear from this definition. If $f: X \to Y$ is a homomorphism of L-algebras, then f(e) = e. It follows that the automorphism group of an L-algebra X of size n is isomorphic to a subgroup of Sym_{n-1} , the symmetric group on n-1 letters.

It can be shown [12, Proposition 2] that for an L-algebra X, the relation \leq given by

$$(2.2) x \leqslant y \Longleftrightarrow x \cdot y = e$$

is a partial order relation. We will call this relation the natural partial order relation on X. Note that from this definition we immediately see that e is the unique maximal element of X.

Given elements $x, y, z \in X$ with $y \leq z$, we can calculate

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z) = (y \cdot x) \cdot e = e,$$

which implies that $x \cdot y \leq x \cdot z$, as well, so the assignments $y \mapsto x \cdot y$ are monotone for all $x \in X$.

One case of natural order is of particular importance in this article.

Definition 2.4. An *L*-algebra *X* is said to be *discrete* if $x \le y$ if and only if x = y or y = e.

This means, an L-algebra is discrete when the order on $X \setminus \{e\}$ is discrete.

2.1. A combinatorial description of finite *L*-algebras. Let $n \ge 2$ and *X* be a finite *L*-algebra of size *n*. Without loss of generality we may assume that $X = \{1, ..., n\}$, where *n* is the logical unit. The *L*-algebra structure on *X* is then described by the matrix $M = (M_{i,j}) \in \mathbb{Z}^{n \times n}$, where

$$M_{i,j} = i \cdot j, \quad i, j \in \{1, \dots, n\}.$$

The matrix M satisfies the following conditions:

- (1) $M_{n,j} = j$ for all $j \in \{1, \dots, n\}$,
- (2) $M_{i,n} = n \text{ for all } i \in \{1, \dots, n\},$
- (3) $M_{k,k} = n$ for all $k \in \{1, ..., n\}$, and
- (4) $M_{M_{i,j},M_{i,k}} = M_{M_{j,i},M_{j,k}}$ holds for all $i,j,k \in \{1,\ldots,n\}$. (5) if $M_{i,j} = n$ and $M_{j,i} = n$, then i = j,

There is a bijective correspondence between L-algebras and matrices satisfying Conditions (1)–(5); see Example 2.5 for a concrete example of a matrix representation of an L-algebra.

Since n is the logical unit, g(n) = n holds for each automorphism g of the Lalgebra. Over the set of $n \times n$ matrices satisfying Conditions (1)–(5) we define the equivalence relation given by

$$M \sim N \iff \text{there exists } g \in \operatorname{Sym}_{n-1} \text{ such that } N_{i,j} = g^{-1}\left(M_{g(i),g(j)}\right)$$

for all $i, j \in \{1, ..., n\}$. It follows that the L-algebras X and Y are isomorphic if and only if $M_X \sim M_Y$.

Example 2.5. Let X be the L-algebra structure over $\{x, y, e\}$ with logical unit e given by

$$e \cdot y = y$$
, $x \cdot y = y \cdot x = e \cdot x = x$.

Let $f: \{1,2,3\} \to \{x,y,e\}$ be the bijective map given by f(1) = x, f(2) = y and f(3) = e. Then X can be represented as the matrix

$$M_X = \begin{pmatrix} 3 & 1 & 3 \\ 1 & 3 & 3 \\ 1 & 2 & 3 \end{pmatrix}.$$

Note that, in this new representation of X, the logical unit is the element 3.

3. Proofs of Theorems 1.1, 1.2 and 1.3

3.1. Proof of Theorems 1.1 and 1.2. To construct all L-algebras of size $n \leq 7$ we need to find all possible matrices $M \in \mathbb{Z}^{n \times n}$ with coefficients in $\{1, 2, \dots, n\}$ that satisfy Conditions (1)-(5). We use Savile Row [9] to find all matrices with entries in $\{1,\ldots,n\}$ satisfying Conditions (1)–(5).

To remove repetitions we proceed as follows. If $g \in \operatorname{Sym}_{n-1}$ and M is a matrix, we denote by M^g the matrix given by

$$(M^g)_{i,j} = g^{-1} (M_{g(i),g(j)})$$

for all $i, j \in \{1, ..., n\}$. We need to find those matrices M such that

$$(3.1) M \leqslant_{\text{lex}} M^g$$

for all $g \in \operatorname{Sym}_{n-1}$, where $A \leq_{\operatorname{lex}} B$ if and only if

$$(A_{1,1}, A_{1,2}, \dots, A_{1,n}, A_{2,1}, A_{2,2}, \dots, A_{n,n})$$

 $\leq (B_{1,1}, B_{1,2}, \dots, B_{1,n}, B_{2,1}, B_{2,2}, \dots, B_{n,n})$

with lexicographical order.

Our method produces a database (159MB) of isomorphism classes of L-algebras of size ≤ 7 in about 6 hours.

Minor variations of our algorithm produce database of particular classes of Lalgebras such as Hilbert algebras. A database (11MB) of isomorphism classes of Hilbert algebras of size \leq 9 was constructed in about two hours. Isomorphism classes of Hilbert algebras of size 10 were only enumerated, not constructed.

- 3.2. **Proof of Theorem 1.3.** The approach used in the proof of Theorem 1.1 does not work for L-algebras of size eight. As is already indicated in the introduction, the main problems we are confronted with are:
 - (1) The vast size of the search space for n=8: from the 64 entries $M_{i,j}$, only 22 (the last row, the last column and the diagonal) follow from the L-algebra axioms. Therefore, the search space contains 8^{42} elements.
 - (2) The lexicographical symmetry breaking is very expensive as there are potentially $7! = |\operatorname{Sym}_{8-1}|$ permutations to consider in order to determine if a matrix is lexicographically least in its conjugacy class. However, there is a significant amount of L-algebras that have a trivial or, at least, small automorphism group. An elimination of the lexicographical symmetry breaking for some of these asymmetric L-algebras is therefore expected to accelerate the search process significantly.

Since each L-algebra has the structure of a poset under (2.2), we can split the problem of constructing all non-isomorphic L-algebras of size n into the construction of all non-isomorphic L-algebras on a given poset (X, \leq) .

Note that a poset of size n on the set $\{1, \ldots, n\}$ with maximal element n can be described by a matrix $P = (P_{i,j}) \in \{0,1\}^{(n-1)\times(n-1)}$, where

$$P_{i,j} = 1 \iff i \leqslant j.$$

Given a matrix P describing the poset (X, \leq) , the condition that a matrix $M_{i,j}$ is an L-algebra on (X, \leq) can then be rewritten as

$$M_{i,j} = n \iff P_{i,j} = 1 \text{ for all } i, j \in \{1, \dots, n-1\},$$

which can easily be translated to a constraint.

To split the search space of all L-algebras of size n, we first compute all nonisomorphic posets on a set of size n-1 and then compute the non-isomorphic L-algebras on each of them separately. Here, we slightly modify the lexicographic condition by not searching for the lexicographically smallest matrix with respect to all symmetries of the set $\{1, \ldots, n-1\}$ but only with respect to automorphisms of the poset (X, \leq) which, in cases of posets with few automorphisms, accelerates the search significantly. Note that this does not necessarily lead to a multiplication table that is the lexicographically least with respect to all of Sym_{n-1} .

Unfortunately, searching by posets is not enough when computing L-algebras of size eight, as in the case of a discrete L-algebra we do not have enough restrictions for the entries in a multiplication table and also there are too many symmetries to

In order to explain our approach, we need to repeat the definition of \land -semibraces [13, Definition 3] and introduce the rather technical notion of a separating kapproximate semilattice.

Definition 3.1. A \land -semibrace is a set X with a distinguished element $e \in X$ and binary operations $\wedge, \cdot : X \times X \to X$, such that the pair (X, \wedge) is a commutative monoid with neutral element e and the following identities hold for all $x, y, z \in X$:

- (1) $e \cdot x = x$,
- $(2) x \cdot e = e,$
- (3) $x \cdot (y \wedge z) = (x \cdot y) \wedge (x \cdot z)$, and (4) $(x \wedge y) \cdot z = (x \cdot y) \cdot (x \cdot z)$.

We remark here that the commutativity of the monoid (X, \wedge) , together with Equation (4), implies that the pair (X, \cdot) fulfils the cycloid equation.

Theorem 3.2 (Rump). Each L-algebra X has a unique \wedge -closure C(X) such that (1) C(X) is a semibrace that is also an L-algebra with logical unit e,

- (2) C(X) contains X as a sub-L-algebra, and
- (3) the monoid $(C(X), \wedge, e)$ is a semilattice generated by the subset X.

Proof. See [13, Theorem 3].

We note here that, as e is the top element of X, the subset $X' = X \setminus \{e\}$ generates the subsemilattice $C(X) \setminus \{e\}$. Using Theorem 3.2, the most obvious approach would now be to first construct all \land -semilattices generated by an (n-1)-element set X' and then search for all L-algebras on $X := X' \cup \{e\}$ with logical unit e that are consistent with the \land -structure in a sense that we will now explain:

Every semilattice is also a poset under the relation

$$x \leqslant y \iff x \land y = x.$$

As C(X) is generated by X, each element $y \in C(X)$ can be written as

$$y = \bigwedge \{ x \in X' : y \leqslant x \},\$$

where $\bigwedge \emptyset = e$. Therefore, each element $y \in C(X)$ can be characterized by the set of all elements above y that are lying in X'.

Note that expressions of the form $(x_1 \wedge x_2 \wedge ... \wedge x_k) \cdot z$ with $x_i \in X$ (i = 1, ..., k) and $z \in X$ can be rewritten in terms of the L-algebra operation only. For example,

$$(x_1 \wedge x_2) \cdot z = (x_1 \cdot x_2) \cdot (x_1 \cdot z),$$

$$(x_1 \wedge x_2 \wedge x_3) \cdot z = ((x_1 \wedge x_2) \cdot x_3) \cdot ((x_1 \wedge x_2) \cdot z)$$

$$= ((x_1 \cdot x_2) \cdot (x_1 \cdot x_3)) \cdot ((x_1 \cdot x_2) \cdot (x_1 \cdot z)).$$

Given a semilattice (C, \wedge) generated by a set X', we call an L-algebra operation \cdot on $X = X' \cup \{e\}$ consistent with the semilattice if the rewritten form of all terms of the form $(x_1 \wedge x_2 \wedge \ldots \wedge x_k) \cdot z$ in terms of \cdot is equal to e if and only if $x_1 \wedge x_2 \wedge \ldots \wedge x_k \leq z$ holds in C.

We have seen that if C is a bounded semilattice with top element e where the subsemilattice $C \setminus \{e\}$ is generated by the subset X', then each element $\bigwedge A \in C$, where $A \subseteq C$, can be identified from the set

$$\overline{A} = \left\{ x \in X' : \left(\bigwedge A \right) \leqslant x \right\}.$$

Note that $\overline{(\bigwedge A) \wedge (\bigwedge B)} = \overline{A} \cup \overline{B}$, so all information about the semilattice C is essentially contained in the operation $A \mapsto \overline{A}$.

This map has the following properties:

- (1) $\overline{\emptyset} = \emptyset$.
- (2) for $A \subseteq X'$, we have $A \subseteq \overline{A}$.
- (3) for $A, B \subseteq X'$, we have the implication $A \subseteq \overline{B} \Rightarrow \overline{A} \subseteq \overline{B}$.
- (4) The mapping $A \mapsto \overline{A}$ is separating points, meaning that for $x, y \in X'$, we have the implication $x \neq y \Rightarrow \overline{\{x\}} \neq \overline{\{y\}}$.

A map $\operatorname{cl}: \mathcal{P}(X') \to \mathcal{P}(X'), A \mapsto \overline{A}$ with these four properties above will be called a separating closure operator on X'. So a possible approach to the enumeration resp. calculation of all L-algebras of size n would be to first construct all semilattices on n-1 generators by determining the different separating closure operators on a set X' of size n-1, up to a relabelling of the generators, and then find, for each semilattice, the non-isomorphic L-algebra operations on $X = X' \cup \{e\}$ which are consistent with the semilattice structure. Note that this approach in fact splits the search case as the \wedge -closure is an invariant of an L-algebra. However, there are two obstacles when trying to use this approach directly:

(1) There are way too many semilattices generated by an n-element set, even for small values of n.




FIGURE 3.1. Another ordering of matrix entries.

(2) The length of the expressions $(x_1 \wedge x_2 \wedge ... \wedge x_k) \cdot z$ contains 2^k terms x_i or z, each one surrounded by k-1 brackets. In particular, these expressions do not have fixed length.

In order to deal with this problem, we decided to only compute part of the semilattice structures and demand consistency with only this part of the semilattice structure. This leads to the notion of a separating k-approximate semilattice. To define this, we clarify some notation: by $\binom{X}{\leqslant k}$, we denote the set of all subsets of a set X whose size is k or less.

Definition 3.3. Let $k \ge 1$ and X be a set. A separating k-approximate semilattice on X is a map $\operatorname{cl}: \binom{X}{\le k} \to \mathcal{P}(X), A \mapsto \overline{A}$ with the following properties:

- $(1) \ \overline{\emptyset} = \emptyset.$
- (2) For $A \in {X \choose \leqslant k}$, we have $A \subseteq \overline{A}$.
- (3) For $A, B \in \binom{X}{\leqslant k}$, we have the implication $A \subseteq \overline{B} \Rightarrow \overline{A} \subseteq \overline{B}$.
- (4) For $x, y \in X$, we have the implication $x \neq y \Rightarrow \overline{\{x\}} \neq \overline{\{y\}}$.

Definition 3.4. If cl_1, cl_2 are two separating k-approximate semilattices on a set X, then they are said to be *conjugate*, if there exists a bijection $\pi: X \to X$, such that $\text{cl}_2(\pi(A)) = \pi(\text{cl}_1(A))$ for all $A \in \binom{X}{< k}$.

Note that each separating 1-approximate semilattice cl on X determines a partial order on X by declaring $x \leq y :\iff y \in \operatorname{cl}(\{x\})$. Also, each partial order on X gives rise to a separating 1-approximate semilattice by defining $\operatorname{cl}(\emptyset) = \emptyset$ and $\operatorname{cl}(\{x\}) = \{y \in X : y \geqslant x\}$. If k = |X|, then a k-approximate semilattice is just a closure operator on X. So a separating k-approximate semilattice on a set K can be seen as something in between a poset on K and a semilattice generated by K.

Given a k-approximate semilattice on a set X' we call an L-algebra operation on the set $X = X' \cup \{e\}$ - with e being the logical unit - consistent with the approximate semilattice if for $l \leq k$ and elements $x_1, \ldots, x_l, y \in X'$, the expression $(x_1 \wedge \ldots \wedge x_l) \cdot y$ evaluates to e (when rewritten in terms of the L-algebra operation) if and only if $y \in \{x_1, \ldots, x_l\}$.

For size 8, we use the lexicographical order of matrices with respect to the ordering $(A_{1,1}, A_{1,2}, A_{2,2}, A_{2,1}, A_{1,3}, A_{2,3}, A_{3,3}, A_{3,2}, A_{3,1}, \ldots, A_{n,n})$, see Figure 3.1. For bigger multiplication tables, this ordering is better suited for checking lexicographical symmetry breaking, as most of its initial segments are contained in smaller squares that are invariant under permutations in Sym_k $(1 \leq k \leq n-1)$.

We now have all tools at our hands to explain how we get our results.

3.3. Discrete *L*-algebras.

Proposition 3.5. Up to isomorphism, there are 149,390,095 discrete L-algebras of size eight.

To deal with the discrete case in size eight, we make use of the following result. For $x, y \in X$, write $x \prec y$ to denote that x < y and there is no $z \in X$ such that x < z < y.

Theorem 3.6 (Rump). If X is a discrete L-Algebra then its \land -closure C(X) has the property that $X \setminus \{e\} = \{y \in C(X) : y \prec e\}$ and is a lower semimodular lattice meaning that C(X) a lattice under the semilattice order the and the implication $a \prec a \lor b \Rightarrow a \land b \prec b$ holds in C(X).

Proof. This is implicitly contained in the proof of [14, Proposition 18]. \Box

In particular, that means that for $e \neq x \in X$ and $y \in C(X)$ with $y \not \leq x$, we have that $x \land y \prec y$. It turns out that splitting the search space by searching for L-algebras consistent with separating 3-approximate semilattices fulfilling a restricted version of lower semimodularity is the best way of constructing all non-isomorphic discrete L-algebras of size eight. Therefore, our approach is as follows:

- (1) Construct all non-conjugate 3-approximate semilattices on $\{1, \ldots, n-1\}$ that fulfill a restricted version of lower semimodularity.
- (2) For each of these 3-approximate semilattices on $\{1, \ldots, n-1\}$, search for all non-isomorphic *L*-algebras on $\{1, \ldots, n\}$ that are consistent with the approximate semilattice structure.

For each step, we now describe the constraints:

3.3.1. Constructing separating 3-approximate semilattices. We model a separating 3-approximate semilattice cl: $\binom{X'}{3} \to \mathcal{P}(X')$ as a map : $\{1, \dots, n-1\}^4 \to \{0, 1\}$, where

$$S(x_1, x_2, x_3, y) = 1 \iff y \in \text{cl}(\{x_1, x_2, x_3\}).$$

This leads to the following set of properties:

- (1) $S(x_1, x_2, x_3, y) = S(x_{\pi(1)}, x_{\pi(2)}, x_{\pi(3)}, y)$ for all $\pi \in \text{Sym}_3$.
- (2) $S(x_1, x_2, x_3, x_i) = 1$ for $i \in \{1, 2, 3\}$.
- (3) If $S(x_1, x_2, x_3, y_i) = 1$ holds for all $i \in \{1, 2, 3\}$ and $S(y_1, y_2, y_3, z) = 1$, then $S(x_1, x_2, x_3, z) = 1$.

By Rump's theorem on the lower semimodularity of C(X), we can add the following property which makes it possible to split the search in a small number of cases:

(4) If $S(x_1, x_2, x_3, y) = 1$, $S(x_1, x_2, x_2, y) = 0$ and $S(x_1, x_2, x_3, z) = 1$ hold, then $S(x_1, x_2, y, z) = 1$.

This means

$$y \in \operatorname{cl}(\{x_1, x_2, x_3\}) \setminus \operatorname{cl}(\{x_1, x_2\}) \implies \operatorname{cl}(\{x_1, x_2, x_3\}) = \operatorname{cl}(\{x_1, x_2, x_3\}).$$

Also, for symmetry breaking, we demand that S has to be lexicographically greatest with respect to relabelling of the variables to ensure that we get as much relations of the form S(i,j,k,l)=1 for small i,j,k,l, as such relations put a lot of restrictions to the first entries of an L-algebra multiplication table.

- 3.3.2. Searching for L-algebras on a fixed 3-approximate semilattice. Given a function $S(x_1, x_2, x_3, y)$ for a separating 3-approximating semilattice on $\{1, \ldots, n-1\}$, we want to construct multiplication tables of L-algebras on $\{1, \ldots, n\}$ that are consistent with the approximate semilattice structure. This means that
 - (1) $M_{i,j} = n \iff (i = j)$ or (j = n). This is just the discreteness condition.
 - (2) For $i, j, k \in \{1, \ldots, n-1\}$, S(i, j, j, k) = 1 holds if and only if $(i \wedge j) \cdot k = e$ which translates to the equation $M_{M_{i,j},M_{i,k}} = n$. As we are looking at discrete L-algebras, this is equivalent to $(M_{i,j} = M_{i,k})$ or (i = k).




FIGURE 3.2. The diamond poset of eight elements.

(3) for
$$i, j, k, l \in \{1, \dots, n-1\}$$
, $S(i, j, k, l) = 1$ holds if and only if $(i \wedge j \wedge k) \cdot l = e$,

which translates to the equation $M_{M_{M_{i,j},M_{i,k}},M_{M_{i,j},M_{i,l}}} = n$. Due to discreteness, this is equivalent to $(M_{M_{i,j},M_{i,k}} = M_{M_{i,j},M_{i,l}})$ or $(M_{i,j} = M_{i,l})$ or (i = l).

(4) For $g \in \operatorname{Sym}_{n-1}$ such that S(i,j,k,l) = S(g(i),g(j),g(k),g(l)) for all $i,j,k,l \in \{1,\ldots,n-1\}$, we have $M \leq_{\operatorname{lex}} M^g$

Proof of Proposition 3.5. The conditions mentioned before can easily be translated to constraints. In total, when distinguishing by separating 3-approximate semilattices of size 7 that fulfill restricted lower semimodularity, we get 73 cases that can be treated separately. This leads to the complete calculation of non-isomorphic discrete L-algebras of size eight.

The construction of the database of isomorphism classes of discrete L-algebras of size eight took about four days, see Table 1.3.

3.4. Non-discrete L-algebras.

Proposition 3.7. Up to isomorphism, there are 89,712,587 non-discrete L-algebras over a poset distinct from the diamond of Figure 3.2.

Proof. In the case that the natural order on $X' = X \setminus \{e\}$ is non-discrete then our approach can be seen as working with separating 1-approximate semilattices, that is, posets:

- (1) Construct all non-isomorphic posets on the set $\{1, \ldots, n-1\}$.
- (2) For each isomorphism type of poset on $\{1, \ldots, n-1\}$, add another element n and search for all non-isomorphic L-algebras on the set $\{1, \ldots, n-1\}$ whose logical unit is n and whose natural order restricts to the given partial order on $\{1, \ldots, n-1\}$.

A few words on the construction of the posets: we use an easy python script to first construct all posets of size n-1 by starting with a poset of size 1 and repeatedly adding new minimal elements by declaring an upwards-closed set from the recent poset as the set of elements lying above the new element. The size of the new constructed upset should be at least as big as the upsets that have already been constructed. We then filter out isomorphic posets by demanding that the constructed poset should be the least with respect to some lexicographic condition. We then encode the poset structure to a zero-one-matrix in a way that will be explained below.

Let X be an L-algebra and $x,y,z\in X.$ If $x\leqslant y$ with respect to the natural order in X, then

$$x \cdot z = e \cdot (x \cdot z) = (x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z).$$

Also recall, that for any elements $x, y, z \in X$, the implication $y \leq z \Rightarrow x \cdot y \leq x \cdot z$ holds.

Therefore, given a partial order relation on $\{1, \ldots, n-1\}$, encoded as a matrix $P \in \{0,1\}^{(n-1)\times(n-1)}$ with

$$P_{i,j} = \begin{cases} 1 & i \leq j, \\ 0 & \text{otherwise,} \end{cases}$$

we use the following constraints, additionally to the L-algebra axioms:

- (1) $M_{i,j} = n \iff P_{i,j} = 1 \text{ for } i, j \in \{1, \dots, n-1\},$

- (1) $M_{i,j} = N$ $\leftarrow N_{i,j} = N_{i$ $M \leqslant_{\text{lex}} M^g$

These conditions can directly translated to constraints that enable us to enumerate almost all non-isomorphic L-algebras whose natural order is non-discrete, except for the case of a diamond order, whose restriction to the set $\{1, \ldots, n-1\}$ is given by $i \leq_{\text{diam}} j \iff i = j \text{ or } i = n - 1.$

Proposition 3.8. Up to isomorphism, there are 138,219,543 L-algebras of size eight on the diamond poset of Figure 3.2.

Proof. This case turns out to be the hardest amongst all non-discrete posets. We split up the case as follows. Let $i, j \in \{1, \dots, n-2\}$. If, in the \land -closure C(X), the condition $i \wedge j \leq k$ holds for some $k \in \{1, \dots, n-2\}$, then this can be expressed as an equation

$$(i \wedge j) \cdot k = e \iff (i \cdot j) \cdot (i \cdot k) = e.$$

We use an approach which can be seen as a really coarse way of using 2approximate semilattices, for as to our knowledge, there is no theorem that describes the semilattice structure of C(X) well enough to restrict the search to a small quantity of 2-approximate semilattices. So we first construct matrices $R \in \{0,1\}^{(n-2)\times(n-2)}$ such that $R_{i,i} = 0$ and $R_{i,j} = R_{j,i}$ for all $i, j \in \{1, \dots, n-2\}$. These matrices parametrize all couples of $i, j \in \{1, ..., n-2\}$ with the property that $i \land j \leqslant k$ holds for all $k \in \{1, \dots, n-2\}$ in the \land -closure of the final L-algebra. If we look at two such matrices as being equivalent if they only differ by a simultaneous relabelling of rows and columns, then we get 156 cases that can be treated as follows:

Given a symmetric zero-one-matrix R with only zeros on the diagonal, we have the following conditions

- (4') the lexicographical symmetry breaking is replaced as follows: if $g \in \operatorname{Sym}_{n-2}$ is such that $R_{i,j} = R_{g(i),g(j)}$ holds for all $i,j \leqslant n-1$, then $M \leqslant_{\operatorname{lex}} M^g$
- (5) for all $i, j \in \{1, \dots, n-2\}$, we have $R_{i,j} = 1$ if and only if $M_{M_{i,j}, M_{i,k}} = n$ holds for all $k \in \{1, \ldots, n-2\}$.

The required 156 cases were treated in about four hours, see Table 1.3 for more information.

4. Linear L-algebras

Recall that under the partial order (2.2) left-multiplications are monotone, i.e. if $y \leq z$, then $x \cdot y \leq x \cdot z$ for all $x \in X$.

Definition 4.1. An L-algebra X is said to be *linear* if $x \leq y$ or $y \leq x$ for all $x, y \in X$.

In the case of linear L-algebras (2.2) is indeed a total order.

For $n \ge 2$ let $\lambda(n)$ be the number of non-isomorphic linear L-algebras with n elements. A look into the sequence A000110 of [18] suggests that $\lambda(n) = B(n-1)$,

where B(m) is the m-th Bell number that counts the number of ways to partition a set with m elements into non-empty subsets.

Definition 4.2. We call an element x in an L-algebra X invariant if $y \cdot x = x$ holds for all y > x.

Lemma 4.3. Let $n \ge 1$ and let X be an L-algebra on the ordered set

$$L_n = \{x_0 > x_1 > \ldots > x_{n-1}\}.$$

Then the following statements hold.

- (1) If $x \ge y > z$, then $x \cdot y > x \cdot z$ for all $x, y, z \in X$.
- (2) $x \cdot y \geqslant y$ holds for all $x, y \in X$.
- (3) If $x \leq y$, then $x \cdot z \geqslant y \cdot z$ for all $x, y, z \in X$.
- (4) If $p \neq e$ is the smallest invariant element of X, then $p \cdot x > x$ for all $x \leq p$.

Proof. (1) Let $x \ge y > z$, then

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z) = e \cdot (y \cdot z) = y \cdot z \neq e.$$

This implies that $x \cdot y \nleq x \cdot z$. As the order on X is total, this implies $x \cdot y > x \cdot z$.

(2) Without loss of generality, we may assume that x > y. By (1), for each $x \in X$ the map $\varphi_x \colon [x_{n-1}, y] \to [x_{n-1}, x \cdot y], z \mapsto x \cdot z$, is a well-defined monotone embedding. Therefore

$$|[x_{n-1},y]| = |\varphi_x([x_{n-1},y])| \leqslant |[x_{n-1},x\cdot y]|.$$

Since these intervals are finite, it follows that $x \cdot y \geqslant y$.

(3) Let $x, y, z \in X$ be such that $x \leq y$. By (2),

$$x \cdot z = e \cdot (x \cdot z) = (x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z) \geqslant y \cdot z.$$

(4) Assume that there exists a maximal x with x < p and $p \cdot x = x$. As x is not invariant, there is a greatest element y such that y > x and $y \cdot x > x$. Then y < p, as $y \ge p$ and (3) imply

$$x = p \cdot x \geqslant y \cdot x > x$$
,

a contradiction. The maximality of x implies that $p \cdot y > y$. Since y is maximal and $p \cdot y > y$, $(p \cdot y) \cdot x = x$. By using the cycloid equation,

$$x = (p \cdot y) \cdot (p \cdot x) = (y \cdot p) \cdot (y \cdot x) = y \cdot x > x,$$

a contradiction. Therefore $p \cdot x > x$.

We can now prove that each linear L-algebra can be built up inductively by adding a new smallest element x' and declaring an admissible value for $p \cdot x'$, where p is the smallest invariant element in the old L-algebra.

Proposition 4.4. Let p be the smallest invariant element in X. Let c be an element in the superset $L_{n+1} \supseteq L_n$ such that $p \cdot x_{n-1} > c$. Then there is a unique L-algebra X' on L_{n+1} such that the canonical embedding $L_n \hookrightarrow L_{n+1}$ embeds X as a sub L-algebra of X' and such that $p \cdot x_n = c$.

As $p \cdot x_n$ must necessarily be smaller than $p \cdot x_{n-1}$ (part (1) of Lemma 4.3), the proposition describes the construction of *each* extension to an *L*-algebra on L_{n+1} .

Proof. We look at two cases, namely $c = x_n$ or $c > x_n$. Note that we must define $a \cdot b$ only for $a \in L_n$ and $b = x_n$, as $x_n \cdot a = e$ is forced by the order of P_{n+1} .

Assume first that $c = x_n$. By Lemma 4.3, setting $a \cdot x_n := x_n$ for $a \neq x_n$ defines the only possible extension to an L-algebra structure on L_{n+1} such that $p \cdot x_n = x_n$. We now check that this definition is indeed compatible with the cycloid equation, meaning that

$$(x \cdot y) \cdot (x \cdot z) = (y \cdot x) \cdot (y \cdot z)$$

holds for all $x, y, z \in L_{n+1}$. Without loss of generality may assume that x, y, z are different. If $z = x_n$, the result is trivially true. Now we may assume that x > y. If $y = x_n$, then

$$(x \cdot x_n) \cdot (x \cdot z) = x_n \cdot (x \cdot z) = e = e \cdot e = (x_n \cdot x) \cdot (x_n \cdot z).$$

This proves that this indeed defines an extension to an L-algebra on L_{n+1} . It remains to show that this is the only possibility. Therefore, let X' be an extension to an L-algebra on L_{n+1} with $p \cdot x_n = x_n$, where p is the smallest invariant element of X.

By Lemma 4.3(1), for a > p, the map $[x_n, p] \to [x_n, a \cdot p] = [x_n, p]$, $x \mapsto a \cdot x$, is a monotone embedding, i.e. an isomorphism of ordered sets. It follows that $a \cdot x_n = x_n$. We now know that $b \cdot x_n = x_n$ for all $b \geqslant p$. Assume now that $p \geqslant a > x_n$ and that $b \cdot x_n = x_n$ holds for all b > a. Since $a \cdot p = e$ and $a \cdot x_n = e$,

$$a \cdot x_n = e \cdot (a \cdot x_n) = (a \cdot p) \cdot (a \cdot x_n)$$

and hence

$$a \cdot x_n = (a \cdot p) \cdot (a \cdot x_n) = (p \cdot a) \cdot (p \cdot x_n) = (p \cdot a) \cdot x_n = x_n,$$

as $p \cdot a > a$ by Lemma 4.3(4).

We now assume that $c > x_n$. Again we need to define $a \cdot x_n$ when $a > x_n$. We claim that the only possible definition is given by

$$a \cdot x_n = \begin{cases} (p \cdot a) \cdot c & a \leqslant p, \\ x_n & a > p. \end{cases}$$

The necessity is easily shown: if p, x and y are elements in a linear L-algebra such that $p \geqslant x$ and $p \geqslant y$, the cycloid equation implies that

$$x \cdot y = e \cdot (x \cdot y) = (x \cdot p) \cdot (x \cdot y) = (p \cdot x) \cdot (p \cdot y).$$

So the first equation follows from demanding $p \cdot x_n = c$. On the other hand, as p is invariant, for all x > p the map given by left-multiplication by x is an automorphism of the ordered set $[x_n, p]$. In particular, $x \cdot y = y$ for all $y \in [x_n, p]$.

Now, we check that these assignments do not violate the cyclic condition. Let $z = x_n$ and x, y > p, then the left side of the equation is

$$(x \cdot y) \cdot (x \cdot x_n) = (x \cdot y) \cdot x_n = x_n,$$

as $x \cdot y \geqslant y > p$ by Lemma 4.3(2). Similarly, $(y \cdot x) \cdot (y \cdot x_n) = x_n$. Now let $z = x_n$ and $x > p \geqslant y$. Then, as before,

$$(x \cdot y) \cdot (x \cdot x_n) = y \cdot x_n,$$

$$(y \cdot x) \cdot (y \cdot x_n) = e \cdot (y \cdot x_n) = y \cdot x_n.$$

For $z = x_n$ and $x, y \leq p$, we use $x \cdot y = (p \cdot x) \cdot (p \cdot y)$ to calculate

$$(x \cdot y) \cdot (x \cdot x_n) = ((p \cdot x) \cdot (p \cdot y)) ((p \cdot x) \cdot c)$$
$$= ((p \cdot y) \cdot (p \cdot x)) ((p \cdot y) \cdot c)$$
$$= (y \cdot x) \cdot (y \cdot x_n).$$

Finally, let $x = x_n$ (the case where $y = x_n$ follows by symmetry). Then

$$(x_n \cdot y) \cdot (x_n \cdot z) = e \cdot e = e.$$

If y > p, then $y \cdot x_n = x_n$ and hence $(y \cdot x_n) \cdot (y \cdot z) = e$. On the other hand, if $y \leq p$, then

$$y \cdot z \geqslant y \cdot x_{n-1} = (p \cdot y) \cdot (p \cdot x_{n-1}) \geqslant (p \cdot y) \cdot c = y \cdot x_n$$

which finally implies that $(y \cdot x_n) \cdot (y \cdot z) = e$.

Theorem 4.5. For all positive integer n, $\lambda(n) = B(n-1)$.

Proof. Note that $\lambda(n)$ is the number of L-algebras on a fixed n-element set L_n . As each isomorphism of L-algebras is also an isomorphism of posets and as finite totally ordered sets do not have any nontrivial automorphisms, any two different L-algebra structures on L_n are non-isomorphic.

We proceed by induction on n. Clearly, $\lambda(1) = B(0) = 1$. Let n > 0 and assume that $\lambda(k) = B(k-1)$ holds for all k < n. For $m \in \{1, \ldots, n-1\}$ let $\lambda(n, m)$ be the number of all linear L-algebras on the ordered set L_n whose smallest invariant element is x_m . We will now show that

$$\lambda(n,m) = \binom{n-1}{m} B(m).$$

To this end, let X be an L-algebra on L_n whose smallest invariant element is x_m . Let

$$X_m = \{x_i \in X : i < m\} \subseteq X$$

and the subset

$$X^m = \{x_m \cdot x_j : j > m\} \subseteq L_{n-1}.$$

We claim that the assignment $X \mapsto (X_m, X^m)$ is a bijective correspondence between L-algebras on L_n with smallest invariant element x_m and pairs (X', A), where X' is an L-algebra on L_m and A is an (n - m - 1)-subset of $L_{n-1} \setminus \{e\}$.

Let X' be an L-algebra on L_m and $A \subseteq L_{n-1}$ be an (n-m-1)-subset of L_{n-1} . By Proposition 4.4, there is a unique L-algebra $X^{(0)}$ on L_{m+1} that has x_m as an invariant element and restricts to X' on L_m . Assume that

$$A = \{a_1 > a_2 > \ldots > x_{n-m-1}\}\$$

and note that from $a_{n-m-1} \geqslant x_{n-2}$ it follows that $a_i \geqslant x_{m-1+i} > x_{m+i}$ for all i. Clearly, x_m is the smallest invariant element of $X^{(0)}$ and $x_m \cdot x_m = e > a_1$. By Proposition 4.4, there is a unique L-algebra $X^{(1)}$ on L_{m+2} such that $p \cdot x_{m+1} = a_1$ and which restricts to $X^{(0)}$ on L_{m+1} . Note that x_m is still the smallest invariant element in $X^{(1)}$ as $a_1 > x_{m+1}$. By repeated use of the proposition, one can show inductively that there is a unique sequence of L-algebras $X^{(i)}$ on L_{m+1+i} for all $i \leqslant n-m-1$ such that $X^{(i)}$ restricts to $X^{(i-1)}$ on L_{m+i} , whose smallest invariant element is x_m and such that $x_m \cdot x_{m+i} = a_i > x_{m+i}$.

We conclude that there is a unique L-algebra on L_n that has x_m as its smallest invariant element, restricts to X' on L_m and fulfils

$$\{x_m \cdot x_j : j > m\} = A.$$

By induction, it follows that

$$\lambda(n,m) = \binom{n-2}{n-m-1} B(m-1) = \binom{n-2}{m-1} B(m-1).$$

Using a well-known recursion formula for Bell numbers (see for example [10]), we conclude that

$$\lambda(n) = \sum_{m=1}^{n-1} \lambda(n,m) = \sum_{m=1}^{n-1} \binom{n-2}{m-1} B(m-1) = B(n-1).$$

Unfortunately, despite using the recursive construction of finite linear L-algebras, this recursion at first glance does not seem to translate into the recursive construction of partitions that leads to the used recursion formula of Bell numbers. We, therefore, pose the following problem:

Problem 4.6. Find an explicit bijection between L-algebras on L_n and partitions of the set $\{0, \ldots, n-1\}$.

5. Regular L-algebras

Definition 5.1. An L-algebra X is called *semiregular* if

$$((x \cdot y) \cdot z) \cdot ((y \cdot x) \cdot z) = ((x \cdot y) \cdot z) \cdot z$$

for all $x, y, z \in X$.

Proposition 5.2 (Rump). If X is a semiregular L-algebra, then $x \leq y \cdot x$ for all $x, y \in X$.

Proof. This is [12, Proposition 14].

The proposition implies that in a semiregular L-algebra X, all upsets

$$x^{\uparrow} = \{ y \in X : y \geqslant x \}$$

are sub-L-algebras.

Definition 5.3. A regular L-algebra is a semiregular L-algebra X such that for all $x, y \in X$ with $x \leq y$ there exists a $z \geq x$ such that $z \cdot x = y$.

A big family of regular L-algebras can be constructed from ℓ -groups. In the following, we will collect the necessary definitions and facts.

First of all, an ordered group is a pair (G, \leq) of a group G and a partial order relation \leq on G that is invariant under multiplication in the sense that for all $x, y, z \in G$, the implication

$$x \leqslant y \Rightarrow (xz \leqslant yz)$$
 and $(zx \leqslant zy)$

holds. As usual, we will just say that G itself is an ordered group whenever it is clear from the context that G is equipped with an appropriate order relation. If G is an ordered group, we call G a lattice-ordered group (ℓ -group, for short) when G is a lattice under \leq . This means that for any elements $x,y\in G$, there is a meet $x\wedge y$, i.e. a greatest element z such that $z\leq x$ and $z\leq y$; also, there is a join $x\vee y$, i.e. a least element z such that $z\geq x$ and $z\geq y$. It can be shown that in an ℓ -group G, for any $x,y,z\in G$, the following equations hold:

$$(x \wedge y)z = xz \wedge yz, \quad z(x \wedge y) = zx \wedge zy$$

 $(x \vee y)z = xz \vee yz, \quad z(x \vee y) = zx \vee zy.$

An important example of a right ℓ -group is the additive group \mathbb{Z} , with \leq being the natural order on the integers. The lattice operations are given by $x \wedge y = \min\{x,y\}$ and $x \vee y = \max\{x,y\}$, respectively, where $x,y \in \mathbb{Z}$.

 ℓ -groups give rise to a special class of L-algebras. Let G be an ℓ -group and define its negative cone as $G^- = \{g \in G : g \leq e\}$. Then G^- has a canonical L-algebra structure given by the binary operation

$$x \cdot y = (x \wedge y)x^{-1} = e \wedge yx^{-1}.$$

In particular, (G^-, \cdot) is a regular L-algebra, see [12, page 2343]. Except for trivial examples, these L-algebras are infinite. However, as $x \cdot y \geqslant y$ holds in any regular L-algebra by Proposition 5.2, it follows that each non-empty, upwards-closed subset $X \subseteq G^-$ is closed under the L-algebra operation and is therefore a regular L-algebra by restriction. Here, a subset $X \subseteq G^-$ is called upwards-closed if for all $x, y \in G^-$ with $x \in X$ and $y \geqslant x$, we also have that $y \in X$. Although G^- is almost always infinite, a sub-L-algebra thereof can be finite in a few more cases.

Before stating Rump's embedding theorem for regular L-algebras, we need the notion of dense element (which can be defined in arbitrary L-algebras).

Definition 5.4. An element x in an L-algebra x is called *dense* whenever there is an $y \in X$ with $x \ge y$ and $x \cdot y = y$.

Theorem 5.5. An L-algebra X can be embedded as a sub-L-algebra in the negative cone of some ℓ -group G if and only if X is regular and has no dense elements aside from e. For such an L-algebra, the ℓ -group G can be chosen in such a way that G^- is generated as a monoid by X, in which case, G is unique up to isomorphism of ℓ -groups.

Proof. See [12, Theorem 4.]. The second part essentially follows from Theorem 3. which implies that the monoid and lattice structures of G^- are determined by X - which implies that those of G are, as well.

If $(G_i)_{i\in I}$ is a (not necessarily finite) family of ℓ -groups, we can form their cardinal sum

$$\bigoplus_{i \in I} G_i := \left\{ (g_i)_{i \in I} \in \prod_{i \in I} G_i \ : \ g_i = 0 \text{ for all but finitely many } i \in I \right\}$$

which again is an ℓ -group under coordinate-wise group and lattice operations. Two important examples of such ℓ -groups are $\mathbb{Z}^m = \bigoplus_{i=1}^m \mathbb{Z}$ and $\mathbb{Z}^\omega = \bigoplus_{i=1}^\infty \mathbb{Z}$.

Definition 5.6. An *infinite-dimensional Young-diagram* is a finite, non-empty and upwards-closed subset in $(\mathbb{Z}^{\omega})^-$.

Definition 5.7. Two infinite-dimensional Young-diagrams are said to be *conjugate* if they can be converted into one another by a permutation of the coordinates of \mathbb{Z}^{ω} .

As an upwards-closed set in the negative cone of an ℓ -group, each infinite-dimensional Young-diagram becomes an instance of a regular L-algebra after restricting the L-algebra operations from $(\mathbb{Z}^{\omega})^-$ to X. Also, conjugate infinite-dimensional Young-diagrams have isomorphic L-algebra structures, for permuting the coordinates of \mathbb{Z}^{ω} is clearly an automorphism of ℓ -groups and therefore also induces an automorphism of the respective L-algebra structure. This implies that conjugate infinite-dimensional Young-diagrams are related by an isomorphism of L-algebras.

When enumerating regular L-algebras we observed that the first numbers are equal to those in sequence A119268 of [18] which counts infinite-dimensional Young-diagrams up to conjugacy. This suggested that infinite-dimensional Young-diagrams might actually be the same as finite regular L-algebras.

Lemma 5.8. Let X be a finite regular L-algebra and $x \in X$. If x is dense, then x = e.

Proof. Let $x \ge y$ be elements in X such that $x \cdot y = y$. By Proposition 5.2, the map $y^{\uparrow} \to y^{\uparrow}$, $x \mapsto x \cdot y$, is well-defined and, due to regularity, surjective. Since X is finite, this map is also injective which implies that the element $y \in y^{\uparrow}$ has exactly one preimage, which must be e.

Theorem 5.9. There exists a bijective correspondence between finite regular L-algebras and infinite-dimensional Young diagrams.

Proof. By Theorem 5.5, there is an ℓ -group G with an embedding $X \hookrightarrow G^-$ that identifies X as a sub-L-algebra of G^- and a generating set thereof. For sake of simplicity, we will identify X with its image in G^- . We want to show that $G \simeq \mathbb{Z}^m$ and that X is upwards-closed in G^- under this embedding.

Recall, that for $x, y \in X$, the relation $x \prec y$ means that x < y and there is no $z \in X$ such that x < z < y. Let $\text{Max}(X) := \{x \in X : x \prec e\}$ be the set of elements lying directly under e.

We claim that G^- is generated by Max(X). Let $M \subseteq G^-$ be the submonoid generated by Max(X). If there was an element in $X \setminus M$, there would also be a

biggest one, say $y \in X \setminus M$. Necessarily, there exists $x \in \operatorname{Max}(X)$ with $x \geqslant y$. By Proposition 5.2 and Lemma 5.8, $x \cdot y > y$. Since $x \in \operatorname{Max}(X)$, $x \cdot y \in M$. Also, we calculate in G^- that $y = x \wedge y = (x \cdot y)x \in M$, a contradiction. We conclude that $X \subseteq M$. As G^- is generated by X, it follows that G^- is generated by $\operatorname{Max}(X)$.

Let now $x,y \in \operatorname{Max}(X)$. By Proposition 5.2, $x \cdot y \geqslant y$. Here, $x \cdot y = e$ is only possible when $x \leqslant y$, i.e. when x = y. Therefore, $x \cdot y = y$ for $x,y \in X$ with $x \neq y$. In this case, one can calculate in G^- that $x \wedge y = (x \cdot y)x = yx$ which is the same as $y \wedge x = xy$. These relations show that we have a decomposition of the ℓ -group G as a cardinal sum

$$G \simeq \bigoplus_{x \in \operatorname{Max}(X)} \mathbb{Z}$$

where Max(X) is identified with the elements in the right group that are -1 in exactly one coordinate and 0 in all others.

It remains to show that X is upwards-closed in G^- . It clearly suffices to show that for $y \in X$ and all $x \in G^-$ with $x \succ y$, we also have $x \in X$. In such a case, $z = yx^{-1} \in \operatorname{Max}(X)$. As we have already seen that G is abelian, we also have $z = x^{-1}y \geqslant y$. We can now calculate in G^-

$$z \cdot y = (z \wedge y)z^{-1} = y(y^{-1}x) = x.$$

As X is a sub-L-algebra of G^- , this implies that $x \in X$.

We have thus shown that each finite regular L-algebra X is isomorphic to an upwards-closed subset in the negative cone of some ℓ -group \mathbb{Z}^m . Under a coordinatewise embedding $\mathbb{Z}^m \hookrightarrow \mathbb{Z}^\omega$, X is identified with an infinite-dimensional Young-diagram. By the uniqueness part of Theorem 5.5, the embedding groups of isomorphic regular L-algebras without nontrivial dense elements are also isomorphic, in the sense of ℓ -groups. Is is easily seen that each isomorphism of two ℓ -groups of the form \mathbb{Z}^m is given by a permutation of coordinates. This implies that each isomorphism class of finite regular L-algebras can be identified with a unique conjugacy class of infinite-dimensional Young-diagrams and that only conjugate Young-diagrams are isomorphic as L-algebras.

ACKNOWLEDGMENTS

We thank Wolfgang Rump for comments and suggestions. This research was supported through the program "Oberwolfach Research Fellows" by the Mathematisches Forschungsinstitut Oberwolfach in 2022. Menchón was partially supported by the National Science Center (Poland), grant number 2020/39/B/HS1/00216; for the purpose of Open Access, the author has applied a CC-BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission. Vendramin was supported in part by OZR3762 of Vrije Universiteit Brussel.

References

- O. Akgün, M. Mereb, and L. Vendramin. Enumeration of set-theoretic solutions to the Yang-Baxter equation. Math. Comp., 91(335):1469-1481, 2022.
- [2] B. Bosbach. Rechtskomplementäre Halbgruppen. Axiome, Polynome, Kongruenzen. Math. Z., 124:273–288, 1972.
- [3] P. Dehornoy. Set-theoretic solutions of the Yang–Baxter equation, RC-calculus, and Garside germs. Adv. Math., 282:93–127, 2015.
- [4] P. Dehornoy, F. Digne, E. Godelle, D. Krammer, and J. Michel. Foundations of Garside theory, volume 22 of EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2015.
- [5] A. Diego. Sur les algèbres de Hilbert. Collection de Logique Mathématique, Sér. A, Fasc. XXI. Gauthier-Villars, Paris; E. Nauwelaerts, Louvain, 1966. Translated from the Spanish by Luisa Iturrioz.

- $[6] \ \ The \ GAP \ Group. \ \textit{GAP-Groups}, \ \textit{Algorithms}, \ \textit{and Programming}, \ \textit{Version 4.11.1}, \ 2021.$
- [7] L. Herman, E. L. Marsden, and R. Piziak. Implication connectives in orthomodular lattices. Notre Dame J. Formal Logic, 16:305–328, 1975.
- [8] D. Mundici. Interpretation of AF C^* -algebras in łukasiewicz sentential calculus. J. Funct. Anal., 65(1):15–63, 1986.
- [9] P. Nightingale, Ö. Akgün, I. P. Gent, C. Jefferson, I. Miguel, and P. Spracklen. Automatically improving constraint models in Savile Row. Artificial Intelligence, 251:35–61, 2017.
- [10] G.-C. Rota. The number of partitions of a set. Amer. Math. Monthly, 71:498-504, 1964.
- [11] W. Rump. A decomposition theorem for square-free unitary solutions of the quantum Yang-Baxter equation. Adv. Math., 193(1):40–55, 2005.
- [12] W. Rump. L-algebras, self-similarity, and l-groups. J. Algebra, 320(6):2328-2348, 2008.
- [13] W. Rump. Semidirect products in algebraic logic and solutions of the quantum Yang-Baxter equation. J. Algebra Appl., 7(4):471–490, 2008.
- [14] W. Rump. Right l-groups, geometric Garside groups, and solutions of the quantum Yang-Baxter equation. J. Algebra, 439:470–510, 2015.
- [15] W. Rump. The L-algebra of Hurwitz primes. J. Number Theory, 190:394-413, 2018.
- [16] W. Rump. Von Neumann algebras, L-algebras, Baer *-monoids, and Garside groups. Forum Math., 30(4):973–995, 2018.
- [17] W. Rump. Symmetric quantum sets and L-algebras. Int. Math. Res. Not. IMRN, (3):1770– 1810, 2022.
- [18] N. J. A. Sloane and T. O. F. Inc. The on-line encyclopedia of integer sequences, 2020.
- [19] T. Traczyk. On the structure of BCK-algebras with $zx \cdot yx = zy \cdot xy$. Math. Japon., 33(2):319–324, 1988.
- [20] Y. Yang and W. Rump. Pseudo-MV algebras as L-algebras. J. Mult.-Valued Logic Soft Comput., 19(5-6):621–632, 2012.

Department of Logic, Nicolaus Copernicus University in Toruń, Fosa Staromiejska 1a, 87-100 Toruń, Poland

Email address: paula.menchon@v.umk.pl

IMAS-CONICET AND DEPTO. DE MATEMÁTICA, FCEN, UNIVERSIDAD DE BUENOS AIRES, PAB. 1, CIUDAD UNIVERSITARIA, C1428EGA, BUENOS AIRES, ARGENTINA

 $Email\ address: {\tt lvendramin@dm.uba.ar}$

Department of Mathematics and Data Science, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel

Email address: carstendietzel@gmx.de
Email address: Leandro.Vendramin@vub.be