

1 **Effects of cadmium and phenanthrene mixtures on aquatic fungi and microbially**
2 **mediated leaf litter decomposition**

3 Catarina Moreirinha, Sofia Duarte, Cláudia Pascoal, Fernanda Cássio

4

5

6 Centre of Molecular and Environmental Biology (CBMA), Department of Biology,

7 University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

8

9

10

11

12 For correspondence:

13 Cláudia Pascoal

14 E-mail cpascoal@bio.uminho.pt

15 Phone +351253604045

16 Fax. +351253678980

17

1 **Abstract**

2 Urbanization and industrial activities have contributed to widespread contamination by
3 heavy-metals and polycyclic aromatic hydrocarbons, but the combined effects of these
4 toxics on aquatic biota and processes are poorly understood. We examined the effects of
5 cadmium (Cd) and phenanthrene (Phe) on the activity and diversity of fungi associated
6 with decomposing leaf litter in streams. Leaves of *Alnus glutinosa* were immersed for
7 10 days in a unpolluted low-order stream in the NW Portugal to allow microbial
8 colonization. Leaves were then exposed in microcosms for 14 days to realistic
9 concentrations of Cd (0.06 - 4.5 mg L⁻¹) and phenanthrene (0.2 mg L⁻¹) either alone or
10 in mixtures. A total of 19 aquatic hyphomycete species were found sporulating on
11 leaves during the whole study. The dominant species was *Articulospora tetracladia*,
12 followed by *Alatospora puchella*, *Clavatospora longibrachiata* and *Tetrachaetum*
13 *elegans*. The exposure to Cd and phenanthrene decreased the contribution of *A.*
14 *tetracladia* to the total conidium production while increased that of *A. pulchella*. Fungal
15 diversity, assessed as DGGE or conidium morphology, was lowered by the exposure to
16 Cd and/or phenanthrene. Moreover, increased Cd concentrations depressed leaf
17 decomposition and fungal reproduction but did not inhibit fungal biomass production.
18 The exposure to phenanthrene potentiated the negative effects of Cd on fungal diversity
19 and activity, suggesting that more studies should be conducted in order to assess the
20 effects of multiple stressors on aquatic biodiversity and stream ecosystem functioning,
21 since it might correspond to more realistic approaches.

22

1 **Introduction**

2 In streams with low autotrophic production, plant-litter decomposition provides carbon
3 and energy for the functioning of aquatic foodwebs (Allan and Castillo 2007). This
4 carbon is first processed through microbial activity increasing the palatability of plant
5 litter for aquatic invertebrates (Bärlocher 2005; Gessner et al. 2007). Among
6 microorganisms, aquatic hyphomycetes play a key role in plant-litter decomposition in
7 streams (Pascoal and Cássio 2004; Pascoal et al. 2005a) due to their ubiquity and ability
8 to produce a variety of extracellular enzymes that degrade the complex plant cell-wall
9 polysaccharides (Bärlocher 2005; Gessner et al. 2007).

10 Human activities from agriculture, mining and industry have contributed to the increase
11 of contaminants entering streams that affect the activity and diversity of biotic
12 communities, including detritus-based food webs (Niyogi et al. 2001; Pascoal et al.
13 2005a,b; Sridhar et al. 2001). Among pollutants, metals can have adverse effects on
14 aquatic biota because of their toxicity and persistence in the environment (Rand et al.
15 1995). Even though aquatic hyphomycetes have been found in metal-polluted streams,
16 the diversity and activity of this group of fungi is constrained by the presence of heavy
17 metals (Bermingham et al. 1996; Niyogi et al. 2002; Pascoal et al. 2005b; Sridhar et al.
18 2005). The exposure of leaf-associated fungal communities to copper and/or zinc has
19 been reported to alter the structure of aquatic hyphomycete communities and reduce
20 fungal sporulation and leaf decomposition (Duarte et al. 2004, 2008a, 2009a).
21 Moreover, metals, including Cd, inhibit the growth and reproduction of several species
22 of aquatic hyphomycetes, but reproduction appears to be more sensitive than growth
23 (Abel and Bärlocher 1984; Azevedo and Cássio 2010; Miersch et al. 1997).

24 Polycyclic aromatic hydrocarbons (PAHs) are a class of toxins containing two or more
25 benzene rings that are known to have carcinogenic and mutagenic properties (Chaudry

1 1994). These compounds are components of coal, crude oil and its derivatives and are
2 ubiquitous in the environment as they are formed during forest fires, combustion of
3 petroleum and incineration of waste. Several studies have shown that microbes have the
4 ability to metabolize and biodegrade PAHs and can potentially be used for
5 bioremediation of contaminated environments (D'Annibale et al. 2006; Johnsen et al.
6 2005; Juhasz and Naidu 2000). For instance, the bacterium *Pseudomonas putida*
7 degraded about 70% of the initial amount of phenanthrene (Phe, 0.47 g L⁻¹) in 27 days
8 (Cuny et al. 2004) and the aquatic hyphomycete *Heliscus lugdunensis* is reported to
9 metabolize metabolites of complex PAHs, such as 1-naphthol (Augustin et al. 2006).
10 PAHs do not dissolve easily in water but can bind strongly to particulate organic matter
11 and accumulate in the sediments of rivers and lakes at high concentrations, affecting
12 aquatic biota and humans through bioaccumulation (Gust and Fleeger 2006; Ribeiro et
13 al. 2005; Scoggins et al. 2007).

14 In aquatic ecosystems, the simultaneous occurrence of PAHs and heavy metals is quite
15 common (Gust and Fleeger 2006). While many studies have investigated the individual
16 effects of these toxins on organisms, relatively few have considered their effects in
17 mixtures, particularly at the community level. In this study, the effects of Cd and
18 phenanthrene on the diversity and activity of aquatic fungal communities associated
19 with decomposing leaves were evaluated. Alder leaves were incubated in a stream to
20 allow microbial colonization and then were exposed to increasing concentrations of Cd
21 (0.06 to 4.5 mg L⁻¹), in the absence or presence of a fixed concentration of phenanthrene
22 (0.2 mg L⁻¹). The effects were evaluated on leaf mass loss and fungal biomass,
23 sporulation and diversity. Although there is no data concerning the co-occurrence of
24 metals and PAHs in Portuguese streams, in the Northwest of Portugal there are several
25 industrial areas and pollution by both kind of toxins are likely to occur in freshwaters

1 (Gust and Fleeger 2006). Although lower concentrations of Cd have been found in the
2 water column in streams of the Northwest Portugal (0.06 mg L^{-1} , Gonçalves 2001), than
3 those used in the current study, in sediments we found concentrations between 0.02 and
4 144 Kg g^{-1} of volatile matter dry weight in the $<63 \mu\text{m}$ sediments fractions (Soares et al.
5 1999). The variation of some physical and chemical characteristics (pH, salinity, redox
6 potential and content of organic chelators) of the water may provoke the release of these
7 metals back to the aqueous phase and higher concentrations of Cd are expected to occur
8 in water column. We expect that Cd and phenanthrene will restrict fungal diversity and
9 activity, and this will be more severe in mixtures with increasing Cd concentrations.
10 However, we also expect that some species might be able to metabolize phenanthrene and
11 thereby survive under the tested conditions.

12

13 **Materials and Methods**

14 **Sampling site**

15 The sampling site is located at Algeriz, a low-order stream in the Northwest of Portugal
16 ($41^{\circ}35' \text{N } 8^{\circ}22' \text{W}$). The riparian vegetation is dominated by *Eucalyptus globulus* Labill.,
17 *Quercus robur* L., *Alnus glutinosa* (L.) Gaertn and *Rubus* sp.

18 Leaves of *A. glutinosa*, collected in October 2006, were air-dried and kept at room
19 temperature until used. The leaves were leached in deionised water for 48 hours and cut
20 into 12 mm diameter disks. Sets of 22 disks were placed into 42 fine-mesh bags (20 x
21 20 cm, 0.5 mm mesh size) to prevent invertebrate colonization. On 20th March 2007,
22 leaf bags were immersed in the stream for 10 days to allow microbial colonization. An
23 additional set of 3 bags were immersed for 15 minutes in the sampling site and the
24 content used to estimate initial leaf dry mass and ergosterol concentration in the leaves.

1 At the time of leaf immersion, stream water had a temperature of 11.8 °C, a pH of 6.9, a
2 redox potential of -8 mV, a concentration of dissolved oxygen of 10.8 mg L⁻¹ and a
3 conductivity of 40 µS cm⁻¹, measured *in situ* with field probes (Multiline F/set 3 n°
4 400327, WTW). Stream water was collected in dark glass bottles, transported on ice to
5 the laboratory and analyzed within 6 h for quantification of inorganic nutrients with a
6 HACH DR/2000 photometer (Hach company, Loveland, CO, USA). Nutrient
7 concentrations were: orthophosphate, 0.06 mg P-PO₄³⁻ L⁻¹ (HACH kit, program 490);
8 nitrate, 0.2 mg N-NO₃⁻ L⁻¹ (HACH kit, program 355); nitrite, 0.005 mg N-NO₂⁻ L⁻¹
9 (HACH kit, program 371); and ammonium <0.01 mg N-NH₃ L⁻¹ (HACH kit, program
10 385). Additional stream water was collected for microcosm experiments.

11

12 Microcosms

13 In the laboratory, sets of 22 leaf disks were placed into 150 mL Erlenmeyer flasks with
14 70 mL of filtered (Macherey-Nagel MN-GF3, glass fiber filter membranes) and
15 sterilized (120 °C, 20 min) stream water. The microcosms were supplemented with
16 cadmium chloride (Sigma) at final concentrations of 0.06, 0.6, 1.2, 3.6 and 4.5 mg L⁻¹
17 Cd and phenanthrene (Fluka) at a final concentration of 0.2 mg L⁻¹, added alone or in
18 mixtures (3 replicates). Phenanthrene was solubilized in ethanol at a final concentration
19 of 0.3 %. Microcosms were incubated on a shaker (110 rpm, Certomat BS 3, Braun,
20 Melsungen, Germany), at 15 °C, in the dark. The microcosm solutions were changed
21 every 4 days, and phenanthrene concentration was checked daily by direct fluorescence
22 spectrophotometry (LS 50 luminescence spectrometer, Perkin-Elmer, Foster city, CA,
23 USA) using the fixed wavelength technique (Watson et al. 2004; Yang et al. 2003).
24 Microcosms without added toxins and microcosms with ethanol, at the concentration
25 used to solubilize phenanthrene, were used as controls (3 replicates).

1 After 14 days of exposure, all microcosms were sacrificed and leaf disks were freeze-
2 dried and weighed (± 0.001 g) for determination of leaf dry mass remaining, and stored
3 at -80 °C for further assays. Sets of non-colonized leaf disks were used to estimate the
4 initial leaf dry mass.

5

6 Fungal biomass

7 Concentration of ergosterol was measured to estimate fungal biomass associated with
8 decomposing leaf disks. Sets of 8 disks, from each replicate microcosm, were heated
9 (80 °C, 30 min) in 0.8 % KOH-methanol for lipid extraction and the extract was purified
10 by solid-phase extraction, according to Gessner (2005). Ergosterol was quantified by
11 high-performance liquid chromatography (HPLC, Beckmann Gold System, Brea, CA,
12 USA) using a LiChrospher RP18 column (Merck). The system was run isocratically
13 with HPLC-grade methanol at 1.4 mL min^{-1} at 33 °C. Peaks of ergosterol were detected
14 at 282 nm and standard series of ergosterol (Sigma) in isopropanol were used to
15 estimate ergosterol concentration in the samples.

16

17 Fungal diversity

18 After 10 and 14 days in microcosms, conidium suspensions were filtered ($5 \mu\text{m}$ pore
19 size; Millipore, Billerica, MA, USA) and the spores were stained with cotton blue in
20 lactic acid. Conidia were counted and identified under a microscope ($400\times$; Leica
21 Biomed, Heerbrugg, Switzerland).

22 DNA was extracted from sets of 3 leaf disks (1 from each replicate) using an UltraClean
23 Soil DNA kit (MoBio Solana Beach, CA, USA). The ITS2 region of fungal ribosomal
24 DNA was amplified by PCR, with ITS3GC and ITS4 primers (White et al. 1990). The
25 reaction mixture contained $4 \mu\text{M}$ of each primer, $1 \mu\text{L}$ (~ 50 ng) of DNA, 1.5 U of Taq

1 polymerase, 3 mM of MgCl₂, 2 mM of dNTPs and 1x of Taq buffer (KCl:(NH₄)₂SO₄)
2 in a final volume of 50 µL. Fungal DNA amplification was carried out in an iCycler
3 Thermal Cycler (BioRad Laboratories, Hercules, CA, USA) and PCR started with a
4 denaturation of 2 min at 95 °C, followed by 36 cycles of denaturation for 30 s at 95 °C,
5 primer annealing for 30 s at 55 °C and extension for 1 min at 72 °C. Final extension was
6 at 72 °C for 5 min (Nikolcheva and Bärlocher 2005; Duarte et al. 2008a).
7 PCR products were analyzed by denaturing gradient gel electrophoresis (DGGE), using
8 a DCode Universal Mutation Detection System (BioRad). Samples of ~750 ng DNA
9 were loaded on 8 % polyacrilamide gel in 1x Tris-acetate-EDTA (TAE) with a
10 denaturing gradient from 30 to 70 % (100 % denaturant corresponds to 40 % formamide
11 and 7 M urea). The gel was run at 55 V, 56 °C for 16 hours and was stained with 1 µg
12 mL⁻¹ of ethidium bromide (BioRad). The images were captured under UV light in a
13 transilluminator Eagle Eye II (Stratagene, La Jolla, CA, USA).

14

15 Data analysis

16 Leaf mass loss, fungal biomass, and sporulation were expressed as percentage of
17 control. To achieve normal distribution and homocedasticity, data were arcsine square
18 root transformed (Zar 1996). Ethanol at the concentration used to solubilize
19 phenanthrene did not significantly inhibit leaf mass loss, fungal biomass and sporulation
20 (t-tests, P > 0.05; Zar 1996). The effects of Cd and phenanthrene on leaf mass loss and
21 ergosterol concentration were tested by a two-way ANOVA (Zar 1996). A three-way
22 ANOVA (Zar 1996) was used to test if Cd, phenanthrene and exposure time affected
23 sporulation rates and the number of leaf-associated aquatic hyphomycete taxa.
24 Dunnett's post-tests were done to test which treatments significantly differed from

1 control (Zar 1996). Statistical analyses were performed using Statistica 7.0 (Statsoft,
2 Inc., Tulsa, OK, USA).
3 The DGGE gel was aligned and normalized using Gelcompar II (Applied Maths, Sint-
4 Martens-Latem, Belgium). Each DGGE band was treated as an individual operational
5 taxonomic unit (OTU) and the number of OTUs was used as a measure of fungal
6 diversity. A correspondence analysis (CA) (Legendre and Legendre 1998) was used to
7 ordinate treatments using fungal communities, based on sporulating species or OTUs
8 (as relative intensity of each band in DGGE fingerprinting). The analyses were done
9 using CANOCO 4.5 (Microcomputer Power, Ithaca, NY, USA).

10

11 **Results**

12 Effects of cadmium and phenanthrene on fungal activity

13 In the absence of Cd and phenanthrene, alder leaves lost approximately 56% of their
14 initial dry mass (0.061 ± 0.009 g) after a 10-day colonization in the stream and 14 days in
15 microcosms. Fungal biomass was $339 \mu\text{g ergosterol g}^{-1}$ dry mass, and fungal sporulation
16 rate on decomposing leaves was 2.94×10^5 and 0.31×10^5 conidia g^{-1} dry mass day^{-1}
17 after 10 and 14 days, respectively.

18 Leaf mass loss was significantly affected by Cd, phenanthrene and the interaction
19 between both toxins (2-way ANOVA, $P < 0.01$ for all factors) (Fig. 1A). Significant
20 inhibition effects were found following the exposure to Cd at high concentrations (≥ 3.6
21 mg L^{-1} ; Dunnett's test, $P < 0.05$) and to phenanthrene alone or in mixtures with Cd
22 (Dunnett's test, $P < 0.01$).

23 Fungal biomass on decomposing leaves was significantly affected by Cd and
24 phenanthrene (2-way ANOVA, $P = 0.001$ and $P = 0.003$, respectively) (Fig. 1B).
25 Significant inhibition effects were found by exposure to mixtures of phenanthrene and

1 Cd at the highest concentration (4.5 mg L⁻¹; Dunnett's test, P<0.05). It seems to exist a
2 pattern that shows that there is a slight stimulation in biomass production with the lower
3 cadmium concentrations, which is significant in one treatment (1.2 mg L⁻¹).
4 Sporulation of aquatic hyphomycetes was significantly affected by the concentration of
5 Cd and phenanthrene (3-way ANOVA, P<0.001, for both comparisons) (Fig. 1C) but
6 not by the exposure time (3-way ANOVA, P=0.1). Moreover, interactions between
7 phenanthrene and exposure time and between the two toxins were significant (3-way
8 ANOVA, P=0.008 P<0.001, respectively; Fig. 1C). At both exposure times, sporulation
9 rate was inhibited by Cd concentrations ≥ 3.6 mg L⁻¹ (Dunnett's test, P<0.05) and by
10 mixtures of Cd and phenanthrene at concentrations of Cd ≥ 0.06 mg L⁻¹ (Dunnett's test,
11 P<0.05).

12

13 Effects of cadmium and phenanthrene on the structure of fungal community

14 During the study, 19 species of aquatic hyphomycetes were found sporulating on alder
15 leaves (Table 1). The exposure to increasing Cd concentrations led to a decrease in the
16 number of fungal species, particularly in microcosms supplemented with the highest Cd
17 concentration (4.5 mg L⁻¹), in which only 8 fungal species were observed (Table 1).
18 *Articulospora tetraccladia* was the dominant species in control microcosms,
19 contributing to 40% to the total conidia production. The exposure to the toxins,
20 especially in mixtures, led to a decrease in the contribution of *A. tetraccladia* (Table 1).
21 On the other hand, the exposure to Cd, alone or in mixtures with phenanthrene,
22 increased the contribution of the co-dominant species *Alatospora pulchella*.
23 *Lemmoniera aquatica* seemed to be tolerant of Cd, since its contribution increased with
24 metal concentration, attaining 41.3% of the total conidia production at the highest Cd
25 concentration (4.5 mg L⁻¹). Phenanthrene increased ca 5-times the contribution of

1 *Tetrachaetum elegans* to conidium production, in the absence or presence of the lowest
2 Cd concentration. An unidentified species became co-dominant in mixtures with high
3 Cd concentrations.

4 Denaturing gradient gel electrophoresis (DGGE) of DNA of fungal communities on
5 decomposing leaves showed 18 OTUs in control microcosms after 14 days in (Table 1,
6 Fig. 2). The exposure to increasing Cd concentrations led to a decrease in the number of
7 OTUs, particularly in the presence of phenanthrene. Generally, higher fungal diversity
8 was found from DGGE than from conidium identification.

9 The CA ordination of fungal communities based on conidium morphology and DGGE
10 OTUs showed that the exposure to Cd and phenanthrene altered the structure of fungal
11 community, with stronger effects for communities exposed to mixtures of both stressors
12 (Fig. 3). The analysis of fungal communities based on conidium morphology showed
13 that the first axis, explaining 58% of the total variance, distributed communities along
14 the gradient of Cd concentration in the absence or presence of phenanthrene (Fig. 3A).
15 The second CA axis explained 37% of the total variance and separated communities
16 exposed to phenanthrene from all the others. The CA ordination of fungal communities
17 based on DGGE OTUs distributed communities along the gradient of Cd concentration
18 defined by the first axis (54% of the total variance), while the second axis (12% of the
19 total variance) separated control communities from all the other treatments, and further
20 separated communities exposed to mixtures of phenanthrene with increasing Cd
21 concentrations (Fig. 3B).

22

23 **Discussion**

24 In the current study, the exposure of freshwater microbial decomposer communities to
25 high Cd concentrations ($\geq 3.6 \text{ mg L}^{-1}$) led to a reduction in leaf decomposition. This

1 might be explained by the reported negative effects of Cd on the activity of fungal
2 extracellular enzymes that degrade plant cell-wall polymers (Baldrian et al. 1996). A
3 poor use of resources were consistent with the observed decrease in fungal activity as
4 sporulation rates in treatments with the highest Cd concentrations, particularly in the
5 presence of phenanthrene. In previous studies concentrations of Cd up to 0.1 mg L⁻¹
6 inhibited both growth and sporulation of aquatic hyphomycetes (Abel and Bärlocher
7 1984) and 1.5 mg L⁻¹ of Cd inhibited leaf decomposition by three strains of aquatic
8 hyphomycetes isolated from a clean site (Fernandes 2008). Although there are no
9 studies in literature about the joint effects of metals and PAHs on aquatic microbially
10 mediated processes, Cd and phenanthrene are reported to have variable effects on
11 organisms depending on the concentration and magnitude of interactions between toxins
12 (Moreau et al. 1999). For instance, the exposure to a mixture of the two toxins resulted
13 in antagonistic effects on the feeding rate of the oligochaete *Ilyodrilus templetoni* (Gust
14 and Fleeger 2006). In another study, Cd combined with phenanthrene decreased the
15 grazing rates of the copepod *Schizopera knabeni* but no interactive effects on feeding
16 were found (Silva et al. 2009). In soils, the addition of heavy metals and PAHs,
17 including Cd and phenanthrene, led to a greater biocidal effect on microbes when toxins
18 were amended together than alone (Maliszewska-Kordybach and Smreczak 2003; Shen
19 et al. 2005). In our study, phenanthrene appeared to accentuate the negative effects of
20 Cd because the highest inhibitions of fungal biomass and sporulation rates were in
21 treatments with both toxins.

22 The inhibitions of fungal biomass and sporulation after exposure to the toxins were
23 probably due to the fact that aquatic fungi were channeling part of the energy available
24 for growth and reproduction to the synthesis of compounds and/or enzymes involved in
25 cellular detoxification processes. Indeed, aquatic hyphomycetes are reported to be able

1 to trigger defense mechanisms against heavy-metal exposure by increasing the activity
2 of antioxidant enzymes (Azevedo et al. 2007, 2009; Braha et al. 2007) or the production
3 of thiol-containing compounds that are able to sequester metal ions or scavenge ROS
4 (Guimarães-Soares et al. 2007; Jaeckel et al. 2005; Miersch et al. 1997). Other possible
5 responses of fungi to heavy metals include the secretion of organic acids,
6 polysaccharides, melanin or proteins capable of complexing and / or precipitate metals,
7 regulation of the transport of metallic, adsorption of metals to the cell wall, and its
8 chemical transformation and cellular compartmentalization (Baldrian 2003, Blaudez et
9 al. 2000). On the other hand, there are studies reporting that white rot fungi are able to
10 metabolize PAHs, including phenanthrene, by the phase I enzymes cytochrome P-450
11 monooxygenase and epoxide hydrolase (Bezalel et al. 1996, 1997; Capotorti et al.
12 2004). Other enzymes, such as manganese peroxidase, are also involved in the
13 degradation of phenanthrene by white-rot fungi (Baborová et al. 2006). However, the
14 addition of phenanthrene led to a more pronounced effect of Cd on the activity of
15 several enzymes produced by soil microorganisms (Shen et al. 2005). We also noticed
16 that Cd alone (1.2 mg L^{-1}) stimulated biomass production. Other studies revealed that
17 the growth of aquatic hyphomycetes was less sensitive than sporulation to metals (Abel
18 and Bärlocher 1984, Duarte et al. 2004, 2008a, 2009). Probably fungi limit their
19 sporulation and increase their growth in these conditions as a defense mechanism.
20 Moreover, bacterial biomass on decomposing leaves is also reported to be severely
21 inhibited in the presence of metals (Duarte et al. 2008a, 2009). Antagonistic interactions
22 between fungi and bacteria are reported to occur during leaf decomposition in streams
23 (Mille-Lindblom and Tranvik 2003); both groups of microorganisms can compete for
24 resources and thus if bacteria are more sensitive to metals, fungi might take advantage
25 and increase its biomass.

1 A decrease in aquatic hyphomycete diversity has been found in metal-polluted streams
2 (Bermingham et al. 1996; Niyogi et al. 2002; Pascoal et al. 2005b; Sridhar et al. 2001).
3 Also, in our study, the exposure to increasing concentrations of Cd caused a decrease in
4 the number of fungal taxa, assessed as conidium morphotypes or DGGE OTUs, with the
5 highest declines in treatments with the highest Cd concentrations and when both toxins
6 were added together (ca. 2- to 3-times lower than control microcosms). However, the
7 effects did not appear to be so drastic when diversity was assessed based on DGGE.
8 This is not surprising since analysis based on DNA offers the advantage of detecting
9 species from non-sporulating mycelia and sporulation has been consistently found to be
10 very sensitive to pollutants (Duarte et al. 2004, 2008a; Niyogi et al. 2002; Sridhar et al.
11 2001, 2005).

12 The exposure to toxins also induced shifts in the structure of fungal community revealed
13 by multivariate analyses based on both sporulating species or OTUs. In this and other
14 studies, microbial communities exposed to high metal concentrations or mixtures of
15 pollutants were most different from control communities (Duarte et al. 2004, 2008a,
16 2009). In our study, the dominance pattern of sporulating fungal species was affected by
17 exposure to both toxins; the dominant species in control, *Articulospora tetracladia*,
18 showed high sensitivity to toxics similar to that found in field observations (Duarte et al.
19 2008b; Pascoal et al. 2005b), while *Alatospora pulchella* was the most tolerant species
20 to Cd, even in the presence of phenanthrene. Some species of aquatic hyphomycetes,
21 such as *Heliscus lugdunensis* and *Flagellospora curta*, are reported to tolerate high
22 levels of Cd (Azevedo and Cássio 2010; Braha et al. 2007; Guimarães-Soares et al.
23 2007) or to metabolize PAHs such as naphthol (Augustin et al. 2006). Interestingly, in
24 our study, some species were even stimulated by phenanthrene, as it was the case of the
25 unknown species, which had increased spore production with

1 exposure to phenanthrene. Overall, results show that the presence of phenanthrene
2 accentuated the deleterious effects of Cd on the diversity and activity of aquatic fungal
3 decomposers. Experiments using multiple stressors might correspond to more realistic
4 approaches to assess the effects of pollutants on aquatic biodiversity and stream
5 ecosystem functioning. In the future, more experiments are needed to better understand
6 the interactions between Cd, phenanthrene and other stressors and to predict their
7 impacts on aquatic detritus foodwebs.

8

9 **Acknowledgements**

10 The Portuguese Foundation for the Science and Technology supported this work
11 (POCI/MAR/56964/2004) and S. Duarte (SFRH/BPD/47574/2008).

12

13 **References**

14 Abel T, Bärlocher F (1984) Effects of cadmium on aquatic hyphomycetes. *Applied and*
15 *Environmental Microbiology* 48: 245-251

16 Alan J, Castillo M (2007) *Stream Ecology* (2nd ed.), Springer, Dordrecht, The
17 Netherlands

18 Augustin T, Schlosser D, Baumbach R, Schmidt J, Grancharov K, Krauss G, Krauss G-
19 K (2006) Biotransformation of 1-Naphtol by a strictly aquatic fungus. *Current*
20 *Microbiology* 52: 216-220

21 Azevedo M, Almeida B, Ludovico P, Cássio F (2009) Metal stress induces programmed
22 cell death in aquatic fungi. *Aquatic Toxicology* 92: 264-270

23 Azevedo M, Carvalho A, Pascoal C, Rodrigues F, Cássio F (2007) Responses of
24 antioxidant defenses to Cu and Zn stress in two aquatic fungi. *Science of the Total*
25 *Environment* 377: 233–243

- 1 Azevedo M, Cássio F (2010) Effects of metals on growth and sporulation of aquatic
2 fungi. *Drug and Chemical Toxicology*. DOI: 10.3109/01480540903431440
- 3 Baborová P, Möder M, Baldrian P, Cajthamlová K, Cajthaml T (2006) Purification of a
4 new manganese peroxidase of the white-rot fungus *Irpex lacteus* and degradation
5 of polycyclic aromatic hydrocarbons by the enzyme. *Research in Microbiology*
6 157: 248-253
- 7 Baldrian P (2003) Interactions of heavy metals with white-rot fungi. *Enzyme and*
8 *Microbial Technology* 32: 78-91
- 9 Baldrian P, Gabriel J, Nerud F (1996) Effect of cadmium on the ligninolytic activity of
10 *Stereum hirsutum* and *Phanerochaete chrysosporium*. *Folia Microbiologica* 41:
11 363-367
- 12 Bärlocher F (2005) Chapter 5: Leaching. In: Graça M, Bärlocher F, Gessner M (eds)
13 *Methods to study litter decomposition*. Springer, Berlin/New York, pp. 33-36
- 14 Bermingham S, Maltby L, Cooke R (1996) Effects of a coal mine effluent on aquatic
15 hyphomycetes. I. Field Study. *Journal of Applied Ecology* 33: 1311-1321
- 16 Bezalel L, Hadar Y, Cerniglia C (1996) Mineralization of polycyclic aromatic
17 hydrocarbons by the white rot fungus *Pleurotus ostreatus*. *Applied and*
18 *Environmental Microbiology* 62: 292–295
- 19 Bezalel L, Hadar Y, Cerniglia C (1997) Enzymatic mechanisms involved in
20 phenanthrene degradation by the white rot fungus *Pleurotus ostreatus*. *Applied*
21 *and Environmental Microbiology* 63: 2495–2501
- 22 Blaudez D, Botton B, Chalot M (2000) Cadmium uptake and subcellular
23 compartmentation in the ectomycorrhizal fungus *Paxillus involutus*. *Microbiology*
24 146: 1109-1117

- 1 Braha B, Tintemann H, Krauss G, Ehrman J, Bärlocher F, Krauss G-J (2007) Stress
2 response in two strains of the aquatic hyphomycete *Heliscus lugdunensis* after
3 exposure to cadmium and copper ions. *Biometals* 20: 3-105
- 4 Capotorti G, Digianvencenzo P, Cesti P, Bernardi A, Guglielmetti G (2004) Pyrene and
5 benzo(a)pyrene metabolism by an *Aspergillus terreus* strain isolated from a
6 polycyclic aromatic hydrocarbons polluted soil. *Biodegradation* 15: 79-85
- 7 Chaudry R (1994) *Biological Degradation and Bioremediation of Toxic Chemicals*.
8 Chapman and Hall, London
- 9 Cuny P, Acquaviva M, Gilewicz M (2004) Phenanthrene degradation, emulsification
10 and surface tension activities of a *Pseudomonas putida* strain isolated from a
11 coastal oil contaminated microbial mat. *Ophelia* 58: 283-287
- 12 D'Annibale A, Rosetto F, Leonardi V, Federici F, Petruccioli M (2006) Role of
13 Autochthonous Filamentous Fungi in Bioremediation of a Soil Historically
14 Contaminated with Aromatic Hydrocarbons. *Applied and Environmental*
15 *Microbiology* 72: 28–36
- 16 Duarte S, Pascoal C, Alves A, Correia A, Cássio F (2008a) Copper and zinc mixtures
17 induce shifts in microbial communities and reduce leaf litter decomposition in
18 streams. *Freshwater Biology* 53: 91-101
- 19 Duarte S, Pascoal C, Cássio F (2004) Effects of zinc on leaf decomposition by fungi in
20 streams: studies in microcosms. *Microbial Ecology* 48: 366-374
- 21 Duarte S, Pascoal C, Cássio F (2008b) High diversity of fungi may mitigate the impact
22 of pollution on plant litter decomposition in streams. *Microbial Ecology* 56: 688-
23 695
- 24 Duarte S, Pascoal C, Cássio F (2009) Functional stability of stream-dwelling microbial
25 decomposers to copper and zinc stress. *Freshwater Biology* 54: 1683-1691

- 1 Fernandes (2008) Effects of fungal diversity and cadmium on leaf litter decomposition
2 in streams: studies in microcosms. M.Sc. Thesis, University of Minho, Braga,
3 Portugal
- 4 Gadd G (1993) Interactions of fungi with toxic metals. *New Phytologist* 124: 25-60
- 5 Gessner M, Gulis V, Kuehn K, Chauvet E, Suberkropp K (2007) Fungal decomposers
6 of plant litter in aquatic ecosystems. *In* Kubicek C, Druzhinina I (2nd ed.) *The*
7 *Mycota: environmental and microbial relationships* (2nd ed.) Springer, Berlin, pp.
8 301-321
- 9 Guimarães-Soares L, Pascoal C, Cássio F (2007) Effects of heavy metals on the
10 production of thiol compounds by the aquatic fungi *Fontanospora fusiramosa* and
11 *Flagellospora curta*. *Ecotoxicology and Environmental Safety* 66: 36-43
- 12 Gonçalves MAP (2001) Determinação de metais pesados em águas superficiais
13 recolhidas no rio Este. M.Sc. thesis, University of Minho, Braga, Portugal
- 14 Gust K, Fleeger J (2006) Exposure to cadmium-phenanthrene mixtures elicits complex
15 toxic responses in the freshwater tubificid oligochaete, *Ilyodrilus templetoni*.
16 *Archives of environmental contamination and toxicology* 51: 54-60
- 17 Jaeckel P, Krauss G-K, Krauss G (2005) Cadmium and zinc response of the fungi
18 *Heliscus lugdunensis* and *Verticillium cf. alboatrum* isolated from highly polluted
19 water. *Science of the Total Environment* 346: 274-279
- 20 Johnsen A, Wick Y, Harms H (2005) Principles of microbial PAH-degradation in soil.
21 *Environmental Pollution* 133: 71-84.
- 22 Juhasz A, Naidu R (2000) Bioremediation of high molecular weight polycyclic aromatic
23 hydrocarbons: a review of the microbial degradation of benzo[a]pyrene.
24 *International Biodeterioration & Biodegradation* 45: 57-88

- 1 Juhasz A, Stanley G, Britz M (2000) Degradation of PAHs in contaminated soil by a
2 bacterial consortium: effects on microtox and mutagenicity bioassays.
3 Biorremediation Journal 4: 271-283
- 4 Legendre P, Legendre L (1998) Numerical ecology (2nd ed.) Elsevier Science BV.
5 Amsterdam. 853 pp
- 6 Maliszewska-Kordybach B, Smreczak B (2003) Habitat function of agricultural soils as
7 affected by heavy metals and polycyclic aromatic hydrocarbons contamination.
8 Environment International 28: 719–728
- 9 Miersch J, Bärlocher F, Bruns I (1997) Effects of cadmium, copper and zinc on growth
10 and thiol content of aquatic hyphomycetes. Hydrobiology 346: 77-84
- 11 Mille-Lindblom C, Tranvik LJ (2003) Antagonism between bacteria and fungi on
12 decomposing aquatic plant litter. Microbial Ecology 45: 173-182
- 13 Moreau C, Klerks P, Haas C (1999) Interaction between phenanthrene and zinc in their
14 toxicity to the sheepshead minnow (*Cyprinodon variegatus*). Environmental
15 Contamination Toxicology 37: 251-257
- 16 Niyogi D, Lewis W, McKnight D (2001) Litter breakdown in mountain streams affected
17 by mine drainage: biotic mediation of abiotic controls. Ecological Applications
18 11: 506-516
- 19 Niyogi D, McKnight D, Lewis W (2002) Fungal communities and biomass in mountain
20 streams affected by mine drainage. Archiv für Hydrobiologie 155: 255–271
- 21 Pascoal C, Cássio F (2004) Contribution of fungi and bacteria to leaf litter
22 decomposition in a polluted river. Applied and Environmental Microbiology 70:
23 5266-5273

- 1 Pascoal C, Cássio F, Marcotegui A, Sanz B, Gomes P (2005a) Role of fungi, bacteria,
2 and invertebrates in leaf litter breakdown in a polluted river. *Journal of the North*
3 *American Benthological Society* 24: 784-797
- 4 Pascoal C, Cássio F, Marvanová L (2005b) Anthropogenic stress may affect aquatic
5 hyphomycete diversity more than leaf decomposition in a low order stream.
6 *Archiv für Hydrobiologie* 162: 481-496
- 7 Rand G, Wells P, McCarty L (1995) Introduction to aquatic toxicology: effects,
8 environmental fate, and risk assessment. In: *Fundamentals of Aquatic Toxicology*
9 (Ed. Rand G.M.) Taylor & Francis, London. pp. 3–66
- 10 Ribeiro C, Vollaire Y, Sanchez-Chardi A, Roche H (2005) Bioaccumulations and the
11 effects of organochlorine pesticides, PAH and heavy metals in the Eel (*Anguilla*
12 *anguilla*) at the Camargue Natural Reserve, France. *Aquatic Toxicology* 74: 53-69
- 13 Scoggins M, McClintock N, Gosselink L, Bryer P (2007) Occurrence of polycyclic
14 aromatic hydrocarbons below coal-tar-sealed parking lots and effects on stream
15 benthic macroinvertebrate communities. *Journal of the North American*
16 *Benthological Society* 26: 694-707
- 17 Shen G, Cao L, Lu Y, Hong J (2005) Influence of phenanthrene on cadmium toxicity to
18 soil enzymes and microbial growth. *Environmental Science and Pollutant*
19 *Resource* 12 (5): 259 – 263
- 20 Silva S, Carman Kevin, Fleeger J, Marshall T, Marlborough S (2009) Effects of
21 phenanthrene and metal-contaminated sediment on the feeding activity of the
22 harpacticoid copepod, *Schizopera knabeni*. *Archives of Environmental*
23 *Contamination and Toxicology* 56: 434-441

- 1 Soares HMVM, Boaventura RAR, Machado ASSC, Esteves da Silva JCG (1999)
2 Sediments as monitors of heavy metal contamination in the Ave river basin
3 (Portugal): multivariate analysis of data. *Environmental Pollution* 105: 311-323
- 4 Sridhar K, Bärlocher F, Krauss G-J, Krauss G (2005) Response of aquatic hyphomycete
5 communities to changes in heavy metal exposure. *International Review of*
6 *Hydrobiology* 90: 21–32
- 7 Sridhar K, Krauss G, Bärlocher F, Raviraja N, Wennrich R, Baumbach R, Krauss G-J
8 (2001) Decomposition of alder leaves in two heavy metal-polluted streams in
9 central Germany. *Aquatic Microbial Ecology* 26: 73-80
- 10 Watson G, Andersen O, Galloway T, Depledge M (2004) Rapid assessment of
11 polycyclic aromatic hydrocarbon (PAH) exposure in decapod crustaceans by
12 fluorimetric analysis of urine and haemolymph. *Aquatic toxicology* 67: 127-142
- 13 White T, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
14 ribosomal RNA genes for phylogenetics. In: *PCR Protocols: a Guide to Methods*
15 *and Applications* (Eds. Innis M, Gelfand D, Sninsky J, White T). Academic Press
16 Inc, New York, pp. 315-322
- 17 Yang X, Peterson D, Baumann P, Lin E (2003) Fish biliary PAH metabolites estimated
18 by fixed-wavelength fluorescence as an indicator of environmental exposure and
19 effects. *Journal of great lakes research* 29: 116-123
- 20 Zar J (1996) *Biostatistical Analysis* (3rd ed.) Prentice Hall. Upper Saddle River, NJ
21 USA, 662 pp
22
23

Table 1. Mean percentage contribution of individual fungal species on decomposing leaves to overall conidia produced after 10 and 14 days of exposure to Cd and Phe alone or in mixtures. Treatments, Cd1: 0.06 mg L⁻¹, Cd2: 0.6 mg L⁻¹, Cd3: 1.2 mg L⁻¹, Cd4: 3.6 mg L⁻¹, Cd5: 4.5 mg L⁻¹, Phe: 0.2 mg L⁻¹, Ct: control. Abb, species abbreviation.

Abb	Species	% of conidia											
		Ct	Cd1	Cd2	Cd3	Cd4	Cd5	Phe	Cd1Phe	Cd2Phe	Cd3Phe	Cd4Phe	Cd5Phe
AA	<i>Alatospora acuminata</i> Ingold	1.8	1.7	3.6	2.7	10.4	1.0	13.0	-	0.1	0.5	-	-
AP	<i>Alatospora pulchella</i> Marvanová	20.4	34.4	63.4	57.7	33.0	31.8	29.8	41.5	72.8	88.6	39.8	54.2
AF	<i>Anguillospora filiformis</i> Greath.	0.5	0.1	-	-	-	-	-	-	-	-	-	-
AT	<i>Articulospora tetracladia</i> Ingold	40.0	19.8	14.7	1.4	4.1	17.7	7.7	15.1	2.1	0.7	-	6.3
CA	<i>Clavariopsis aquatica</i> De Wild.	2.0	0.1	<0.1	0.5	-	0.2	-	-	-	-	-	-
CL	<i>Clavatospora longibrachiata</i> (Ingold) Marvanová & Sv. Nilsson	15.9	36.6	2.8	0.1	0.6	-	-	-	-	-	-	-
DF	<i>Dimorphospora foliicola</i> Tubaki	-	-	-	-	-	-	0.8	0.1	-	-	-	-
LA	<i>Lemonniera aquatica</i> De Wild.	0.9	0.9	1.7	5.8	28.5	41.3	-	1.5	7.1	2.1	22.2	4.2
LC	<i>Lunulospora curvula</i> Ingold	0.2	0.3	<0.1	-	-	-	-	-	-	-	-	-
TE	<i>Tetrachaetum elegans</i> Ingold	6.6	4.1	12.2	27.5	12.8	-	33.7	38.5	16.3	5.4	2.3	4.2
TB	<i>Tetracladium breve</i> A. Roldán	0.1	0.1	0.4	0.1	-	-	0.6	-	-	-	-	-
TSt	<i>Tetracladium setigerum</i> (Grove) Ingold	-	-	-	-	-	0.2	0.8	-	-	-	-	-
TSp	<i>Tricladium splendens</i> Ingold	1.3	1.0	0.5	0.6	1.6	0.5	2.2	0.8	0.1	1.3	4.0	-
TA	<i>Triscelophorus cf. acuminatus</i> Nawawi	0.3	-	-	-	-	-	-	-	-	-	-	-
Un	Unknown branched species (50-30 µm)	0.4	0.2	0.7	-	-	7.5	0.1	2.4	1.1	1.5	31.8	31.3
S1	Sigmoid 1 (80-2 µm)	0.3	0.1	-	0.1	0.2	-	-	-	-	-	-	-
S2	Sigmoid 2 (50-1.5 µm)	0.2	0.1	-	-	-	-	-	-	-	-	-	-
S3	Sigmoid 3 (20-4 µm)	9.4	0.1	-	3.7	9.0	-	11.4	0.2	0.2	-	-	-
S4	Sigmoid 4 (110-1.5 µm)	<0.1	0.2	0.1	-	-	-	-	-	0.1	-	-	-
	N° of conidial morphotypes	17	16	12	11	9	8	10	8	9	7	5	5
	N° of DGGE OTUs	18	19	16	14	14	13	13	12	12	10	10	10

Figure legends

Figure 1. Effects of Cd and / or Phe on leaf mass loss (A), fungal biomass (B) and fungal sporulation (C). Leaf mass loss and fungal biomass were measured after 14 days of exposure, while fungal sporulation was measured after 10 and 14 days. n=3, error bars indicate ± 1 SEM; *: $p < 0.05$. Treatments, Cd1: 0.06 mg L^{-1} , Cd2: 0.6 mg L^{-1} , Cd3: 1.2 mg L^{-1} , Cd4: 3.6 mg L^{-1} , Cd5: 4.5 mg L^{-1} , Phe: 0.2 mg L^{-1} . Control: $0,061 \pm 0,009 \text{ g}$

Figure 2. DGGE fingerprints of the ITS2 region of rDNA of fungal communities on decomposing leaves. M, mixture of DNA from pure cultures. AT, *Articulospora tetracladia*; VE, *Varicosporium elodeae*; AA, *Alatospora acuminata*, LA, *Lemonniera aquatica* and TC, *Tricladium chaetocladium*. Treatments, Cd1: 0.06 mg L^{-1} , Cd2: 0.6 mg L^{-1} , Cd3: 1.2 mg L^{-1} , Cd4: 3.6 mg L^{-1} , Cd5: 4.5 mg L^{-1} , Phe: 0.2 mg L^{-1} .

Figure 3. Correspondence Analysis (CA) diagrams for ordination of fungal communities based on the 19 sporulating species (A) and on the 24 DGGE OTUs (B) exposed to Cd and/or Phe for 14 days in microcosms. See Table 1 for species names and abbreviations in A. Treatments, Cd1: 0.06 mg L^{-1} , Cd2: 0.6 mg L^{-1} , Cd3: 1.2 mg L^{-1} , Cd4: 3.6 mg L^{-1} , Cd5: 4.5 mg L^{-1} , Phe: 0.2 mg L^{-1}