

# Phytochemistry Reviews

## Hypericum sp.: Essential oil composition and biological activities

--Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Full Title:</b>	Hypericum sp.: Essential oil composition and biological activities
<b>Article Type:</b>	Review Article
<b>Keywords:</b>	Antibacterial, antifungal, antiviral, antiangiogenic, antitumor activities; biotechnology
<b>Corresponding Author:</b>	Manuel Fernandes-Ferreira, Ph.D. CITAB - Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Department of Biology, Faculty of Sciences, University of Porto Porto, PORTUGAL
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	CITAB - Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Department of Biology, Faculty of Sciences, University of Porto
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Ana Patrícia Guedes, Ph.D.
<b>First Author Secondary Information:</b>	
<b>All Authors:</b>	Ana Patrícia Guedes, Ph.D. Franklin Gregory, Ph.D. Manuel Fernandes-Ferreira, Ph.D.
<b>All Authors Secondary Information:</b>	
<b>Abstract:</b>	Phytochemical characterization of several species of Hypericum genus has been studied for a long time. Several reviews, most of them on <i>H. perforatum</i> , have already been published concerning the characterization of alcoholic and water extracts as well as their biological activities. Studies on the essential oils of <i>H. perforatum</i> and other species of this genus have already been published, some of them reporting the positive biological activities of these essential oils. Additionally, variations on the essential oils of Hypericum species induced by seasonal variation, geographic distribution, phenological cycle and type of the organ in which essential oils are produced and/or accumulated have also been reported. However, so far, no review paper has been published gathering all the reported data on Hypericum essential oils and respective biological activities. Thus in this chapter we collect and summarize as many information as possible concerning composition and biological activities of essential oils and essential oil containing crude extracts of Hypericum species, as well the biotechnology approaches envisaging their improvement.

## ***Hypericum* sp.: Essential oil composition and biological activities**

Ana P. Guedes<sup>1,2</sup>, Franklin Gregory<sup>1,2</sup> and Manuel Fernandes-Ferreira<sup>1,3\*</sup>

<sup>1</sup>*CITAB - Centre for the Research and Technology of Agro-Environmental and Biological Sciences,*

<sup>2</sup>*Department of Biology, University of Minho, 4710-057 Braga, Portugal*

<sup>3</sup>*Department of Biology, Faculty of Science, University of Porto, FC4, 4169-007 Porto, Portugal.*

### **\* To whom correspondence should be addressed:**

*Manuel Fernandes Ferreira*

*Department of Biology, Faculty of Science, University of Porto, FC4, 4169-007 Porto, Portugal*

Phone: 00351 220402732 / 00351 220402000; fax: 00351 220402 009

Email: [manuel.ferreira@fc.up.pt](mailto:manuel.ferreira@fc.up.pt)

### **Abstract**

Phytochemical characterization of several species of *Hypericum* genus has been studied for a long time. Several reviews, most of them on *H. perforatum*, have already been published concerning the characterization of alcoholic and water extracts as well as their biological activities. Studies on the essential oils of *H. perforatum* and other species of this genus have already been published, some of them reporting the positive biological activities of these essential oils. Additionally, variations on the essential oils of *Hypericum* species induced by seasonal variation, geographic distribution, phenological cycle and type of the organ in which essential oils are produced and/or accumulated have also been reported. However, so far, no review paper has been published gathering all the reported data on *Hypericum* essential oils and respective biological activities. Thus in this chapter we collect and summarize as many information as possible concerning composition and biological activities of essential oils and essential oil containing crude extracts of *Hypericum* species, as well the biotechnology approaches envisaging their improvement.

## Key Words

Antibacterial, antifungal, antiviral, antiangiogenic, antitumor activities; biotechnology

## Introduction

*Hypericum* is an important genus of the family Clusiaceae (Guttiferae), which includes about 450 species of trees, shrubs and herbs. Some of the general morphological characteristics of this genus are: (i) opposite simple entire exstipulated leaves; (ii) (4-)5-merous flowers consisting of green sepals and free yellow petals (occasionally, the sepals as well as the petals can be red-tinged); (iii) free of more or less united ovary and; (iv) stamens organized in 5 fascicles. The main feature of some of the important species is the presence of translucent and often black or red schizogenous hypericin glands. Although, these glands are present in the plant parts like stem, flowers and leaves, their distribution differ between species. Hypericin was first identified in *Hypericum* species and serves as a chemotaxonomic marker of the genera. For more details about the taxonomy and distribution of this genus, refer Robson (2006) (Robson, 2006).

Many species of the *Hypericum* genus have a long traditional value as medicinal plants. Ancient Greeks knew about the medicinal properties of this genus. Greek herbalist Dioscorides recommends four species of *Hypericum* (Upericon, Askuron, Androsaimon, and Koris) for sciatica and burns. He referred that *H. crispum* and *H. barbatum* have antidiuretic and antimalarial properties. Theophrastus recommends *H. lanuginosum* for external application, while Pliny says it should be taken in wine against poisonous reptiles. *H. coris*, another Greek species, was mentioned by Hippocrates and Pliny. Yet another species, *H. perforatum* (St. John's wort) has been appreciated for its medicinal value since at least as early as the 1st century A.D. and was known by Hippocrates, Pliny, and Dioscorides who included it in the *De materia medica* (Bombardelli *et al.*, 1995; Upton, 1997)

Currently, several species of this genus has been used in ailments as knowledge-based medicine in many countries. In Portuguese folk medicine, some *Hypericum* species are used the major of them being *H. androsaemum*, *H. perforatum*, and *H. undulatum*. *H. androsaemum* (Hipericão do Gerês) is known for its diuretic effect and infusions are used for kidney and bladder ailments, whereas *H. undulatum* extract is used for liver problems. Flowers decoction of *H. undulatum* is also traditionally used for migraine, bladder ailments; and as renal antispasmodic and hepatic protector (Ferreira *et al.*, 2006). Moreover, decoctions of the flowering aerial parts of *H.*

*perforatum* have been used to treat liver troubles (Camejo-Rodrigues *et al.*, 2003), depression, liver disorders and rheumatism (Nogueira *et al.*, 1998). In the Canary Islands (Spain), infusions prepared from the flowers, leaves and fruits of various species of the genus *Hypericum* have been used as a vermifuge, diuretic, as well as a wound healing, sedative, antihysterical and antidepressive agent (Prado *et al.*, 2002; Rabanal *et al.*, 2002). In Italy, leaves and stems of *H. hircinum* have been widely used against cough (Pieroni *et al.*, 2004). *H. perforatum*, a well-known species of this genus, has been included in the traditional pharmacopeia of many countries. For instance, in the 1997 German List of Prescription Drugs described more than 50 preparations from *H. perforatum*. In the German Drug Prescription Report 1996, about 131 million of daily doses of products prepared from this species were prescribed. In Serbia, aerial parts of *H. perforatum* were cited to be used internally and externally for all ailments (Jaric *et al.*, 2007). Different parts of the plant and different preparations are externally applied or orally administrated for several ailments, including digestive, urinary, respiratory and cardiac diseases in Turkey (Kültür, 2007). In Bulgarian phytotherapy, aerial parts of *H. perforatum* have been recommended for treatment of gastric and duodenal ulcer; as regenerative and anti-inflammatory agent in digestive tract diseases and as epithelotonic agent (Ivanova *et al.*, 2005). In Italy, leaves and stems of *H. hircinum* have been widely used against cough (Pieroni *et al.*, 2004).

In the last decade, several pharmacological studies have been performed using crude extracts to evaluate the traditional knowledge. Results of those studies have revealed that extract of *Hypericum* exert several pharmacological properties including antidepressant, antimicrobial and wound healing effects (Ishiguro *et al.*, 1990; Decosterd *et al.*, 1991; Jayasuriyab *et al.*, 1991; Rocha *et al.*, 1995; Öztürk *et al.*, 1996; Sokmen *et al.*, 1999; Daudt *et al.*, 2000; Mukherjee *et al.*, 2000; Pistelli *et al.*, 2000; Avato *et al.*, 2004).

Plants produce and accumulate different types of compounds. Molecules with an important role in basic life functions such as cell division, growth, respiration, storage, and reproduction were described as primary metabolites (Bougard *et al.*, 2001). Several other compounds which are not essential for the above functions are known as secondary metabolites. For a long time, secondary metabolites have been considered as waste by-products of plant metabolism. Recent improvement of biochemical techniques and the rise of molecular biology have shown that these molecules play a major role in the adaptation of plants to their environment including biotic and abiotic challenges. They also act on animals (anti-feeding properties), pathogens (phytoalexins) and other plants (allelopathy) (Bougard *et al.*, 2001). Plant secondary metabolites include a vast array of compounds that to date sum up to more than 200,000 defined structures.

Studies on medicinal plants are rapidly increasing because of the search for new active molecules, and for the improvement in the production of plants or molecules for the herbal pharmaceutical industries (Poutaraud *et al.*, 2005). As a genus with several medicinal plant species, *Hypericum* has recently drawn the attention of phytochemists and pharmacologists. However, it is clear from the literature that *H. perforatum* is the only most characterized species of this genus in terms of phytochemistry and pharmacology. Under the stimulus of great scientific interest and economic value acquired by *H. perforatum*, studies with other plants of *Hypericum* genus have also been carried out revealing their antidepressant, analgesic, anti-inflammatory, antioxidant, antimicrobial and wound healing properties (Öztürk *et al.*, 1996; Apaydin *et al.*, 1999; Daudt *et al.*, 2000; Mukherjee *et al.*, 2001; Trovato *et al.*, 2001; Öztürk *et al.*, 2002; Prado *et al.*, 2002; Rabanal *et al.*, 2002; Sánchez-Mateo *et al.*, 2002; Cakir *et al.*, 2003; Couladis *et al.*, 2003; Heilmann *et al.*, 2003; Rabanal *et al.*, 2005).

Phytochemical characterisation of several species of this genus have revealed the presence of a variety of compounds, but not limited to phenylpropanes, flavonol derivatives, biflavones, proanthocyanidines, xanthenes, phloroglucinols and naphthodianthrones (for review see (Bombardelli *et al.*, 1995; Nahrstedt *et al.*, 1997; Hölzl *et al.*, 2003)) and essential oils. In spite of the large size of *Hypericum* genus, the composition of volatile compounds is known in only a small number of species with the exception of *H. perforatum* (Saroglou *et al.*, 2007). The present review summarizes information available on the chemical composition of essential oils isolated from *Hypericum* genus. Additionally, the biological activities of this genus with special emphasis on essential oil composition are also discussed.

## **Chemical Composition of *Hypericum* Essential Oils**

Interest in essential oils has revived in recent decades, with the popularity of aromatherapy, a branch of alternative medicine which claims that the specific aromas carried by essential oils have curative effects. Oils are volatilized or diluted in a carrier oil and used in massage or burned as incense. About 300 essential oils out of 3000 known are commercially important mainly for their flavours and fragrances (Burt, 2004).

Essential oils are aromatic oily extracts obtained from the plants. Composition of essential oils can differ between different parts of the plants. Steam distillation is the most commonly used method of extraction for commercial production of essential oils (Burt, 2004), although other extraction methods such as expression, fermentation, enfleurage or extraction are also in practice. In

the steam distillation, the water is heated and the steam passes through the plant material, thus vaporizing the volatile compounds. The vapors flow through a coil where they condense back to liquid, which is then collected in the receiving vessel. Because of the mode of extraction, mostly by distillation from aromatic plants, the extract contains a variety of volatile molecules such as terpenes and terpenoids, phenol-derived aromatic components and aliphatic components (Bakkali *et al.*, 2008).

Essential oils are very complex natural mixtures which can contain about 20–60 components at quite different concentrations. Hence, they are generally characterized by two or three major components at fairly high concentrations compared to other components that are present in trace amount (Bakkali *et al.*, 2008).

The main essential oil constituents identified in *Hypericum* species are summarised in Table 1. Characterization of essential oils from species of *Hypericum* genus revealed the presence of monoterpenoid and sesquiterpenoid compounds, as well as alkanes and aldehydes as the main compounds in the most of them (Mathis *et al.*, 1964a; Mathis *et al.*, 1964b; Mathis *et al.*, 1964c). In the essential oils of *H. ericoides*,  $\alpha$ -curcumene,  $\alpha$ -pinene,  $\gamma$ -muurolene and  $\delta$ -cadinene were the most represented compounds (Cardona *et al.*, 1983).

In general, species of this genus have higher amount of sesquiterpenoids in their essential oils (Table 1). Indeed, *E*-caryophyllene was one of the most represented compounds in most of the studied *Hypericum* essential oils. Germacrene D was another major sesquiterpene hydrocarbon common to several species. Caryophyllene oxide, spathulenol and globulol, are the oxygenated-sesquiterpenes present in the top three most represented compounds of some *Hypericum* species. Likewise,  $\alpha$ -pinene and  $\beta$ -pinene were the two mostly represented monoterpene hydrocarbons in essential oils from *Hypericum*. Interestingly, in *H. barbatum* the three most represented compounds are monoterpene hydrocarbons ( $\alpha$ -pinene,  $\beta$ -pinene and limonene) (Saroglou *et al.*, 2007). *n*-Alkanes were found as the major compounds in few species of the genus. In *H. foliosium*, *H. hirsutum*, *H. myrianthum*, *H. richeri* and *H. triquetrifolium*, *n*-nonane is among the three most represented compounds. Some of these species also had high proportions of *n*-undecane (Santos *et al.*, 1999; Bertoli *et al.*, 2003; Ferraz *et al.*, 2005; Ferretti *et al.*, 2005; Saroglou *et al.*, 2007).

**Table 1 – Main essential oil constituents of *Hypericum* species**

Species	Origin	Main constituents	Reference
<i>H. alpinum</i>	Serbia	$\beta$ -Pinene (13.3%)	(Saroglou <i>et al.</i> , 2007)

Species	Origin	Main constituents	Reference
		$\gamma$ -Terpinene (7.7%) (-)-( <i>E</i> )-Caryophyllene (6.5%)	
<i>H. acmosepalum</i>	China	$\beta$ -Selinene (16.3%) <i>ar</i> -Curcumene (12.6%)	(Demirci <i>et al.</i> , 2005)
<i>H. aegypticum</i> <i>ssp. aegypticum</i>	Libya	Ishwarane (14.4%) Eudesm-11-en-4-ol stereoisomer (9.6%) Eudesm-11-en-4-ol stereoisomer (10.7%)	(Crockett <i>et al.</i> , 2007)
<i>H. aegypticum</i> <i>ssp. marrocanum</i>	Northwestern Africa	Caryophyllene oxide (29.2%) $\beta$ -Caryophyllene (15.1%) Caryophylladienol-II (9.7%)	(Crockett <i>et al.</i> , 2007)
<i>H. androsaemum</i>	Portugal	C <sub>15</sub> H <sub>24</sub> (27.6%) Germacrene D (12.3%) $\beta$ -Caryophyllene (14.0%)	(Nogueira <i>et al.</i> , 1998)
	Portugal	( <i>E</i> )-Caryophyllene (9.4- 15.1%) $\gamma$ -Elemene (8.0-17.9%) $\beta$ -Gurjunene (6.1-15.5%)	(Guedes <i>et al.</i> , 2003)
	Portugal	( <i>E</i> )-Caryophyllene (9.0- 17.0%) $\gamma$ -Elemene (9.3-17.3%) $\beta$ -Gurjunene (7.9-14.8%)	(Guedes <i>et al.</i> , 2004)
<i>H. balearicum</i>	Balearic Islands	$\alpha$ -Pinene (28.5%) $\beta$ -Pinene (20.4%) $\beta$ -Eudesmol (11.2%)	(Crockett <i>et al.</i> , 2007)
<i>H. barbatum</i>	Serbia	(-)- $\alpha$ -Pinene (17.1%) (-)- $\beta$ -Pinene (17.0%) (-)-Limonene (6.0%)	(Saroglou <i>et al.</i> , 2007)
<i>H. beanii</i>	China	Caryophyllene oxide (18.7%) $\beta$ -Selinene (16.3%) $\gamma$ -Muurolene (11.3%)	(Demirci <i>et al.</i> , 2005)
<i>H. brasiliense</i>	Brazil	$\beta$ -Caryophyllene (29.5%)	(Abreu <i>et al.</i> , 2004)

Species	Origin	Main constituents	Reference
		$\alpha$ -Humulene (12.7%) Caryophyllene oxide (9.9%)	
<i>H. bupleuroides</i>	Turkey	$\beta$ -Sesquiphellandrene (33.2%) $\beta$ -Caryophyllene (20.2%) Selina-3,7(11)-diene (7.0%)	(Demirci <i>et al.</i> , 2006)
<i>H. calcynum</i>	China	$\beta$ -Pinene (29.2%) $\alpha$ -Terpineol (11.5%)	(Demirci <i>et al.</i> , 2005)
<i>H. carinatum</i>	Brazil	$\beta$ -Caryophyllene (21.0%) $\alpha$ - <i>trans</i> -Bergamotene (10.0%) Caryophyllene oxide (9.5%)	(Ferraz <i>et al.</i> , 2005)
<i>H. choisyanum</i>	China	Cis-eudesma-6,11-diene (11.4%)	(Demirci <i>et al.</i> , 2005)
<i>H. connatum</i>	Brazil	Caryophyllene oxide (40.1%) $\beta$ -Caryophyllene (13.1%) Humulene oxide II (10.5%)	(Ferraz <i>et al.</i> , 2005)
<i>H. coris</i>	France	$\alpha$ -Curcumene (40.1%) $\gamma$ -Cadinene (14.7%) $\delta$ -Cadinene (6.6%)	(Schwob <i>et al.</i> , 2002)
<i>H. delphicum</i>	Arabian peninsula	Caryophyllene oxide (31.5%) $\beta$ -Caryophyllene (18.2%) <i>n</i> -Undecane (17.5%)	(Crockett <i>et al.</i> , 2007)
<i>H. foliosum</i>	Portugal	<i>n</i> -Nonane (28.7-72.6%) Limonene (6.9-45.8%) Terpinolene (0.5-18.8%) $\beta$ -Caryophyllene (1.1-6.9%) $\beta$ -Pinene (0.3-6.3%)	(Santos <i>et al.</i> , 1999)
<i>H. forrestii</i>	China	Caryophyllene oxide (12.7%) $\alpha$ -Pinene (10.4%)	(Demirci <i>et al.</i> , 2005)
<i>H. heterophyllum</i>	Turkey	Isocaryophyllene (17.1%) $\alpha$ -Pinene (11.6%)	(Cakir <i>et al.</i> , 2004)



Species	Origin	Main constituents	Reference
		$\delta$ -Cadinene (9.5%)	
<i>H. hirsutum</i>	Serbia	Nonane (24.8%) Undecane (13.3%) (-)-( <i>E</i> )-Caryophyllene (5.4%)	(Saroglou <i>et al.</i> , 2007)
	Serbia	<i>n</i> -Undecane (32.2%) Patchoulene (11.8%) Caryophyllene oxide (9.3%)	(Gudžic <i>et al.</i> , 2007)
<i>H. humifusum</i>	Portugal	$\alpha$ -Pinene (80.6%) $\beta$ -Pinene (4.7%) Germacrene D (2.1%)	(Nogueira <i>et al.</i> , 1998)
	Portugal	$\alpha$ -Pinene (44.7-77.2%) $\beta$ -Pinene (4.7-7.7%) $\beta$ -Caryophyllene (1.2-9.3%) Germacrene D (1.9-6.1%)	(Nogueira <i>et al.</i> , 2008)
<i>H. kouytchense</i>	China	<i>cis</i> - $\beta$ -Guaiene (10.7%) $\gamma$ -Muuroolene (12.4%)	(Demirci <i>et al.</i> , 2005)
<i>H. lancasteri</i>	China	$\beta$ -Selinene (11.4%) Eudesmadienone (10.8%)	(Demirci <i>et al.</i> , 2005)
<i>H. leschenaultii</i>	China	Cuparene (24.8%) $\gamma$ -Muuroolene (16.8%)	(Demirci <i>et al.</i> , 2005)
<i>H. linarifolium</i>	Portugal	$\alpha$ -Pinene (31.1%) $\beta$ -Caryophyllene (11.6%) <i>n</i> -Undecane (7.0%)	(Nogueira <i>et al.</i> , 1998)
	Portugal	$\alpha$ -Pinene (19.9-31.2%) $\beta$ -Pinene (5.0-11.0%) $\beta$ -Caryophyllene (6.6-11.6%)	(Nogueira <i>et al.</i> , 2008)
<i>H. linarioides</i>	Turkey	$\delta$ -Cadinene (6.9%) $\gamma$ -Muuroolene (5.5%) ( <i>Z</i> )- $\beta$ -Farnesene (5.2%)	(Cakir <i>et al.</i> , 2005)
<i>H. lysimachioides</i>	South eastern Anatolia	Caryophyllene oxide (30.8%) $\alpha$ -Longifolene (6.4%)	(Toker <i>et al.</i> , 2006)

Species	Origin	Main constituents	Reference
		$\beta$ -Selinene (6.7%)	
<i>H. maculatum</i>	Serbia	Spathulenol (6.8%) Globulol (10.2%) Nonane (5.5%)	(Saroglou <i>et al.</i> , 2007)
<i>H. monogynum</i>	China	Tricosane (13.3%) Myrcene (10.4%)	(Demirci <i>et al.</i> , 2005)
<i>H. myrianthum</i>	Brazil	Undecane (20.7%) Nonane (17.5%) Dehydro-aromadendrene (8.6%)	(Ferraz <i>et al.</i> , 2005)
<i>H. olympicum</i>	Serbia	( <i>E</i> )-Anethole (30.7%) $\beta$ - Farnesene (12.4%) $\delta$ -Cadinene (8.7%)	(Gudžić <i>et al.</i> , 2001)
	Greece	Germacrene D (16.0%) ( <i>E</i> )-Caryophyllene (7.4%) Spathulenol (6.7%)	(Pavlović <i>et al.</i> , 2006)
<i>H. patulum</i>	China	$\beta$ -selinene (14.7%)	(Demirci <i>et al.</i> , 2005)
<i>H. polyanthemum</i>	Brazil	HP1 Benzopyrans (26.7%) HP2 Benzopyrans (13.2%) Undecane (7.9%)	(Ferraz <i>et al.</i> , 2005)
<i>H. pseudohenryi</i>	China	$\beta$ -Selinene (18.5%)	(Demirci <i>et al.</i> , 2005)
<i>H. pulchrum</i>	Portugal	$\alpha$ -Pinene (35.7-49.8%) $\beta$ -Pinene (9.0-12.5%) Germacrene D (2.4-5.4%)	(Nogueira <i>et al.</i> , 1998)
	Portugal	$\alpha$ -Pinene (49.8%) $\beta$ -Pinene (12.5%) Germacrene D (5.4%)	(Nogueira <i>et al.</i> , 2008)
<i>H. richeri</i>	Italy	( <i>Z</i> )- $\beta$ -Ocimene (19.5%) <i>n</i> -Nonane (13.8%) $\beta$ -Bisabolene (8.7%)	(Ferretti <i>et al.</i> , 2005)
<i>H. roeperanum</i>	East Africa	$\gamma$ -Curcumene (15.6%)	(Crockett <i>et al.</i> ,

Species	Origin	Main constituents	Reference
		(2 <i>E</i> ,6 <i>E</i> )-Farnesol (7.8%) <i>ar</i> -Curcumene (7.7%)	2007)
<i>H. scabrum</i>	Uzbekistan	$\alpha$ -Pinene (11.2%) Spathulenol (7.2%) <i>p</i> -Cymene (6.1%)	(Baser <i>et al.</i> , 2002)
<i>H. ternum</i>	Brazil	$\beta$ -Caryophyllene (12.0%) Bicyclogermacrene (10.0%) $\beta$ -Cadinene (5.0%)	(Ferraz <i>et al.</i> , 2005)
<i>H. tetrapterum</i>	Greece	$\alpha$ -Copaene (11.3%) $\alpha$ -Longipinene (9.7%) Caryophyllene oxide (8.9%)	(Pavlović <i>et al.</i> , 2006)
<i>H. tomentosum</i>	Portugal	$\beta$ -Caryophyllene (12.6%) <i>n</i> -Undecane (7.5%) $\alpha$ -Humulene (5.2%)	(Nogueira <i>et al.</i> , 1998)
<i>H. xmoserianum</i>	China	$\gamma$ -Muurolene (10.7%) $\delta$ -Cadinene (10.2%)	(Demirci <i>et al.</i> , 2005)
<i>H. tomentosum</i>	Tunisia	Menthone (17.0%) <i>n</i> -Octane (9.9%) $\beta$ -Caryophyllene (5.3%) $\alpha$ -Pinene (5.2%)	(Hosni <i>et al.</i> , 2008)
<i>H. triquetrifolium</i>	Italy	$\alpha$ -Humulene <i>cis</i> -Calamene $\delta$ -Cadinene $\alpha$ -Pinene (10.33%) Caryophyllene oxide (1.38%)	(Karim <i>et al.</i> , 2007)

Essential oils content and composition can be greatly affected by several parameters including seasonal variation (Guedes *et al.*, 2004), phenological cycle (Schwob *et al.*, 2004) and geographic distribution. We have summarised the most represented compounds of the essential oils of *H. perforatum*, *H. perfoliatum* and *H. hyssopifolium* based on their geographical distribution in Table 2.

**Table 2 - Variation in the essential oil composition of a few *Hypericum* species based on their geographical distribution**

Species	Origin	Main constituents of essential oil	Reference
<i>H. perforatum</i>	Serbia	<i>cis</i> -Caryophyllene (48.5%) $\beta$ - Farnesene (12.1%) 2-Methyl-dodecane (5.7%)	(Gudžić <i>et al.</i> , 1997)
	Serbia	$\beta$ -Caryophyllene (14.2%) 2-Methyl-octane (13.1%) 2-Methyl-decane (7.9%)	(Gudžić <i>et al.</i> , 2001)
	Serbia	$\alpha$ -Pinene (8.6%) ( <i>Z</i> )- $\beta$ - Farnesene (6.6%) Germacrene D (6.8%)	(Saroglou <i>et al.</i> , 2007)
	Portugal	Germacrene D (20.0%) $\beta$ -Caryophyllene (10.9%) 2-Methyl-octane (9.7%)	(Nogueira <i>et al.</i> , 1998)
	Uzbekistan	$\beta$ -Caryophyllene (11.7%) Caryophyllene oxide (6.3%) Spathulenol (6.0%)	(Baser <i>et al.</i> , 2002)
	Lithuania	Caryophyllene oxide (6.1-35.8%) Germacrene D (4.5- 31.5%) $\beta$ -Caryophyllene (5.1-19.1%)	(Mockutė <i>et al.</i> , 2003)
	Italy	2-Methyl-octane (21.1%) Germacrene D (17.6%) $\alpha$ -Pinene (15.8%)	(Pintore <i>et al.</i> , 2005)
	Greece	$\alpha$ -Pinene (21.0%) 2-Methyl-octane (12.6%) $\gamma$ -Muurolene (6.9%)	(Pavlović <i>et al.</i> , 2006)
	India	Ishwarane $\alpha$ -cuprenene	(Weyerstahl <i>et al.</i> , 1995)
<i>H. perforatum</i>	Portugal	$\alpha$ -Pinene (50.0%) <i>n</i> -Nonane (16.8%) <i>n</i> -Undecane (8.8%)	(Nogueira <i>et al.</i> , 1998)

Species	Origin	Main constituents of essential oil	Reference
	Portugal	$\alpha$ -Pinene (39.4-64.3%) <i>n</i> -Nonane (11.9-23.8%) $\beta$ -Pinene(1.9-3.2%)	(Nogueira <i>et al.</i> , 2008)
	Algeria	Thymol (22.1%) $\tau$ -Cadinol (18.5%) 4,5-Dimethyl-2-ethylphenol (13.0%)	(Touafek <i>et al.</i> , 2005)
	Greece	$\alpha$ -Pinene (48.6%) <i>n</i> -Nonane (8.5%) $\delta$ -Cadinene (4.6%)	(Couladis <i>et al.</i> , 2001)
	Greece	$\alpha$ -Pinene (34.2%) $\beta$ -Pinene (9.2%) $\delta$ -Cadinene (8.1%)	(Couladis <i>et al.</i> , 2001)
	Greece	$\alpha$ -Pinene (41.3%) $\beta$ -Pinene (6.5%) $\delta$ -Cadinene (6.2%) <i>n</i> -Nonane (6.1%)	(Petrakis <i>et al.</i> , 2005)
	Tunisia	$\alpha$ -Pinene (13.1%) <i>allo</i> -Aromadendrene (11.4%) Germacrene-D (10.6%)	(Hosni <i>et al.</i> , 2008)
<b><i>H. hyssopifolium</i></b>	South eastern Anatolia	Caryophyllene oxide (20.4%) Spathulenol (13.4%) Caryophyllene alcohol (9.0%)	(Toker <i>et al.</i> , 2006)
	Turkey	$\alpha$ -Pinene (57.3%) $\beta$ -Pinene (9.0%) Limonene (6.2%)	(Cakir <i>et al.</i> , 2004)
	France	Spathulenol (19.5 %) Tetradecanol (10.2%) Dodecanol (9.3%)	(Schwob <i>et al.</i> , 2006)

$\beta$ -Caryophyllene and Germacrene D were abundant in essential oils of *H. perforatum* from Portugal and Lithuania, but not in the one from Greece. Gudžić and co-workers (1997; 2001) reported  $\beta$ -Caryophyllene as the most represented compound in essential oils of *H. perforatum* from Rujan mountain and Vlasina, both in Serbia (Gudžić *et al.*, 1997; Gudžić *et al.*, 2001). However this sesquiterpene was not one of the three major compounds in the essential oils of this species in Barelic, another region of Serbia (Saroglou *et al.*, 2007).  $\alpha$ -Pinene predominance was registered in the essential oils from plants cultivated in Italy, Greece and Serbia (Barelic). Essential oils from Lithuania and Uzbekistan were the only ones in which oxygenated-sesquiterpenes were the major compounds, caryophyllene oxide and spathulenol for Uzbekistan and caryophyllene oxide for Lithuania. In the essential oils of *Hypericum perforatum* L. from North Indian, ishwarane and  $\alpha$ -cuprenene were identified (Weyerstahl *et al.*, 1995). In essential oils of *H. perforatum* from Portugal, Greece and Tunisia some homogeneity was found in the major compounds. In all of them  $\alpha$ -pinene was one of the major compounds, with *n*-nonane and  $\beta$ -pinene common to some of them. However, data on essential oils from plants cultivated in Algeria show a different composition regarding the three most represented compounds, without any mono-or sesquiterpene hydrocarbon between them. Essential oils of *H. hyssopifolium* from south eastern Anatolia and France shared the predominance of the oxygenated-sesquiterpene spathulenol. However, in Turkey the three most represented compounds in this species were monoterpene hydrocarbons, with  $\alpha$ -pinene as the main constituent. Differences in the essential oils profile of plants cultivated in different locations can be attributed to different climatic and pedological conditions.

As shown in Table 3, *Hypericum* essential oil profile could also be dependent on the plant organs from which they are extracted. For example, essential oils extracted from flowers and leaves of *H. triquetrifolium* vary greatly in their composition. Flowers have a high representation of monoterpene hydrocarbons, whereas they are absent in the essential oils of leaves. On the other hand, although the essential oils composition of leaves and flowers were similar, the main constituent, caryophyllene oxide, varied in the concentration in both *H. androsaemum* and *H. perforatum*. Similarly, leaves and flowers of *H. caprifoliatum* varied in the nonane and  $\beta$ -caryophyllene concentrations.

**Table 3 - Variation in the essential oil composition of some *Hypericum* species based, on the plant part from which they were extracted**

Species	Main constituents of essential oil	Reference
---------	------------------------------------	-----------

	Leaves/ vegetative part	Flowers/ flowering top	
<i>H. androsaemum</i>	Caryophyllene oxide (35.8%)	$\alpha$ -Guaiene (40.2%)	(Morteza-Semnani <i>et al.</i> , 2005))
	Ishwarane (30.5%)	Caryophyllene oxide (28.0%)	
	Humulene epoxide II (5.6%)	Khusinol (4.2%)	
<i>H. caprifoliatum</i>	Nonane (44.6%)	Nonane (55.8%)	(Ferraz <i>et al.</i> , 2005)
	$\beta$ -Caryophyllene (11.2%)	$\beta$ -Caryophyllene (5.9%)	
	Bicyclogermacrene (5.6%)	Undecane (5.0%)	
<i>H. perforatum</i>	Caryophyllene oxide (9.3-25.9%)	Caryophyllene oxide (7.7-34%)	(Radusiene <i>et al.</i> , 2005)
	Spathulenol (6.4-15.7%)	$\beta$ -caryophyllene (4.2-14.2%)	
	Tetradecanol (1.1-24.5%)	Viridiflorol (4.5-11%)	
<i>H. triquetrifolium</i>	Nonane (14.7%)	Myrcene (16.4%)	(Bertoli <i>et al.</i> , 2003)
	Germacrene D (12.7%)	$\alpha$ -Pinene (13.3%)	
	$\beta$ -caryophyllene (10.9%)	Sabinene (13.1%)	

### Biological Activities of *Hypericum* Essential Oils

Since the middle ages, essential oils have been widely used for bactericidal, virucidal, fungicidal, antiparasitical, insecticidal, medicinal and cosmetic applications (Bakkali *et al.*, 2008). Nowadays, essential oils are used in pharmaceutical, sanitary, cosmetic, agricultural and food industries (Burt, 2004).

Although *Hypericum* extracts have been studied in terms of their biological activities, very few studies have so far been performed with essential oils. Antimicrobial activities have been the most reported biological activities for the essential oils of *Hypericum* species (Tables 4a. 4b. 4c).

**Table 4a - Antimicrobial activities of *Hypericum* essential oils on standard micro organisms in different *in vitro* assays evaluated by the Minimum Inhibitory Concentration (MIC)**

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (MIC)	Reference
<i>H. alpinum</i>	<i>Escherichia coli</i>	50.0 µg/ml	(Saroglou <i>et al.</i> , 2007)
	<i>Proteus mirabilis</i>	no activity	
	<i>Agrobacterium tumefaciens</i>	25.0 µg/ml	
	<i>Pseudomonas aeruginosa</i>	no activity	
	<i>Pseudomonas tolaasii</i>	50.0 µg/ml	
	<i>Salmonella enteritidis</i>	50.0 µg/ml	
	<i>Staphylococcus aureus</i>	12.5 µg/ml	
	<i>Micrococcus luteus</i>	12.5 µg/ml	
	<i>Sarcina lutea</i>	12.5 µg/ml	
	<i>Bacillus cereus</i>	12.5 µg/ml	
	<i>Candida albicans</i>	no activity	
<i>H. barbatum</i>	<i>Escherichia coli</i>	25.0 µg/ml	(Saroglou <i>et al.</i> , 2007)
	<i>Proteus mirabilis</i>	50.0 µg/ml	
	<i>Agrobacterium tumefaciens</i>	25.0 µg/ml	
	<i>Pseudomonas aeruginosa</i>	50.0 µg/ml	
	<i>Pseudomonas tolaasii</i>	25.0 µg/ml	
	<i>Salmonella enteritidis</i>	25.0 µg/ml	
	<i>Staphylococcus aureus</i>	6.25 µg/ml	
	<i>Micrococcus luteus</i>	6.25 µg/ml	
	<i>Sarcina lutea</i>	6.25 µg/ml	
	<i>Bacillus cereus</i>	6.25 µg/ml	
	<i>Candida albicans</i>	25.0 µg/ml	
<i>H. canariense</i>	<i>Bacillus cereus</i> var. <i>mycoides</i>	0.05 mg/ml	(Rabanal <i>et al.</i> , 2005)
	<i>Micrococcus luteus</i>	0.05 mg/ml	
	<i>Staphylococcus aureus</i>	0.05 mg/ml	



<i>Hypericum</i> sps	Test organism	Antimicrobial activity (MIC)	Reference
	<i>Staphylococcus epidermis</i>	0.11 mg/ml	
	<i>Bordetella bronchiseptica</i>	0.11mg/ml	
<b><i>H. coris</i></b>	<i>Escherichia coli</i>	no activity	(Schwob <i>et al.</i> , 2002)
	<i>Enterococcus hirae</i>	no activity	
	<i>Staphylococcus aureus</i>	CI 100µg/ml	
	<i>Candida albicans</i>	no activity	
	<i>Saccharomyces cerevisiae</i>	CI 100µg/ml	
<b><i>H. glandulosum</i></b>	<i>Bacillus cereus</i> var. <i>mycoides</i>	0.22 mg/ml	(Rabanal <i>et al.</i> , 2005)
	<i>Micrococcus luteus</i>	0.15 mg/ml	
	<i>Staphylococcus aureus</i>	0.09 mg/ml	
	<i>Staphylococcus epidermis</i>	0.22 mg/ml	
	<i>Bordetella bronchiseptica</i>	0.11mg/ml	
<b><i>H. grandifolium</i></b>	<i>Bacillus cereus</i> var. <i>mycoides</i>	0.05 mg/ml	(Rabanal <i>et al.</i> , 2005)
	<i>Micrococcus luteus</i>	0.11 mg/ml	
	<i>Staphylococcus aureus</i>	0.09 mg/ml	
	<i>Staphylococcus epidermis</i>	0.15 mg/ml	
	<i>Bordetella bronchiseptica</i>	0.18 mg/ml	
<b><i>H. hirsutum</i></b>	<i>Escherichia coli</i>	50.0 µg/ml	(Saroglou <i>et al.</i> , 2007)
	<i>Proteus mirabilis</i>	no activity	
	<i>Agrobacterium tumefaciens</i>	50.0 µg/ml	
	<i>Pseudomonas aeruginosa</i>	no activity	
	<i>Pseudomonas tolaasii</i>	50.0 µg/ml	
	<i>Salmonella enteritidis</i>	50.0 µg/ml	
	<i>Staphylococcus aureus</i>	25.0 µg/ml	
	<i>Micrococcus luteus</i>	25.0 µg/ml	
	<i>Sarcina lutea</i>	12.5 µg/ml	

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (MIC)	Reference
	<i>Bacillus cereus</i>	12.5 µg/ml	
	<i>Candida albicans</i>	no activity	
<b><i>H. maculatum</i></b>	<i>Escherichia coli</i>	25.0 µg/ml	(Saroglou <i>et al.</i> , 2007)
	<i>Proteus mirabilis</i>	50.0 µg/ml	
	<i>Agrobacterium tumefaciens</i>	25.0 µg/ml	
	<i>Pseudomonas aeruginosa</i>	25.0 µg/ml	
	<i>Pseudomonas tolaasii</i>	25.0 µg/ml	
	<i>Salmonella enteritidis</i>	25.0 µg/ml	
	<i>Staphylococcus aureus</i>	12.5 µg/ml	
	<i>Micrococcus luteus</i>	12.5 µg/ml	
	<i>Sarcina lutea</i>	12.5 µg/ml	
	<i>Bacillus cereus</i>	12.5 µg/ml	
	<i>Candida albicans</i>	50.0 µg/ml	
	<i>Escherichia coli</i>	25.0 µg/ml	(Saroglou <i>et al.</i> , 2007)
	<i>Proteus mirabilis</i>	50.0 µg/ml	
	<i>Agrobacterium tumefaciens</i>	25.0 µg/ml	
	<i>Pseudomonas aeruginosa</i>	50.0 µg/ml	
	<i>Pseudomonas tolaasii</i>	25.0 µg/ml	
	<i>Salmonella enteritidis</i>	25.0 µg/ml	
<b><i>H. perforatum</i></b>	<i>Staphylococcus aureus</i>	12.5 µg/ml	
	<i>Micrococcus luteus</i>	12.5 µg/ml	
	<i>Sarcina lutea</i>	12.5 µg/ml	
	<i>Bacillus cereus</i>	12.5 µg/ml	
	<i>Candida albicans</i>	50.0 µg/ml	
	<i>Aspergillus niger</i>	15.0 µg/ml	(Rančić <i>et al.</i> , 2005)
	<i>Aspergillus flavus</i>	15.0 µg/ml	

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (MIC)	Reference
	<i>Cladosporium cladosporioides</i>	15.0 µg/ml	
	<i>Penicillium funiculosum</i>	15.0 µg/ml	
	<i>Trichoderma viride</i>	15.0 µg/ml	
<b><i>H. rumeliacum</i></b>	<i>Staphylococcus aureus</i>	7.83 mg/ml	(Couladis <i>et al.</i> , 2003)
	<i>Staphylococcus epidermidis</i>	11.2 mg/ml	
	<i>Escherichia coli</i>	3.8 mg/ml	
	<i>Enterobacter cloacae</i>	17.2 mg/ml	
	<i>Klebsiella pneumoniae</i>	9.3 mg/ml	
	<i>Pseudomonas aeruginosa</i>	7.35 mg/ml	
	<i>Candida albicans</i>	6.34 mg/ml	
	<i>C. tropicalis</i>	5.25 mg/ml	
	<i>C. glabrata</i>	4.75 mg/ml	
	<i>Escherichia coli</i>	25.0 µg/ml	(Saroglou <i>et al.</i> , 2007)
	<i>Proteus mirabilis</i>	50.0 µg/ml	
	<i>Agrobacterium tumefaciens</i>	25.0 µg/ml	
	<i>Pseudomonas aeruginosa</i>	25.0 µg/ml	
	<i>Pseudomonas tolaasii</i>	25.0 µg/ml	
	<i>Salmonella enteritidis</i>	25.0 µg/ml	
	<i>Staphylococcus aureus</i>	6.25 µg/ml	
	<i>Micrococcus luteus</i>	12.5 µg/ml	
	<i>Sarcina lutea</i>	6.25 µg/ml	
	<i>Bacillus cereus</i>	12.5 µg/ml	
	<i>Candida albicans</i>	25.0 µg/ml	

**Table 4b - Antimicrobial activities of *Hypericum* essential oils on standard micro organisms in different *in vitro* assays evaluated by the zone of inhibition (mm).**

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (mm)	Reference
<i>H. perforatum</i>	<i>Candida albicans</i>	5 mm (2.5µl)	(Rančić <i>et al.</i> , 2005)
	<i>Escherichia coli</i>	7 mm (1µl)	
	<i>Micrococcus luteus</i>	6 mm (1µl)	
	<i>Pseudomonas tolaasii</i>	3 mm (1µl)	
	<i>Salmonella enteritidis</i>	4 mm (1µl)	
	<i>Salmonella typhimurium</i>	4 mm (1µl)	
	<i>Staphylococcus aureus</i>	5 mm (1µl)	
	<i>Staphylococcus epidermidis</i>	4 mm (1µl)	
<i>H. scabrum</i>	<i>Escherichia coli</i> K12	18 mm (40µg/disc)	(Kızıllı <i>et al.</i> , 2004)
	<i>Escherichia coli</i> PBR322	10 mm (40µg/disc)	
	<i>Escherichia coli</i> PUC9	14 mm (60µg/disc)	
	<i>Bacillus brevis</i> ATCC	10 mm (40µg/disc)	
	<i>Bacillus cereus</i> DM65	14 mm (60µg/disc)	
	<i>Streptococcus pyogenes</i> DM41	10 mm (60µg/disc)	
	<i>Pseudomonas aeruginosa</i> DMC66	16 mm (40µg/disc)	
	<i>Staphylococcus aureus</i> DMC70	16 mm (40µg/disc)	
<i>Candida albicans</i> DM31	18 mm (60µg/disc)		
<i>H. scabroides</i>	<i>Escherichia coli</i> K12	16 mm (40µg/disc)	(Kızıllı <i>et al.</i> , 2004)
	<i>Escherichia coli</i> PBR322	16 mm (40µg/disc)	
	<i>Escherichia coli</i> PUC9	20 mm (40µg/disc)	
	<i>Bacillus brevis</i> ATCC	12 mm (40µg/disc)	
	<i>Bacillus cereus</i> DM65	10 mm (60µg/disc)	
	<i>Streptococcus pyogenes</i> DM41	10 mm (40µg/disc)	
	<i>Pseudomonas aeruginosa</i> DMC66	14 mm (40µg/disc)	

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (mm)	Reference
	<i>Staphylococcus aureus</i> DMC70	10 mm (40µg/disc)	
	<i>Candida albicans</i> DM31	16 mm (40µg/disc)	
<i>H. triquetrifolium</i>	<i>Escherichia coli</i> K12	12 mm (60µg/disc)	(Kızıllı <i>et al.</i> , 2004)
	<i>Escherichia coli</i> PBR322	10 mm (40µg/disc)	
	<i>Escherichia coli</i> PUC9	10 mm (40µg/disc)	
	<i>Bacillus brevis</i> ATCC	10 mm (40µg/disc)	
	<i>Bacillus cereus</i> DM65	12 mm (40µg/disc)	
	<i>Streptococcus pyogenes</i> DM41	12 mm (60µg/disc)	
	<i>Pseudomonas aeruginosa</i> DMC66	12 mm (40µg/disc)	
	<i>Staphylococcus aureus</i> DMC70	16 mm (40µg/disc)	
<i>H. hyssopifolium</i> var. <i>microcalcynum</i>	<i>Candida albicans</i> DM31	12 mm (40µg/disc)	
	<i>Escherichia coli</i> K12	14 mm (40µg/disc)	(Toker <i>et al.</i> , 2006)
	<i>Escherichia coli</i> PBR322	10 mm (40µg/disc)	
	<i>Escherichia coli</i> PUC9	8 mm (40µg/disc)	
	<i>Bacillus brevis</i> ATCC	16 mm (60µg/disc)	
	<i>Bacillus cereus</i> DM65	10 mm (40µg/disc)	
	<i>Streptococcus pyogenes</i> DM41	14 mm (60µg/disc)	
	<i>Pseudomonas aeruginosa</i> DMC66	12 mm (40µg/disc)	
	<i>Staphylococcus aureus</i> DMC70	12 mm (40µg/disc)	
	<i>Candida albicans</i> DM31	12 mm (40µg/disc)	

**Table 4c - Antimicrobial activities of *Hypericum* essential oils on standard micro organisms in different *in vitro* assays evaluated by % of inhibition**

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (% Inhibition)	Reference
<i>H. linarioides</i>	<i>Alternaria solani</i>	0% (5 mg/ml)	(Cakir <i>et al.</i> , 2005)
	<i>Fusarium acuminatum</i>	0% (5 mg/ml)	
	<i>Fusarium culmorum</i>	13.6% (5 mg/ml)	
	<i>Fusarium equiseti</i>	0% (5 mg/ml)	
	<i>Fusarium oxysporum</i>	0% (5 mg/ml)	
	<i>Fusarium sambucinum</i>	9.6% (5 mg/ml)	
	<i>Fusarium solani</i>	0% (5 mg/ml)	
	<i>Verticillium albo-atrum</i>	36.0% (5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-5	0% (5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-9	87.5% (5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-11	2.8% (5 mg/ml)	
<i>H. heterophyllum</i>	<i>Rhizoctonia solani</i> AG-3	0% (5 mg/ml)	(Cakir <i>et al.</i> , 2004)
	<i>Rhizoctonia solani</i> AG-4	0% (5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-5	6% (0.5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-9	12% (0.5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-11	54% (0.5 mg/ml)	
	<i>Fusarium oxysporum</i>	0% (5 mg/ml)	
	<i>Fusarium culmorum</i>	0% (5 mg/ml)	
	<i>Fusarium sambucinum</i>	0% (5 mg/ml)	
	<i>Fusarium solani</i>	0% (5 mg/ml)	
	<i>Fusarium acuminatum</i>	0% (5 mg/ml)	
<i>H. hyssopifolium</i>	<i>Rhizoctonia solani</i> AG-3	0% (5 mg/ml)	(Cakir <i>et al.</i> , 2004)
	<i>Rhizoctonia solani</i> AG-4	11% (0.5 mg/ml)	

<i>Hypericum</i> sps	Test organism	Antimicrobial activity (% Inhibition)	Reference
	<i>Rhizoctonia solani</i> AG-5	28% (0.5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-9	6% (0.5 mg/ml)	
	<i>Rhizoctonia solani</i> AG-11	46% (0.5 mg/ml)	
	<i>Fusarium oxysporum</i>	6% (0.5 mg/ml)	
	<i>Fusarium culmorum</i>	6% (0.5 mg/ml)	
	<i>Fusarium sambucinum</i>	0% (5 mg/ml)	
	<i>Fusarium solani</i>	0% (5 mg/ml)	
	<i>Fusarium acuminatum</i>	0% (5 mg/ml)	

It is likely that the antimicrobial properties of the *Hypericum* essential oils are based in a great part due to the presence of  $\alpha$ -pinene,  $\beta$ -pinene and (*E*)-caryophyllene as these compounds are known for their antimicrobial effects (Aliγιannis *et al.*, 2001; Mourey *et al.*, 2002; Cakir *et al.*, 2004; Costa *et al.*, 2008). As these compounds are widely found in the essential oils of *Hypericum*, it is not surprising to see many reports on antimicrobial activities of species of this genus (Gudžić *et al.*, 1997; Schwob *et al.*, 2002b; Couladis *et al.*, 2003; Cakir *et al.*, 2004; Cakir *et al.*, 2005; Schwob *et al.*, 2006; Toker *et al.*, 2006; Saroglou *et al.*, 2007). Anti-inflammatory and anticarcinogenic activities have been attributed to  $\beta$ -caryophyllene (Zheng *et al.*, 1992; Martin *et al.*, 1993; Kubo *et al.*, 1996; Tambe *et al.*, 1996). The monoterpene hydrocarbon,  $\alpha$ -pinene have also shown to exert anti-inflammatory activity in rats (Martin *et al.*, 1993).

### ***Antibacterial activity***

Essential oils extracted from the flowers of *H. perforatum* grown in the Vlasina region, Serbia had antibacterial activity against *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, *Sarcina lutea*, *Bacillus subtilis* 841, *Salmonella enteritidis* and *Klebsiella pneumoniae* (Gudžić *et al.*, 1997). However, no activity was found against *Pseudomonas aeruginosa*. (Schwob *et al.*, 2002b) The volatile fraction of *H. coris* aerial parts, consisting mainly of  $\alpha$ -curcumene showed moderate antibacterial activity against *Staphylococcus aureus* whereas, no activity was found against *Escherichia coli* and *Enterococcus hirae* (Schwob *et al.*, 2002b). On the other hand,

moderate growth inhibitory activity was found in the essential oils of *H. hyssopifolium* against *E. hirae* and *S. aureus*, while no activity was detected against *E. coli* (Schwob *et al.*, 2006). Essential oils of *H. lysimachioides* and *H. hyssopifolium* were highly effective in growth inhibition of nine microorganisms (*Escherichia coli* K12, *E. coli* PBR 322, *E. coli* PUC 9, *Bacillus brevis* ATCC, *B. cereus* DMC65, *Streptococcus pyogenes* DMC41, *Pseudomonas aeruginosa* DMC66, and *Staphylococcus aureus* DMC70) at a concentration of 60 to 80 µg (Toker *et al.*, 2006). Essential oil isolated from the aerial parts of *H. rumeliacum* exhibited moderate activities against all the tested bacteria (*S. aureus*, *S. epidermidis*, *E. coli*, *Enterobacter cloacae*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*), with a minimal inhibitory concentration (MIC) of 3.80–17.20 mg/ml (Couladis *et al.*, 2003). The *H. rumeliacum* essential oil showed its highest activity against a Gram-negative strain of *E. coli*, while *E. cloacae* appeared to be the most resistant. According to the authors, the antibacterial properties of *H. rumeliacum* oil could be associated with the high percentage of  $\alpha$ -pinene and  $\beta$ -pinene.

In a recent study, essential oils of six *Hypericum* species (*H. alpinum*, *H. barbatum*, *H. rumeliacum*, *H. hirsutum*, *H. maculatum* and *H. perforatum*) were tested against several bacteria (*Escherichia coli* strain ATCC 35218, *Proteus mirabilis*, *Agrobacterium tumefaciens*, *Pseudomonas aeruginosa*, *Pseudomonas tolaasii*, *Salmonella enteritidis* strain ATCC 13076, *Staphylococcus aureus* strain ATCC 6538, *Micrococcus luteus*, *Sarcina lutea* strain ATCC 9341 and *Bacillus cereus*) (Saroglou *et al.*, 2007). They found that *H. barbatum* essential oil was proved to be the most active, while the essential oils of *H. alpinum* and *H. hirsutum* were inactive against the clinical species of *P. mirabilis* and *P. aeruginosa*.

### ***Antifungal activity***

Essential oil of *H. perforatum* flowers was found to be effective against the fungus *Aspergillus niger*, but not against *Candida albicans* (Gudžić *et al.*, 1997). Similarly, *H. coris* essential oil consisting mainly of  $\alpha$ -curcumene did not show any antimicrobial activity against *Candida albicans* (Couladis *et al.*, 2003), whereas antimicrobial activity was found against *Saccharomyces cerevisiae*. Couladis and co-workers (Couladis *et al.*, 2003) tested the activity of essential oil isolated from the aerial parts of *H. rumeliacum* against three fungi (*Candida albicans*, *C. tropicalis*, and *C. glabrata*) and reported the MIC value of 4.75–6.34 mg/ml. Saroglou *et al.* (Couladis *et al.*, 2003) tested essential oils of six *Hypericum* species against *Candida albicans*. Among them, *H. barbatum*, *H. rumeliacum*, *H. maculatum* and *H. perforatum* essential oils were effective. The MIC for *H. barbatum* and *H. rumeliacum* was 25 µg/ml whereas, *H. maculatum* and



*H. perforatum* showed an MIC of 50 µg/ml, whereas *H. alpinum* and *H. hirsutum* essential oils were inactive.

Cakir and co-workers tested the essential oils of three species of *Hypericum* against several phytopathogenic fungi using agar diffusion method for the possible use in agricultural pest and disease control. A moderate antifungal activity of *H. hyssopifolium* and *H. heterophyllum* was found against *Fusarium acuminatum*, and *Rhizoctonia solani* (strains AG-5, AG-9 and AG-11) (Couladis *et al.*, 2003). *H. linarioides* essential oils was active against *Rhizoctonia solani* strain AG-9 and *Verticillium alboatrum* (Couladis *et al.*, 2003).

Although the antimicrobial activity of essential oils from many plant species has been extensively surveyed, their accurate mechanism of action has not been reported in great details. However, it is thought that it might involve membrane disruption by lipophilic compounds, such as terpenoids (Cowan, 1999). It has been demonstrated that  $\alpha$ -pinene and  $\beta$ -pinene are able to destroy cellular integrity, and thereby, inhibit respiration and ion transport processes, and they can also increase the membrane permeability in yeast cells and isolated mitochondria (Andrews *et al.*, 1980; Uribe *et al.*, 1985). Griffin *et al.* (1999) postulated a relative inactivity of hydrocarbons, which is correlated to their limited hydrogen capacity and water solubility. On the other hand, besides being active, ketones, aldehydes and alcohols have different specificity and activity levels. These facts seem to be associated to the functional group and with hydrogen-bonding parameters. Indeed, it seems that there is a relationship between the compounds' chemical structure and the antimicrobial activities they exert (Griffin *et al.*, 1999). Several data suggest that a great antimicrobial potential could be ascribed to the oxygenated terpenes (Panizi *et al.*, 1993; Adam *et al.*, 1998; Saroglou *et al.*, 2007).

### ***Antiangiogenic activity***

Recently, antiangiogenic activity of essential oil of *Hypericum perforatum* has been demonstrated using the chicken chorio allantoic membrane (CAM) assay by Demirci *et al.* (2008) (Demirci *et al.*, 2008). The essential oils at various concentrations (5-50 microgram/pellet) remarkably prevented new blood vessel growth in the *in vivo* chicken embryo compared to standards. Antiangiogenic effect of *H. perforatum* has also been attributed to hyperforin, a phenolic type compound that do not takes part of the essential oil (Martínez-Poveda B *et al.*, 2005). This compound has shown to inhibit angiogenesis both *in vitro* and *in vivo*. Thus, hyperforin and essential oil may have potential use in cancer and metastasis inhibition, as well as in treatment of angiogenesis-related pathologies.

### ***Antioxidant activity***

Antioxidant and inhibitory activity of acetylcholinesterase were shown in assays performed with essential oils of *Hypericum undulatum* (Ferreira *et al.*, 2006). According to the authors, further investigations should be done with this essential oils to evaluate its potential use in preventing or alleviating patients suffering from Alzheimer's Disease (Ferreira *et al.*, 2006).

### **Biological Activities of *Hypericum* Extracts**

The most of the pharmacological properties described in the previous section related with *Hypericum* essential oils coincides well with biological activities reported to *Hypericum* crude extracts. Hence, we speculate that at least some of the biological activities exhibited by the crude extract should also be present in the essential oils. Here we summarize some of the important biological activities of *Hypericum* crude extract, which have not so far been demonstrated with essential oils, but it would be worth to be done.

### ***Antitumor activity***

Hartwell (1970) reported 14 entries referring to *Hypericum* as a folk remedy for various neoplastic conditions in a survey on the use of plants for the treatment of cancer (Hartwell, 1970). Thus, in the last years, *Hypericum* extracts have been investigated for their potential use as antitumor agent. Antiproliferative activity of crude methanolic extracts of *H. caprifoliatum* Cham. & Schlecht., *H. carinatum* Griseb., *H. connatum* Lam., *H. myrianthum* Cham. & Schlecht., *H. polyanthemum* Klotzsch ex Reichardt and *H. ternum* A. St. Hil. were tested against two cell lines (HT-29 human colon carcinoma cells and H-460 non-small cell lung carcinoma). The most active crude methanolic extracts were those from *H. caprifoliatum*, *H. myrianthum* and, to a lesser extent, from *H. connatum* (Ferraz *et al.*, 2005a). *H. perforatum* extracts (commercially available St. John's wort preparations) showed potential anticarcinogenic effect, since they functioned as potent inhibitors of the major human procarcinogen-activating enzyme, the isoform CYP1A1 (Schwarz *et al.*, 2003). These extracts should be also evaluated for cancer chemopreventive potential. Different antiproliferative effects of *Hypericum* methanolic extracts on leukemia cell lines (K562 and U937), human colon carcinoma cells (HT-29) and non-small cell lung carcinoma (H-460) were also reported (Hostanska *et al.*, 2003a; Ferraz *et al.*, 2005a).

### ***Antioxidant activity***

The actual knowledge that reactive oxygen species are involved in the genesis of several diseases, such as arteriosclerosis, rheumatism, some cancers and ageing has prompted the search for new antioxidants (Heilmann *et al.*, 2003). Thus, there is an increasing demand for phytochemicals with antioxidative activity, not only plant extracts but also isolated compounds. The high amount of phenolic compounds produced by species of the genus *Hypericum* makes its extract very interesting from the potential antioxidant activity point of view. *H. barbatum* Jacq., *H. hirsutum* L., *H. linarioides* Bosse, *H. maculatum* Crantz, *H. olympicum* L., *H. perforatum* L., *H. richeri* Vill., *H. rumeliacum* Boiss. and *H. tetrapterum* Fries crude methanolic extracts have been tested for their antioxidant activity (Radulovic *et al.*, 2007). Even though, all the extracts of the *Hypericum* species studied possess a significant antioxidant activity, it was higher in the crude methanolic extracts of the flowers of *H. perforatum*, followed closely by aerial parts of *H. barbatum*. The antioxidant activity of ethanolic extracts of *H. perforatum* has also been demonstrated (Silva *et al.*, 2005). Methanolic extracts of *Hypericum rumeliacum* Boiss. also revealed antioxidant activity (Galati *et al.*, 2008).

*H. triquetrifolium*, is another species of the genus, which water and methanol extracts had significant antioxidant activities (Conforti *et al.*, 2002; Tawaha *et al.*, 2007). Ethanol and water extracts of the flowers of *H. venustum* showed a strong reducing power, free radicals and hydrogen peroxide scavenging activity (Spiteller *et al.*, 2008). Amongst 21 species of medicinal plants widely used in Bulgaria, extracts of *H. perforatum* prepared as infusions with water, showed considerable antioxidant activity (Ivanova *et al.*, 2005). Various extracts from *H. undulatum*, such as essential oils, decoctions and ethanolic extracts showed inhibitory activity of acetylcholinesterase and antioxidant activity, demonstrating that it may help to prevent or alleviate patients suffering from Alzheimer's Disease (Ferreira *et al.*, 2006). Recently, water and methanol extracts of *H. triquetrifolium*, from Jordania, revealed high antioxidant activities, concomitant with high amounts of phenolic compounds. Significant antioxidant activity has also been attributed to methanolic extracts of *H. triquetrifolium* (Conforti *et al.*, 2002).

### ***Antiviral activity***

Schmitt and co-workers (2001) tested aqueous and methanolic extracts of *H. connatum*, *H. caprifoliatum* and *H. polyanthemum* for their antiviral activity against feline immunodeficiency

virus (FIV). Feline immunodeficiency virus (FIV) is a causative agent of acquired immunodeficiency syndrome (AIDS)-like disease in cats. The fact that FIV has similarity to HIV-1 in many molecular and biochemical properties, and the pathogenesis of FIV infection is thought to be similar to that of HIV infection makes it an attractive model for AIDS research. From the three plants studied only *H. connatum*, used in traditional medicine, showed activity against FIV (Schmitt *et al.*, 2001). Antiviral activity against herpes simplex viruses (HSV) was also shown in crude methanolic extracts of *H. connatum* (Fritz *et al.*, 2007).

### ***Wound-healing and anti-inflammatory activities***

Besides the above mentioned, several other biological activities have been attributed to the *Hypericum* extracts. Wound-healing activity of *H. perforatum* alcoholic extracts was recently shown, when extracts were incubated with chicken embryonic fibroblasts from fertilized eggs (Öztürk *et al.*, 2007). The authors found that the mechanism of action of these extracts was similar to that of titrated extract of *Centella asiatica*, which is used in Europe in wound healing drugs (Maquart *et al.*, 1999). Indeed, both extracts seemed to stimulate collagen synthesis in fibroblast cultures and increase the tensile strength of tissues.

Analgesic and topical anti-inflammatory activities in mice were also reported for methanol extracts of three other *Hypericum* species, *H. reflexum*, *H. canariense* L., and *H. glandulosum* Ait. (Rabanal *et al.*, 2005; Sánchez-Mateo *et al.*, 2006). Furthermore, the methanol extracts of *H. perforatum*, *H. empetrifolium*, *H. triquetrifolium* and *H. rumeliacum* have been shown to exhibit anti-inflammatory activity (Apaydin *et al.*, 1999; Öztürk *et al.*, 2002; Galati *et al.*, 2008). Analgesic activity was also attributed to the *H. empetrifolium* extracts (Trovato *et al.*, 2001).

### **Biotechnology in *Hypericum* Essential Oil Improvement**

Recent data have shown a complex composition of the essential oils as well as the influence of different parameters in their quality. Indeed, as mentioned above, the composition of essential oils depends on climatic and pedological conditions, plant organ and vegetative cycle stage. Thus, it is of utmost importance to characterize the essential oils composition as well as the influence of the referred parameters on its quality, in order to obtain essential oils of constant composition. According to Bakkali (2008), this could only be possible if essential oils are extracted under the

same conditions from the same organ of the plant which has been growing on the same soil, under the same climate and has been picked in the same season.

However, one possible alternative to conventional agriculture practices for the production of essential oils could be the use of *in vitro* cultures (Rao *et al.*, 2002; Lila, 2005). This technology mostly offers the possibility of having a controlled system of production, ensuring a continuous supply of products with uniform quality and yield (Lila, 2005). To the best of our knowledge, essential oils composition upon *in vitro* cultures has been reported only in *H. androsaemum* (Guedes *et al.*, 2003). In the essential oils isolated from shoots of *H. androsaemum*, sesquiterpene hydrocarbons was the major group of compounds, representing more than 80% of the total essential oil. Its major constituent was  $\gamma$ -muurolene (15% of the total essential oil) (Guedes *et al.*, 2003).

Genetic transformation methods can also be used to improve the essential oils content. Transformation is currently used for genetic manipulation of major economic crops, vegetables, ornamental, medicinal, fruit, tree and pasture plants, using *Agrobacterium*-mediated or direct transformation methods (Riva *et al.*, 1998). Genetic transformation on species of *Hypericum* has only been performed in *H. perforatum*. A successful protocol of particle bombardment-mediated transformation for *H. perforatum* L. has been reported (Franklin *et al.*, 2007). According to the authors, this species remains highly recalcitrant towards *Agrobacterium*-mediated transformation. Further data have shown a reduction of *Agrobacterium*-viability when in contact with *H. perforatum* cells. Possibly, this can be due to the activation of defence mechanisms of the plant cells against the bacteria (Franklin *et al.*, 2008). The establishment of successful transformation procedures and the knowledge of biosynthetic pathways of useful secondary metabolites can make possible the modulation of its production. In the last decades, great progress has been made in the elucidation of plant terpenoids biosynthetic pathways at the gene and enzyme levels. Indeed, Metabolic Engineering is seen as a powerful tool to improve secondary metabolites production. For *Hypericum* essential oils, the knowledge of the enzymes involved in the production of its compounds as well as their regulatory mechanisms can possibly lead to the increase in the production of essential oils with higher amount of the desirable compounds. For instance,  $\beta$ -caryophyllene, one of the major constituents of essential oils of some *Hypericum* species, should be an interesting molecule to increase production due to its abovementioned pharmacological activities. Moreover, the enzyme responsible for its synthesis,  $\beta$ -caryophyllene synthase, has already been isolated and characterized in several species (*Artemisia annua* (Cai *et al.*, 2002), *Arabidopsis thaliana* (Chen *et al.*, 2003) and *Cucumis sativus* (Mercke *et al.*, 2004). Different approaches can be used in order to modulate its production, such as over-expression of genes

involved in the production of the desirable compounds, manipulation of transcription factors controlling the expression of cascades of genes, or even the identification of tissue/organ-specific promoters (Lange *et al.*, 1999; Mahmoud *et al.*, 2002).

## Conclusion

Without doubt, *Hypericum* is an important genus containing a vast array of secondary metabolites distributed by water, alcoholic extracts and essential oils. Although essential oil composition of several *Hypericum* species has been reported, its pharmacological studies are scarce. Essential oils characterization of *H. perforatum*, *H. androsaemum* and *H. undulatum*, are under way in our laboratory.

## Acknowledgements

This work was sponsored by EU (FSE/FEDER) and Portuguese Government (FCT) through the Project PTDC/AGR-AAM/70418/2006 and the Grant SFRH/BD/13283/2003.

## References

- Abreu, I., M. Reis, A. Marsaioli and P. Mazzafera (2004). "Essential oil composition of *Hypericum brasiliense* choise." *Flavour and Fragrance Journal*(19): 80-82.
- Adam, K., A. Sivropoulou, S. Kokkini, T. Lanaras and M. Arsenakis (1998). "Antifungal Activities of *Origanum vulgare* subsp. *hirtum*, *Mentha spicata*, *Lavandula angustifolia*, and *Salvia fruticosa* Essential Oils against Human Pathogenic Fungi." *J. Agric. Food Chem.* **46**(5): 1739-1745.
- Aligiannis, N., E. Kalpoutzakis, I. B. Chinou, S. Mitakou, E. Gikas and A. Tsaibopoulos (2001). "Composition and Antimicrobial Activity of the Essential Oils of Five Taxa of Sideritis from Greece." *J. Agric. Food Chem.* **49**(2): 811-815.
- Andrews, R., L. Parks and K. Spence (1980). "Some effects of Douglas fir terpenes on certain microorganisms." *Applied and Environmental Microbiology* **40**: 301-304.
- Apaydin, S., U. Zeybek and I. Ince (1999). "*Hypericum triquetrifolium* Turra. extract exhibits antinociceptive activity in the mouse." *J Ethnopharmacol* **67**: 307-312.
- Avato, P., F. Raffo, G. Guglielmi, C. Vitali and A. Rosato (2004). "Extracts from St John's wort and their antimicrobial activity." *Phytotherapy Research* **18**(3): 230-232.
- Bakkali, F., S. Averbeck, D. Averbeck and M. Idaomar (2008). "Biological effects of essential oils - A review." *Food and Chemical Toxicology* **46**(2): 446-475.

- Baser, K. H. C., T. Ozek, H. R. Nuriddinov and A. B. Demirci (2002). "Essential oils of two *Hypericum* species from Uzbekistan." *Chemistry of Natural Compounds* **38**(1): 54-57.
- Bertoli, A., F. Menichini, M. Mazzetti, G. Spinelli and I. Morelli (2003). "Volatile constituents of the leaves and flowers of *Hypericum triquetrifolium* Turra." *Flavour and Fragrance Journal* **18**(2): 91-94.
- Bombardelli, E. and P. Morazzoni (1995). "*Hypericum perforatum*." *Fitoterapia* **66**: 43-68.
- Bougard, F., A. Gravot, S. Milesi and E. Grontier (2001). "Production of Plant Secondary Metabolites: a Historical Perspective." *Plant Science* **161**: 839-851.
- Burt, S. (2004). "Essential oils: their antibacterial properties and potential applications in foods—a review." *International Journal of Food Microbiology* **94**: 223– 253.
- Cai, Y., J. Jia, J. Crock, Z. Lin, C. XY and C. R (2002). " A cDNA clone for  $\beta$ -caryophyllene synthase from *Artemisia annua*." *Phytochemistry* **61**: 523-529.
- Cakir, A., S. Kordali, H. Kilic and E. Kaya (2005). "Antifungal properties of essential oil and crude extracts of *Hypericum linarioides* Bosse." *Biochemical Systematics and Ecology* **33**(3): 245-256.
- Cakir, A., S. Kordali, H. Kilic and E. Kaya (2005). "Antifungal properties of essential oil and crude extracts of *Hypericum linarioides* Bosse." *Biochemical Systematics and Ecology* **33**(3): 245-256.
- Cakir, A., S. Kordali, H. Zengin, S. Izumi and T. Hirata (2004). "Composition and antifungal activity of essential oils isolated from *Hypericum hyssopifolium* and *Hypericum heterophyllum*." *Flavour and Fragrance Journal* **19**(1): 62-68.
- Cakir, A., A. Mavi, A. Yildirim, M. E. Duru, M. Harmandar and C. Kazaz (2003). "Isolation and characterization of antioxidant phenolic compounds from the aerial parts of *Hypericum hyssopifolium* L. by activity-guided fractionation." *Journal of Ethnopharmacology* **87**(1): 73-83.
- Camejo-Rodrigues, J., L. Ascensão, M. À. Bonet and J. Vallès (2003). "An ethnobotanical study of medicinal and aromatic plants in the Natural Park of "Serra de São Mamede" (Portugal)." *Journal of Ethnopharmacology* **89**(2-3): 199-209.
- Cardona, M., J. Marco, J. Sendra, E. Seoane and J. Ibanez (1983). "Waxes and volatile oils in *Hypericum-ericoides* (Guttiferae)." *Lipids* **18**(6): 439-442.
- Chen, F., D. Tholl, J. C. D'Auria, A. Farooq, E. Pichersky and J. Gershenzon (2003). "Biosynthesis and Emission of Terpenoid Volatiles from Arabidopsis Flowers." *Plant Cell* **15**(2): 481-494.
- Conforti, F., G. A. Statti, R. Tundis, F. Menichini and P. Houghton (2002). "Antioxidant activity of methanolic extract of *Hypericum triquetrifolium* Turra aerial part." *Fitoterapia* **73**(6): 479-483.
- Costa, E. V., S. D. Teixeira, F. A. Marques, M. C. T. Duarte, C. Delarmelina, M. L. B. Pinheiro, J. R. Trigo and B. H. L. N. Sales Maia (2008). "Chemical composition and antimicrobial activity of the essential oils of the Amazon Guatteropsis species." *Phytochemistry* **69**(9): 1895-1899.
- Couladis, M., P. Baziou, P. V. Petrakis and C. Harvala (2001). "Essential oil composition of *Hypericum perforatum* L. growing in different locations in Greece." *Flavour and Fragrance Journal* **16**(3): 204-206.
- Couladis, M., I. B. Chinou, O. Tzakou and P. V. Petrakis (2003). "Composition and antimicrobial activity of the essential oil of *Hypericum rumeliacum* subsp. *apollinis* (Boiss. & Heldr.)." *Phytotherapy Research* **17**(2): 152-154.
- Cowan, M. (1999). "Plant Products as Antimicrobial Agents." *Clinical Microbiology Reviews* **12**(4): 564-582.

- Crockett, S. L., B. Demirci, K. H. C. BaÅYer and I. A. Khan (2007). "Analysis of the Volatile Constituents of Five African and Mediterranean *Hypericum* L. (Clusiaceae, Hypericoideae) Species." *Journal of Essential Oil Research* **19**(4): 302-306.
- Daudt, R., G. von Poser, G. Neves and S. Rates (2000). "Screening for the antidepressant activity of some species of *Hypericum* from South Brazil." *Phytotherapy Research* **14**(5): 344-346.
- Decosterd, L., E. Hoffmann, R. Kyburz, D. Bray and K. Jostettmann (1991). "A new phloroglucinol derivative from *Hypericum calycinum* with antifungal and *in vitro* antimalarial activity." *Planta Medica* **57**: 548-551.
- Demirci, B., K. H. C. Baser, S. L. Crockett and I. A. Khan (2005). "Analysis of the Volatile Constituents of Asian *Hypericum* L. (Clusiaceae, Hyperidoideae) Species." *Journal of Essential Oil Research* **17**(6): 659-663.
- Demirci, B., T. Kıyan, F. Demirci and K. BaÅer (2008). "Antiangiogenic effects of *Hypericum perforatum* L. essential oil." *PSE-natural products in cancer therapy*.
- Demirci, F. and K. H. C. Baser (2006). "Volatiles of *Hypericum bupleuroides* Griseb." *The Journal of essential oil research* **18**(6): 650-651.
- Ferraz, A., D. H. Faria, M. N. Benneti, A. Brondani da Rocha, G. Schwartzmann, A. Henriques and G. L. von Poser (2005). "Screening for antiproliferative activity of six southern Brazilian species of *Hypericum*." *Phytomedicine* **12**(1-2): 112-115.
- Ferraz, A., D. H. Faria, M. N. Benneti, A. Brondani da Rocha, G. Schwartzmann, A. Henriques and G. L. von Poser (2005a). "Screening for antiproliferative activity of six southern Brazilian species of *Hypericum*." *Phytomedicine* **12**(1-2): 112-115.
- Ferraz, A. B. F., R. P. Limberger, S. A. L. Bordignon, G. L. v. Poser and A. T. Henriques (2005). "Essential oil composition of six *Hypericum* species from southern Brazil." *Flavour and Fragrance Journal* **20**(3): 335-339.
- Ferreira, A., C. Proena, M. L. M. Serralheiro and M. E. M. Araujo (2006). "The *in vitro* screening for acetylcholinesterase inhibition and antioxidant activity of medicinal plants from Portugal." *Journal of Ethnopharmacology* **108**(1): 31-37.
- Ferretti, G., F. Maggi and B. Tirillini (2005). "Essential oil composition of *Hypericum richeri* Vill. from Italy." *Flavour and Fragrance Journal* **20**(3): 295-298.
- Franklin, G., L. Conceio, E. Kombrink and A. Dias (2008). "*Hypericum perforatum* plant cells reduce *Agrobacterium* viability during co-cultivation." *Planta* **227**: 1401-1408.
- Franklin, G., M. Oliveira and A. C. P. Dias (2007). "Production of transgenic *Hypericum perforatum* plants via particle bombardment-mediated transformation of novel organogenic cell suspension cultures." *Plant Science* **172**(6): 1193-1203.
- Fritz, D., C. R. Venturi, S. Cargnin, J. Schripsema, P. M. Roehe, J. A. Montanha and G. L. von Poser (2007). "Herpes virus inhibitory substances from *Hypericum connatum* Lam., a plant used in southern Brazil to treat oral lesions." *Journal of Ethnopharmacology* **113**(3): 517-520.
- Galati, E., G. Contartese, N. Miceli, M. Taviano, V. Sdrafkakis, M. Couladis, O. Tzakou and F. Lanuzza (2008). "Antiinflammatory and antioxidant activity of *Hypericum rumeliacum* Boiss. subsp. *apollinis* (Boiss. & Heldr.) Robson & Strid methanol extract." *Phytotherapy Research* **22**(6): 766-771.
- Griffin, S., S. Wyllie, J. Markham and D. Leach (1999). "The role of structure and molecular properties of terpenoids in determining their antimicrobial activity." *Flavour and Fragrance Journal* **14**(5): 322-332.
- Gudzić, B., S. Đorđević, R. Palić and G. Stojanovi (2001). "Essential oils of *Hypericum olympicum* L. and *Hypericum perforatum* L." *Flavour and Fragrance Journal* **16**(3): 201-203.



- Gudžić, B., J. M. Nedeljković, S. Đorđević and J. J. Čomor (1997). "Composition and antimicrobial activity of essential oil of Hyperici herb (*Hypericum perforatum* L.) from Vlasina Region." *Facta Universitatis* **1**(4): 47-51.
- Gudžić, B., A. Šmelcerović, S. Dorđević, N. Mimica-Dukić and M. Ristić (2007). "Essential oil composition of *Hypericum hirsutum* L." *Flavour and Fragrance Journal* **22**(1): 42-43.
- Guedes, A. P., L. R. Amorim, A. Vicente and M. Fernandes-Ferreira (2004). "Variation of the essential oil content and composition in leaves from cultivated plants of *Hypericum androsaemum* L." *Phytochemical Analysis* **15**: 146-151.
- Guedes, A. P., L. R. Amorim, A. M. S. Vicente, G. Ramos and M. Fernandes-Ferreira (2003). "Essential Oils from Plants and *in vitro* Shoots of *Hypericum androsaemum* L." *J. Agric. Food Chem.* **51**(5): 1399-1404.
- Hartwell, J. L. (1970). "Plants used against cancer. A survey." *Lloydia* **33**: 288-392.
- Heilmann, J., K. Winkelmann and O. Sticher (2003). "Studies on the antioxidative activity of phloroglucinol derivatives isolated from *Hypericum* species." *Planta medica* **69**(3): 202-206.
- Hölzl, J. and M. Petersen (2003). Culture and biotechnology of *Hypericum*. *Chemical constituents of Hypericum ssp.* E. Ernst. London and New York, Taylor & Francis: 77-93.
- Hosni, K., K. Msaâda, M. Ben Taârit, O. Ouchikh, M. Kallel and B. Marzouk (2008). "Essential oil composition of *Hypericum perforatum* L. and *Hypericum tomentosum* L. growing wild in Tunisia." *Industrial Crops and Products* **27**(3): 308-314.
- Hostanska, K., S. Bommer, M. Weber, B. Krasniqi and R. Saller (2003a). "Comparison of the growth-inhibitory effect of *Hypericum perforatum* L. extracts, differing in the concentration of phloroglucinols and flavonoids, on leukaemia cells." *Journal of Pharmacy and Pharmacology* **55**(7): 973-980.
- Ishiguro, K., M. Yamaki, M. Kashihara, S. Takagi and K. Isoi (1990). "Sarothralin G: a new antimicrobial compound from *Hypericum japonicum*." *Planta Medica* **56**: 274-276.
- Ivanova, D., D. Gerova, T. Chervenkov and T. Yankova (2005). "Polyphenols and antioxidant capacity of Bulgarian medicinal plants." *Journal of Ethnopharmacology* **96**(1-2): 145-150.
- Jaric, S., Z. Popovic, M. Macukanovic-Jocic, L. Djurdjevic, M. Mijatovic, B. Karadzic, M. Mitrovic and P. Pavlovic (2007). "An ethnobotanical study on the usage of wild medicinal herbs from Kopaonik Mountain (Central Serbia)." *Journal of Ethnopharmacology* **111**(1): 160-175.
- Jayasuriyab, H., A. M. Clark and J. D. McChesney (1991). "New antimicrobial filicinic acid derivatives from *Hypericum drummondii*." *Journal of Natural Products* **54**(5): 1314-1320.
- Karim, H., M. Kamel, B. Mouna, C. Thouraya and M. Brahim (2007). "Essential oil composition of *Hypericum triquetrifolium* Turra. aerial parts." *Ital J Biochem* **56**(1): 40-56.
- Kızıllı, G., Z. Toker, H. Özen and Ç. Aytekin (2004). "The Antimicrobial Activity of Essential Oils of *Hypericum scabrum*, *Hypericum scabroides* and *Hypericum triquetrifolium*." *Phytotherapy Research* **18**: 339-341.
- Kubo, I., S. Chaudhuri, Y. Kubo, Y. Sanchez, T. Ogura, T. Saito, H. Ishikawa and H. Haraguchi (1996). "Cytotoxic and Antioxidative Sesquiterpenoids from *Heterotheca inuloides*." *Planta Medica*(5): 427-430.
- Kültür, S. (2007). "Medicinal plants used in Kırklareli Province (Turkey)." *Journal of Ethnopharmacology* **111**(2): 341-364.
- Lange, B. and R. Croteau (1999). "Genetic engineering of essential oil production in mint." *Current Opinion in Plant Biology* **2**: 139-144.
- Lila, M. (2005). Valuable Secondary Products from *In vitro* Culture. *Plant development and Biotechnology*. R. Trigiano and D. Gray, CRC Press: 285-289.

- Mahmoud, S. and R. Croteau (2002). "Strategies for transgenic manipulation of monoterpene biosynthesis in plant." *Trends Plant Science* **7**: 366–373.
- Maquart, F. X., F. Chastang, A. Simeon, P. Birembaut, P. Gillery and Y. Wegrowski (1999). "Triterpenes from *Centella asiatica* stimulate extracellular matrix accumulation in rat experimental wounds." *European Journal of Dermatology* **9**(4): 289-96.
- Martin, S., E. Padilla, M. A. Ocete, J. Galvez, J. Jiménez and A. Zarzuelo (1993). "Anti-Inflammatory Activity of the Essential Oil of *Bupleurum fruticosens*." *Planta Medica*(6): 533-536.
- Martínez-Poveda B, Quesada AR and M. MA (2005). "Hyperforin, a bio-active compound of St. John's Wort, is a new inhibitor of angiogenesis targeting several key steps of the process." *International Journal of Cancer* **117**(5): 775–780.
- Mathis, C. and G. Ourisson (1964a). "Étude chimio-taxonomique du genre *Hypericum* : II. Identification de constituants de diverses huiles essentielles d'*Hypericum*." *Phytochemistry* **3**(1): 115-131.
- Mathis, C. and G. Ourisson (1964b). "Étude chimio-taxonomique du genre *Hypericum* : III. Répartition des carbures saturés et des monoterpènes dans les huiles essentielles d'*Hypericum*." *Phytochemistry* **3**(1): 133-141.
- Mathis, C. and G. Ourisson (1964c). "Etude chimio-taxonomique du genre *Hypericum*--IV. : Repartition des sesquiterpenes, des alcools monoterpéniques et des aldehydes satures dans les huiles essentielles d'*Hypericum*." *Phytochemistry* **3**(3): 377-378.
- Mercke, P., I. F. Kappers, F. W. A. Verstappen, O. Vorst, M. Dicke and H. J. Bouwmeester (2004). "Combined Transcript and Metabolite Analysis Reveals Genes Involved in Spider Mite Induced Volatile Formation in Cucumber Plants." *Plant Physiol.* **135**(4): 2012-2024.
- Mockutė, D., G. Bernotienė and A. Judžentienė (2003). "Volatile compounds of the aerial parts of wild St. John's wort (*Hypericum perforatum* L.) plants." *Chemija* **14**(3): 108-111.
- Morteza-Semnani, K. and M. Saedi (2005). "The essential oil composition of *Hypericum androsaemum* L. leaves and flowers from Iran." *Flavour and Fragrance Journal* **20**(3): 332-334.
- Mourey, A. and N. Canillac (2002). "Anti-listeria monocytogenes activity of essential oils components of conifers." *Food Control* **13**: 289–292.
- Mukherjee, P., R. Verpoorte and B. Suresh (2000). "Evaluation of in-vivo wound healing activity of *Hypericum patulum* (family: Hypericaceae) leaf extract on different wound model in rats." *Journal of Ethnopharmacology* **70**: 315–321.
- Mukherjee, P. K., G. S. Saritha and B. Suresh (2001). "Antibacterial spectrum of *Hypericum hookerianum*." *Fitoterapia* **72**(5): 558-560.
- Nahrstedt, A. and V. Butterweck (1997). "Biologically active and other chemical constituents of the herb of *Hypericum perforatum* L." *Pharmacopsychiatry* **30**: 129-134.
- Nogueira, T., F. Duarte, F. Venâncio, R. Tavares, M. Lousã, C. Bicchi and P. Rubiolo (1998). "Aspectos Quimiotaxonómicos do Género *Hypericum* L. em Portugal." *Silva Lusitana* **6**(1): 55-61.
- Nogueira, T., M. J. Marcelo-Curto, A. C. Figueiredo, J. G. Barroso, L. G. Pedro, P. Rubiolo and C. Bicchi (2008). "Chemotaxonomy of *Hypericum* genus from Portugal: Geographical distribution and essential oils composition of *Hypericum perforatum*, *Hypericum humifusum*, *Hypericum linarifolium* and *Hypericum pulchrum*." *Biochemical Systematics and Ecology* **36**(1): 40-50.
- Öztürk, B., S. Apaydin, E. Goldeli, I. Ince and U. Zeybek (2002). "*Hypericum triquetrifolium* Turra. extract exhibits antiinflammatory activity in the rat." *J Ethnopharmacol* **80**: 207-209.

- Öztürk, N., S. Korkmaz and Y. Öztürk (2007). "Wound-healing activity of St. John's Wort (*Hypericum perforatum* L.) on chicken embryonic fibroblasts." *Journal of Ethnopharmacology* **111**(1): 33-39.
- Öztürk, Y., S. Aydin, R. Beis, K. Baser and H. Berberolu (1996). "Effects of *Hypericum calycinum* L. Extract on the Central Nervous System in Mice." *Phytotherapy Research* **10**(8): 700-702.
- Panizi, L., G. Flamini, P. L. Cioni and I. Morelli (1993). "Composition and antimicrobial properties of essential oils of four Mediterranean Lamiaceae." *Journal of Ethnopharmacology* **39**: 167-170.
- Pavlović, M., O. Tzakou, P. V. Petrakis and M. Couladis (2006). "The essential oil of *Hypericum perforatum* L., *Hypericum tetrapterum* Fries and *Hypericum olympicum* L. growing in Greece." *Flavour and Fragrance Journal* **21**(1): 84-87.
- Petrakis, P. V., M. Couladis and V. Roussis (2005). "A method for detecting the biosystematic significance of the essential oil composition: The case of five Hellenic *Hypericum* L. species." *Biochemical Systematics and Ecology* **33**(9): 873-898.
- Pieroni, A., C. L. Quave and R. F. Santoro (2004). "Folk pharmaceutical knowledge in the territory of the Dolomiti Lucane, inland southern Italy." *Journal of Ethnopharmacology* **95**(2-3): 373-384.
- Pintore, G., M. Chessa, G. Boatto, R. Cerri, M. Usai and B. Tirillini (2005). "Essential oil composition of *Hypericum perforatum* L. var. *angustifolium* DC growing wild in Sardinia (Italy)." *The Journal of essential oil research* **17**(5): 533-535
- Pistelli, L., A. Bertoli, S. Zucconelli, I. Morelli, L. Panizzi and F. Menichini (2000). "Antimicrobial activity of crude extracts and pure compounds of *Hypericum hircinum*." *Fitoterapia* **71**(Supplement 1): S138-S140.
- Poutaraud, A. and P. Girardin (2005). "Improvement of medicinal plant quality: a *Hypericum perforatum* literature review as an example." *Plant Genetic Resources* **3**(2): 178-189.
- Prado, B., R. Rabanal and C. Sánchez-Mateo (2002). "Evaluation of the central properties of several *Hypericum* species from the Canary Islands." *Phytotherapy Research* **16**(8): 740-744.
- Rabanal, R., C. Bonkanka, M. Hernández-Pérez and C. Sánchez-Mateo (2005). "Analgesic and topical anti-inflammatory activity of *Hypericum canariense* L. and *Hypericum glandulosum* Ait. ." *Journal of Ethnopharmacology* **96**(3): 591-596.
- Rabanal, R. M., A. Arias, B. Prado, M. Hernández-Pérez and C. C. Sánchez-Mateo (2002). "Antimicrobial studies on three species of *Hypericum* from the Canary Islands." *Journal of Ethnopharmacology* **81**(2): 287-292.
- Radulovic, N., V. Stankov-Jovanovic, G. Stojanovic, A. Smelcerovic, M. Spitteller and Y. Asakawa (2007). "Screening of *in vitro* antimicrobial and antioxidant activity of nine *Hypericum* species from the Balkans." *Food Chemistry* **103**(1): 15-21.
- Radusiene, J., A. Judzentiene and G. Bernotiene (2005). "Essential oil composition and variability of *Hypericum perforatum* L. growing in Lithuania." *Biochemical Systematics and Ecology* **33**(2): 113-124.
- Rančić, A., M. Soković, J. Vukojević, A. Simić, P. Marin, S. Duletić-Laušević and D. Djoković (2005). "Chemical Composition and Antimicrobial Activities of Essential Oils of *Myrrhis odorata* (L.) Scop, *Hypericum perforatum* L and *Helichrysum arenarium* (L.) Moench." *Journal of Essential Oil Research* **17**(3): 341-345.
- Rao, S. R. and G. A. Ravishankar (2002). "Plant cell cultures: Chemical factories of secondary metabolites." *Biotechnology Advances* **20**: 101-153.

- Riva, G. d. I., J. González-Cabrera, R. Vázquez-Padrón and C. Ayra-Pardo (1998). "Agrobacterium tumefaciens: a natural tool for plant transformation." *EJB Electronic Journal of Biotechnology* **1**(3): 1-16.
- Robson, N. (2006). "Studies in the genus *Hypericum* L. (Clusiaceae). 1 Section 9. *Hypericum sensu lato* (part 3): subsection 1. *Hypericum* series 2. *Senanensia*, subsection 2. *Erecta* and section 9b. *Graveolentia*." *Systematics and Biodiversity* **4**: 19-98
- Rocha, L., A. Marston, O. Potterat, M. A. C. Kaplan, H. Stoeckli-Evans and K. Hostettmann (1995). "Antibacterial phloroglucinols and flavonoids from *Hypericum brasiliense*." *Phytochemistry* **40**(5): 1447-1452.
- Sánchez-Mateo, C. C., C. X. Bonkanka, M. Hernández-Pérez and R. M. Rabanal (2006). "Evaluation of the analgesic and topical anti-inflammatory effects of *Hypericum reflexum* L. fil." *Journal of Ethnopharmacology* **107**(1): 1-6.
- Sánchez-Mateo, C. C., B. Prado and R. M. Rabanal (2002). "Antidepressant effects of the methanol extract of several *Hypericum* species from the Canary Islands." *Journal of Ethnopharmacology* **79**(1): 119-127.
- Santos, P. A. G., A. C. Figueiredo, J. G. Barroso, L. G. Pedro and J. J. C. Scheffer (1999). "Composition of the essential oil of *Hypericum foliosum* Aiton from five Azorean islands." *Flavour and Fragrance Journal* **14**(5): 283-286.
- Saroglou, V., P. D. Marin, A. Rancic, M. Veljic and H. Skaltsa (2007). "Composition and antimicrobial activity of the essential oil of six *Hypericum* species from Serbia." *Biochemical Systematics and Ecology* **35**(3): 146-152.
- Schmitt, A. C., A. P. Ravazzolo and G. L. von Poser (2001). "Investigation of some *Hypericum* species native to Southern of Brazil for antiviral activity." *Journal of Ethnopharmacology* **77**(2-3): 239-245.
- Schwarz, D., P. Kisselev and I. Roots (2003). "St. John's Wort Extracts and Some of Their Constituents Potently Inhibit Ultimate Carcinogen Formation from Benzo[a]pyrene-7,8-Dihydrodiol by Human CYP1A1." *Cancer Res* **63**(22): 8062-8068.
- Schwob, I., J.-M. Bessiere, V. Masotti and J. Viano (2004). "Changes in essential oil composition in Saint John's wort (*Hypericum perforatum* L.) aerial parts during its phenological cycle." *Biochemical Systematics and Ecology* **32**(8): 735-745.
- Schwob, I., J. M. Bessiere, M. Dherbomez and J. Viano (2002). "Composition and antimicrobial activity of the essential oil of *Hypericum coris*." *Fitoterapia* **73**(6): 511-513.
- Schwob, I., J. M. Bessiere, M. Dherbomez and J. Viano (2002b). "Composition and antimicrobial activity of the essential oil of *Hypericum coris*." *Fitoterapia* **73**(6): 511-513.
- Schwob, I., J. Viano, G. Jann-Para, J.-M. Bessiere and M. Dherbomez (2006). "Composition and antimicrobial activity of the essential oil of *Hypericum hyssopifolium* ssp. *hyssopifolium* from southeast France." *The Journal of essential oil research* **18**(4): 469-471.
- Silva, B. A., F. Ferreres, J. O. Malva and A. C. P. Dias (2005). "Phytochemical and antioxidant characterization of *Hypericum perforatum* alcoholic extracts." *Food Chemistry* **90**(1-2): 157-167.
- Sokmen, A., B. M. Jones and M. Erturk (1999). "The *in vitro* antibacterial activity of Turkish medicinal plants." *Journal of Ethnopharmacology* **67**: 79-86.
- Spiteller, M., T. Özen, A. Smelcerovic, S. Zuehlke and N. Mimica-Dukic (2008). "Phenolic constituents and the *in vitro* antioxidant activity of the flowers of *Hypericum venustum*." *Fitoterapia* **79**(3): 191-193.

- Tambe, Y., H. Tsujiuchi, G. Honda, Y. Ikeshiro and S. Tanaka (1996). "Gastric Cytoprotection of the Non-Steroidal Anti-Inflammatory Sesquiterpene, beta-Caryophyllene." *Planta Medica* **5**(5): 469-470.
- Tawaha, K., F. Q. Alali, M. Gharaibeh, M. Mohammad and T. El-Elimat (2007). "Antioxidant activity and total phenolic content of selected Jordanian plant species." *Food Chemistry* **104**(4): 1372-1378.
- Toker, Z., G. Kizil, H. Ç. Özen, M. Kizil and S. Ertekin (2006). "Compositions and antimicrobial activities of the essential oils of two *Hypericum* species from Turkey." *Fitoterapia* **77**(1): 57-60.
- Touafek, O., A. Nacer, A. Kabouche and Z. Kabouche (2005). "Analysis of the essential oil of algerian *Hypericum perforatum* (L)." *Flavour and Fragrance Journal* **20**(6): 669-670.
- Trovato, A., E. Raneri, M. Kouladis, O. Tzakou, M. Taviano and E. Galati (2001). "Anti-inflammatory and analgesic activity of *Hypericum empetrifolium* Willd. (Guttiferae)." *Farmaco* **56**: 455-457.
- Upton, R. (1997). *American Herbal Pharmacopoeia and Therapeutic Compendium, St. John's Wort Monograph*. Santacruz, CA.
- Uribe, S., T. Ramirez and A. Pena (1985). "Effects of  $\beta$ -pinene on yeast membrane functions." *Journal of Bacteriology* **161**: 195-200.
- Weyerstahl, P., U. Splittgerber, H. Marschall and V. K. Kaul (1995). "Constituents of the leaf essential oil of *Hypericum perforatum* L. from India." *Flavour and Fragrance Journal* **10**(6): 365-370.
- Zheng, G.-Q., P. M. Kenney and L. K. T. Lam (1992). "Sesquiterpenes from Clove (*Eugenia caryophyllata*) as Potential Anticarcinogenic Agents." *J. Nat. Prod.* **55**(7): 999-1003

