

Effects of NR1I2 and ABCB1 Genetic Polymorphisms on Everolimus Pharmacokinetics in Japanese Renal Transplant Patients

著者	Yagishita Hironobu, Kagaya Hideaki, Saito Mitsuru, Numakura Kazuyuki, Yamamoto Ryohei, Sagehashi Ryuichiro, Habuchi Tomonori, Satoh Shigeru, Miura Masatomo
journal or publication title	INTERNATIONAL JOURNAL OF MOLECULAR SCIENCES
volume	23
number	19
year	2022
出版者	MDPI
関連リンク	http://dx.doi.org/10.3390/ijms231911742 (http://dx.doi.org/10.3390/ijms231911742)
著作権等	(C) 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).
URL	http://hdl.handle.net/10295/00006225

doi: 10.3390/ijms231911742



Article

Effects of *NR1I2* and *ABCB1* Genetic Polymorphisms on Everolimus Pharmacokinetics in Japanese Renal Transplant Patients

Hironobu Yagishita ¹, Hideaki Kagaya ¹, Mitsuru Saito ², Kazuyuki Numakura ², Ryohei Yamamoto ², Ryuichiro Sagehashi ², Tomonori Habuchi ², Shigeru Satoh ³ and Masatomo Miura ^{1,4,*} 

¹ Department of Pharmacy, Akita University Hospital, 1-1-1 Hondo, Akita 010-8543, Japan

² Department of Urology, Akita University Graduate School of Medicine, 1-1-1 Hondo, Akita 010-8543, Japan

³ Center for Kidney Disease and Transplantation, Akita University Hospital, 1-1-1 Hondo, Akita 010-8543, Japan

⁴ Department of Pharmacokinetics, Akita University Graduate School of Medicine, 1-1-1 Hondo, Akita 010-8543, Japan

* Correspondence: m-miura@hos.akita-u.ac.jp; Tel.: +81-18-884-6439

Abstract: The purpose of this study was to evaluate the effects of *NR1I2* (7635G>A and 8055C>T) and *ABCB1* (1236C>T, 2677G>T/A, and 3435C>T) genetic polymorphisms on everolimus pharmacokinetics in 98 Japanese renal transplant patients. On day 15 after everolimus administration, blood samples were collected just prior to and 1, 2, 3, 4, 6, 9, and 12 h after administration. The dose-adjusted area under the blood concentration–time curve (AUC₀₋₁₂) of everolimus was significantly lower in patients with the *NR1I2* 8055C/C genotype than in those with other genotypes ($p = 0.022$) and was significantly higher in male patients than female patients ($p = 0.045$). Significant correlations between the dose-adjusted AUC₀₋₁₂ of everolimus and age ($p = 0.001$), aspartate transaminase ($p = 0.001$), and alanine transaminase ($p = 0.005$) were found. In multivariate analysis, aging ($p = 0.008$) and higher alanine transaminase levels ($p = 0.032$) were independently predictive of a higher dose-adjusted everolimus AUC₀₋₁₂. Aging and hepatic dysfunction in patients may need to be considered when evaluating dose reductions in everolimus. In renal transplant patients, management using everolimus blood concentrations after administration may be more important than analysis of *NR1I2* 8055C>T polymorphism before administration.

Keywords: everolimus; polymorphism; pregnane X-receptor; P-glycoprotein



Citation: Yagishita, H.; Kagaya, H.; Saito, M.; Numakura, K.; Yamamoto, R.; Sagehashi, R.; Habuchi, T.; Satoh, S.; Miura, M. Effects of *NR1I2* and *ABCB1* Genetic Polymorphisms on Everolimus Pharmacokinetics in Japanese Renal Transplant Patients. *Int. J. Mol. Sci.* **2022**, *23*, 11742. <https://doi.org/10.3390/ijms231911742>

Academic Editor: Anthony M George

Received: 3 August 2022

Accepted: 28 September 2022

Published: 3 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Everolimus, a mammalian target of rapamycin inhibitor, has been approved for the prophylaxis of acute rejection in renal transplant recipients [1–5]. Individual variability in blood concentrations of everolimus involves several factors, including genetic factors and drug interactions [6]. Everolimus is metabolized by cytochrome P450 (CYP) 3A4 and CYP3A5 in the gut and liver, although CYP3A4 contributes more to this process than CYP3A5 [7]. In several studies, the pharmacokinetics of everolimus have been reported to independent of CYP3A5 polymorphism [6–10].

Everolimus is also a substrate of P-glycoprotein (encoded by *ABCB1*) in the intestines. The calcineurin inhibitor cyclosporine, an inhibitor of P-glycoprotein, significantly increases the area under the blood concentration–time curve (AUC) and the maximum blood concentration (C_{max}) of everolimus but does not affect the elimination half-life [11], and these pharmacokinetic parameters support that drug interactions between cyclosporine and everolimus occur in the intestines [11]. Thus, P-glycoprotein inhibitors primarily affect the oral bioavailability of everolimus rather than everolimus clearance, highlighting the key role of intestinal P-glycoprotein. The three most common single nucleotide polymorphisms

(SNPs) identified in the *ABCB1* transporter are 1236C>T, 2677G>T/A, and 3435C>T [12]; however, in a previous study using a population pharmacokinetic model of 53 renal transplant recipients, *ABCB1* 1236C>T, 2677G>T/A, and 3435C>T polymorphisms did not affect the apparent oral clearance of everolimus [9]. P-glycoprotein and CYP3A is regulated by activated pregnane X-receptor (PXR, *NR1I2*), an important nuclear receptor [13,14], and the 7635G>A (rs6785049) and 8055C>T (rs2276706) polymorphisms in the *NR1I2* gene for human PXR are associated with altered CYP3A4 regulation [15]. However, in this previous study using a pharmacokinetic model [9], *NR1I2* 7635G>A and 8055C>T also did not affect the apparent oral clearance of everolimus. Until now, the effects of *NR1I2* and *ABCB1* genetic polymorphisms on pharmacokinetic parameters such as the AUC and C_{max} to evaluate the involvement of intestinal P-glycoprotein have been unclear. Therefore, the clinical study using pharmacokinetic parameters for the absorption process, but not the apparent clearance, of everolimus is necessary.

Although everolimus can be utilized after renal transplantation to reduce nephrotoxicity induced by tacrolimus [1–3], the blood concentrations of everolimus are not influenced by tacrolimus but are affected by cyclosporine [10,16–18]. Therefore, the dose of everolimus that achieves equivalent exposure is 1.5- or 2-fold higher in the presence of tacrolimus than in the presence of cyclosporine [11,17,18]. Accordingly, when everolimus is administered in combination with cyclosporine, the influence of genetic polymorphisms, such as polymorphisms in *NR1I2* or *ABCB1*, on everolimus pharmacokinetics cannot be sufficiently assessed. In a previous study using a population pharmacokinetic model [9], the blood concentrations of everolimus after beginning treatment at an initial dose of 3 mg twice daily were adjusted based on the target trough blood concentration (C_0) of 6–8 ng/mL when a calcineurin inhibitor-free monotherapy regimen was switched from an immunosuppressive regimen including cyclosporine [9]. Moreover, when everolimus was used in combination with other immunosuppressive drugs, such as calcineurin inhibitors and glucocorticoids, the C_0 range was generally set to 3–8 ng/mL [6]. Until now, in renal transplant recipients, the effects of *NR1I2* and *ABCB1* genetic polymorphisms on pharmacokinetics of everolimus in combination with tacrolimus have remained unclear. In addition, we cannot evaluate the influence of P-glycoprotein on an absorption process of everolimus using only one point of the C_0 . Many studies have investigated the influence of ABC transporter polymorphisms by using only one point of the everolimus C_0 [19–21].

Accordingly, in this study, we evaluated the effects of *NR1I2* and *ABCB1* genetic polymorphisms on everolimus pharmacokinetics in 98 Japanese renal transplant patients in combination with tacrolimus.

2. Results

The clinical characteristics of the patients prior to initiation of everolimus therapy are listed in Table 1. The median age was 54.5 years, and the median body weight was 58.7 kg. The genotype frequencies for the *NR1I2* (7635G>A and 8055C>T) and *ABCB1* (1236C>T, 2677G>T/A, and 3435C>T) genetic polymorphisms in 98 Japanese renal transplant patients are shown in Table 1.

On day 15 at a steady-state after beginning treatment at an initial everolimus dose of 0.75 mg twice daily (1.5 daily dose), the everolimus C_0 was significantly correlated with the AUC_{0-12} (slope = 10.978, intercept = 15.710, $r^2 = 0.862$, $p < 0.001$). The dose-adjusted C_0 and AUC_{0-12} of everolimus in patients with the *NR1I2* 8055C/C genotype were significantly lower than those in patients with the 8055C/T or 8055T/T genotype ($p = 0.011$ and 0.022, respectively); however, there were no significant differences in the elimination half-life among the three groups (Figure 1 and Table 2). The geometric mean dose-adjusted C_0 and AUC_{0-12} of everolimus in patients with the *NR1I2* 8055C/C, 8055C/T and 8055T/T genotype were 3.7, 5.2, and 4.4 ng/mL/mg, respectively ($p = 0.011$, one-way ANOVA test), and 60.0, 76.9, and 71.7 ng·h/mL/mg, respectively ($p = 0.027$, one-way ANOVA test). In addition, there were significant differences in the dose-adjusted trough blood concentrations at 12 h after everolimus administration (C_{12}) and the elimination half-life of

everolimus among the three genotype groups of *ABCB1* 2677G>T/A ($p = 0.042$ and 0.035 , respectively); however, blood concentrations of everolimus in heterozygous carriers of the *ABCB1* 2677 T or A allele were the highest (Table 2).

Table 1. Clinical characteristics of patients before beginning everolimus therapy.

Numbers of Patients (Female: Male)	98	(34: 64)
Age (years)	54.5	(44.0–63.3)
Body weight (kg)	58.7	(50.4–66.5)
Laboratory test values		
Aspartate transaminase (IU/L)	14.0	(11.0–16.0)
Alanine transaminase (IU/L)	14.0	(10.0–20.0)
Serum albumin (g/dL)	3.8	(3.5–4.0)
Total bilirubin (mg/dL)	0.4	(0.4–0.6)
Serum creatinine (mg/dL)	1.3	(1.0–1.7)
<i>NR1I2</i> 7635G>A (rs6785049) G/G: G/A: A/A	32: 48: 18	
<i>NR1I2</i> 8055C>T (rs2276707) C/C: C/T: T/T	20: 52: 26	
<i>ABCB1</i> 1236C>T (rs1128503) C/C: C/T: T/T	10: 44: 44	
<i>ABCB1</i> 2677G>T/A (rs2032582) G/G: G/T+G/A: T/T+T/A	27: 53: 18	
<i>ABCB1</i> 3435C>T (rs1045642) C/C: C/T: T/T	36: 42: 20	

Data are presented as the median (quartile 1–quartile 3) or number.




Figure 1. Mean (\pm standard deviation) plasma concentration–time profiles of everolimus in renal transplant recipients with the *NR1I2* 8055C/C (closed circles, $n = 20$), 8055C/T (open squares, $n = 52$), or 8055T/T genotype (open circles, $n = 26$) on day 15 after initiation of everolimus treatment at a dose of 0.75 mg twice daily (1.5 mg daily dose) in combination with tacrolimus.

The dose-adjusted C_0 and AUC_{0-12} values of everolimus in male patients were significantly higher than those in female patients ($p = 0.004$ and 0.045 , respectively; Table 3). Significant correlations were observed between the dose-adjusted C_0 of everolimus on day 15 after beginning therapy and age ($p < 0.001$), body weight ($p = 0.028$), aspartate transaminase ($p < 0.001$), alanine transaminase ($p < 0.001$), and total bilirubin ($p = 0.039$; Table 3). Furthermore, significant correlations were found between the dose-adjusted AUC_{0-12} of everolimus and age ($p = 0.001$), aspartate transaminase ($p = 0.001$), and alanine transaminase ($p = 0.005$; Table 3).

Table 2. Pharmacokinetic parameters of everolimus in *NR1I2* and *ABCB1* genotype groups.

<i>NR1I2</i> 7635G>A (rs6785049)	G/G	G/A	A/A	<i>p</i> -value
Numbers of patients	32	48	18	
C ₀ /D (ng/mL/mg)	4.6 (3.4–6.8)	5.0 (3.9–6.2)	3.6 (2.9–5.6)	0.245
C ₁₂ /D (ng/mL/mg)	4.0 (2.9–6.2)	4.2 (3.4–5.6)	3.4 (2.4–5.3)	0.385
C _{max} /D (ng/mL/mg)	10.1 (6.9–11.9)	9.0 (7.9–11.6)	8.6 (6.1–11.4)	0.716
Half-life (h)	7.9 (5.8–9.0)	7.7 (6.1–10.0)	7.0 (6.0–10.6)	0.119
AUC ₀₋₁₂ /D (ng·h/mL/mg)	76.3 (54.9–100)	75.5 (63.6–89.9)	63.2 (48.5–80.9)	0.371
<i>NR1I2</i> 8055C>T (rs2276707)	C/C	C/T	T/T	<i>p</i> -value
Numbers of patients	20	52	26	
C ₀ /D (ng/mL/mg)	3.4 (2.7–5.1)	5.3 (3.9–6.7)	4.5 (3.1–7.0)	0.011
C ₁₂ /D (ng/mL/mg)	3.2 (2.2–5.3)	4.4 (3.5–5.9)	3.9 (3.0–6.1)	0.069
C _{max} /D (ng/mL/mg)	7.6 (6.0–9.7)	9.9 (7.9–12.0)	10.0 (8.0–11.4)	0.057
Half-life (h)	7.0 (5.4–9.8)	8.0 (6.4–10.5)	7.0 (5.7–8.3)	0.119
AUC ₀₋₁₂ /D (ng·h/mL/mg)	62.0 (43.5–75.0)	82.7 (64.1–97.9)	66.4 (59.7–90.8)	0.022
<i>ABCB1</i> 1236C>T (rs1128503)	C/C	C/T	T/T	<i>p</i> -value
Numbers of patients	10	44	44	
C ₀ /D (ng/mL/mg)	5.0 (4.2–6.9)	4.3 (3.2–6.4)	5.0 (3.4–6.5)	0.646
C ₁₂ /D (ng/mL/mg)	4.2 (3.9–5.9)	3.6 (2.6–5.3)	4.3 (3.1–6.0)	0.321
C _{max} /D (ng/mL/mg)	9.6 (8.3–12.7)	8.5 (6.7–9.5)	9.8 (7.8–11.4)	0.341
Half-life (h)	6.9 (5.7–9.8)	7.2 (6.0–9.5)	7.8 (6.3–9.8)	0.526
AUC ₀₋₁₂ /D (ng·h/mL/mg)	82.4 (67.7–89.1)	69.3 (54.1–87.6)	74.6 (57.3–98.2)	0.321
<i>ABCB1</i> 2677G>T/A (rs2032582)	G/G	G/T+G/A	T/T+T/A	<i>p</i> -value
Numbers of patients	27	53	18	
C ₀ /D (ng/mL/mg)	4.4 (2.8–5.3)	5.1 (3.7–7.0)	4.4 (3.7–6.3)	0.109
C ₁₂ /D (ng/mL/mg)	3.5 (2.6–4.6)	4.4 (3.3–6.2)	3.8 (2.7–5.6)	0.042
C _{max} /D (ng/mL/mg)	8.9 (6.6–11.1)	9.7 (7.8–12.0)	8.8 (7.2–10.5)	0.583
Half-life (h)	6.4 (5.4–8.4)	8.0 (6.5–10.0)	7.5 (6.1–9.7)	0.035
AUC ₀₋₁₂ /D (ng·h/mL/mg)	64.0 (52.9–84.0)	76.4 (61.9–99.3)	67.7 (54.4–87.4)	0.110
<i>ABCB1</i> 3435C>T (rs1045642)	C/C	C/T	T/T	<i>p</i> -value
Numbers of patients	36	42	20	
C ₀ /D (ng/mL/mg)	4.8 (3.7–6.7)	4.2 (3.3–6.5)	5.0 (3.9–6.2)	0.530
C ₁₂ /D (ng/mL/mg)	4.0 (3.2–5.6)	3.8 (2.7–6.7)	4.6 (3.2–5.9)	0.660
C _{max} /D (ng/mL/mg)	10.0 (6.9–12.1)	9.2 (7.0–11.2)	8.8 (7.9–12.1)	0.654
Half-life (h)	6.9 (5.5–10.0)	7.7 (6.2–9.6)	7.8 (6.5–9.3)	0.728
AUC ₀₋₁₂ /D (ng·h/mL/mg)	77.8 (60.0–88.0)	68.1 (55.9–94.3)	74.6 (63.6–96.4)	0.659

Data are presented as the median (quartile 1–quartile 3) or number. C₀, trough blood concentration at morning time; C₁₂, trough blood concentration at nighttime of 12 h after administration; C_{max}, maximum blood concentration; AUC₀₋₁₂, area under the blood concentration–time curve from 0 to 12 h; D, single dose. Kruskal–Wallis test.

The results of multiple regression analyses, including covariate analyses, are listed in Table 4. Aging ($p = 0.001$), higher alanine transaminase value ($p = 0.019$), and body weight ($p = 0.027$) were independently predictive of a higher dose-adjusted everolimus C₀. In addition, aging ($p = 0.008$) and higher alanine transaminase value ($p = 0.032$) were independently predictive of a higher dose-adjusted everolimus AUC₀₋₁₂. However, the determination coefficients for the everolimus C₀ and AUC₀₋₁₂ were low (0.164 and 0.100, respectively; Table 4).

There were significant differences in sex; everolimus C₀ and AUC₀₋₁₂ on day 15; single dose of everolimus at 1 year; and patient age between patients with dose reduction in everolimus within 1 year based on the target C₀ range and patients with no change in dose (Table 5). However, there were no significant differences between genotypes of *NR1I2* and *ABCB1* (Table 5). On the other hand, there were no significant differences in the everolimus C₀ or AUC₀₋₁₂ on day 15 between patients with dose reduction in everolimus within 1 year based on the onset of everolimus-induced side effects, such as stomatitis and leukopenia and patients with no change in dose (Table 5). Although everolimus

was used in combination with tacrolimus, the changes in everolimus dose within 1 year after beginning everolimus therapy were caused by being above the target concentration range of tacrolimus. There were no significant differences in tacrolimus C_0 and AUC_{0-24} between each group. The dose reductions in everolimus by the onset of side effects or above the target C_0 range could not be predicted from information of *NR112* and *ABCB1* genetic polymorphisms.

Table 3. Comparison of everolimus dose-adjusted C_0 and AUC_{0-12} values and clinical characteristics of patients.

Clinical Characteristics	<i>n</i>	Dose-Adjusted C_0 (ng/mL/mg)		<i>p</i> Value
		Median	Quartile 1–3	
Sex				
Female	34	3.7	(2.7–5.5)	0.004 ^a
Male	64	5.1	(4.2–6.6)	
Correlation coefficient (<i>r</i>)				
Age (years)		0.359		<0.001
Body weight (kg)		0.223		0.028
Laboratory test values				
Aspartate transaminase		0.364		<0.001
Alanine transaminase		0.356		<0.001
Serum albumin		−0.034		0.740
Total bilirubin		0.209		0.039
Serum creatinine		0.055		0.592
Clinical Characteristics	<i>n</i>	Dose-Adjusted AUC_{0-12} (ng·h/mL/mg)		<i>p</i> Value
		Median	Quartile 1–3	
Sex				
Female	34	63.3	(51.1–87.0)	0.045 ^a
Male	64	77.2	(63.2–96.0)	
Correlation coefficient (<i>r</i>)				
Age (years)		0.327		0.001
Body weight (kg)		0.159		0.118
Laboratory test values				
Aspartate transaminase		0.328		0.001
Alanine transaminase		0.283		0.005
Serum albumin		−0.067		0.512
Total bilirubin		0.170		0.094
Serum creatinine		0.041		0.686

^a Mann–Whitney test.

Table 4. Stepwise multiple regression analysis of explanatory variables for everolimus dose-adjusted C_0 and AUC_{0-12} values.

Explanatory Variable for Everolimus C_0	Slope	SE	SRC	<i>p</i> Value	R^2
Age (years)	0.055	0.016	0.318	0.001	0.164
Alanine transaminase (IU/L)	0.047	0.020	0.224	0.019	
Body weight (kg)	0.031	0.014	0.216	0.027	
Intercept =	−0.500	1.336			
Explanatory Variable for Everolimus AUC_{0-12}	Slope	SE	SRC	<i>p</i> value	R^2
Age (years)	0.548	0.202	0.262	0.008	0.100
Alanine transaminase (IU/L)	0.537	0.246	0.210	0.032	
Intercept =	38.115	11.518			

SE, standard error; SRC, standardized regression coefficient.

Table 5. Relationships between changes in everolimus dose within 1 year after beginning everolimus therapy and genotypes of *NR1I2* and *ABCB1*.

Change of Everolimus Dose Within 1 Year	Onset of Side Effects		Dose Adjusted Based on Target Range of Everolimus C ₀			
	Dose Reduction or Withdrawal		Dose Reduction	No Change	Increase in Dose	
Numbers of patients (Female: male)	27 (9: 18)		22 (3: 19) *	37 (16: 21)	12 (6: 6)	
C ₀ on day 15 (ng/mL)	3.9 (3.1–5.4)		4.3 (3.6–6.2) **	3.4 (2.7–3.9)	2.7 (2.1–3.2)	
AUC ₀₋₁₂ on day 15 (ng·h/mL)	59.1 (48.4–74.4)		70.8 (53.1–78.6) **	51.8 (45.8–63.3)	40.9 (34.2–46.8) *	
Starting single dose (mg, baseline)	0.75		0.75	0.75	0.75	
Single dose at 1 year (mg)	0.5 (0–0.5) ***		0.5 (0.25–0.5) ***	0.75	1.0 (1.0–1.2) ***	
Tacrolimus C ₀	7.6 (6.7–9.1)		7.8 (5.6–8.5)	7.0 (5.7–8.8)	8.2 (6.7–9.0)	
Tacrolimus AUC ₀₋₂₄	266 (220–297)		273 (225–315)	269 (215–320)	271 (245–338)	
Age (years)	54.0 (44.0–63.0)		58.5 (55.5–65.0) *	51.0 (39.5–63.0)	49.5 (40.0–60.0)	
Body weight (kg)	59.7 (46.6–76.5)		59.9 (55.4–65.7)	54.5 (46.9–64.0)	57.9 (51.7–64.6)	
Alanine transaminase (IU/L)	15.0 (12.0–22.0)		15.0 (10.0–25.3)	14.0 (8.5–20.0)	11.5 (9.3–13.0)	
<i>NR1I2</i> 7635G>A, G/G: G/A: A/A	9: 14: 4		5: 11: 6	11: 21: 5	7: 2: 3	
<i>NR1I2</i> 8055C>T, C/C: C/T: T/T	3: 17: 7		8: 10: 4	6: 21: 10	3: 4: 5	
<i>ABCB1</i> 1236C>T, C/C: C/T: T/T	3: 8: 16		2: 11: 9	4: 19: 14	1: 6: 5	
<i>ABCB1</i> 2677G>T/A, G/G: G/T+G/A: T/T+T/A	6: 14: 7		4: 14: 4	14: 18: 5	3: 7: 2	
<i>ABCB1</i> 3435C>T, C/C: C/T: T/T	9: 11: 7		8: 9: 5	13: 17: 7	6: 5: 1	

Data are presented as the median (quartile 1–quartile 3) or number. Target range of everolimus C₀: 3–5 ng/mL. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared with the no change group.

3. Discussion

To the best of our knowledge, this is the first study to report the effects of *NR1I2* and *ABCB1* genetic polymorphisms on the actual AUC₀₋₁₂ of everolimus calculated using many sampling points with larger numbers of patients. In the clinical concentration range of everolimus, a univariate analysis of 98 renal transplant recipients showed that the dose-adjusted C₀ and AUC₀₋₁₂ values of everolimus in patients with the *NR1I2* 8055C/C genotype were significantly lower than those in patients with the 8055C/T or 8055T/T genotype; however, in the multivariate analysis, *NR1I2* and *ABCB1* polymorphisms did not affect interindividual variability in everolimus blood concentrations. In multivariate analyses, age and alanine transaminase values had major effects on everolimus C₀ and AUC₀₋₁₂. In addition, the age of patients who underwent dose reductions in everolimus within 1 year was significantly higher than that in patients who did not undergo dose changes. Therefore, everolimus dose reductions should be considered as patients age and in patients with hepatic dysfunction.

In patients with mild and moderate hepatic impairment, the dose of everolimus should initially be reduced [22]. After the initial dose reduction, the dose of everolimus may be adjusted based on the target blood concentration of everolimus [6,22]. Therefore, for renal transplant recipients, based on alanine transaminase values of recipients before everolimus administration, the clinician adjust the initial dose of everolimus (0.25 or 0.5 mg twice daily) and reduce it from the standard dose (0.75 mg twice daily). In the univariate analysis in the current study, the dose-adjusted C₀ and AUC₀₋₁₂ values of everolimus in male patients were significantly higher than those in female patients; however, in the multivariate analysis, sex difference did not affect variability in everolimus blood concentrations. In our study, the alanine transaminase values of male patients were significantly higher than that of female patients ($p < 0.001$). Thus, patient backgrounds seem to be a cause of sex difference, and in the multivariate analysis, sex difference was excluded. Consequently, careful monitoring of alanine transaminase values for renal transplant recipients, especially for elderly patients, is necessary.

In a previous study, patient age and body weight did not contribute to interindividual variability in everolimus blood concentrations [23], in contrast to the results of our current study. Additionally, in this previous study, the mean patient age was 44.4 years [23], which was much younger than that (52.8 years, median 54.5 years) in the current study. Moreover, the mean body weight of patients in the previous study was higher than that in the current study (76.7 versus 60.7 kg, respectively) [23]. Thus, the Japanese patients included in our study were older and had lower body weights, and these factors may have resulted in differences in everolimus exposure. Similar to our study, another previous

study in Japanese renal transplant recipients (median age, 51 years) showed that the dose-adjusted everolimus C_0 was affected by patient age [24]. Furthermore, the expression of P-glycoprotein in intestinal tissue was not correlated with patient age (in patients 21–67 years old) [25]. Therefore, reductions in the drug-metabolizing capacity of the liver observed during aging may increase everolimus exposure.

Four previous clinical studies demonstrated the effects of *ABCB1* genetic polymorphisms on everolimus pharmacokinetics in 53 renal [9], 24 renal [19], 37 cardiac [20], and 65 lung transplant recipients [21]. Although only everolimus C_0 was used in these studies [19–21], in all of these studies, *ABCB1* genetic polymorphisms did not affect everolimus blood concentrations. Therefore, *ABCB1* genotyping prior to the initiation of everolimus therapy is not recommended [6], consistent with the findings of our current study. Because the activation of PXR induces the expression of drug-metabolizing enzymes, such as CYP3A and ABC transporters (e.g., P-glycoprotein) [13,14,26], differences in PXR activation may influence interindividual variability in everolimus pharmacokinetics because everolimus is a substrate of both CYP3A and P-glycoprotein. In the univariate analysis, the dose-adjusted C_0 and AUC_{0-12} values of everolimus in patients with the *NR1I2* 8055C/C genotype were significantly lower than those in patients with other genotypes; however, no changes in the elimination half-life were observed among genotypes. This phenomenon suggests that intestinal P-glycoprotein affects everolimus absorption. However, in the multivariate analysis, *NR1I2* polymorphisms did not affect the blood concentrations of everolimus. By contrast, blood concentrations of everolimus seemed to be more strongly influenced by aging and liver function than by the *NR1I2* 8055C>T polymorphism. Consequently, our current results using actual pharmacokinetic parameters of everolimus obtained at eight time points were consistent with the results obtained from a previous study using a population model [9]. Thus, the pharmacokinetics of everolimus in renal transplant recipients cannot be predicted based on drug metabolism and transport-related SNPs. Management using everolimus blood concentrations after administration may be more important than an analysis of drug metabolism and transport-related SNPs before everolimus administration. Similar to the previous reports [9,27], the development of a population pharmacokinetic mode will be necessary to improve the precision of the therapeutic drug monitoring of everolimus. Further study for population pharmacokinetic mode development using our everolimus pharmacokinetic data will be necessary.

4. Materials and Methods

4.1. Patients and Protocols

Ninety-eight Japanese renal transplant recipients (34 women and 64 men) who received renal grafts at Akita University Hospital between October 2013 and June 2021 were enrolled in the retrospective study. The study protocol was approved by the Ethics Committee of Akita University Graduate School of Medicine (approval no. 1140), and all patients provided written informed consent. The study was carried out during hospitalization.

The criteria for eligibility for the study were as follows: (1) patients were treated with an immunosuppressive regimen based on tacrolimus (Graceptor; Astellas, Tokyo, Japan), mycophenolate mofetil (MMF; Cellcept; Chugai Pharmaceutical, Tokyo, Japan), and steroids, and on day 15 after renal transplantation, everolimus (Certican; Novartis Pharma, Tokyo, Japan) was added; (2) patients received tacrolimus every 24 h at the designated time (09:00 AM), and MMF and everolimus in equally divided doses every 12 h at designated times (09:00 AM and 21:00 PM); (3) patients without serious hepatic dysfunction, renal dysfunction, or gastrointestinal motility; (4) patients who were not taking concomitant drugs, supplements, or foods that may affect CYP3A or P-glycoprotein function; (5) nonsmokers; and (6) patients with an ABO compatible blood type.

All patients received everolimus 0.75 mg twice daily (1.5 mg daily dose) as the initial dose. The target C_0 of everolimus was 3–5 ng/mL after the second week [17]. The target C_0 values of tacrolimus were 10–12 ng/mL during the first week, 8–10 ng/mL during the second to fourth weeks after renal transplantation, and 5–8 ng/mL thereafter. Methyl-

prednisolone was given concomitantly at a dose of 500 mg intravenously (i.v.) on the day of surgery and was tapered to 40 mg/day i.v. during the first week. Subsequently, 10–15 mg/day oral prednisolone was administered in the second to third weeks and 7.5–10 mg/day oral prednisolone was administered thereafter.

Everolimus dose reductions within 1 year were carried out based on the grades of reported side effects, such as stomatitis and leukopenia, and on C_0 values of above the target range of 5.0 ng/mL. By contrast, increased everolimus doses were administered based on the target everolimus C_0 of 3.0 ng/mL.

4.2. Sample Collection and Analytical Methods

On day 15 after everolimus administration (namely, day 29 after renal transplantation), whole blood samples were collected by venipuncture just prior to (C_0) and at 1, 2, 3, 4, 6, 9, and 12 h (C_{12}) after everolimus and tacrolimus administration at 09:00 AM. In addition, for tacrolimus, whole blood samples were also collected at 24 h after administration. Thereafter, blood concentrations of everolimus and tacrolimus were determined by electrochemiluminescence immunoassay using a Cobas e411 system (Roche, Tokyo, Japan) and chemiluminescence magnetic microparticle immunoassays on an Architect-i1000 system (Abbott Laboratories, Abbott Park, IL, USA), respectively, according to the manufacturers' instructions.

4.3. Genotyping

DNA was extracted from peripheral blood samples with a QIAamp Blood Kit (Qiagen, Hilden, Germany) and stored at $-80\text{ }^\circ\text{C}$ until analysis. Genotyping procedures identifying the C and T alleles in exon 12 (1236C>T, rs1128503), the G and T/A alleles in exon 21 (2677G>T/A, rs2032582), and the C and T alleles in exon 26 (3435C>T, rs1045642) of the *ABCB1* gene [28–30]; the G and A alleles in intron 5 (7635G>A, rs6785049) and the C and T alleles in intron 6 (8055C>T, rs2276707) of the *NR1I2* gene [31,32] were identified using polymerase chain reaction–restriction fragment length polymorphism.

4.4. Pharmacokinetic Analysis

Pharmacokinetic analyses of everolimus were carried out using a standard noncompartmental method with Phoenix WinNonlin 6.4 (Pharsight Co., Mountain View, CA, USA). The AUC_{0-12} was calculated using the linear trapezoidal rule. The C_{\max} and C_0 were obtained directly from the profile. The elimination half-life was obtained using the log-linear regression of the terminal phase of the concentration–time data with at least 3 sampling points (elimination half-life = $\ln 2/k_e$; where k_e = elimination rate constant).

4.5. Statistical Procedures

Kolmogorov–Smirnov tests were used to assess distributions. The clinical characteristics of renal transplant recipients were expressed as medians (quartile 1–quartile 3) or numbers. Kruskal–Wallis tests or Mann–Whitney U tests were used to elucidate differences between groups. Spearman's rank correlation coefficient test was used to assess correlations in continuous values between groups, and all results were expressed as correlation coefficients (r values). The effects of factors in univariate analysis were evaluated using stepwise multiple linear regression analysis. Variables with borderline significance ($p < 0.2$) on the univariate analysis were subjected to multivariate regression analyses. Dummy variables were used to replace the groups (1 and 0 in 2 groups; 1 and 0, 0 and 0, and 0 and 1 in 3 groups). Results with p values of less than 0.05 were considered significant, and SPSS 20.0 for Windows (SPSS IBM Japan Inc., Tokyo, Japan) was used for all statistical analyses.

5. Conclusions

Age and alanine transaminase values had major effects on everolimus C_0 and AUC_{0-12} . Therefore, aging and hepatic dysfunction should be considered when evaluating the need for everolimus dose reduction. Management using blood concentrations of everolimus after

administration may be more important than analysis of drug metabolism and transport-related genetic polymorphisms before everolimus administration. Especially for elderly renal transplant recipients, careful monitoring of everolimus blood concentrations and alanine transaminase values is necessary.

Author Contributions: H.Y.: H.K., M.S., S.S. and M.M. participated in the design of the study and reviewed the results. M.S., K.N., R.Y., R.S. and S.S. were responsible for patient enrollment and were involved in data acquisition. H.Y. and H.K. carried out genotyping. H.Y. and H.K. were involved in acquisition of data. H.K. analyzed blood concentrations. H.Y. and M.M. were responsible for the statistical analysis. H.Y., H.K. and M.M. drafted the manuscript. M.S., K.N., M.S., S.S. and T.H. helped to draft the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by a grant (no. 20K07150) from the Japan Society for the Promotion of Science, Tokyo, Japan.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board (or Ethics Committee) of Akita University Graduate School of Medicine (approval no. 1140, 25 December 2013).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data and materials are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

Disclosure: All authors report that they have no relevant relationships to disclose.

References

1. Shihab, F.; Qazi, Y.; Mulgaonkar, S.; McCague, K.; Patel, D.; Peddi, V.R.; Shaffer, D. Association of clinical events with everolimus exposure in kidney transplant patients receiving low doses of tacrolimus. *Am. J. Transplant.* **2017**, *17*, 2363–2371. [[CrossRef](#)] [[PubMed](#)]
2. Sageshima, J.; Ciancio, G.; Chen, L.; Dohi, T.; El-Hinnawi, A.; Paloyo, S.; Misawa, R.; Ekwenna, O.; Yatawatta, A.; Burke, G.W., 3rd. Everolimus with low-dose tacrolimus in simultaneous pancreas and kidney transplantation. *Clin. Transplant.* **2014**, *28*, 797–801. [[CrossRef](#)] [[PubMed](#)]
3. Langer, R.M.; Hene, R.; Vitko, S.; Christiaans, M.; Tedesco-Silva, H., Jr.; Ciechanowski, K.; Cassuto, E.; Rostaing, L.; Vilatoba, M.; Machein, U.; et al. Everolimus plus early tacrolimus minimization: A phase III, randomized, open-label, multicentre trial in renal transplantation. *Transpl. Int.* **2012**, *25*, 592–602. [[CrossRef](#)] [[PubMed](#)]
4. Chan, L.; Hartmann, E.; Cibrik, D.; Cooper, M.; Shaw, L.M. Optimal everolimus concentration is associated with risk reduction for acute rejection in de novo renal transplant recipients. *Transplantation* **2010**, *90*, 31–37. [[CrossRef](#)] [[PubMed](#)]
5. Chan, L.; Greenstein, S.; Hardy, M.A.; Hartmann, E.; Bunnapradist, S.; Cibrik, D.; Shaw, L.M.; Munir, L.; Ulbricht, B.; Cooper, M. Multicenter, randomized study of the use of everolimus with tacrolimus after renal transplantation demonstrates its effectiveness. *Transplantation* **2008**, *85*, 821–826. [[CrossRef](#)] [[PubMed](#)]
6. Shipkova, M.; Hesselink, D.A.; Holt, D.W.; Billaud, E.M.; van Gelder, T.; Kunicki, P.K.; Brunet, M.; Budde, K.; Barten, M.J.; De Simone, P.; et al. Therapeutic drug monitoring of everolimus: A consensus report. *Ther. Drug Monit.* **2016**, *38*, 143–169. [[CrossRef](#)]
7. Picard, N.; Rouguieg-Malki, K.; Kamar, N.; Rostaing, L.; Marquet, P. CYP3A5 genotype does not influence everolimus in vitro metabolism and clinical pharmacokinetics in renal transplant recipients. *Transplantation* **2011**, *91*, 652–656. [[CrossRef](#)]
8. Moes, D.J.; Swen, J.J.; den Hartigh, J.; van der Straaten, T.; van der Heide, J.J.; Sanders, J.S.; Bemelman, F.J.; de Fijter, J.W.; Guchelaar, H.J. Effect of CYP3A4*22, CYP3A5*3, and CYP3A combined genotypes on cyclosporine, everolimus, and tacrolimus pharmacokinetics in renal transplantation. *CPT Pharmacomet. Syst. Pharmacol.* **2014**, *3*, e100. [[CrossRef](#)]
9. Moes, D.J.; Press, R.R.; den Hartigh, J.; van der Straaten, T.; de Fijter, J.W.; Guchelaar, H.J. Population pharmacokinetics and pharmacogenetics of everolimus in renal transplant patients. *Clin. Pharmacokinet.* **2012**, *51*, 467–480. [[CrossRef](#)]
10. Kagaya, H.; Niioka, T.; Saito, M.; Inoue, T.; Numakura, K.; Yamamoto, R.; Akamine, Y.; Habuchi, T.; Satoh, S.; Miura, M. Prediction of tacrolimus exposure by CYP3A5 genotype and exposure of co-administered everolimus in Japanese renal transplant recipients. *Int. J. Mol. Sci.* **2018**, *19*, 882. [[CrossRef](#)]
11. Kovarik, J.M.; Kalbag, J.; Figueiredo, J.; Rouilly, M.; Frazier, O.L.; Rordorf, C. Differential influence of two cyclosporine formulations on everolimus pharmacokinetics: A clinically relevant pharmacokinetic interaction. *J. Clin. Pharmacol.* **2002**, *42*, 95–99. [[CrossRef](#)] [[PubMed](#)]
12. Brambila-Tapia, A.J. MDR1 (ABCB1) polymorphisms: Functional effects and clinical implications. *Rev. Invest. Clin.* **2013**, *65*, 445–454. [[PubMed](#)]

13. Lv, C.; Huang, L. Xenobiotic receptors in mediating the effect of sepsis on drug metabolism. *Acta. Pharm. Sin. B.* **2020**, *10*, 33–41. [[CrossRef](#)] [[PubMed](#)]
14. Niu, X.; Wu, T.; Li, G.; Gu, X.; Tian, Y.; Cui, H. Insights into the critical role of the PXR in preventing carcinogenesis and chemotherapeutic drug resistance. *Int. J. Biol. Sci.* **2022**, *18*, 742–759. [[CrossRef](#)] [[PubMed](#)]
15. Zhang, J.; Kuehl, P.; Green, E.D.; Touchman, J.W.; Watkins, P.B.; Daly, A.; Hall, S.D.; Maurel, P.; Relling, M.; Brimer, C.; et al. The human pregnane X receptor: Genomic structure and identification and functional characterization of natural allelic variants. *Pharmacogenetics* **2001**, *11*, 555–572. [[CrossRef](#)] [[PubMed](#)]
16. Christians, U.; Jacobsen, W.; Benet, L.Z.; Lampen, A. Mechanisms of clinically relevant drug interactions associated with tacrolimus. *Clin. Pharmacokinet.* **2002**, *41*, 813–851. [[CrossRef](#)] [[PubMed](#)]
17. Rostaing, L.; Christiaans, M.H.; Kovarik, J.M.; Pascual, J. The pharmacokinetics of everolimus in de novo kidney transplant patients receiving tacrolimus: An analysis from the randomized ASSET study. *Ann. Transplant.* **2014**, *19*, 337–345.
18. Felipe, C.; Ferreira, A.; Bessa, A.; Abait, T.; Perez, J.D.; Casarini, D.E.; Medina-Pestana, J.; Tedesco, H. Adequacy of initial everolimus dose, with and without calcineurin inhibitors, in kidney transplant recipients. *Ther. Drug Monit.* **2018**, *40*, 52–58. [[CrossRef](#)]
19. Sridharan, K.; Shah, S.; Jassim, A.; Hammad, M.; Ebrahim, A.L.; Al Gadhban, J.E.; Al Segai, O. Evaluation of pharmacogenetics of drug-metabolizing enzymes and drug efflux transporter in renal transplants receiving immunosuppressants. *J. Pers. Med.* **2022**, *19*, 823. [[CrossRef](#)]
20. Lesche, D.; Sigurdardottir, V.; Setoud, R.; Englberger, L.; Fiedler, G.M.; Largiader, C.R.; Mohacsi, P.; Sistonen, J. Influence of CYP3A5 genetic variation on everolimus maintenance dosing after cardiac transplantation. *Clin. Transplant.* **2015**, *29*, 1213–1220. [[CrossRef](#)]
21. Schoeppler, K.E.; Aquilante, C.L.; Kiser, T.H.; Fish, D.N.; Zamora, M.R. The impact of genetic polymorphisms, diltiazem, and demographic variables on everolimus trough concentrations in lung transplant recipients. *Clin. Transplant.* **2014**, *28*, 590–597. [[CrossRef](#)] [[PubMed](#)]
22. Kovarik, J.M.; Sabia, H.D.; Figueiredo, J.; Zimmermann, H.; Reynolds, C.; Dilzer, S.C.; Lasseter, K.; Rordorf, C. Influence of hepatic impairment on everolimus pharmacokinetics: Implications for dose adjustment. *Clin. Pharmacol. Ther.* **2001**, *70*, 425–430. [[CrossRef](#)]
23. Kovarik, J.M.; Kahan, B.D.; Kaplan, B.; Lorber, M.; Winkler, M.; Rouilly, M.; Gerbeau, C.; Cambon, N.; Boger, R.; Rordorf, C. Everolimus Phase 2 Study Group. Longitudinal assessment of everolimus in de novo renal transplant recipients over the first post-transplant year: Pharmacokinetics, exposure-response relationships, and influence on cyclosporine. *Clin. Pharmacol. Ther.* **2001**, *69*, 48–56. [[CrossRef](#)] [[PubMed](#)]
24. Geka, Y.; Hamada, Y.; Fuchinoue, S.; Kimura, T. Evaluation of factors influencing the ratio of the trough blood concentration to dose level of everolimus in Japanese kidney transplant recipients. *Transpl. Immunol.* **2022**, *73*, 101609. [[CrossRef](#)] [[PubMed](#)]
25. Peeters, L.E.J.; Andrews, L.M.; Hesselink, D.A.; de Winter, B.C.M.; van Gelder, T. Personalized immunosuppression in elderly renal transplant recipients. *Pharmacol. Res.* **2018**, *130*, 303–307. [[CrossRef](#)] [[PubMed](#)]
26. Geick, A.; Eichelbaum, M.; Burk, O. Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin. *J. Biol. Chem.* **2001**, *276*, 14581–14587. [[CrossRef](#)]
27. Robertsen, I.; Debord, J.; Asberg, A.; Marquet, P.; Woillard, J.B. A limited sampling strategy to estimate exposure of everolimus in whole blood and peripheral blood mononuclear cells in renal transplant recipients using population pharmacokinetic modeling and Bayesian estimators. *Clin. Pharmacokinet.* **2018**, *57*, 1459–1469. [[CrossRef](#)]
28. Wu, L.; Xu, X.; Shen, J.; Xie, H.; Yu, S.; Liang, T.; Wang, W.; Shen, Y.; Zhang, M.; Zheng, S. MDR1 gene polymorphisms and risk of recurrence in patients with hepatocellular carcinoma after liver transplantation. *J. Surg. Oncol.* **2007**, *96*, 62–68. [[CrossRef](#)] [[PubMed](#)]
29. Tanaka, H.; Imamura, N.; Oguma, N.; Shintani, T.; Tanaka, K.; Hyodo, H.; Oda, K.; Kimura, A. Acute myelogenous leukemia with PIG-A gene mutation evolved from aplastic anemia-paroxysmal nocturnal hemoglobinuria syndrome. *Int. J. Hematol.* **2001**, *73*, 206–212. [[CrossRef](#)] [[PubMed](#)]
30. Cascorbi, I.; Gerloff, T.; Johne, A.; Meisel, C.; Hoffmeyer, S.; Schwab, M.; Schaeffeler, E.; Eichelbaum, M.; Brinkmann, U.; Roots, I. Frequency of single nucleotide polymorphisms in the P-glycoprotein drug transporter MDR1 gene in white subjects. *Clin. Pharmacol. Ther.* **2001**, *69*, 169–174. [[CrossRef](#)]
31. Dring, M.M.; Goulding, C.A.; Trimble, V.I.; Keegan, D.; Ryan, A.W.; Brophy, K.M.; Smyth, C.M.; Keeling, P.W.; O'Donoghue, D.; O'Sullivan, M.; et al. The pregnane X receptor locus is associated with susceptibility to inflammatory bowel disease. *Gastroenterology.* **2006**, *130*, 341–348. [[CrossRef](#)] [[PubMed](#)]
32. Kimura, Y.; Selmi, C.; Leung, P.S.; Mao, T.K.; Schauer, J.; Watnik, M.; Kuriyama, S.; Nishioka, M.; Ansari, A.A.; Coppel, R.L.; et al. Genetic polymorphisms influencing xenobiotic metabolism and transport in patients with primary biliary cirrhosis. *Hepatology* **2005**, *41*, 55–63. [[CrossRef](#)] [[PubMed](#)]