

# Clonal hematopoiesis in adult pure red cell aplasia

|                              |  |
|------------------------------|--|
| 著者                           | Fujishima Naohito, Kohmaru Junki, Koyota Souichi, Kuba Keiji, Saga Tomoo, Omokawa Ayumi, Moritoki Yuki, Ueki Shigeharu, Ishida Fumihiro, Nakao Shinji, Matsuda Akira, Ohta Akiko, Tohyama Kaoru, Yamasaki Hiroshi, Usuki Kensuke, Nakashima Yasuhiro, Sato Shinya, Miyazaki Yasushi, Nannya Yasuhito, Ogawa Seishi, Sawada Kenichi, Mitani Kinuko, Hirokawa Makoto |
| journal or publication title | SCIENTIFIC REPORTS   |
| volume                       | 11   |
| year                         | 2021   |
| 出版者                          | Nature Research  |
| 関連リンク                        | <a href="http://dx.doi.org/10.1038/s41598-021-81890-5">http://dx.doi.org/10.1038/s41598-021-81890-5</a> ( <a href="http://dx.doi.org/10.1038/s41598-021-81890-5">http://dx.doi.org/10.1038/s41598-021-81890-5</a> )  |



|      |   |
|------|---|
| 著作權等 | Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> .<br>(C) The Author(s) 2021 |
| URL  | <a href="http://hdl.handle.net/10295/00006287">http://hdl.handle.net/10295/00006287</a>   |

doi: 10.1038/s41598-021-81890-5



OPEN

## Clonal hematopoiesis in adult pure red cell aplasia

Naohito Fujishima<sup>1,17</sup>, Junki Kohmaru<sup>2,17</sup>, Souichi Koyota<sup>2</sup>, Keiji Kuba<sup>2,3</sup>, Tomoo Saga<sup>4</sup>, Ayumi Omokawa<sup>4</sup>, Yuki Moritoki<sup>4</sup>, Shigeharu Ueki<sup>4</sup>, Fumihiro Ishida<sup>5</sup>, Shinji Nakao<sup>6</sup>, Akira Matsuda<sup>7</sup>, Akiko Ohta<sup>8</sup>, Kaoru Tohyama<sup>9</sup>, Hiroshi Yamasaki<sup>10</sup>, Kensuke Usuki<sup>11</sup>, Yasuhiro Nakashima<sup>12</sup>, Shinya Sato<sup>13</sup>, Yasushi Miyazaki<sup>13</sup>, Yasuhito Nannya<sup>14</sup>, Seishi Ogawa<sup>14</sup>, Kenichi Sawada<sup>15</sup>, Kinuko Mitani<sup>16</sup> & Makoto Hirokawa<sup>4</sup>

Idiopathic pure red cell aplasia (PRCA) and secondary PRCA associated with thymoma and large granular lymphocyte leukemia are generally considered to be immune-mediated. The PRCA2004/2006 study showed that poor responses to immunosuppression and anemia relapse were associated with death. PRCA may represent the prodrome to MDS. Thus, clonal hematopoiesis may be responsible for treatment failure. We investigated gene mutations in myeloid neoplasm-associated genes in acquired PRCA. We identified 21 mutations affecting amino acid sequences in 11 of the 38 adult PRCA patients (28.9%) using stringent filtering of the error-prone sequences and SNPs. Four PRCA patients showed 7 driver mutations in *TET2*, *DNMT3A* and *KDM6A*, and 2 PRCA patients carried multiple mutations in *TET2*. Five PRCA patients had mutations with high VAFs exceeding 0.3. These results suggest that clonal hematopoiesis by stem/progenitor cells might be related to the pathophysiology of chronic PRCA in certain adult patients.

Idiopathic pure red cell aplasia (PRCA) and secondary PRCA not responding to treatments for the underlying diseases in adults are generally considered to be immune-mediated and are treated by immunosuppressive therapy<sup>1,2</sup>. We previously conducted the PRCA2004/2006 study and reported that poor responses to induction therapy and anemia relapse were associated with death<sup>3–6</sup>. Principal causes of death were infections and organ failure. Different outcomes in adult PRCA patients depending on their responses to immunosuppression suggest the heterogeneity of chronic PRCA in adults. Based on previous findings, idiopathic PRCA may be the prodrome to myelodysplastic syndromes<sup>7,8</sup>. In some cases, erythroid hypoplasia/aplasia has been observed in patients with myelodysplastic syndrome (MDS)<sup>9–11</sup>.

Theoretically, there are two potential mechanisms of unresponsiveness to immunosuppression: clonal changes in autoaggressive lymphocytes reacting against erythroid progenitors and clonal hematopoiesis by stem/progenitor cells that have undergone somatic mutations during the disease progression of PRCA. Regarding the former, mutations in the signal transducer and activator of the transcription 3 gene (*STAT3*) were detected in 40% of patients with large granular lymphocyte (LGL) leukemia<sup>12</sup> and have been found in PRCA<sup>13</sup>, aplastic anemia, and MDS patients<sup>14</sup>. Kawakami et al. recently reported that *STAT3* mutations were detected in 43% of PRCA

<sup>1</sup>Division of Blood Transfusion, 44-2 Hiroomote Aza Hasunuma, Akita 010-8543, Japan. <sup>2</sup>Akita University Bioscience Education and Research Center, Akita, Japan. <sup>3</sup>Department of Biochemistry and Metabolic Science, Akita University Graduate School of Medicine, Akita, Japan. <sup>4</sup>Department of General Internal Medicine and Clinical Laboratory Medicine, Akita University Graduate School of Medicine, Akita, Japan. <sup>5</sup>Department of Biomedical Laboratory Sciences, Shinshu University School of Medicine, Matsumoto, Japan. <sup>6</sup>Department of Cellular Transplantation Biology, Kanazawa University Graduate School of Medicine, Kanazawa, Japan. <sup>7</sup>Department of Hemato-Oncology, International Medical Center, Saitama Medical University, Saitama, Japan. <sup>8</sup>Division of Public Health, Department of Social Medicine, Saitama Medical University, Saitama, Japan. <sup>9</sup>Department of Laboratory Medicine, Kawasaki Medical School, Okayama, Japan. <sup>10</sup>Department of Hematology and Oncology, Kumamoto City Hospital, Kumamoto, Japan. <sup>11</sup>Department of Hematology, NTT Medical Center Tokyo, Tokyo, Japan. <sup>12</sup>Department of Medicine and Bioregulatory Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan. <sup>13</sup>Department of Hematology, Atomic Bomb Disease and Hibakusha Medicine Unit, Atomic Bomb Disease Institute, Nagasaki University, Nagasaki, Japan. <sup>14</sup>Department of Pathology and Tumor Biology, Graduate School of Medicine, Kyoto University, Kyoto, Japan. <sup>15</sup>Department of Hematology, Nephrology and Rheumatology, Akita University Graduate School of Medicine, Akita, Japan. <sup>16</sup>Department of Hematology and Oncology, Dokkyo Medical University, Tochigi, Japan. <sup>17</sup>These authors contributed equally: Naohito Fujishima and Junki Kohmaru. ✉email: naofuji@doc.med.akita-u.ac.jp; makihiroka08@gmail.com

patients, including LGL leukemia-associated PRCA, idiopathic PRCA and thymoma-associated PRCA<sup>15</sup>. They also reported that *STAT3* mutation-positive patients were less responsive to cyclosporine than those with wild-type *STAT3*. These results appear to support the hypothesis that clonal changes in autoaggressive lymphocytes may be related to the refractoriness of PRCA to immunosuppression.

To test the working hypothesis that clonal hematopoiesis by stem/progenitor cells might be related to the poor responses and outcomes of PRCA patients, we investigated how often clonal hematopoiesis is detected in adult chronic PRCA. We examined gene mutations in myeloid neoplasm-associated genes and found that rare variants or mutations in myeloid neoplasm-associated genes were detected in adult chronic PRCA.

## Results

**Landscape of somatic mutations in PRCA, aplastic anemia, myeloid neoplasm.** After stringent filtering of the error-prone sequences and excluding SNPs, the remaining variant sequences were curated as described in the Methods section. We identified at least one genomic mutation in 11 out of 38 patients with PRCA (28.9%), and we observed one mutation in only 1 out of 13 patients with aplastic anemia ( $p=0.151$ , Fisher test) and at least one mutation in 7 out of 9 patients with myeloid neoplasms ( $p=0.0178$ , Fisher test) (Fig. 1). No gene mutations were detected in 13 healthy control donors (data not shown). The prevalence of gene mutations in idiopathic, thymoma/thymic cancer-associated, LGL-associated, and other causes of PRCA was 28% (5 of 18), 56% (5 of 9), 20% (1 of 5), and 0% (0 of 6), respectively.

**Interpretation of gene mutations in PRCA.** We classified the curated mutations into three categories: driver, potential driver, and nondriver. The driver mutations were judged based on the previously reported gene mutations implicated in myeloid malignancies in public databases. With this strategy, we identified 21 mutations affecting amino acid sequences in 11 patients with PRCA (Table 1). Four patients with PRCA had 7 driver mutations, including *TET2*, *DNMT3A*, and *KDM6A*. The affected genes were *TET2*, *DNMT3A*, and *KDM6A* with VAFs of 0.03915 to 0.32235. Nine potential driver mutations were found in 8 patients, and the mutated genes were *DNMT3A*, *SMC3*, *CUX1*, *ETV6*, *BCOR*, and *IKZF1* with VAFs of 0.0415 to 0.47025. The five mutations that had been considered irrelevant to myeloid malignancy were found in 4 patients with VAFs of 0.04125–0.501. Collectively, 5 PRCA patients had mutations with very high VAFs exceeding 0.3. Five of 11 patients carried multiple mutations (2–3 genes), and 2 of these 5 patients carried 2 mutations in the same *TET2* gene. *TET2*, *DNMT3A*, *KDM6A*, *CUX1*, and *BCOR1* mutations were detected frequently in study participants.

We observed only one mutation of the *ASXL1* gene in a patient with aplastic anemia (Table 2). We also detected frequent gene mutations in 8 of 9 patients with myeloid neoplasms, including multiple driver mutations.

## Discussion


The present results demonstrate that adult chronic PRCA patients frequently have genomic alterations in myeloid neoplasm-associated genes, which are derived from heterogeneous origins, including common SNPs, rare variant SNPs, and mutations. With regard to the mutations in PRCA, we made two important observations in this study. First, we found 21 distinct mutations of myeloid neoplasm-associated genes in 11 out of 38 adult patients (28.9%) with chronic PRCA, and the 4 patients had driver mutations in *TET2*, *DNMT3A* or *KDM6A*. Second, despite its unknown relevance to disordered hematopoiesis, we found ambiguous and unlikely driver mutations in 4 patients. Interestingly, 5 patients had mutations with high VAFs exceeding 0.3. These two findings suggest the working hypothesis that clonal hematopoiesis by stem/progenitor cells might be related to the pathophysiology of chronic PRCA, at least in a certain proportion of adult patients.

In the present study, we focused on genes implicated in myeloid malignancies and found that mutations in myeloid neoplasm-associated genes in PRCA patients were widely distributed in epigenetics, cohesin, transcription, and cell cycle-related genes. Although *STAT3* mutations are frequently detected in PRCA patients<sup>13,15</sup>, we did not identify *STAT3* mutations in the present study because this gene was not included in the target sequencing panel.

The mutational patterns of PRCA resembled those of myeloid neoplasms. The median age of PRCA patients in this study was 65 years. Jaiswal S, et al. previously reported that age-related clonal hematopoiesis (CHIP) in individuals in their 60 s has been reported in 5.6% (95% confidence interval, 5.0 to 6.3) of the general population, and the majority of these variants occur in epigenetic-related genes, such as *DNMT3A*, *TET2*, and *ASXL1*<sup>16</sup>. This has been confirmed by Genovese et al.<sup>17</sup>. Therefore, it is possible to assume that the mutations of *TET2* and *DNMT3A* in our PRCA patient cohort might reflect the age-related clonal hematopoiesis. The prevalence of genomic alterations in the present PRCA cohort appears to exceed the rate of age-related clonal hematopoiesis, although this remains to be clarified by examining the age-matched healthy controls, since the median age of the healthy control donors in this study was 25.

The limitations of our study were mostly derived from its design. First, we did not have germline controls for analyses. Thus, we could not correctly determine the origins of mutations. Second, a large portion of the samples were analyzed in a retrospective manner, the sample size was too small, and clinical information was limited. Thus, we could not explore the association of the mutations with clinical responses to treatment. Third, we only examined the targeted mutations of 54 myeloid neoplasm-associated genes but not of other genes involved in hematopoiesis. Finally, we could not determine the identified mutations in myeloid lineage cells or lymphoid cells or both in PRCA patients because of the analyzed samples.

The reason why immunosuppression is effective in erythropoietic disorders with genetic alterations in myeloid neoplasm-associated genes is still being debated. Although genomic alterations may become driver mutations that directly affect hematopoiesis, mutational events may also produce mutant proteins that are recognized as altered self<sup>18</sup>. Recently, chronic inflammatory responses have been proposed to reflect the pathophysiology of



**Figure 1.** Landscape of gene mutation in PRCA. Overall genomic alterations are shown here. Each row indicates a distinct genomic alteration, and each column shows the results of the respective individual. Upper and lower panels show mutations in coding and noncoding regions, respectively. Mutations with high VAFs (> 0.40) and low VAFs ( $\leq 0.40$ ) are depicted in red and blue, respectively. None of the 13 healthy control donors showed any genomic alterations.

| Interpretation             | Patient/Age   | Classification | VAF           | Gene     | Functional consequence | Amino acid change | Sample     |
|----------------------------|---------------|----------------|---------------|----------|------------------------|-------------------|------------|
| Driver mutations           | PR-17-12/73 y | Idiopathic     | 0.03915       | TET2     | Frameshift             | p.G641fs          | PBMCs      |
|                            |               |                | 0.1119        | TET2     | SNV                    | p.H1904R          |            |
|                            | PR-17-55/81 y | Idiopathic     | 0.0458        | TET2     | Frameshift             | p.T759fs          | PBMCs      |
|                            |               |                | PR-17-50/50 y | Thymoma  | 0.0486                 | KDM6A             |            |
|                            | 0.06955       | DNMT3A         |               |          | Splicing               | c.1429+1G>A       |            |
|                            | PR-18-76/82 y | LGL-L          |               |          | 0.32235                | TET2              | Frameshift |
| 0.0617                     |               |                | TET2          | Stopgain | p.R1465X               |                   |            |
| Potential driver mutations | PR-17-12/73 y | Idiopathic     | 0.4336        | ETV6     | SNV                    | p.H308N           | PBMCs      |
|                            | PR-17-47/45 y | Idiopathic     | 0.07685       | DNMT3A   | SNV                    | p.T808I           | PBMCs      |
|                            | PR-17-9/60 y  | Idiopathic     | 0.47025       | BCORL1   | SNV                    | p.G1391R          | PBMCs      |
|                            | PR-17-55/81 y | Idiopathic     | 0.10555       | SMC3     | SNV                    | p.K684I           | PBMCs      |
|                            |               |                | 0.0415        | CUX1     | SNV                    | p.S1028P          |            |
|                            | PR-17-13/63 y | Thymoma        | 0.04455       | DNMT3A   | SNV                    | p.P799S           | PBMCs      |
|                            | PR-17-50/50 y | Thymoma        | 0.0903        | CUX1     | SNV                    | (c.A1585T)        | PBMCs      |
|                            | PR-18-81/72 y | Thymoma        | 0.05355       | DNMT3A   | SNV                    | p.L373V           | WB         |
|                            | PR-17-51/78 y | Thymic cancer  | 0.0567        | IKZF1    | SNV                    | p.D488N           | PBMCs      |
| Nondriver mutations        | PR-17-15/56 y | Idiopathic     | 0.501         | KDM6A    | SNV                    | p.A694T           | PBMCs      |
|                            |               |                | 0.36885       | KDM6A    | SNV                    | p.Y80C            |            |
|                            | PR-17-53/48 y | Thymoma        | 0.40885       | BCORL1   | SNV                    | p.E1094G          | PBMCs      |
|                            | PR-17-51/78 y | Thymic cancer  | 0.04125       | CDKN2A   | SNV                    | (c.A217T)         | PBMCs      |
|                            |               |                | 0.09305       | CUX1     | SNV                    | p.Q329R           |            |

**Table 1.** Driver and nondriver mutations in pure red cell aplasia. LGL-L, large granular lymphocyte leukemia; PBMCs, peripheral blood mononuclear cells; WB, whole blood.

| Interpretation             | Patient/Age   | Disease         | VAF         | Gene      | Functional consequence | Amino acid change | Sample |
|----------------------------|---------------|-----------------|-------------|-----------|------------------------|-------------------|--------|
| Driver mutations           | PR-17-25/72 y | Aplastic anemia | 0.1105      | ASXL1     | Frameshift             | p.P808fs          | PBMCs  |
|                            | PR-17-2/82 y  | MDS-MLD         | 0.07735     | U2AF1     | SNV                    | p.S34F            | WB     |
|                            |               |                 | 0.49555     | SRSF2     | SNV                    | p.P95L            |        |
|                            |               |                 | PR-17-39/NA | MDS-MLD   | 0.0864                 | CBL               |        |
|                            | 0.1591        | GNAS            |             |           | SNV                    | p.R201H           |        |
|                            | PR-17-40/NA   | MDS-MLD         | 0.3232      | U2AF1     | SNV                    | p.Q157R           | WB     |
|                            | PR-17-41/NA   | AML-MRC         | 0.04445     | ZRSR2     | Stopgain               | p.K106X           | WB     |
|                            | PR-18-77/70 y | MPN/MDS         | 0.0324      | ASXL1     | Frameshift             | p.G642fs          | WB     |
|                            | PR-18-78/73 y | MDS-SLD         | 0.7453      | EZH2      | SNV                    | p.D659G           | WB     |
|                            |               |                 | 0.4073      | SF3B1     | SNV                    | p.K700E           |        |
|                            |               |                 | 0.40385     | RUNX1     | Stopgain               | p.R201X           |        |
|                            |               |                 | 0.3342      | PTPN11    | SNV                    | p.N58Y            |        |
|                            |               |                 | MDS-L2007   | Cell line | 0.67895                | CEBPA             |        |
|                            | 0.6548        | NRAS            |             |           | SNV                    | p.G12A            |        |
| 0.9569                     | TP53          | Splicing        |             |           | c.672+1G>A             |                   |        |
| Potential driver mutations | PR-18-77/70 y | MPN/MDS         | 0.74965     | CUX1      | Frameshift             | p.R254fs          | WB     |
|                            | PR-18-78/73 y | MDS-SLD         | 0.0345      | BCOR      | Frameshift             | p.E1182fs         | WB     |
| Nondriver mutations        | PR-18-70/NA   | MDS-EB1         | 0.04745     | NOTCH1    | SNV                    | p.L1531M          | WB     |

**Table 2.** Driver and nondriver mutations in aplastic anemia and myeloid neoplasms. MDS myelodysplastic syndrome, MDS-MLD MDS with multilineage dysplasia, AML-MRC acute myeloid leukemia with myelodysplasia-related changes, MPN, myeloproliferative neoplasm, MDS-SLD MDS with single lineage dysplasia, MDS-EB MDS with excess blasts, NA not available, PBMCs peripheral blood mononuclear cells, WB whole blood.

MDS<sup>19–23</sup>. Moreover, Balasubramanian et al. reported that the genetic alterations found in MDS were detected in 5 typical idiopathic PRCA patients in their cohort<sup>24</sup>, suggesting that our results are consistent with their findings. Another possibility is that immunosuppressants induce the erythroid differentiation of hematopoietic stem/progenitor cells in vitro, as described previously by Sawafuji et al.<sup>25</sup>.

It is generally believed that idiopathic PRCA and secondary PRCA not responding to treatments for the underlying diseases in adults are generally considered to be immune-mediated and are treated with immunosuppressive therapy<sup>1,2</sup>. Our present study has demonstrated that adult chronic PRCA is a heterogeneous hematopoietic disorder with various genetic backgrounds. The pathogenicity or functional significance of the somatic mutations found in the present PRCA patient cohort remains to be elucidated by future experiments.

Molecular genetic testing with next-generation sequencing is a candidate tool for the classification and prognostic evaluation of bone marrow failure, such as MDS and aplastic anemia<sup>26–32</sup>. A prospective study is needed to confirm the present results and clarify the diagnostic and predictive values of genetic variations in myeloid neoplasm-associated genes in acquired PRCA. This project is ongoing in collaboration with the prospective cohort study PRCA2016 currently being conducted in Japan.

## Methods

**Patients.** The present study included 38 patients with chronic acquired PRCA (18 idiopathic, 9 thymoma-, 5 LGL leukemia-, 2 systemic lupus erythematosus-, 2 viral infection-, one autoimmune hepatitis-, and one end-stage renal disease-associated PRCA); patients had a median age of 65 years (range: 34–82 years) and were recruited from multiple collaborative centers. The disease status varied among patients. Thirteen patients with idiopathic aplastic anemia had a median age of 68 years (range: 35–88 years), 9 with myeloid neoplasms had a median age of 75 years (range: 70–82 years), and 13 healthy control donors had a median age of 25 years (range: 23–34 years). The diagnostic criteria and classification of PRCA as well as the response criteria for PRCA were based on previous findings<sup>1,3–5</sup>. Aplastic anemia was diagnosed and classified according to previous findings<sup>33,34</sup>. MDS was diagnosed and classified according to the WHO 2017 classification of myeloid neoplasms<sup>35</sup>. This study was approved by the institutional review board of Akita University Graduate School of Medicine and was performed in accordance with the Declaration of Helsinki. Written informed consent was obtained from patients and healthy controls. This study was registered as an observational study at the UMIN Clinical Trials Registry (UMIN-CTR) (UMIN000033866).

**Cell line.** The established cell line MDS-L2007 was used to assess the sensitivity of targeted sequencing for myeloid malignancy-associated genes. This cell line was established from a patient with MDS by Matsuoka et al.<sup>36</sup> and harbored mutations including *NRAS* G12A, *CEBPA* Q311X, and *TET2* F1901fs.

**Mutation analysis.** Extracted genomic DNA samples from peripheral blood were subjected to targeted sequencing for 54 myeloid malignancy-associated genes using a MiSeq system and a TruSight Myeloid Sequencing Panel kit according to the manufacturer's instructions (Illumina). To address the clonal hematopoiesis by stem/progenitor cells that might be related to the pathogenesis and pathophysiology of PRCA, the NGS panel primarily included myeloid malignancy-associated genes. *STAT3* gene mutations were not included. In some cases, DNA from peripheral blood mononuclear cells (PBMCs) obtained by density gradient centrifugation was used in the analyses. Germline controls were not included in the present study.

The sensitivity of the mutation analysis was assessed using serial dilutions of the MDS-L2007 cells in healthy control donor cells (Supplementary Fig. 1). To avoid false-positive results, mutations were identified only by the presence of identical gene alterations in two different sequencing runs for each sample because certain gene mutations were detected in one sequencing run but not in the other, particularly when VAFs were not high. Other researchers have established a cutoff value of higher than 0.02 for positive VAFs in age-related clonal hematopoiesis<sup>16</sup>. Thus, we employed a cutoff value for positive variant allele frequencies (VAFs) of higher than 0.03 for assessing the global genetic alterations in patients at the first step of mutation analysis.

Synonymous single nucleotide variants (SNVs) and ambiguous SNVs that were also found in control specimens were excluded. SNVs that were registered in public databases—including the dbSNP (the National Center for Biotechnology Information), jMorp (Japanese Multi Omics Reference Panel), 1000 Genomes Project (IGSR: The International Genome Sample Resource), and ExAC (Exome Aggregation Consortium)—were considered putative SNPs. To obtain the mutations affecting amino acid sequences (potentially relevant to the pathophysiology in chronic PRCA), exonic variants were selected and curated with public databases, including CLINVAR, COSMIC v90, and InterVar<sup>26,27,37</sup> (Supplementary Data).

**Statistical analysis.** Comparisons between different groups were carried out using Fisher's exact test as appropriate.  $P < 0.05$  was considered to indicate statistical significance. Fisher's exact test p-value calculations were double-checked for accuracy by two independent coresearchers. All statistical analyses were performed using the EZR software program<sup>38</sup>.

Received: 27 September 2020; Accepted: 12 January 2021

Published online: 26 January 2021



## References

- Means, R. T. Jr. Pure red cell aplasia. *Blood* **128**, 2504–2509 (2016).
- Sawada, K., Fujishima, N. & Hirokawa, M. Acquired pure red cell aplasia: updated review of treatment. *Br. J. Haematol.* **142**, 505–514 (2008).
- Sawada, K. *et al.* Long-term outcome of patients with acquired primary idiopathic pure red cell aplasia receiving cyclosporine A. A nationwide cohort study in Japan for the PRCA Collaborative Study Group. *Haematologica* **92**, 1021–1028 (2007).
- Hirokawa, M. *et al.* Long-term response and outcome following immunosuppressive therapy in thymoma-associated pure red cell aplasia: a nationwide cohort study in Japan by the PRCA collaborative study group. *Haematologica* **93**, 27–33 (2008).
- Fujishima, N. *et al.* Long-term responses and outcomes following immunosuppressive therapy in large granular lymphocyte leukemia-associated pure red cell aplasia: a Nationwide Cohort Study in Japan for the PRCA Collaborative Study Group. *Haematologica* **93**, 1555–1559 (2008).
- Hirokawa, M. *et al.* Long-term outcome of patients with acquired chronic pure red cell aplasia (PRCA) following immunosuppressive therapy: a final report of the nationwide cohort study in 2004/2006 by the Japan PRCA collaborative study group. *Br. J. Haematol.* **169**, 879–886 (2015).
- Wang, S. A. *et al.* Myelodysplastic syndrome with pure red cell aplasia shows characteristic clinicopathological features and clonal T-cell expansion. *Br. J. Haematol.* **138**, 271–275 (2007).
- Inui, Y. *et al.* Isolated isochromosome 17q in myelodysplastic syndromes with pure red cell aplasia and basophilia. *Int. Med.* **51**, 1579–1584 (2012).
- Williamson, P. J., Oscier, D. G., Bell, A. J. & Hamblin, T. J. Red cell aplasia in myelodysplastic syndrome. *J. Clin. Pathol.* **44**, 431–432 (1991).
- García-Suarez, J., Pascual, T., Muñoz, M. A., Herrero, B. & Pardo, A. Myelodysplastic syndrome with erythroid hypoplasia/aplasia: a case report and review of the literature. *Am. J. Hematol.* **58**, 319–325 (1998).
- Goyal, R., Varma, N. & Marwaha, R. K. Myelodysplastic syndrome with erythroid hypoplasia. *J. Clin. Pathol.* **58**, 320–321 (2005).
- Koskela, H. L. *et al.* Somatic STAT3 mutations in large granular lymphocytic leukemia. *N. Engl. J. Med.* **366**, 1905–1913 (2012).
- Ishida, F. *et al.* STAT3 gene mutations and their association with pure red cell aplasia in large granular lymphocyte leukemia. *Cancer Sci.* **105**, 342–246 (2014).
- Jerez, A. *et al.* STAT3 mutations indicate the presence of subclinical T-cell clones in a subset of aplastic anemia and myelodysplastic syndrome patients. *Blood* **122**, 2453–2459 (2013).
- Kawakami, T. *et al.* Frequent STAT3 mutations in CD8<sup>+</sup> T cells from patients with pure red cell aplasia. *Blood Adv.* **2**, 2704–2712 (2018).
- Jaiswal, S. *et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N. Engl. J. Med.* **371**, 2488–2498 (2014).
- Genovese, G. *et al.* Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N. Engl. J. Med.* **371**, 2477–2487 (2014).
- Finn, O. J. Cancer immunology. *N. Engl. J. Med.* **358**, 2704–2715 (2008).
- Barreyro, L., Chlon, T. M. & Starczynowski, D. T. Chronic immune response dysregulation in MDS pathogenesis. *Blood* **132**, 1553–1560 (2018).
- Ganan-Gomez, I. *et al.* Deregulation of innate immune and inflammatory signaling in myelodysplastic syndromes. *Leukemia* **29**, 1458–1469 (2015).
- de Vries, A. C. *et al.* T-cell receptor Vbeta CDR3 oligoclonality frequently occurs in childhood refractory cytopenia (MDS-RC) and severe aplastic anemia. *Leukemia* **22**, 1170–1174 (2008).
- Kiladjian, J. J. *et al.* Activation of cytotoxic T-cell receptor gammadelta T lymphocytes in response to specific stimulation in myelodysplastic syndromes. *Haematologica* **93**, 381–389 (2008).
- Kordasti, S. Y. *et al.* IL-17-producing CD4(+) T cells, pro-inflammatory cytokines and apoptosis are increased in low risk myelodysplastic syndrome. *Br. J. Haematol.* **145**, 64–72 (2009).
- Balasubramanian, S. K. *et al.* Rational management approach to pure red cell aplasia. *Haematologica* **103**, 221–230 (2018).
- Sawafuji, K., Miyakawa, Y., Kizaki, M. & Ikeda, Y. Cyclosporin A induces erythroid differentiation of K562 cells through p38 MAPK and ERK pathways. *Am. J. Hematol.* **72**, 67–69 (2003).
- Makishima, H. *et al.* Dynamics of clonal evolution in myelodysplastic syndromes. *Nat. Genet.* **49**, 204–212 (2017).
- Yoshizato, T. *et al.* Somatic mutations and clonal hematopoiesis in aplastic anemia. *N. Engl. J. Med.* **373**, 35–47 (2015).
- Cazzola, M., Della Porta, M. G. & Malcovati, L. The genetic basis of myelodysplasia and its clinical relevance. *Blood* **122**, 4021–4034 (2013).
- Papaemmanuil, E. *et al.* Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood* **122**, 3616–3627 (2013).
- Bejar, R. *et al.* Validation of a prognostic model and the impact of mutations in patients with lower-risk myelodysplastic syndromes. *J. Clin. Oncol.* **30**, 3376–3382 (2012).
- Gerstung, M. *et al.* Combining gene mutation with gene expression data improves outcome prediction in myelodysplastic syndromes. *Nat. Commun.* **6**, 5901. <https://doi.org/10.1038/ncomms6901> (2015).
- Steensma, D. P. How I use molecular genetic tests to evaluate patients who have or may have myelodysplastic syndromes. *Blood* **132**, 1657–1663 (2018).
- Killick, S. B. *et al.* Guidelines for the diagnosis and management of adult aplastic anaemia. *Br. J. Haematol.* **172**, 187–207 (2016).
- Young, N. S. Aplastic anemia. *N. Engl. J. Med.* **379**, 1643–1656 (2018).
- Hasserjian R. P., *et al.* Myelodysplastic syndromes: overview. In *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Revised 4th Edition* (ed. Swerdlow SH *et al.*) 97–106 (International Agency for Research on Cancer, 2017).
- Matsuoka, A. *et al.* Lenalidomide induces cell death in an MDS-derived cell line with deletion of chromosome 5q by inhibition of cytokinesis. *Leukemia* **24**, 748–755 (2010).
- Yoshizato, T. *et al.* Genetic abnormalities in myelodysplasia and secondary acute myeloid leukemia: impact on outcome of stem cell transplantation. *Blood* **129**, 2347–2358 (2017).
- Kanda, Y. Investigation of the freely available easy-to-use software “EZ” for medical statistics. *Bone Marrow Transpl.* **48**, 452–458 (2013).

## Acknowledgements

The authors are grateful to Noriko Tan and Kaori Kato for their technical and secretarial assistance. This study was supported by a grant from the Japan Agency for Medical Research and Development (AMED) under grant number 17929782 (MH) and a Japanese Society of Hematology Research Grant.

## Author contributions

M.H. designed and performed the experiments. M.H. and N.F. wrote the manuscript. J.K., S.K., T.S., A.O., Y.M., S.U., and K.T. performed the experiments. N.F., F.I., A.M., H.Y., K.U., Y.N., S.S., and Y.M. collected clinical



samples. K.K., S.N., A.O., K.S., Y.N., S.O., and K.M. contributed to scientific discussions. All authors read and approved the manuscript.

### Competing interests

The authors declare no competing interests.

### Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-81890-5>.

**Correspondence** and requests for materials should be addressed to N.F. or M.H.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021