

BIB



CENTRE OF BIOLOGICAL ENGINEERING UNIVERSIDADE DO MINHO

INSTITUTE FOR BIOTECHNOLOGY AND BIOENGINEERING

## Integrated system for macroalgae production and conversion into biogas

Ana Nobre, Patrícia Gonçalves, José Carlos Costa, Madalena Alves

IBB-Institute for Biotechnology and Bioengineering, Centre of Biological Engineering, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal

Recently research on energy from macroalgae is being reconsidered driven by the following factors: fuel price increase,  $CO_2$  mitigation policies and interest on renewable energy sources after the Kyoto protocol, and need for energy crops not competing with land for food production. However, the commercial expansion of this energy source is limited by its economic feasibility. In this presentation we analyse the development of integrated systems that promote synergies between macroalgae/biogas production and activities such as aquaculture and urban wastewater treatment. The recycling of nutrients and  $CO_2$  by macroalgae can be an opportunity to reduce the biomass-biogas production cost. Other advantage is the proximity between biomass production, conversion into energy and its consumption, thus avoiding energy losses and pollution in transportation.

Experimental work is underway and includes batch tests to evaluate biodegradability of *Ulva* spp. and *Gracilaria* sp. and co-digestion of these macroalgae with sewage sludge from Beirolas wastewater treatment plant (WWTP). Furthermore, continuous co-digestion experiments are planned to be carried out in a 30 L experimental reactor to be placed in the same WWTP. The objectives are to reproduce sludge digestion in Beirolas WWTP and evaluate algae/sludge co-digestion performance.

Preliminary results indicate methane yields in accordance with similar batch experiments, whereby our  $CH_4$  yields per volatile solids added range between 0.14-0.20 m<sup>3</sup> CH4 kg<sup>-1</sup>  $VS_{added}$  against 0.16-0.27 m<sup>3</sup> CH4 kg<sup>-1</sup>  $VS_{added}$  [1,2] depending on the algae species and pretreatment. Overall, *Ulva* sp. shows a better performance over *Gracilaria* sp. as also confirmed in other experiments [1].

## References

- [1] Habig, C., DeBusk, T. A. and Ryther, J. H., 1984. The effect of nitrogen content on methane production by the marine algae Gracilaria tikvahiae and Ulva sp. Biomass, 4(4): 239-251.
- [2] Nielsen, H.B. and Heiske, S., 2010. Anaerobic digestion of macroalgae: methane potentials, pretreatment, inhibition and co-digestion. 10 th IWA World Symposium on Anaerobic Digestion.