

Complete Genome Sequence of the *Pseudomonas aeruginosa* Bacteriophage phiIBB-PAA2

Diana P. Pires,^a Andrew M. Kropinski,^{b,c} Joana Azeredo,^a Sanna Sillankorva^a

Institute for Biotechnology and Bioengineering, Centre of Biological Engineering, Universidade do Minho, Campus de Gualtar, Braga, Portugal^a; Public Health Agency of Canada, Laboratory for Foodborne Zoonoses, Guelph, ON, Canada^b; Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON, Canada^c

***Pseudomonas aeruginosa* phage phiIBB-PAA2 is a broad-host-range virus isolated from raw hospital sewage (Porto, Portugal). This phage has a terminally redundant (183 bp), 45,344-bp double-stranded DNA (dsDNA) genome encoding 66 coding sequences (CDSs) and 3 tRNAs. It belongs to the family *Podoviridae* and the genus *Luz24likevirus*.**

Received 19 November 2013 Accepted 15 January 2014 Published 6 February 2014

Citation Pires DP, Kropinski AM, Azeredo J, Sillankorva S. 2014. Complete genome sequence of the *Pseudomonas aeruginosa* bacteriophage phiIBB-PAA2. *Genome Announc.* 2(1):e01102-13. doi:10.1128/genomeA.01102-13.

Copyright © 2014 Pires et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Diana P. Pires, priscilapires@deb.uminho.pt.

Pseudomonas aeruginosa is a significant opportunistic pathogen frequently associated with nosocomial infections. The infections caused by this bacterium are generally related to its ability to form biofilms on a variety of surfaces. These infections are difficult to treat due to the low susceptibility of the biofilms to antibiotics (1, 2).

The strictly lytic phage phiIBB-PAA2 was isolated from hospital wastewater in Porto, Portugal, using a reference strain (ATCC 10145) as the host. Our group previously showed that this phage has a broad lytic spectrum even against multidrug-resistant *P. aeruginosa* clinical strains (3).

The transmission electron microscopy (TEM) morphological characterization of phiIBB-PAA2 showed that it belongs to the family *Podoviridae*. The phage genome was sequenced using Roche/454-recommended procedures at the Plateforme d'analyses of the Institut de Biologie Intégrative et des Systèmes (Laval University, Québec, QC, Canada). Shotgun reads were assembled using the gsAssembler module of Newbler v 2.5.3. The potential coding sequences (CDSs) were initially annotated using myRAST (4). Each of the predicted proteins was screened against the National Center for Biotechnology Information (NCBI) protein databases, using BLASTP (5). Promoter sequences were predicted using PHIRE 1.0 (6) and BPROM (7), while predicted terminators were annotated using ARNold (8). tRNAscan-SE (9) was used for tRNA annotation.

The genome of the phage phiIBB-PAA2 consists of a linear double-stranded DNA (dsDNA) of 45,344 bp with a GC content of 52% and two 183-bp-long direct terminal repeats (DTRs) determined by Sanger DNA sequencing. The genome was scanned for CDSs, resulting in 66 predicted genes ranging from 141 bp to 3,168 bp, 45 of which are rightward oriented while the others are leftward oriented. The initiation codon of 89% of the genes is ATG, while 8% start with GTG and 3% with TTG. Based on the BLASTP analyses, only 25% of the proteins have assigned functions.

Our analyses further reveal that phage phiIBB-PAA2 has 8 host-dependent promoters, 3 terminators, and 3 tRNAs. The

closest relative of phage phiIBB-PAA2 is *Pseudomonas* phage LUZ24 (10), with which it shares 83.2% nucleotide identity.

Nucleotide sequence accession number. The complete genome of the *P. aeruginosa* phage phiIBB-PAA2 has been deposited in GenBank under the accession number [KF856712](http://www.ncbi.nlm.nih.gov/GenBank/FASTA/KF856712).

ACKNOWLEDGMENTS

D.P.P. and S.S. acknowledge the financial support from the Portuguese Foundation for Science and Technology (FCT) through the grants SFRH/BD/764407/2011 and SFRH/BPD/48803/2008. This work was supported by the FEDER fund through the COMPETE program and through the National fund of FCT in the scope of the project PTDC/EBB-BIO/114760/2009 (FCOMP-01-0124-FEDER-014759).

REFERENCES

- Mah TF, Pitts B, Pellock B, Walker GC, Stewart PS, O'Toole GA. 2003. A genetic basis for *Pseudomonas aeruginosa* biofilm antibiotic resistance. *Nature* 426(6964):306–310. <http://dx.doi.org/10.1038/nature02122>.
- Ma L, Conover M, Lu H, Parsek MR, Bayles K, Wozniak DJ. 2009. Assembly and development of the *Pseudomonas aeruginosa* biofilm matrix. *PLOS Pathog.* 5(3):e1000354. <http://dx.doi.org/10.1371/journal.ppat.1000354>.
- Pires D, Sillankorva S, Faustino A, Azeredo J. 2011. Use of newly isolated phages for the control of *Pseudomonas aeruginosa* PAO1 and ATCC 10145 biofilms. *Res. Microbiol.* 162:798–806. <http://dx.doi.org/10.1016/j.resmic.2011.06.010>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
- Lavigne R, Sun WD, Volckaert G. 2004. PHIRE, a deterministic approach to reveal regulatory elements in bacteriophage genomes. *Bioinformatics* 20:629–635. <http://dx.doi.org/10.1093/bioinformatics/btg456>.
- Solovyev V, Salamov A. 2011. Automatic annotation of microbial genomes and metagenomic sequences, p 61–78. *In* Li RW (ed), *Metagenomics and its applications in agriculture, biomedicine and environmental studies*. Nova Science Publishers, Hauppauge, NY.

8. Naville M, Ghullot-Gaudeffroy A, Marchais A, Gautheret D. 2011. ARNold: a web tool for the prediction of Rho-independent transcription terminators. *RNA Biol.* 8:11–13. <http://dx.doi.org/10.4161/rna.8.1.13346>.
9. Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res.* 33(Web Server issue):W686–W689. <http://dx.doi.org/10.1093/nar/gki366>.
10. Ceysens P-J, Hertveldt K, Ackermann H-W, Noben J-P, Demeke M, Volckaert G, Lavigne R. 2008. The intron-containing genome of the lytic *Pseudomonas* phage LUZ24 resembles the temperate phage PaP3. *Virology* 377(2):233–238. <http://dx.doi.org/10.1016/j.virol.2008.04.038>.