

MEtop – a top FCNC event generator

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2013 J. Phys.: Conf. Ser. 447 012031

(<http://iopscience.iop.org/1742-6596/447/1/012031>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 193.137.16.114

This content was downloaded on 16/12/2014 at 16:31

Please note that [terms and conditions apply](#).

MEtop - a top FCNC event generator

Rita Coimbra¹, António Onofre², Rui Santos^{3,4} and Miguel Won^{1,4}

¹ LIP / Departamento de Física, Universidade de Coimbra, 3004-516 Coimbra, Portugal

² Departamento de Física, Universidade do Minho, 4710-057 Braga, Portugal

³ Instituto Superior de Engenharia de Lisboa - ISEL, Rua Conselheiro Emídio Navarro 1, 1959-007 Lisboa, Portugal

⁴ Centro de Física Teórica e Computacional, Faculdade de Ciências, Universidade de Lisboa, Av. Prof. Gama Pinto 2, 1649-003 Lisboa, Portugal

E-mail: rita.coimbra@coimbra.lip.pt, Antonio.Onofre@cern.ch, rsantos@cii.fc.ul.pt, miguel.won@coimbra.lip.pt

Abstract. In this work we present a new Monte Carlo generator for Direct top and Single top production via flavour-changing neutral currents (FCNC). This new tool calculates the cross section and generates events with Next-to-Leading order precision for the Direct top process and Leading-Order precision for all other FCNC single top processes. A set of independent dimension six FCNC operators has been implemented - including four-fermion operators - where at least one top-quark is present in the interaction.

1. Introduction

With the new experimental data from the Large Hadron Collider (LHC) one will be able to scrutinise the Standard Model (SM) boundaries with unprecedented precision. Flavour-changing neutral currents (FCNC) top decays such as $t \rightarrow V q$, where $V = Z, \gamma, g$ and $q = u, c$ are highly suppressed in the SM due to the well known Glashow-Iliopoulos-Maiani (GIM) mechanism [1]. top-quark FCNC branching ratios are of the order of 10^{-14} and 10^{-12} for the electroweak and strong case respectively. However, some versions of two-Higgs doublet models (2HDMs) predict quite larger values. In fact, for some regions of the 2HDM parameter space the $\text{BR}(t \rightarrow g q)$ could reach values of the order of 10^{-4} [2], which is eight orders of magnitude above the SM value. The high statistics expected in LHC data forces an increase in the precision of all relevant theoretical calculations. Hence, at least Next-to-Leading (NLO) calculations should be made available whenever possible. The experimental searches performed both at LHC and at the Tevatron on the direct top FCNC process have already used NLO cross sections from [3]. However, the NLO calculation was used only as a normalization factor and the events were generated using LO generators like TopRex [4] or Protos [5]. In this work we present a new Monte Carlo tool that generates FCNC direct top events with NLO precision. The FCNC interactions were implemented via an effective lagrangian [6] where a complete set of dimension six operators was used. The events are generated in the Les-Houches Event (LHE) format [7] and can therefore be easily interfaced with shower algorithms such as the one in Pythia [8]. Since the generator calculates the total inclusive cross section, the FCNC single top process was also included and can be generated independently. Not only strong FCNC interactions, but also electroweak and four-fermions dimensions six operators were included. Each operator can be turned on independently allowing for independent studies of the different Lorentz structures. In



section 2 the effective formalism is presented while in section 3 we show the physical processes already included in MEtop. In section 4 we give a description of the algorithm used by the Monte Carlo generator in order to obtain a final NLO result. In section 5 we discuss a few selected results and in section 6 we sum up our conclusions.

2. Effective operators

In the effective lagrangian formalism we assume that the SM is no more than a low energy limit of some more general theory to be revealed at TeV scale. In this context, the top-quark FCNC interaction is seen at low energies as a contact interaction that can be parametrized with higher dimension effective operators. The complete lagrangian is written as

$$\mathcal{L} = \mathcal{L}^{SM} + \frac{1}{\Lambda} \mathcal{O}^5 + \frac{1}{\Lambda^2} \mathcal{O}^6 + \mathcal{O}\left(\frac{1}{\Lambda}\right) \quad (1)$$

where we have truncated the expansion series at dimension $d = 6$. We follow the formalism in [6], where the SM symmetries are imposed and all operators are built using the already known SM fields. This results in a reduced list of the possible dimension six operators. Additionally, baryon and lepton number conservation is imposed, which restricts the operators to dimension six only. Even imposing the SM symmetries, the number of all possible FCNC operators is quite vast. However, the list can be reduced by using appropriate equations of motion and Fierz transformations applied to the physical operators we are dealing with. The final minimal complete set of FCNC operators for top-quark physics was obtained in [9, 10, 11, 12]. With this procedure, a single operator survives in the strong sector [11]. Following the notation in [6] this operator can be written as

$$\mathcal{O}_{uG\phi}^{ij} = \bar{q}_L^i \lambda^a \sigma^{\mu\nu} u_R^j \tilde{\phi} G^{a\mu\nu} \quad (2)$$

where $G_{\mu\nu}^a$ is the gluonic field tensor, u^i right-handed u or c -quark field and q^i is the left-handed quark doublet. The same operator also appears in the literature as a dimension 5 operator

$$i\kappa_i \frac{g_s}{\Lambda} \bar{q}_i \lambda^a \sigma^{\mu\nu} (f_i + h_i \gamma_5) t G_{\mu\nu}^a \quad , \quad (3)$$

where κ_i is a real parameter, $i = u, c$, g_s is the strong coupling constant and f_i and h_i are complex numbers which respect $|f_i|^2 + |h_i|^2 = 1$. This operator is responsible for the effective FCNC vertices tgu and tgc . The minimum set of operators in the electroweak sector is

$$\mathcal{O}_{uB\phi}^{ij} = \bar{q}_L^i \sigma^{\mu\nu} u_R^j \tilde{\phi} B_{\mu\nu} \quad , \quad \mathcal{O}_{uW\phi}^{ij} = \bar{q}_L^i \tau_I \sigma^{\mu\nu} u_R^j \tilde{\phi} W_{\mu\nu}^I \quad , \quad (4)$$

$$\mathcal{O}_{\phi u}^{ij} = i(\phi^\dagger D_\mu \phi) (\bar{u}_R^i \gamma^\mu u_R^j) \quad , \quad (5)$$

$$\mathcal{O}_{\phi q}^{(1),ij} = i(\phi^\dagger D_\mu \phi) (\bar{q}_L^i \gamma^\mu q_L^j) \quad , \quad \mathcal{O}_{\phi q}^{(3),ij} = i(\phi^\dagger \tau_I D_\mu \phi) (\bar{q}_L^i \gamma^\mu \tau_I q_L^j) \quad , \quad (6)$$

$$\mathcal{O}_{u\phi}^{ij} = (\phi^\dagger \phi) (\bar{q}_L^i u_R^j \tilde{\phi}) \quad , \quad (7)$$

where $B^{\mu\nu}$ and $W_{\mu\nu}^I$ are the $U(1)_Y$ and $SU(2)_L$ field tensors, respectively. The field ϕ is the usual Higgs boson doublet. In addition to this electroweak set of operators, we have included subsets of four-fermion operators which are described in [13].

3. Physical Processes

The physical processes included in METop are FCNC direct top production and FCNC single top production. Direct top is available both at LO and at NLO while FCNC single top is available only at LO. All processes can be generated for both pp and $p\bar{p}$ hadron colliders. Furthermore, the full top decay chain is included so that spin correlations are preserved. Also, in order to perform spin correlations studies, a on/off switch for the top decay chain was made available. In figures 1 and 2 we show the generic Feynman diagrams for the physical processes implemented in METop. The production process depicted in figure 1 has only strong FCNC operator contributions. Born level FCNC direct top is shown in figure 1 (left) together with the $t+g$ process (right). In figure 2 we present the diagrams for the LO $t+q$ production process from gluon fusion (right) and with quarks in the initial state (left). The former is only affected by strong FCNC operators while the later has contributions from the strong, electroweak and four-fermion sectors.




Figure 1. FCNC leading order direct top production and top + gluon production at the parton level. Only FCNC strong operators contribute to the process.




Figure 2. FCNC leading order top + quark production at the parton level. FCNC strong and electroweak operators contribute to the process together with 4F operators.

4. NLO approximation

As mentioned in the previous section, METop is able to generate FCNC direct top events with NLO precision. The implementation was performed by adopting an NLO effective approximation [14]. A general NLO calculation of an n -particles final state process can be represented by

$$d\sigma_{NLO} = d\sigma_B d\Phi_n + d\sigma_V d\Phi_n + d\sigma_R d\Phi_{n+1}, \quad (8)$$

where B stands for Born, V for Virtual and R for Real. It is well known that the virtual term has infrared divergences that will only be cured by the inclusion of the real radiation term σ_R . We show in figure 3 the class of diagrams contributing to NLO direct top production. The first process $qg \rightarrow t$ correspond to the Born term. The second, $qg \rightarrow t$, corresponds to the Virtual term while the third, $qg \rightarrow gt$, is the real radiation term. This last term gives rise to an infrared divergence which correspond to a low P_T and/or to a collinear external gluon. Kinematically, the divergence can be parametrized by the gluon transverse momentum, P_T , and




Figure 3. Inclusive FCNC direct top production at NLO in QCD.

it obviously occurs in the limit $P_T \rightarrow 0$. In order to avoid the divergence one could just impose a P_T cut. The resulting region has to be excluded from the event space generation with some care - since the problem arises only in the collinear/soft limit, we can fill this phase space region with a branching mechanism. By doing so, we assume a collinear factorization, where the QCD radiation is emitted by one of the external legs of the Born configuration. This approximation can be translated by

$$|M_{t+g}|^2 d\Phi_2 \rightarrow |M_t|^2 d\Phi_1 \frac{\alpha_S}{2\pi} \frac{dt}{t} P_{q,gg}(z) dz \frac{d\phi}{2\pi} \quad (9)$$

where M_{t+g} and M_t are the real radiation for $(t+g)$ and direct top amplitudes, respectively; $d\Phi_i$ is the phase space for the i th body processes and $P_{q,gg}(z)$ is the Altarelli-Parisi splitting function. The variable t is a resolution parameter which in our case is the top quark transverse momentum P_T^{top} . The hard P_T region will then be filled by using the appropriate transition amplitude and the low P_T region by the use of a shower mechanism. In this NLO effective approximation [14], the P_T phase-space is then divided in two regions parametrized by P_T^{top} , which we call P_T^{match} . This parameter plays the role of a matching parameter where the criterion for a good transition is the smoothness of the final P_T distribution. The final cross section free of divergences can be written as

$$\sigma_{NLO} = K\sigma_B(P_T^{PS} < P_T^{match}) + \sigma_R(P_T > P_T^{match}), \quad (10)$$

where P_T^{PS} is the transverse momentum of the parton shower emission. The contribution from the virtual term $d\sigma_V$ in equation 8 is included via a K-factor applied to the born term and because born and virtual terms share the same $2 \rightarrow 1$ topology we assume they have similar kinematics. It is from equation 10 that events in METop are generated. Since METop is a parton level generator, the events must then be submitted to a parton shower simulator in order to include initial (ISR) and final (FSR) state radiation. Therefore, since these parton showers are based on a branching mechanism, we leave the collinear and/or soft P_T phase space region to be filled by the corresponding parton shower. In order to avoid double counting we just have to ensure that the parton shower will not populate the $P_T > P_T^{match}$ region in the first emission. To prevent it, the shower mechanism must be performed with the PT-ordered scheme with the first emission starting at $P_T = P_T^{match}$. For the NLO direct top process, METop assumes that the Pythia shower will be used with its PT-ordered scheme [15] flag on. Finally, we note that because this is a $2 \rightarrow 1$ process, the shower mechanism implemented in Pythia will not add FSR but only ISR. This does not pose any problems because FSR is negligible when compared to ISR due the large top-quark mass.

5. The generation process and some results

In figure 4 we show the diagram flow of the MEtop generation process. It was drawn specifically for the NLO direct top case. The remaining processes follow the same flow except for the K-factor calculation step. The amplitudes used to evaluate the cross sections were generated with CalcHEP [16], whit Feynman rules derived with LanHEP [17] and all integrations were performed using the Cuba Library [18]. The K-factors are calculated "on the fly" and therefore each sub process has its own normalization factor.




Figure 4. MEtop flow cart for NLO Direct top generation.

In figure 5 we present the top-quark P_T distribution for direct top after the first emission and with the starting scale shower at m_t (black solid line). The blue dashed line represents the top P_T distribution of the real radiation process. As explained before, both regions overlap and therefore a matching must be introduced in order to avoid double counting. The matching process must be chosen so that the transition between the rescaled direct top distribution and the real radiation process is smooth. In figure 6 we show the top P_T distribution for $P_T^{match} = 10, 15$ Gev where direct top is the gray dashed line and $t + g$ is the blue dashed line. The first one corresponds to the top distribution after the first ISR emission, where the PT-shower starting scale was set to 10 and 15 Gev, respectively; the second one is the top distribution from the 't+g' process where a P_T^{cut} of 10 and 15 Gev was imposed. We have studied several P_T^{match} scenarios in the 5 Gev to 20 Gev range finding no significant differences. We therefore recommend that this matching value should be included in any analysis as a systematic error.

In figure 7 we show the final NLO direct top quark P_T and η distribution (solid line), as well as the LO result (dashed line) after full ISR, FSR and Multiple Interactions (MP). It is clear that in both distributions the final NLO result does not amount to simply multiplying the LO result by a K-factor. In fact, the distributions show that the NLO contribution concentrates more events in the low P_T region and produces top quarks at higher angles. In figure 8 we show the equivalent distribution for the lepton coming from the decaying W. The b -quark and neutrino distributions show similar differences between the NLO and LO results. For the top decay products, the NLO contribution does not result in a major change in distributions making a K-factor adjustment suitable. However, one must pay special attention to analysis where the top quark momentum is reconstructed. For example, analysis which include the study of the top spin polarisation usually requires full reconstruction of its momentum. In this case, figure 7 show that NLO events should be considered. Finally, when studying the inclusive NLO direct top production one must add the $t + jet$ (single top FCNC) process as well. The distribution of




Figure 5. P_T distribution of the top quark for $\sqrt{s} = 7$ TeV. The black solid line is for direct top production after the first branching in ISR, with starting scale of m_t . The blue dashed line is for the hard process top+gluon production.




Figure 6. P_T distribution of top quark after the first ISR branching with a P_T^{match} of 10 GeV (left) and 15 GeV (right).




Figure 7. Comparison of the LO and NLO P_T (left) and η (right) distributions of the top quark at the partonic level after the full shower (ISR+FSR) and Multiple Interaction.

the sum of the two processes is shown in figure 9 for the top P_T (right) and the top η (left). The FCNC Single top process was generated with a $P_T^{cut} = 10$ GeV and only subprocesses where a




Figure 8. Comparison of the LO and NLO P_T (left) and η (right) distributions of the lepton from $t \rightarrow bW \rightarrow bl\nu$ at the partonic level after the full shower (ISR+FSR) and Multiple Interaction.

FCNC interaction takes place were included. Processes like $u\bar{d} \rightarrow t\bar{b}$ which are pure SM single top production were discarded. In tables 1 and 2 we show the total contributions from each LO and NLO processes. They were calculated for $k_{u,c} = 0.01$ and for LHC @ 7 TeV. As we can see from figure 9, the FCNC single top contribution does not produce considerable shape modifications, although its contribution to the total cross section is approximately 13%.




Figure 9. P_T (left) and η (right) distributions of the top quark for NLO direct top (solid line) and NLO direct top plus $pp \rightarrow tq$ with $P_T^{\text{match}} = 10$ GeV and jet $p_T > 10$ GeV.

Table 1. FCNC Direct top (anti-top) LO and NLO cross sections for $k_{u,c} = 0.01$ and LHC @ 7 TeV.

Subprocess	LO	NLO
$ug \rightarrow t$	6.12	8.74
$cg \rightarrow t$	0.91	1.67

6. Conclusions

We have presented a new Monte Carlo generator dedicated to top FCNC physics. We have included the main FCNC top production channels: direct top and single top. The direct top channel is available at NLO and the single top at LO. A complete set of dimension six operators

Table 2. FCNC $t + q$ with $k_{u,c} = 0.01$ and LHC @ 7 Tev and $PT_{cut} = 10$ Gev.

Subprocess	$k_u = 0.01$	$k_c = 0.01$
$p, p \rightarrow t, q$	1.12	0.40

are already included. We have shown that the NLO top distributions obtained for the direct top cannot be built via a K-factor from the LO ones. Therefore NLO events should be used especially in the cases where top-quark momentum reconstruction is required.

Acknowledgments

This work is partially supported by the Portuguese *Fundação para a Ciência e a Tecnologia* (FCT) under contracts CERN/FP/123619/2011 and PTDC/FIS/117951/2010. RS is also partially supported by an FP7 Reintegration Grant, number PERG08-GA-2010-277025 and by PEst-OE/FIS/UI0618/2011. RC is funded by FCT through the grant SFRH/BPD/45198/2008.

References

- [1] Glashow S L, Iliopoulos J and Maiani L 1970 Weak Interactions with Lepton-Hadron Symmetry *Phys. Rev. D* **2** 1285
- [2] Aguilar-Saavedra J A 2004 Top flavor-changing neutral interactions: Theoretical expectations and experimental detection *Acta Phys. Polon. B* **35** 2695 [arXiv:hep-ph/0409342].
- [3] Liu J J, Li C S, Yang L L and Jin L G 2005 Next-to-leading order QCD corrections to the direct top quark production via model-independent FCNC couplings at hadron colliders *Phys. Rev. D* **72** 074018
- [4] Slabospitsky S R and Sonnenschein L 2002 TopReX generator (version 3.25): Short manual *Comput. Phys. Commun.* **148** 87
- [5] Aguilar-Saavedra J A 2010 Zt, gamma t and t production at hadron colliders via strong flavour-changing neutral couplings *Nucl. Phys. B* **837** 122
- [6] Buchmüller W and Wyler D 1986 Effective Lagrangian Analysis of New Interactions and Flavor Conservation *Nucl. Phys. B* **268** 621
- [7] Allwall J *et al.* 2007 A standard format for Les Houches event files *Comput. Phys. Commun.* **176** 300
- [8] Sjostrand T, Mrenna S and Skands P 2006 PYTHIA 6.4 Physics and Manual *J. High Energy Phys.* JHEP0605(2006)026
- [9] Ferreira P M, Oliveira O and Santos R 2006 Flavour changing strong interaction effects on top quark physics at the LHC *Phys. Rev. D* **73** 034011
- [10] Ferreira P M and Santos R 2006 Strong flavour changing effective operator contributions to single top quark production *Phys. Rev. D* **73** 054025
- [11] Aguilar-Saavedra J A 2009 A Minimal set of top anomalous couplings *Nucl. Phys. B* **812**, 181
- [12] Grzadkowski B, Iskrzynski M, Misiak M and Rosiek J 2010 Dimension-Six Terms in the Standard Model Lagrangian *J. High Energy Phys.* JHEP1010(2010)085
- [13] Coimbra R, Onofre A, Santos R and Won M 2012 METop - a generator for single top production via FCNC interactions *Eur. Phys. J. C* **72** 2222
- [14] Boos E E, Bunichev V E, Dudko L V, Savrin V I and Sherstnev A V 2006 Method for simulating electroweak top-quark production events in the NLO approximation: SingleTop event generator, *Phys. Atom. Nucl.* **69** 1317
- [15] Sjostrand T and Skands P 2005 Transverse-momentum-ordered showers and interleaved multiple interactions *Eur. Phys. J. C* **39** 129 [hep-ph/0408302].
- [16] Pukhov A 2004 CalcHEP 2.3: MSSM, structure functions, event generation, batches, and generation of matrix elements for other packages *Preprint* arXiv:hep-ph/0412191
- [17] Semenov A 1998 LanHEP: A package for automatic generation of Feynman rules from the Lagrangian *Comput. Phys. Commun.* **115** 124
Semenov A 2009 LanHEP - a package for the automatic generation of Feynman rules in field theory. Version 3.0 *Comput. Phys. Commun.* **180** 431
- [18] Hahn T 2005 CUBA: A library for multidimensional numerical integration *Comput. Phys. Commun.* **168** 78