

BIODEGRADATION OF THE FUNGICIDE METALAXYL BY ZYGOMYCETES

P77

**M. R. Martins¹, R. S. Ribeiro¹, C. Santos²,
J. Cruz-Morais¹ and N. Lima²**

¹ ICAAM & Chemistry Department, School of Sciences and Technology, University of Évora, Évora, Portugal

² IBB/Biological Engineering Centre, University of Minho, Braga, Portugal

e-mail: mrm@uevora.pt

A large number of fungicides, such as metalaxyl [methyl N-(methoxyacetyl)-N-(2,6-xylyl)-DL-alaninate], are used against Oomycete species that cause downy mildew in agricultural crops, including grapes. However, these compounds are potentially harmful for terrestrial and aquatic environments, and to human health due to carcinogenic and mutagenic properties. There is an increase in interest in using filamentous fungi, such as members of the *Zygomycetes* which degrade xenobiotic compounds using nonspecific extracellular enzymes for the bioremediation of pesticides in polluted soils.

In this study, several *Zygomycetes* including *Gongronella*, *Absidia*, *Circinella* and *Rhizopus* species, were used to screen degradation of metalaxyl using solid and liquid cultures. In order to identify and characterise these taxa a polyphasic approach including morphology characterization, molecular fingerprint M13-PCR and Matrix Assisted Laser Desorption Ionization Time of Flight Intact Cell Mass Spectrometry (MALDI-TOF ICMS) was used. After selective enrichment on solid medium containing a fungicide gradient concentration of 0-100 mg.L⁻¹, *Absidia glauca* (CBS 101.08) and *Rhizopus orizae* (CCMI 900) showed a capacity to tolerate high metalaxyl concentrations. These resistant strains were selected to perform liquid assays using Yeast Nitrogen liquid cultures supplemented with sucrose (5 g.L⁻¹) and metalaxyl (100 mg.L⁻¹). Biomass concentration was determined by dry weight. The *A. glauca* and *R. orizae* strains showed specific growth rates of 0.774 h⁻¹ and 0.999 h⁻¹, respectively. Sucrose was completely consumed within 5 d with a sucrose consumption rate of 0.93 and 0.84 g.L⁻¹.day⁻¹ for *A. glauca* and *R. orizae*, respectively. The non-degraded metalaxyl in liquid cultures was determined by UV-HPLC and evaluated periodically for 21 d. The metalaxyl degradation rate for *A. glauca* and *R. orizae* was 2.22 mg.L⁻¹.day⁻¹ and 2.29 mg.L⁻¹.day⁻¹, respectively.

Results suggest that *A. glauca* (CBS 101.08) and *R. orizae* (CCMI 900) can be used in soil bioremediation experimentation for metalaxyl degradation. These strains are now under study to determine the presence of extracellular enzymes involved in the process.

BIOLOGICAL RESOURCE CENTRES

Closing the gap
between science
and society

ECCO XXXI MEETING

ABSTRACTS BOOK

Edited by
Russell Paterson
Marta F. Simões
Leonel Pereira
Cledir Santos
Nelson Lima

Biological Resource Centres

Closing the gap between
science and society

Edited by

Russell Paterson
Marta F. Simões
Leonel Pereira
Cledir Santos
Nelson Lima

ECCO XXXI Meeting
Abstracts Book



Abstracts of the 31st European Culture Collections'
Organization Meeting, Universidade do Minho, Braga, Portugal,
14-15 June 2012.



Biological Resource Centres **Closing the gap between science and society**

Editors: Russell Paterson,
Marta Simões,
Leonel Pereira,
Cledir Santos,
Nelson Lima

Published by: Micoteca da Universidade do Minho

Printed: Candeias Artes Gráficas | Braga | Portugal

Depósito Legal: 345229/12

ISBN: 978-972-97916-5-9

Production run: 200 copies

All rights are reserved to the Micoteca da Universidade do Minho, however the abstracts may be freely reviewed, abstracted, reproduced or translated, in part or in whole, but not for sale or use in conjunction with commercial purposes. The views expressed or implied in this publication, unless otherwise noted, should not be interpreted as official positions of the Micoteca da Universidade do Minho