

Use of Matrix-assisted Laser Desorption/Ionization-Time of Flight Mass Spectrometry (MALDI-TOF MS) for the identification of beer spoilage bacteria.

Anneleen WIEME^{1,2},

Doctoraatspromotoren: Prof. Dr. Anita VAN LANDSCHOOT¹ and Prof. Dr. Peter VANDAMME²

¹ University College Ghent (Association University Ghent), Faculty Applied Bioscience Engineering, Laboratory of Biochemistry and Brewing, Valentin Vaerwyckweg 1, 9000 Ghent, Belgium, 092424236 – Anneleen.Wieme@Hogent.be

² Ghent University, Department of Biochemistry and Microbiology, Laboratory of Microbiology, K.L. Ledeganckstraat 35, BE-9000 Ghent, Belgium, 092645114

Beer is a beverage with good microbiological stability because it contains almost no oxygen and nutrients for bacterial growth. In addition, low pH, high CO₂-content and the presence of ethanol and antibacterial hop compounds ensure microbial stability. Nevertheless, beer spoilage induced by bacteria is a common problem in the brewing industry and these spoilage bacteria typically cause visible turbidity, acidity and off-flavours. Currently, these bacteria are detected with culture-dependent methods using selective media or with faster identification methods such as DNA-typing, ribotyping and other PCR-based techniques. These approaches are notoriously laborious, expensive, time-consuming and moreover, lack specificity and sensitivity. The present study aims to develop a quick, specific and inexpensive method to detect and identify beer spoilage bacteria in the brewing industry. To achieve this, an extensive database comprising MALDI-TOF MS-profiles of more than 260 established and accurately identified contaminants and beer spoilage strains was built. In addition to these strains, strains of the same species originating from other niches, besides spoiled beer, were also included in order to encompass the phenotypic diversity of the spoilage species. Among others, strains of *Lactobacillus brevis* (29), *Lb. lindneri* (3), "*Lb. brevisimilis*" (1), *Lb. buchneri* (5), *Lb. coryniformis* (1), *Lb. plantarum* (8), *Lb. parabuchneri* (15), *Lb. paracollinoides* (2), *Lb. perolens* (10), *Pediococcus damnosus* (9), *P. inopinatus* (10), *Pectinatus cerevisiiphilus* (1), *P. frisingensis* (2), *Selenomonas lacticifex* (1), *Megasphaera cerevisiae* (2), and *Zymophilus raffinovorans* (2) were included in the database. The resulting set of profiles (\pm 6500 good quality profiles) allowed the assignment of reproducible species-specific biomarker peaks for all spoilage species. All strains were not only cultured at species-specific conditions (type medium, growth temperature, oxygen requirements, growth time), but also on selective and non-selective media. Different media are used to enable the exclusion of medium-associated peaks from the species-specific biomarker peaks. Consequently, identification of novel beer spoilage isolates can be easily and rapidly performed. Nevertheless, the final aim of this research is to detect and identify these bacteria in a spoiled sample with minimal time-consuming culture steps.

Financial support was granted by the Research Fund of University College Ghent.