

University of Groningen

Real-world Data Reveal Long Drug Survival for Guselkumab in Patients with Plaque Psoriasis

BioCAPTURE consortium; van Muijen, Marloes E.; Thomas, Sarah E.; Vellinga, Douwe; Bouwman, Silke; van Doorn, Martijn B.A.; Politiek, Klazien; Otero, Marisol E.; van den Reek, Juul M.P.A.; de Jong, Elke M.G.J.

Published in:
Acta dermato-venereologica

DOI:
[10.2340/actadv.v102.685](https://doi.org/10.2340/actadv.v102.685)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

BioCAPTURE consortium, van Muijen, M. E., Thomas, S. E., Vellinga, D., Bouwman, S., van Doorn, M. B. A., Politiek, K., Otero, M. E., van den Reek, J. M. P. A., & de Jong, E. M. G. J. (2022). Real-world Data Reveal Long Drug Survival for Guselkumab in Patients with Plaque Psoriasis. *Acta dermato-venereologica*, 102, [adv00755]. <https://doi.org/10.2340/actadv.v102.685>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Real-world Data Reveal Long Drug Survival for Guselkumab in Patients with Plaque Psoriasis

Marloes E. VAN MUIJEN^{1,2}, Sarah E. THOMAS^{1,2}, Douwe VELLINGA³, Silke BOUWMAN⁴, Martijn B. A. VAN DOORN⁵, Klazien POLITIEK⁶, Marisol E. OTERO^{1,2}, Juul M. P. A. VAN DEN REEK^{1,2}, Elke M. G. J. DE JONG^{1,2,7} and the BioCAPTURE consortium
¹Department of Dermatology and ²Radboud Institute for Health Sciences, Radboud University Medical Centre, Geert Grooteplein Zuid 10, Postbus 9101, NL-6525 GA Nijmegen, Departments of Dermatology: ³Alrijne Ziekenhuis, Leiderdorp, ⁴University Medical Centre Groningen, Groningen, ⁵Erasmus University Medical Centre, Rotterdam and ⁶Medisch Centrum Leeuwarden, Leeuwarden, and ⁷Radboud University, Nijmegen, The Netherlands. E-mail: Marloes.vanMuijen@radboudumc.nl

Accepted Jul 5, 2022; Epub ahead of print Jul 5, 2022

Acta Derm Venereol 2022; 102: adv00755. DOI: 10.2340/actadv.v102.685

Guselkumab has been registered as the first interleukin-23 (IL-23) inhibitor for treatment of psoriasis. Randomized controlled trials (RCTs) have shown a favourable efficacy and safety profile for guselkumab (1, 2). However, RCTs may not adequately reflect the real-world situation (3). The primary objective of this real-world observational multicentre study was to evaluate 1- and 2-year drug survival (DS) of guselkumab, split for discontinuation due to ineffectiveness or side-effects. A further aim was to elucidate predictors for a shorter guselkumab DS.

METHODS AND RESULTS

A detailed description of the methods is given in Appendix S1. Data from patients with plaque psoriasis treated with guselkumab were collected from the prospective BioCAPTURE registry (www.biocapture.nl) and retrospective data from 4 other centres in the Netherlands (time-frame 2020 to 2021). Temporary treatment interruptions for any reason were allowed if <90 days. This 90-day gap was prolonged up to 1 year if patients discontinued due to fear of COVID-19 or due to remission. In the Kaplan–Meier analyses, 3 separate DS curves were created with an event for discontinuation in general (all reasons), due to ineffectiveness or to side-effects. Discontinuation due to an increase in musculoskeletal complaints in patients with psoriatic arthritis (PsA) was considered as an event in side-effect analyses. Univariable and multivariable Cox regression models were used to identify factors affecting DS.

Participating centres and patient and treatment characteristics are shown in Tables SI and SII, respectively. A total of 195 patients

(288.4 actively-treated patient years) were included; 110 (56.4%) were male, and 58 (29.7%) were biologic naive at guselkumab initiation. Forty (20.5%) patients had a rheumatologist-confirmed diagnosis of PsA. Six (3.1%) patients shortened the dosing interval, and 27 (13.8%) lengthened the interval.

Overall guselkumab DS rates after 1 and 2 years were 85.5% and 77.8%, respectively. One- and 2-year DS rates for discontinuation related to ineffectiveness were 92.8% and 88.7%, and for discontinuation related to side-effects were 94.3% and 92.1%, respectively (**Fig. 1**). The outputs of the Cox regression analyses are shown in Table SIII.

The multivariable model showed a significant association between diabetes mellitus type 2 (DMt2) and a shorter DS (hazard ratio (HR) 3.69 (95% confidence interval (95% CI) 1.14–11.98) ($p=0.030$) due to ineffectiveness. Multivariable analyses for predictors of side-effect-related DS showed a significant association for a shorter DS in patients with PsA (HR 7.51 (95% CI 2.26–24.95) ($p=0.001$)).

DISCUSSION

This study shows that 1- and 2-year DS for guselkumab was high, both for discontinuation due to side-effects and ineffectiveness. The latter finding is notable, as in previous literature higher discontinuation rates due to ineffectiveness have been described for other types of biologics (4). Previous studies on guselkumab DS in real-world settings have also reported high first-year DS (ranging from 68.0% (5) to 95.0% (6)), although sample size was often small, and the event definition and duration of follow-up varied (5–11).

A substantial number of patients in this study ($n=27$, 13.8%) used a lengthened dosing interval, which suggests that, for guselkumab, high therapeutic effectiveness can be maintained even on a lower dose. In ongoing studies on guselkumab for psoriasis, the use of a prolonged dosing interval is currently being evaluated (12, 13).

Having PsA was associated with a shorter DS due to side-effects. It should be noted that the association between side-effect-related discontinuation and PsA was largely explained by patients with pre-existent PsA who experienced an increase in musculoskeletal complaints. In contrast, a systematic review on predictors of persistence for other biologics, described having PsA as predictive for longer survival (14). Furthermore, we found an association between DMt2 and a higher risk of discontinuation due to ineffectiveness. In support of our




Fig. 1. Kaplan–Meier drug survival analysis of guselkumab during 2 years, split for reason of discontinuation.

findings, the Corrona psoriasis registry has previously reported that diabetes reduced the risk of achieving various biologic treatment goals (15).

A strength of this study is the large study population, and high external validity due to the multicentre design. Due to the COVID-19 pandemic, there were fewer clinical visits during the study period and more treatment interruptions due to fear of COVID-19. These interruptions were handled differently (see Appendix S1), leading to a more realistic reflection of DS in non-COVID-19 time-frames.

In conclusion, this study found a high 1- and 2-year DS for guselkumab. Reassuringly, discontinuation due to ineffectiveness or side-effects was very uncommon. Having DMt2 was associated with a shorter DS due to ineffectiveness, whereas having PsA was associated with a shorter DS due to side-effects. A substantial proportion of patients (14%) was able to prolong their dosing interval.

ACKNOWLEDGEMENTS

The authors would like to thank Eldrid Schoonhoven, Janneke Huizinga and Barbara Horváth for their contribution to data entry.

Conflicts of interest: MEvM carries out clinical trials for AbbVie, Celgene, Janssen and Novartis, and has received a speaking fee from Janssen. All funding is not personal, but goes to the independent Research Fund of the Department of Dermatology of the Radboud University Medical Centre Nijmegen (Radboudumc), The Netherlands. DV carried out clinical trials for Novartis and attended advisory boards from AbbVie, Almirall, Janssen, Novartis, Leo Pharma and UCB. MBAvD has received consulting fees or honorarium from Novartis, AbbVie, Pfizer, Leo Pharma, Sanofi, Lilly, Janssen and Celgene, has received a grant and payment for lectures including service on speakers bureaus from Novartis, Sanofi and Janssen. KP has attended advisory boards for Sanofi, Leo Pharma and AbbVie, and has received reimbursement for organizing a symposium from AbbVie. MEO has acted as consultant for Lilly. SRPD has attended advisory boards for AbbVie, Janssen and Leo Pharma, and has received a congress fee from AbbVie. RAT has attended advisory boards from Leo Pharma, Lilly, and Novartis. PPMvL has received funding from Wyeth for research and carried out clinical trials for Abbott and Janssen. P. M. van Lümig has received speaking and consulting fees from Wyeth and Schering-Plough and has received reimbursement for attending a symposium from Schering-Plough and Pfizer. P. M. van Lümig has attended advisory boards for AbbVie, Leo Pharma, Novartis and UCB. JMPAvdR carried out clinical trials for AbbVie, Celgene and Janssen and has received speaking fees/attended advisory boards from AbbVie, Janssen, BMS, Almirall, LEO Pharma, Novartis, UCB and Eli Lilly and reimbursement for attending a symposium from Janssen, Pfizer, Celgene and AbbVie. All funding is not personal, but goes to the independent research fund of the Department of Dermatology of Radboudumc Nijmegen, the Netherlands. EMGJdJ has received research grants for the independent research fund of the department of dermatology of the Radboudumc Nijmegen, the Netherlands from AbbVie, Novartis, Janssen Pharmaceutica and Leo Pharma. Has acted as consultant and/or paid speaker for and/or participated in research sponsored by companies that manufacture drugs used for the treatment of psoriasis including AbbVie, Janssen Pharmaceutica, Novartis, Lilly, Celgene, Leo Pharma, UCB and Almirall. All funding is not personal, but goes to the independent research fund

of the Department of Dermatology of Radboudumc, Nijmegen, the Netherlands.

REFERENCES

1. Reich K, Griffiths CEM, Gordon KB, Papp KA, Song M, Ranzazzo B, et al. Maintenance of clinical response and consistent safety profile with up to 3 years of continuous treatment with guselkumab: results from the VOYAGE 1 and VOYAGE 2 trials. *J Am Acad Dermatol* 2020; 82: 936–945.
2. Reich K, Papp KA, Armstrong AW, Wasfi Y, Li S, Shen YK, et al. Safety of guselkumab in patients with moderate-to-severe psoriasis treated through 100 weeks: a pooled analysis from the randomized VOYAGE 1 and VOYAGE 2 studies. *Br J Dermatol* 2019; 180: 1039–1049.
3. Mason KJ, Barker J, Smith CH, Hampton PJ, Lunt M, McElhone K, et al. Comparison of drug discontinuation, effectiveness, and safety between clinical trial eligible and ineligible patients in BADBIR. *JAMA Dermatol* 2018; 154: 581–588.
4. Lin PT, Wang SH, Chi CC. Drug survival of biologics in treating psoriasis: a meta-analysis of real-world evidence. *Sci Rep* 2018; 8: 16068.
5. Schwensen JFB, Nielsen VW, Nissen CV, Sand C, Gniadecki R, Thomsen SF. Effectiveness and safety of guselkumab in 50 patients with moderate to severe plaque psoriasis who had previously been treated with other biologics: a retrospective real-world evidence study. *J Eur Acad Dermatol Venereol* 2021; 35: e341–e343.
6. Iznardo H, Vilarrasa E, López-Ferrer A, Puig L. Real world drug survival of guselkumab, ixekizumab and secukinumab for psoriasis. *Br J Dermatol* 2021; 185: 660–662.
7. Lee EB, Reynolds KA, Pithadia DJ, Egeberg A, Wu JJ. Drug survival of guselkumab for psoriasis in a real-world setting: a single-center retrospective chart review. *J Dermatolog Treat* 2020; 31: 342–343.
8. Dapavo P, Siliquini N, Mastorino L, Avallone G, Merli M, Agostini A, et al. Efficacy, safety and drug survival of IL-23, IL-17 and TNF-alpha inhibitors for psoriasis treatment: a retrospective study. *J Dermatolog Treat* 2022; 33: 2352–2357.
9. Ruiz-Villaverde R, Rodriguez-Fernandez-Freire L, Armario-Hita JC, Pérez-Gil A, Galán-Gutiérrez M. Guselkumab: mid-term effectiveness, drug survival, and safety in real clinical practice. *Dermatol Ther* 2021; 34: e14798.
10. Torres T, Puig L, Vender R, Lynde C, Piaserico S, Carrascosa JM, et al. Drug survival of IL-12/23, IL-17 and IL-23 inhibitors for psoriasis treatment: a retrospective multi-country, multicentric cohort study. *Am J Clin Dermatol* 2021; 22: 567–579.
11. Lytvyn Y, Zaaroura H, Mufti A, AlAbdulrazzaq S, Yeung J. Drug survival of guselkumab in patients with plaque psoriasis: a 2 year retrospective, multicenter study. *JAAD Int* 2021; 4: 49–51.
12. van der Schoot LS, van den Reek J, Grine L, Schots L, Kievit W, Lambert JLW, et al. Dose reduction of the new generation biologics (IL-17 and IL-23 inhibitors) in psoriasis: study protocol for an international, pragmatic, multicenter, randomized, controlled, non-inferiority study—the BeNeBio study. *Trials* 2021; 22: 707.
13. Eyerich K, Weisenseel P, Pinter A, Schäkel K, Asadullah K, Wegner S, et al. IL-23 blockade with guselkumab potentially modifies psoriasis pathogenesis: rationale and study protocol of a phase 3b, randomised, double-blind, multicentre study in participants with moderate-to-severe plaque-type psoriasis (GUIDE). *BMJ Open* 2021; 11: e049822.
14. Mourad A, Straube S, Armijo-Olivo S, Gniadecki R. Factors predicting persistence of biologic drugs in psoriasis: a systematic review and meta-analysis. *Br J Dermatol* 2019; 181: 450–458.
15. Enos CW, Ramos VL, McLean RR, Lin TC, Foster N, Dube B, et al. Comorbid obesity and history of diabetes are independently associated with poorer treatment response to biologics at 6 months: a prospective analysis in Corrona Psoriasis Registry. *J Am Acad Dermatol* 2022; 86: 68–76.