

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Acinetobacter baumannii

Whiteway, Clémence; Breine, Anke; Philippe, Chantal; Van der Henst, Charles

Published in: Trends in Microbiology

DOI: 10.1016/j.tim.2021.11.008

Publication date: 2022

Document Version Publisher's PDF, also known as Version of record

Link to publication

Citation for pulished version (HARVARD): Whiteway, C, Breine, A, Philippe, C & Van der Henst, C 2022, 'Acinetobacter baumannii', Trends in Microbiology, vol. 30, no. 2, pp. 199-200. https://doi.org/10.1016/j.tim.2021.11.008

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Trends in Microbiology | Microbe of the Month

Acinetobacter baumannii

Clémence Whiteway, ^{1,2} Anke Breine, ^{1,2} Chantal Philippe, ³ and Charles Van der Henst ^{1,2,*}

¹Microbial Resistance and Drug Discovery, VIB-VUB Center for Structural Biology, VIB, Flanders Institute for Biotechnology, Brussels, Belgium ²Structural Biology Brussels, Vrije Universiteit Brussel (VUB), Brussels, Belgium

³Research Unit in the Biology of Microorganisms (URBM), NARILIS, University of Namur (UNamur), Namur, Belgium



Acinetobacter baumannii is a ubiquitous, Gram-negative, nonflagellated coccobacillus bacterium commonly isolated from the environment. In human medicine, this opportunistic pathogen is responsible for hospitaland community-acquired infections. Resistance to last-resort antibiotics, such as colistin, tigecycline, and carbapenems, earns this bacterium a place among the most problematic nosocomial ESKAPE pathogens, being classified as a bacterium for which research and development of new antibiotics is critically needed by the World Health Organization (WHO) and as an urgent threat to public health by the Centers for Disease Control and Prevention (CDC). Its ability to form biofilms and resist desiccation and disinfectants is additionally alarming as these characteristics allow *A. baumannii* to thrive within hospital settings. The antibiotic resistance, the environmental persistence, along with the absence of identified host-damaging toxins in its genome, suggest that the virulence potential of *A. baumannii* is based on a 'persist and resist' strategy wherein these bacteria also resist complement-mediated killing and oxidative stress. At the genetic level, due to a plastic genome, we observe a high heterogeneity amongst isolates, adding complexity to the study of *A. baumannii* as an entity. Therefore, *A. baumannii* represents a modern and worldwide challenge which requires constant surveillance by the public health community.

KEY FACTS:

Isolation of extensively resistant and pan-drug-resistant strains ranked as 'high priority' by the WHO (global priority list of antibiotic-resistant bacteria to guide research, discovery and development of new antibiotics, 2017) and the CDC (antibiotic resistance threats in the USA, 2019).

Belongs to the 'ESKAPE' group of most problematic nosocomial pathogens.

Resistant to desiccation, disinfectant, oxidative stress, and complementmediated killing.

First described as a fast-growing extracellular bacterium; several strains show a potential for intracellular growth in eukaryotic cells.

High genetic diversity; rapidly evolving bacteria, with some isolates competent for natural transformation.

Lipooligosaccharide instead of classical lipopolysaccharide, with 14 variants identified so far.

At least 128 polysaccharide capsule types identified so far.

Presence of several secretion systems (SSs): T1SS, T2SS, and T6SS, but absence of T3SS and T4SS usually found in Gram-negative pathogens.

Phenotypic heterogeneity by 'phase variation' within clonal populations, with virulent opaque and capsulated (VIR-O) and avirulent translucent and less capsulated (AV-T) types identified.

DISEASE FACTS:

Nosocomial, community-acquired, and opportunistic infections.

Broad infection sites [skin and soft tissue, urinary tract, digestive and respiratory tracts, blood, central nervous system (meningitis), etc.].

*Correspondence: charles.vanderhenst@vub.vib.be (C. Van der Henst).




Trends in Microbiology, February 2022, Vol. 30, No. 2

© 2021 The Author(s). Published by Elsevier Ltd. https://doi.org/10.1016/j.tim.2021.11.008 This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Trends in Microbiology | Microbe of the Month

Acinetobacter baumannii (AB5075)



Trends in Microbiology

Declaration of interests

No interests were declared.

Literature

- 1. Sarshar, M. et al. (2021) Acinetobacter baumannii: an ancient commensal with weapons of a pathogen. Pathogens 10, 387
- 2. De Oliveira, D.M.P. et al. (2020) Antimicrobial resistance in ESKAPE pathogens. Clin. Microbiol. Rev. 33, e00181-19
- 3. Wong, D. et al. (2017) Clinical and pathophysiological overview of Acinetobacter infections: a century of challenges. Clin. Microbiol. Rev. 30, 409-447
- 4. Nowak, J. et al. (2017) High incidence of pandrug-resistant Acinetobacter baumannii isolates collected from patients with ventilator-associated pneumonia in Greece, Italy and Spain as part of the MagicBullet clinical trial. J. Antimicrob. Chemother. 72, 3277-3282
- 5. Chin, C.Y. et al. (2018) A high-frequency phenotypic switch links bacterial virulence and environmental survival in Acinetobacter baumannii. Nat. Microbiol. 3, 563-569
- 6. Wyres, K.L. et al. (2020) Identification of Acinetobacter baumannii loci for capsular polysaccharide (KL) and lipooligosaccharide outer core (OCL) synthesis in genome assemblies using curated reference databases compatible with Kaptive. Microb. Genom. 6, e000339
- 7. Gebhardt, M.J. et al. (2015) Joint transcriptional control of virulence and resistance to antibiotic and environmental stress in Acinetobacter baumannii, mBio 6, e01660-15
- 8. Gallagher, L.A. et al. (2015) Resources for genetic and genomic analysis of emerging pathogen Acinetobacter baumanii. J. Bacteriol. 197, 2027-2035
- 9. Harding, C.M. et al. (2018) Uncovering the mechanisms of Acinetobacter baumannii virulence. Nat. Rev. Microbiol. 16, 91-102

Can thrive in hospital settings, with a peculiar virulence based on a 'persist and resist' strategy.

Infection routes and factors favoring infections: colonization of mechanical devices such as catheters and ventilation equipment, open wounds, major trauma or burns, prolonged hospital stays, and immunocompromisation.

Therapeutic dead end because of last-resort antibiotic resistance.

TAXONOMY AND CLASSIFICATION: **KINGDOM:** Bacteria **PHYLUM:** Proteobacteria **CLASS:** Gammaproteobacteria **ORDER:** Pseudomonadales FAMILY: Moraxellaceae **GENUS:** Acinetobacter SPECIES: Acinetobacter baumannii



200 Trends in Microbiology, February 2022, Vol. 30, No. 2 © 2021 The Author(s). Published by Elsevier Ltd. https://doi.org/10.1016/j.tim.2021.11.008 This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/bv-nc-nd/4.0/).

