

1 Abstract

2 *Campylobacter* enteritis is the most reported zoonotic disease in many developed
3 countries where it imposes a serious health burden. *Campylobacter* transmission
4 to humans occurs primarily through the chicken vector. Chicks are regarded as a
5 natural host for *Campylobacter* species and are ~~mostly~~ colonized with *C. jejuni* in
6 particular. But despite carrying a very high bacterial load in their gastrointestinal
7 tract these birds, in contrast to humans, do not develop pathological signs. It
8 seems that in chickens *C. jejuni* principally harbours in the cecal mucosal crypts,
9 where an inefficient inflammatory response fails to clear the bacterium from the
10 gut. Recent intensive research resulted in an increased insight into the crosstalk
11 between *C. jejuni* and its avian host. This review discusses the chicken intestinal
12 mucosal immune response upon *C. jejuni* entrance, leading to tolerance and
13 persistent cecal colonization. It might in addition provide a solid base for further
14 research regarding this topic aiming to fully understand the host-bacterium
15 dynamics of *C. jejuni* in chicks and to develop effective control measures to clear
16 this zoonotic pathogen from poultry lines.

17

18 **Keywords:** *Campylobacter jejuni*; broiler chicken; immune response; tolerance;
19 persistent colonization

20

1 **Introduction**

2 From 2005 onwards, *Campylobacter* enteritis has been the most reported zoonotic
3 disease in many developed countries (EFSA, 2011). ~~And although mostly self-~~
4 ~~limiting, several sequelae might be developed, such as Guillain-Barré syndrome,~~
5 ~~reactive arthritis, irritable bowel syndrome and inflammatory bowel disease,~~
6 ~~which can eventually result in mortality (EFSA, 2010). Thus, campylobacteriosis~~
7 ~~poses a serious health burden in developed countries, where disease in humans is~~
8 ~~mostly caused by pathogenic *C. jejuni* strains (EFSA, 2011). Chickens are a~~
9 ~~natural host for *Campylobacter* spp. and are often colonized with *C. jejuni* in~~
10 ~~particular (EFSA, 2011). This review will therefore mainly focus on the~~
11 ~~interaction of *C. jejuni* with the chicken host. Despite being colonized in their~~
12 ~~ceca at a high degree, broiler chickens do not show typical signs of pathology and~~
13 ~~carry a high *C. jejuni* load until slaughter. As a consequence, slaughter and~~
14 ~~carcass processing of such animals results in the contamination of their meat~~
15 ~~products (Rosenquist et al., 2006), which are major sources for transmitting this~~
16 ~~pathogen to humans (EFSA, 2010). In addition, *C. jejuni* is frequently found in~~
17 ~~the intestines of broiler roosters and laying hens too (Cox et al., 2009). A decline~~
18 ~~in human cases of campylobacteriosis is not bound to happen as long as the poor~~
19 ~~understanding on the host bacterium interactions of *C. jejuni* in its chicken host~~
20 ~~hampers. Until recently to now,~~ the knowledge on the chicken immune response
21 in general ~~is~~ has been poor, hampering the development of control strategies to
22 eradicate *C. jejuni* from poultry animals (Hermans et al., 2011a). However,
23 intensive research in the past few years has resulted in an increased insight into

1 | the chicken immune response toward *C. jejuni* ~~entraneecolonization~~. This review
2 | discusses the ~~dual~~ interaction between *C. jejuni* and its chicken host, focussing on
3 | immune responses, leading to persistent, high-level cecal colonization. At the end
4 | of this review the mechanisms that are potentially responsible for the redirection
5 | ~~of~~ this response toward tolerance, and thus for the different disease outcome
6 | compared to humans, are handled.

7

8 | **Colonization pattern and antigenic variation of *Campylobacter*** 9 | ***jejuni* in chicks**

10 | Despite some reports of *Campylobacter*-induced diarrhea, systemic invasion,
11 | growth retardation and jejunal villus atrophy (Ruiz-Palacios et al., 1982; Sanyal et
12 | al., 1984; Sang et al., 1989; Lam et al., 1992; Lamb-Rosteski et al., 2008) it is
13 | generally accepted that *C. jejuni* colonizes the avian gut as a commensal.
14 | Colonization of chickens with *C. jejuni* does not cause clinical illness nor changes
15 | in cecal mucosa morphology even though large numbers of the bacterium reside
16 | ~~in their eecathere~~ (generally around 10^6 to 10^8 cfu/g), the predominant site for
17 | colonization (Beery et al., 1988; Van Deun et al., 2008; Meade et al., 2009b).
18 | Commensal bacteria in general do not colonize outside the gastrointestinal (GI)
19 | tract, but strangely enough *C. jejuni* can readily be found in various extraintestinal
20 | organs of broilers too. Up to seven days after oral and cloacal inoculation, the
21 | bacterium was found in the thymus, spleen, liver/gallbladder and bursa of
22 | Fabricius (Cox et al., 2005; Van Deun et al., 2008; Meade et al., 2009b). In a
23 | study with two-week-old chicks that were inoculated with *C. jejuni* at day-of

1 hatch, high bacterial numbers ($> 5 \log \text{CFU/g}$) were isolated from spleen and liver
2 of most of the birds (Lamb-Rosteski et al., 2008). In addition, *C. jejuni* was
3 isolated from the reproductive tract and ovarian follicles of laying hens (Cox et
4 al., 2009). The dissemination of *C. jejuni* to other organs seems to be correlated
5 with the invasive potential in primary cecal epithelial cells of chicks (Van Deun et
6 al., 2008), suggesting that *C. jejuni* translocates the epithelial barrier
7 transcellularly (through the chicken crypt epithelium) rather than paracellularly
8 (between cells).

9 Upon ingestion, *C. jejuni* reaches the cecum and multiplies, resulting in an
10 established colonizing population within 24 h after infection (Coward et al., 2008;
11 Smith et al., 2008). Most broiler flocks become colonized only at an age of two to
12 four weeks after which the infection rapidly spreads to almost all birds ($>95\%$),
13 which remain colonized until slaughter (Jacobs-Reitsma et al., 1995; Stern et al.,
14 2001; Stern, 2008; van Gerwe et al., 2009). Although not all birds in a flock were
15 colonized, it was demonstrated that *C. jejuni* can be isolated from laying hens
16 until an age of 42 weeks (Lindblom et al., 1986) and probably longer, since
17 experimental periods exceeding one year are not documented. This implies that *C.*
18 *jejuni* can evade the chicken host immune system. However, in a study by
19 Cawthraw and Newell (2010) colonization of breeder birds decreased over time,
20 indicating ~~and resist elimination by some~~ mucosal clearance. In addition, with
21 older birds it cannot be ruled out that replacement of one strain by an
22 immunologically distinct strain (strain succession) occurred, disguising mucosal
23 clearance of the former *C. jejuni* strain.

1 *Campylobacter*-positive flocks are often colonized with more than one sero-
2 or genotype at the same time (referred to as co-colonization), which may be
3 explained by recurring environmental exposure to the bacterium but also by
4 genetic changes within the *C. jejuni* population (van de Giessen et al., 1992;
5 Jacobs-Reitsma et al., 1995). The dominating strains are replaced throughout the
6 colonization period, probably due to strain-specific immune responses, and it
7 seems that this colonization pattern is mainly determined by the chicken host and
8 not by the host microbiota (Skanseng et al., 2007; Ridley et al., 2008). Indeed,
9 different breeds of chicken may differ in their susceptibility to colonization by *C.*
10 *jejuni* (Stern et al., 1990; Boyd et al., 2005). It has been suggested that a paternal
11 effect might be an important genetic factor influencing resistance to *C. jejuni*
12 colonization in broilers (Li et al., 2008). However, there are also other lines of
13 evidence suggesting that external factors are responsible for the *Campylobacter*
14 colonization pattern in broilers. It has been found in artificially inoculated birds
15 that different *C. jejuni* genotypes may compete for colonization leading to a *C.*
16 *jejuni* succession in broilers (Konkel et al., 2007).

17 *C. jejuni* isolates often show increased colonization potential after passage
18 through the chicken gut (Ringoir & Korolik, 2003). Chicken intestinal
19 colonization may favour genetic recombination in *C. jejuni*, resulting in different
20 *flaA* types, ribo- and PFGE patterns (Hanninen et al., 1999; Van Deun et al., 2007;
21 Hanel et al., 2009). Interstrain genetic exchange and intragenomic alterations were
22 shown to occur *in vivo*, even in the absence of selective pressure (de Boer et al.,
23 2002). It has been demonstrated that bacteriophage genes are known to be present

1 in the genome of *C. jejuni* and that phages can alter PFGE patterns of this
2 bacterium (Barton et al., 2007; Clark & Ng, 2008). Both phage-dependent and -
3 independent rearrangements of the genome result in an enormous antigenic
4 variation among *C. jejuni* isolates with the former resulting in phage-resistant *C.*
5 *jejuni* types (Scott et al., 2007a, 2007b). Besides protection against phage
6 predation, this generation of antigenic diversity may also play an important role in
7 immune evasion and thus in chicken gut colonization. However, *C. jejuni* strains
8 that underwent rearrangements leading to phage-resistance were demonstrated to
9 be inefficient colonizers of the chick intestine (Scott et al., 2007b). There is still
10 some controversy regarding the genomic instability of *C. jejuni* since Nielsen et
11 al. (2001) concluded that many strains were genetically stable as tested by
12 ribotyping, PFGE, RAPD and Penner heat-stable serotyping after *in vitro* and *in*
13 *vivo* (through mice) passage. Moreover, Manning et al. (2001) concluded that this
14 stability could be maintained despite exposure to various environmental
15 conditions over long time periods and covering large distances. Also, it has been
16 suggested that subtype pattern variations in *C. jejuni* leading to phenotypic
17 changes, occur only occasionally during *in vivo* passage (Konkel et al., 2007). On
18 the other hand, Ridley et al. (2008) observed that, although stable during single
19 cecal colonization of one individual strain, the *C. jejuni* genome can undergo
20 changes upon competitive stress (i.e. during co-colonization) in the avian gut,
21 leading to PFGE type variants with different colonization capacities from a single
22 parent clone. This genetic and phenotypic diversity might play a role in the
23 improved fitness of certain *C. jejuni* strains to survive and colonize another host.

1

2 **Crosstalk between *C. jejuni* and the chicken gut mucosa**

3 **Colonization mechanism**

4 Although *C. jejuni* is likely to encounter environmental stressors compromising
5 optimal growth in its chicken host (Murphy et al., 2006), the bacterium
6 persistently colonizes the chicken gut. This indicates that the bacterium harbours
7 regulatory systems conferring protection toward a hostile environment inside its
8 host. Although it is clear that successful colonization of the chicken GI tract is a
9 multifactorial process (Newell, 2002), the mechanism by which *C. jejuni* is able to
10 persistently evade the chicken immune response is poorly understood.

11 Upon entering the chicken GI tract, *C. jejuni* moves toward the intestinal
12 epithelial border, probably mediated by chemotaxis. *C. jejuni* is attracted by
13 intestinal mucins, as well as several amino acids, carbohydrates and salts of
14 organic acids, while the chemoattractive properties of L-fucose are controversial
15 (Vegge et al., 2009). *C. jejuni* responds to these chemicals via methyl-accepting
16 chemotaxis proteins (MCP) (Vegge et al., 2009), of which the most important are
17 the determinant of colonization proteinB (DocB) and the chemoreceptor
18 transducer-like protein1 (Tlp1), [while](#) the chemotaxis regulatory proteinY (CheY)
19 [shuttles between these MCP and the flagellar motor](#) (Hendrixson & DiRita, 2004;
20 Hartley-Tassell et al., 2010). The putative adaptation system CheBR is believed to
21 be involved in the response of *C. jejuni* to these environmental signals by
22 modifying its chemoreceptors (including Tlp1) (Kanungpean et al., 2011). DocB
23 and Tlp1 truncation, however, does not alter the chemotactic behaviour of *C.*

1 | *jejuni in vitro*, ~~indicating that they either serve partly redundant chemotactic~~
2 | ~~functions or a different function. Indeed, these MCP proteins were shown to but~~
3 | ~~rather reduces its~~ invasive potential in chicken embryo intestinal cells (Vegge et
4 | al., 2009). In any case, there is no doubt that DocB and Tlp1 are indispensable for
5 | *C. jejuni* to colonize chicks ~~the *in vivo* function of these proteins, as well as~~
6 | ~~chemotaxis regulation *in vivo* in general, remain somewhat obscure~~. For moving
7 | toward the most favourable conditions for growth *C. jejuni* needs intact flagella
8 | and it seems that especially *flaA*, *flgK*, *cj1324* and the motility accessory factor
9 | *maf5* gene are crucial for colonizing the chicken gut (Hermans et al. 2011b).

10 | The host intestinal mucus layer that lines the epithelial cells prevents most
11 | commensal bacteria to make direct contact with the epithelial surface by
12 | constituting a viscous physical barrier and by harbouring secretory IgA and
13 | antimicrobial peptides (Ivanov & Littman, 2011). And although increased
14 | viscosity has been associated with down-regulation of *flaA* promoter activity
15 | (Allen et al., 2001), the modified flagellum of *C. jejuni* allows the bacterium to
16 | penetrate the viscous mucus layer (Guerry, 2007) and ~~to reach and from~~ making
17 | directe contact with the intestinal epithelial cells. Although *C. jejuni* is not found
18 | to be attached to chicken cecal crypt microvilli *in vivo* (Beery et al., 1988) the
19 | bacterium has been observed intracellularly in intestinal epithelial cells of three-
20 | day-old experimentally inoculated chickens and in chicken primary cecal
21 | epithelial crypt cells *in vitro* (Van Deun et al., 2008). Moreover, several adhesins
22 | of *C. jejuni* have been implicated to be important for chick colonization.
23 | Therefore, upon entering the chicken gut it is believed that *C. jejuni* adheres to the

1 epithelial cells, mediated by intact flagella and surface-exposed proteins. In
2 particular CadF (*Campylobacter* adhesion to fibronectin) ~~(CadF)~~ and FlpA
3 ~~(fibronectin-like protein A (FlpA))~~ were identified as important adhesins for
4 colonization, while the potential contribution of *Campylobacter* adhesion protein
5 A (CapA) is less clear ~~*Campylobacter* adhesion protein A (CapA) has been~~
6 ~~implicated as a putative adhesion (Ashgar et al., 2007). In contrast, in another~~
7 ~~study no reduced colonization in chicks was observed for a *capA* mutant, although~~
8 ~~*C. jejuni* adherence to chicken LMH cells was attenuated (Flanagan et al., 2009).~~
9 ~~This study also revealed that *capA* is not conserved among *C. jejuni* isolates,~~
10 ~~suggesting only a limited role for CapA during chicken colonization.~~ (Hermans et
11 al. 2011b) Also several surface-accessible carbohydrate structures of *C. jejuni*,
12 such as lipooligosaccharide (LOS) and an intact capsule, are involved in adhesion
13 with in particular the capsular polysaccharide transporter gene *kpsM* and the *N*-
14 linked general protein glycosylation pathway gene *pglH* being important for
15 colonization of the chicken intestinal tract (Karlyshev et al., 2004; Hermans et al.,
16 2011b).

17 Adhesion of *C. jejuni* to gut epithelial cells ~~is probably~~ may be followed by
18 marginal invasion in these cells. Upon exposure to chicken mucus, the flagellar
19 apparatus increases the secretion of *Campylobacter* invasion antigens (Cia),
20 important for *in vitro* cell invasion and chick colonization (Ziprin et al., 2001;
21 Konkel et al., 2004; Biswas et al., 2007). Also *C. jejuni* LOS is important for
22 epithelial cell invasion as well as for immune evasion in humans and sialylation of
23 the LOS outer core further enhances these traits (Louwen et al., 2008; Habib et al.,

1 2009). *C. jejuni* is not able to survive for long periods in primary chicken cecal
2 epithelial cells, nor is it able to multiply in cultured human intestinal epithelial
3 cells. Therefore, intracellular replication in these cells is probably not important
4 for persistent *in vivo* colonization. Rather, invasion of cecal crypt epithelial cells
5 would be followed by evading these cells allowing *C. jejuni* to replicate in the
6 mucus, which seems to provide all necessary nutrients for optimal growth, and re-
7 invasion to escape mucosal clearance (Van Deun et al., 2008). Strangely, in
8 contrast to Caco-2 invasion, the invasion capacity of *C. jejuni* in primary chicken
9 cecal epithelial cells *in vitro* is not correlated with *in vivo* gut colonization, but is
10 with systemic dissemination (Van Deun et al., 2008). Therefore, the genuine
11 contribution of epithelial cell invasion during cecal colonization of chicks with *C.*
12 *jejuni* is not clearly definable and can only be speculated on.

13 Next In addition to these three key events (chemotaxis, adhesion and possibly
14 invasion), ~~also~~ a plethora of additional mechanisms, including several stress
15 responses, multidrug and bile resistance regulation, iron regulation and energy
16 metabolism are definitely important for initial and persistent high-level
17 colonization of the avian GI tract with *C. jejuni* (Hermans et al. 2011b).

18

19 **Chicken intestinal immune response upon to C. jejuni entranecolonization**

20 *Protection of young chicks against C. jejuni colonization*

21 Day-of-hatch chicks have no established gut flora and possess an immature
22 mucosal immune system. In the cecum, it is only after four to seven days post-
23 hatch that an increase in cecal pro-inflammatory chemo- (such as interleukin-8

1 (IL-8)) and cytokine expression and heterophil numbers can be observed, upon
2 exposure to feed and microflora (Bar-Shira & Friedman, 2006). Hatchlings are
3 also unprotected by adaptive immunity, which only starts to develop after a few
4 days of life (Friedman et al., 2003). Nevertheless, colonization of chickens with
5 *C. jejuni* during this critical period seems not to occur. Instead, maternally-derived
6 antibodies generated against flagellin proteins (such as FlaA), adhesins (such as
7 CadF) and other *C. jejuni* surface components are important in protecting young
8 chickens from *C. jejuni* colonization during the first two weeks, the so called lag-
9 phase (Sahin et al., 2001, 2003; Shoaf-Sweeney et al., 2008; Zeng et al., 2009).
10 Killing of *C. jejuni* by maternal antibodies happens in a complement-mediated,
11 strain-specific way (Young et al., 2007). These antibodies confer enhanced
12 protection against challenge with a homologous strain compared to a heterologous
13 strain, probably because they retard motility of a homologous, but not that of a
14 heterologous strain, as shown *in vitro* (Sahin et al., 2003). After the lag-phase,
15 chickens show an increased susceptibility to colonization with *C. jejuni* which
16 coincides with a loss of maternally derived, circulating anti-*Campylobacter* IgY
17 antibodies, suggesting that adaptive immunity is not critical in protecting broilers
18 from colonization (Cawthraw et al., 2010). Interestingly, day-of-hatch broilers
19 have been shown to be very susceptible to *C. jejuni* colonization, which again
20 diminished over the first few days of life (Cawthraw et al., 2010; Conlan et al.,
21 2011), while transmission of *C. jejuni* between co-housed birds is lower in day-
22 old chicks compared to two-week-old birds (Conlan et al., 2011). This indicates

1 | that a lack of exposure of broiler flocks to *C. jejuni* and/or reduced transmission
2 | during the early stages of rearing may also contribute to the observed lag-phase.

3 Developing chicken embryos have increased expression levels of several
4 avian β -defensins, a group of antimicrobial peptides important in innate and
5 adaptive immune responses that might contribute to the observed protection
6 toward *C. jejuni* infection *in ovo* and post-hatch (Meade et al., 2009a). For the β -
7 defensin gallinacin-6, for instance, *in vitro* antibacterial activity against *C. jejuni*
8 has been demonstrated (van Dijk et al., 2007).

9

10 *Innate immune response*

11 | The chicken intestinal innate immune system ~~is built up by~~ comprises several
12 tissues, cells (such as epithelial cells, monocytes/macrophages, dendritic cells,
13 natural killer cells and neutrophils) and germline-encoded molecules (such as
14 chemo- and cytokines, antimicrobial peptides and nitric oxide) that can limit both
15 commensal and pathogenic invading bacteria (Brisbin et al., 2008). Some *in vitro*
16 studies with macrophages and epithelial cells, both primary and cultured,
17 contributed to the insight into the chicken immune response toward *C. jejuni*
18 infection. *C. jejuni* has been shown to be adhesive to, invasive in and to stimulate
19 inflammatory responses from these cells (Smith et al., 2005; Byrne et al., 2007;
20 Larson et al., 2008; Van Deun et al., 2008). Evidence of both *in vitro* uptake of *C.*
21 *jejuni* by chicken peritoneal macrophages (Myszewski & Stern, 1991) and *in vivo*
22 presence of *C. jejuni* within chicken epithelial cells and macrophages (Ruiz-
23 Palacios et al., 1991) exists.

1 A crucial step in the host innate immune response to bacterial entrance in the
2 GI tract is the activation of Toll-like receptors (TLRs), expressed on a variety of
3 cells of the GI mucosa including macrophages and epithelial cells, the latter
4 forming the first borderline defence against invading pathogens (He et al., 2006;
5 Linde et al., 2008). TLRs are recognized by specific bacterial ligands and, once
6 activated, promote the expression of effector molecules such as antimicrobial
7 peptides, NO and inflammatory cytokines. Although knowledge on avian TLR
8 biology is only starting to unravel, very recently several chicken TLRs have been
9 implicated to play a role in *C. jejuni* recognition. The chicken TLR4/myeloid
10 differentiation protein-2 (chTLR4/chMD-2) complex and cell-surface expressed
11 chTLR2 recognize *Campylobacter* LOS and lipopeptides, respectively. Both
12 receptors are potently activated by lysed *Campylobacter* bacteria. However, loss
13 of bacterial cell wall integrity does not seem to play a critical role in TLR
14 activation, because also live *Campylobacter* bacteria are able to elicit a marked
15 inflammatory response in chickens (de Zoete et al., 2010). TLR5 specifically
16 recognizes conserved regions of bacterial flagellins, thereby preventing intestinal
17 pathology. *C. jejuni*, however, lacks these TLR5-recognition sites and is therefore
18 unable to activate chTLR5, indicating that TLR5 signaling does not play a critical
19 role in the chick immune response against *C. jejuni* (Guerry, 2007; de Zoete et al.,
20 2010). Finally, TLR21, which is unique to avian, amphibian and fish species,
21 enables recognition of unmethylated single stranded microbial 2'-
22 deoxyribo(cytidine-phosphateguanosine) (CpG) DNA motifs with a broad ligand
23 specificity. *C. jejuni* CpG DNA is internalized through endocytosis and most

1 likely interacts with chTLR21 intracellularly, similar to the interaction of CpG
2 DNA with the functional homologue (TLR9) in mammals (de Zoete et al., 2010,
3 Kestra et al., 2010).

4 Activation of chTLR2, chTLR4 and chTLR21 results in an innate immune
5 response through myeloid differentiation primary response gene 88 (MyD88)-
6 dependent activation of nuclear transcription factor kappaB (NF- κ B) and
7 subsequent production of inflammatory cytokines and chemokines (Brownlie et
8 al., 2009; de Zoete et al., 2010; Kestra et al., 2010). Additionally, chTLR4 and
9 chTLR21 ligands can induce the production of inducible nitric oxide synthase-
10 mediated NO from chicken monocytes (He et al., 2006). In mammals, TLR-
11 signaling also involves a TLR4-mediated MyD88-independent pathway
12 associated with the induction of late phase NF- κ B and interferon (IFN)-inducible
13 genes, such as IFN- β , involved in natural killer cell activation, and maturation of
14 dendritic cells (Yamamoto et al., 2004). Chickens, however, lack this pathway and
15 therefore have an aberrant response to *C. jejuni* LOS compared to mammalian
16 species, rendering them much more resistant to the toxic effects of these TLR4
17 agonists. Although the TLR4-mediated MyD88-dependent pathway, leading to
18 early phase activation of NF- κ B, is intact, this explains in part the absence of
19 pathological signs in chicks in response to infection with *C. jejuni*, despite cell
20 adhesion and invasion (Kestra & van Putten, 2008; Shaughnessy et al., 2009; de
21 Zoete et al., 2010).

22 Upon *Campylobacter* infection, primary chick kidney cells and the avian
23 macrophage cell line HD11 express NO and pro-inflammatory cyto- (IL-6 and IL-

1 1 β) and chemokines (chIL-8) (Larson et al., 2008). Production of NO by activated
2 macrophages is important for their bactericidal activity (Linde et al., 2008). IL-1 β
3 and IL-6 are both major mediators of the innate immune system, while IL-6 is
4 also involved in the immunological switch from innate to adaptive immunity
5 (Smith et al., 2005). IL-1 β is primarily produced by monocytes/macrophages and
6 is involved in the inflammatory response of chickens against microbial products
7 (such as lipopolysaccharide (LPS)) by instructing epithelial cells and
8 macrophages to produce chemokines (Bar-Shira & Friedman, 2006). The chicken
9 orthologue of mammalian IL-8 (CXCLi1 and CXCLi2, but here referred to as
10 chIL-8) (Kaiser et al., 1999; Smith et al., 2005) attracts heterophils and, unlike its
11 mammalian counterpart, also monocytes to the site of infection (Martins-Green,
12 2001). It has been demonstrated that the *N*-terminus of chIL-8, where the
13 chemotactic activity resides, is structurally homologous to that of monocyte
14 chemotactic protein-1 (Borrmann et al., 2007). This human chemokine is
15 chemotactic for monocytes, probably explaining the chemotactic movement of
16 monocytes toward chIL-8. A marked chIL-8 response is induced in chicken LMH
17 and primary intestinal cells upon inoculation with *C. jejuni* (Brisbin et al., 2008;
18 Li et al., 20010). Finally, also in chicken embryo intestinal cells *C. jejuni* is
19 capable of inducing a pro-inflammatory response (Smith et al., 2008; Li et al.,
20 2010).

21 Despite the lack of association of *C. jejuni* with chicken crypt epithelium *in vivo*,
22 some recent reports demonstrate the initiation of a mild inflammatory response in
23 chickens upon exposure to the bacterium. *C. jejuni* colonization in chickens is

1 accompanied by infiltration of proinflammatory cells in mucosal tissues, although
2 overt signs of cell invasion or pathology were not found (Larson et al., 2008;
3 Smith et al., 2008). Upon inoculation of four-week-old broilers, an early increase
4 (six h post- inoculation (pi)) in circulating monocytes/macrophages was observed
5 and increased numbers were maintained after 48 h (Meade et al., 2009b).
6 Strikingly, heterophil numbers remained unaltered during [this](#) time course.
7 Absence of a heterophil infiltrate was also observed in cecal mucosal tissues of
8 three-week-old hens [24 h after directly injection of their cecum](#) with *C. jejuni*
9 (Van Deun et al., 2008). In contrast, another study (Smith et al., 2008) showed a
10 minor, although significant induction of heterophil infiltration in cecal tissues one
11 day and four days after inoculating two-week-old broiler chicks, as well as in the
12 ileum at four days post-inoculation. [It cannot be ruled out that also in the studies](#)
13 [by Meade et al. \(2009b\) and Van Deun et al. \(2008\) a heterophil influx could have](#)
14 [been observed after four days, but the discrepancy in heterophil influx after one](#)
15 [day between these studies is not clear. Possibly, the different *C. jejuni* strains](#)
16 [used in these studies may have accounted for this. But more likely, differences in](#)
17 [chicken lines and bird age were responsible in the differential host response](#)
18 [because in the study by Smith et al. \(2008\) an out-bred flock was used and the](#)
19 [heterophil influx observed in two-week-old birds was absent in day-of-hatch](#)
20 [chicks.](#) In one-day-old birds, however, this induction was not observed.
21 Expression of both TLR4 and TLR21, but not TLR2, is readily increased (six h pi)
22 in cecal tissues in response to *C. jejuni* inoculation (Meade et al., 2009b;
23 Shaughnessy et al., 2009). In two- and four-week-old broiler chicks this is

1 accompanied, however, by only a limited cytokine gene expression except for a
2 marked increase in *chIL-8* expression already after 6-12 h pi which is maintained
3 over 48 h after inoculation (Shaughnessy et al., 2009) and longer (Smith et al.,
4 2008). *IL-1 β* expression levels are moderately increased after 20-24 h and
5 decrease afterwards, while increased *IL-6* expression is evident only after 48 h at
6 the earliest (Keestra & van Putten, 2008; Smith et al., 2008). A similar response
7 can be observed in ileal tissues although a marked induction of *IL-6* expression
8 levels was already evident in these tissues at six h pi after which they started to
9 drop again. In one-day-old chicks, these responses are less pronounced or absent
10 although also in these animals *IL-8* expression in cecal tissues is induced. Overall,
11 induction of cytokines is most evident within 24 h after inoculation after which
12 the expression levels drop again. Because the intestinal bacterial load in these
13 *Campylobacter*-colonized chicks did not lower during the examined time-course,
14 there clearly exist some mechanisms that are responsible for controlling this pro-
15 inflammatory response (Smith et al., 2008). Expression levels of anti-
16 inflammatory *IL-10*, *IL-13* and transforming-growth factor β 4 (*TGF- β 4*) were not
17 detected in cecum, ileum and spleen, and the signals modulating the pro-
18 inflammatory response, resulting in sustained and unaffected *C. jejuni*
19 colonization, are yet unknown (Smith et al., 2008; Shaughnessy et al., 2009). *C.*
20 *jejuni* colonization in chicks significantly reduces expression levels of several
21 antimicrobial peptide genes (Meade et al., 2009a). This downregulation might
22 represent one mechanism whereby *C. jejuni* modulates the immune response,
23 limiting the efficacy of these antimicrobial factors and enabling itself to

1 persistently colonize its host at high levels. As stated above, gallinacin-6 has a
2 bactericidal effect on *C. jejuni* (van Dijk et al., 2007). Based on mRNA levels,
3 expression of this defensin is low in the avian intestinal tract, and no detailed
4 studies have been done upon the time of writing this review that indicate an
5 inducible upregulation of gallinacin-6 after exposure to *C. jejuni*. In a recent study
6 by Shaughnessy et al. (2011) 270 genes were found to be significantly ($P < 0.01$)
7 differentially expressed after 20 h in four-week-old chicks colonized with *C.*
8 *jejuni* compared to *C. jejuni* free chicks. These genes corresponded to the
9 activation of several biological processes, including immune responses. Although
10 differences in expression were only marginal, this response was hypothesized to
11 point toward an innate T-cell response in the ceca of chickens 20 h after
12 inoculation with *C. jejuni* (Shaughnessy et al., 2011).

13

14 *Adaptive immune response*

15 The type of immune response generated against *C. jejuni* depends on the cytokine
16 microenvironment induced by the chick innate defence cells. This in turn is
17 determined by the interaction of TLRs and other pathogen recognition receptors
18 expressed on these cells with their respective ligands. In chickens, not all of these
19 receptors and cytokines are fully identified yet, making the switch from innate to
20 adaptive immunity in this species not completely understood (Brisbin et al.,
21 2008).

22 In chickens, intestinal antigens are capable of entering the bursa of Fabricius,
23 the site of primary B cell development (Brisbin et al., 2008). Chickens have an

1 incomplete antibody response toward T-cell independent type 2 antigens which
2 activate B cells independently of T cells (Jeurissen et al., 1998). Because these
3 antigens are usually of polysaccharide nature, an insufficient humoral response
4 toward certain surface-accessible carbohydrate structures (SACS) of *C. jejuni*
5 might contribute to the inability of the chicken immune system to clear this
6 microorganism, despite the antigenic potential of *C. jejuni* LOS and its capsule
7 (Oza et al., 2002) and the marked immunogenicity of *C. jejuni* flagellin (Widders
8 et al., 1998). Moreover, an outer membrane protein extract of *C. jejuni* has been
9 shown to cause apoptosis of chicken blood and spleen lymphocytes, probably
10 promoting immune evasion of *C. jejuni* in the chick (Zhu et al., 1999). An
11 antibody response to *C. jejuni* might, however, contribute to protection against
12 intestinal colonization of chickens, which show a significant increase in specific
13 mucosal and circulating IgG (IgY) and IgA and circulating IgM antibody titres
14 when colonized with *Campylobacter* (Cawthraw et al., 1994; Widders et al.,
15 1998). In these studies flagellin was shown to be the immunodominant antigen,
16 which is rather peculiar due to the lack of functional TLR5-recognition sites in *C.*
17 *jejuni* flagellin, permitting TLR5 evasion (Guerry, 2001; de Zoete et al., 2010).
18 Nevertheless, vaccinating chicks with a hybrid protein based on *C. jejuni* FlaA
19 induced a specific response against this protein and reduced colonization in these
20 birds (Khoury and Meinersmann, 1995). An antibody response specific for native
21 flagellin was also induced in the serum of chickens immunized with purified *C.*
22 *jejuni* flagellin. Serum and GI secretion antibodies specific for *C. jejuni* whole
23 cells were, however, only induced ~~But only~~ when ~~the~~ this protein was

1 | complemented with killed *C. jejuni* whole cells, ~~which moreover result~~inged in
2 | reduced cecal *C. jejuni* counts in these birds (Widders et al., 1998). This might
3 | indicate that the epitopes of *C. jejuni* flagella are not accessible for these
4 | antibodies in intact bacteria and that possibly other antigens, not detected in this
5 | study, were responsible for the induction of anti-*C. jejuni* antibodies reducing the
6 | cecal bacterial load. Recent studies gave more insight into this enigma and
7 | identified additional immunogens of *C. jejuni* promoting the humoral immune
8 | respons in chicks. Amongst others the *C. jejuni* ferric enterobactin receptor CfrA
9 | (involved in iron regulation), the outer membrane chanel CmeC (involved in
10 | multidrug resistance), Cj0091 (belonging to a lipoprotein-encoding operon), the
11 | lipoprotein CjaA and CjaC (mediating amino acid transport), CadF and LOS were
12 | shown to be immunogenic and expressed during *in vivo* colonization (Shoaf-
13 | Sweeney et al., 2008; Zeng et al., 2009; Oakland et al., 2011). Both the sera of
14 | young chicks free of *C. jejuni* and older birds colonized with the bacterium were
15 | reactive against recombinant CfrA, indicating that they are not only passed from
16 | the mature hen to the hatchling but are also induced during colonization of
17 | broilers after the lag-phase (Zeng et al., 2009). It was speculated that antibodies
18 | directed to CfrA hinder the interaction of FeEnt with its receptor. Proper
19 | functioning of CfrA is crucial for *C. jejuni* colonization in chicks, indicating that
20 | CfrA antibodies are potentially protective. Also *C. jejuni* CjaA-based vaccines
21 | were shown to induce specific serum IgY and mucosal IgA antibody responses
22 | against CjaA and reduced cecal colonization of vaccinated chickens (Buckley et
23 | al., 2010).

1 Intestinal epithelial cells might contribute to a mucosal IgA response by the
2 GALT, located beneath the epithelial cell border in the lamina propria, in a T-cell
3 dependent manner by producing IL-6 after contact with *C. jejuni* (Faragasan,
4 2008). Secretory IgA is the major immunoglobulin isotype in mucosal secretions
5 and generally responsible for preventing sub-epithelial translocation of
6 commensal bacteria by preventing their adhesion to epithelial cells or returning
7 bacteria that already reached the basolateral site, without eliciting an
8 inflammatory response (Brisbin et al., 2008). Moreover, by its resistance to
9 normal intestinal proteases, through dimerization on the surface of mucosal
10 epithelial cells, IgA is ideally suited for host defences at the mucosal surface of
11 the GI tract (Phalipon et al., 2002). IgA might thus play an important role in
12 limiting the mucosal immune response to *C. jejuni* in chickens and redirecting it
13 toward tolerance.

14 Most *C. jejuni* strains possess genes encoding a cell death-promoting
15 cytolethal distending toxin (CDT) of which the expression is induced in both the
16 avian and human gut (Abuoun et al., 2005). During human infection with *C.*
17 *jejuni*, neutralizing antibodies against CDT are induced, but not during
18 colonization in chickens and it seems that production of this toxin in general is not
19 important for chick colonization as opposed to its suspected role during
20 pathogenesis in humans (Abuoun et al., 2005; Biswas et al., 2006).

21 As mentioned above, genetically distinct chicken lines may differ in their
22 susceptibility toward cecal *C. jejuni* colonization (Stern et al., 1990). Further
23 research in this area revealed insulin receptor signaling and metabolism process

1 pathways to be key players of this differential response (Li et al., 2010). In a more
2 resistant line, lymphocyte activation, lymphoid organ development functions and
3 circadian rhythm were important in the cecal host defence upon *C. jejuni*
4 inoculation. In a more susceptible line, cell differentiation, communication and
5 signaling pathways were important during host defence, with a marked
6 upregulation in lipid, glucose and amino acid metabolism.

7

8 Chicken systemic immune response to *C. jejuni*

9 The frequently observed systemic colonization of *C. jejuni* in chicks indicates that
10 the bacterium, despite the induction of secretory IgA by the GALT, is capable of
11 breaching the gut epithelial barrier. As in the GI tract this happens without
12 developing pathology or inducing excessive inflammation, although chicks can
13 mount an adaptive T cell response to *C. jejuni* when it reaches and colonizes the
14 liver (Jennings et al., 2011). In colonized flocks, almost all birds carry *C. jejuni* in
15 their ceca but significantly less birds harbour the bacteria in their liver tissues
16 (Jennings et al., 2011). Whether host-specific differences decide over *C. jejuni*
17 dissemination, or a T cell response is responsible for the eradication of *C. jejuni*
18 from the host liver in some animals, is not known. In any case, *C. jejuni*-specific
19 antibody responses are apparently not capable of clearing the bacterium from the
20 chicken gut, but nevertheless do indicate that there indeed must have been a
21 preceding close interaction between *C. jejuni* and the host epithelial cells.

22 The two chicken lines used in the study of Li et al. (2010) also differed in their
23 systemic response to *C. jejuni* (Li et al., 2011). In the spleen, a secondary

1 lymphoid organ of the avian immune system important for lymphocyte activation,
2 proliferation and differentiation, the response to *C. jejuni* in the more resistant line
3 was characterized, as in the cecum, by lymphocyte activation and differentiation.
4 In addition, splenic host genes for humoral responses and Ig heavy and light chain
5 were upregulated. These responses initiate adaptive immune responses to *C. jejuni*
6 and are probably responsible for an increased genetic resistance to systemic *C.*
7 *jejuni* colonization. In the susceptible line, genes for regulation of erythrocyte
8 differentiation, hemopoiesis and RNA biosynthesis processes were
9 downregulated. This study also revealed distinct innate defense mechanisms
10 against *C. jejuni* by the two chicken lines. Apoptosis and cytochrome *c* release
11 from mitochondria was associated with increased resistance against *C. jejuni*
12 colonization. Probably, these events induce increased apoptosis of infected host
13 cells, thereby destroying the habitat of the bacteria and contributing to the
14 increased resistance to splenic colonization with *C. jejuni*.

15

16 **Interaction with the host microbiota**

17 Little is known currently about the effect of the natural avian gut microbiota on
18 the level of *C. jejuni* colonization. In general, host microbiota imposes a
19 colonization barrier for intruding pathogens by competing for nutrients (such as
20 carbon) and host receptors. Their composition, however, can alter the outcome of
21 invading enteric bacteria (by e.g. altering the virulence properties of these
22 bacteria), resulting in either clearance or colonization (Keeney & Finlay, 2011).
23 And although it has been suggested that the colonization pattern of *C. jejuni* in

1 chicks is mainly determined by the chicken host but not by the host microbiota
2 (Ridley et al., 2008), also the composition of the latter might contribute to the
3 observed colonization pattern. Changes in *C. jejuni* loads in the commercial
4 turkey intestine seemed to correlate to, but are not dependent on, two acute
5 transitions in the cecal microbiota composition during the turkey development
6 phases (Scupham, 2009). With an approach called antibiotic dissection, day-old
7 turkey poults were inoculated with cecal contents of *Campylobacter*-free adult
8 turkeys after which the microbial communities in these poults were modified by
9 different antibiotic treatments. Molecular examination of the constituents of these
10 communities detected that a subtype I of *Megamonas hypermegale* correlated with
11 decreased colonization ability of *C. jejuni*, while a virginiamycin-derived cecal
12 microbiota seemed to be correlated with enhanced colonization ability (Scupham
13 et al., 2010). These results indicate that *C. jejuni* may respond to the presence of
14 specific subsets of the avian gut microbiota. It has, however, to be examined if the
15 effect of these gut microbiota alterations on *C. jejuni* in turkeys also applies to
16 chicks.

17

18 **Hypothetical mechanism of the interaction between *C. jejuni* and the chicken** 19 **gut mucosa**

20 The interaction of *C. jejuni* with its avian host is very complex, evidenced by the
21 extensive interplay between several key mediators important in successful and
22 persistent colonization in the chicken GI tract. In chicks, this dual interaction is
23 clearly influenced by both the *C. jejuni* strain and the chicken line involved. The

1 information reviewed above suggests that, despite the lack of a developed
2 pathology, a pro-inflammatory response is developed in the chicken intestinal
3 mucosa during asymptomatic colonization with *C. jejuni*. Upon *Campylobacter*
4 entrance in the avian GI tract, an early induced production of chIL-8 by intestinal
5 epithelial cells is observed, followed by macrophage recruitment and production
6 of proinflammatory cytokines. This is, however, not accompanied by the
7 recruitment of heterophils (the avian equivalent of mammalian neutrophils) to the
8 site of infection. In a later stage, a specific mucosal IgA response is mounted
9 against *C. jejuni*, but this induction is not capable of clearing the bacterium from
10 the gut. This humoral response is moreover not capable to prevent *C. jejuni*
11 from further interacting with and to translocate across the gut epithelium and
12 to disseminate systemically. Also the specific T cell response that is triggered
13 upon *C. jejuni* entrance in the extra-intestinal organs does neither not result in
14 clearance from these tissues, nor pathology. Because *C. jejuni* colonizes the
15 chicken gut persistently, it is thus must be capable of somehow evading this the
16 inefficient host immune response and. But also the chicken host might be
17 involved in maintaining homeostasis during persistent colonization (see further).
18 In **Figure 1**, a schematic overview is given of a simplified hypothetical
19 mechanism involved in the interaction of *C. jejuni* with the chicken gut, after lag-
20 phase, leading to successful and persistent colonization of the GI tract, without
21 developing pathology.

22

23 **Commensal *C. jejuni* colonization in chicks: immunological tolerance?**

1 In mammals commensal infections are characterized by the absence of a
2 neutrophil infiltrate or a classical inflammation as seen during pathogenic
3 infection (MacPherson & Uhr, 2004), indicating that the interaction between *C.*
4 *jejuni* and its chicken host is indeed of commensal nature. Intestinal homeostasis
5 during commensal colonization requires that a proinflammatory response is
6 rapidly controlled. In mammals not much is known about the host regulatory
7 mechanisms that contribute to tolerance without reducing bacterial numbers, but,
8 restricting the bacteria to the lumen (so they cannot reach the epithelial cells and
9 the immune system) and inducing an anti-inflammatory response are believed to
10 induce a state of “immunological ignorance” (Ivanov & Littman, 2011). Due to a
11 lack of knowledge about the interaction between *C. jejuni* and the chicken
12 immune system it remains unclear how homeostasis is maintained in chickens
13 colonized with *C. jejuni*. An apparent induction of a mild intestinal pro-
14 inflammatory response, the inability to demonstrate upregulation of anti-
15 inflammatory cytokines, occasional invasion of cecal crypt epithelial cells and
16 regular dissemination to extra-intestinal organs upon *C. jejuni* colonization of the
17 chicken host, suggests that their interaction is not a tale of ignorance but rather a
18 cohort of active processes, exerted by the two partners, resulting in
19 “immunological tolerance”. *C. jejuni* itself might escape or alter the inflammatory
20 response by, for instance, down-regulating antimicrobial peptide gene expression
21 in the chicken gut, but other potential mechanism(s) or bacterial factor(s) of *C.*
22 | *jejuni* involved in immune evasion are currently not known. Alternatively, [also](#)

1 the chicken host might support tolerance to maintain homeostasis during
2 persistent, asymptomatic colonization (Pédron & Sansonetti, 2004).

3 First of all, the differential composition of the chicken intestinal mucus layer,
4 compared to its human counterpart, probably plays an important role in promoting
5 homeostasis during *C. jejuni* colonization. Chicken intestinal mucins have been
6 shown to reduce the adhesive and especially the invasive capacity of *C. jejuni* in
7 human primary and cultured intestinal epithelial cells (Byrne et al., 2007; Alemka
8 et al., 2010). In contrast, human-derived mucus promotes adhesion and entrance
9 (Byrne et al., 2007). Moreover, MUC2, the most abundantly secreted mucin in the
10 human intestine, is a major chemoattractant for *C. jejuni* and induces the
11 expression of several colonization- and virulence-associated genes (Tu et al.,
12 2008). To date, no such properties have been assigned to chicken mucins. Host
13 intestinal mucins can be either secreted or expressed at the apical surface of the
14 (cecal) mucosal epithelial cells and are readily found to be coated with
15 fucosylated glycans in terminal positions (Stahl et al., 2011). Although the
16 chemotactic properties of L-fucose were not validated by Vegge et al. (2009), it is
17 believed that *C. jejuni* is attracted to, and binds with both mucin and L-fucose.
18 Presence of the latter at certain concentrations might moreover increase *C. jejuni*
19 *flaA* promoter activity (Allen et al., 2001). Therefore, fucosylated glycans may
20 function as adherence factors for *C. jejuni*. In addition, although it was believed
21 until now that *C. jejuni* is an asaccharolytic organism, very recent evidence
22 indicates that some strains are able to use L-fucose as a substrate for growth (Stahl
23 et al., 2011). Thus, chemotaxis toward, adhesion to and subsequent utilization of

1 L-fucose by *C. jejuni* strains possessing a functional L-fucose uptake and
2 metabolism pathway provides them with a competitive advantage. This seems,
3 however, to be only the case during pathogenic (in human), but not during
4 commensal (in chick) colonization (Stahl et al., 2011). Probably, next to
5 decreasing the intestinal barrier permeability to *C. jejuni*, the highly sulfated
6 fucosylated *O*-glycan mucin structures found in chickens decrease the
7 accessibility of, and thus the responsiveness of *C. jejuni* to L-fucose. Indeed, upon
8 feeding young chicks with an excess of free L-fucose also here a competitive
9 colonization advantage was observed for wild-type *C. jejuni* over a mutant lacking
10 a functional fucose permease gene, important for L-fucose transport into the
11 bacterial cell (Stahl et al., 2011). Thus, a high degree of L-fucose masking
12 through increased sulfation might give further explanation to the lack of
13 association of *C. jejuni* with the chicken crypt epithelium *in vivo*. To conclude,
14 there is increasing evidence that the composition of the chicken mucus layer is
15 involved in the hindered contact between *C. jejuni* with the chicken intestinal
16 epithelial surface. Indeed, *C. jejuni* is not closely associated with chicken crypt
17 epithelium *in vivo* but rather resides in the mucus within the lumen of the crypts
18 (Beery et al., 1988). However, the effect of chicken mucus on *C. jejuni* invasion
19 in primary chicken epithelial cells has not yet been examined. Moreover, as the
20 bacterium can be frequently detected in extra-intestinal organs of chicks, the
21 mucus layer is not likely to be an efficient barrier to prevent close interaction with
22 *C. jejuni* and the intestinal epithelial lining. In contrast, it seems that it indirectly
23 promotes *C. jejuni* invasion through the secretion of Cia proteins (Biswas et al.,

2007). Further research will therefore have to reveal the genuine contribution of the mucus layer to GI and systemic colonization of *C. jejuni* in chicks.

Also the adaptive immune system of the chick might participate in the tolerogenic response to *C. jejuni*. Upon intestinal colonization, specific IgA against *C. jejuni* is induced. IgA is believed to induce the modulation of epitope expression by bacteria and to reduce intestinal proinflammatory signalling (Peterson et al., 2007). This indicates that the induction of IgA could lead to immune evasion, but whether the induction of IgA in chickens colonized with *C. jejuni* might be responsible for the noninflammatory *C. jejuni*-chicken gut relationship is not clear.

Next, murine intestinal epithelial cells are tolerized to LPS early after birth by exposure to exogenous LPS, facilitating microbial colonization and the establishment of a stable intestinal host-microbe homeostasis (MacPherson & Uhr, 2004). Whether in chickens LOS tolerance in the gut is involved in a tolerance-oriented integrated mucosal immune system, allowing commensal colonization of *C. jejuni*, is not clear.

Finally, chickens have an aberrant response to *C. jejuni* LOS and ~~are unresponsive to *C. jejuni* flagellin~~, due to the absence of a late phase NF- κ B response and TLR5 recognition sites, respectively. Only the first is likely to contribute to the differential *C. jejuni* response in humans and chicks because *C. jejuni* escapes TLR5 recognition in humans too (de Zoete et al., 2010). Next to these responses, colonized chickens might further induce tolerance by expressing factors that blunt *C. jejuni* components which could induce inflammation

1 (MacPherson & Uhr, 2004). However, potential candidates have not yet been
2 identified.

3

4 **Concluding remarks**

5

6 Chickens are often colonized by the zoonotic pathogen *Campylobacter jejuni* and
7 broiler meat products are considered to be the main source of campylobacteriosis
8 in humans. In humans, *C. jejuni* is capable of causing severe inflammatory
9 disease, while chickens are colonized asymptotically. How *C. jejuni* shapes the
10 mucosal immune system of the gut during health and disease is, however, poorly
11 understood. Upon entering the chicken GI tract, *C. jejuni* establishes a complex
12 interaction with its host, resulting in persistent high-level cecal colonization.
13 Although evidence is emerging suggesting that *C. jejuni* poorly invades the GI
14 tract of chicks and inefficiently elicits the chick's immune system, no pathology is
15 observed. Moreover, it seems that *C. jejuni* is capable of evading the immune
16 response and to even colonize systemically. This inefficient, controlled
17 inflammatory response is not capable of clearing *C. jejuni* from the chicken gut
18 and many processes might be involved in redirecting the response toward
19 tolerance. The underlying mechanisms of the crosstalk between *C. jejuni* and
20 chicks are just now starting to unravel and further research is warranted.
21 Especially the mechanisms allowing this bacterium to persistently evade the
22 immune response should deserve full attention. After all, a better understanding of
23 the chick immune response upon *C. jejuni* entrance, as well as further elucidation

1 of the colonization mechanism of the bacterium in this host might promote the
2 development of effective control measures to clear this human pathogen from
3 poultry lines. For this purpose it might be of particular interest to identify chicken
4 factors, if any, involved in blunting *C. jejuni* virulence factors, while *C. jejuni*
5 | colonization factors identified to da-te might hold promise for effective subunit
6 | vaccines. Moreover, the differential disease outcome in chicks and humans upon
7 | exposure to *C. jejuni* might be explained. Could it be due to the differences in
8 | mucin composition, TLR signalling, effect of CDT or humoral responses in these
9 | hosts, or are there other, yet to defined, mechanisms that ~~deside~~determine ~~over~~
10 | the commensal or pathogenic nature of *C. jejuni*. Answering these questions,
11 | ~~based on what is currently known and described in this review,~~ could explain why
12 | and how a single bacterium is capable of causing severe inflammatory disease in
13 | one host while being (seemingly?) completely harmless in another.

14

15 **Acknowledgements**

16 In the authors' research groups, research on *Campylobacter* is financially
17 supported by the Federal Public Service of Health, Food Chain Safety and
18 Environment (FOD, Brussels, Belgium): project RT08/8-CAMPOUL. The authors
19 would like to thank Isabel De Smet for the graphical design of Figure 1.

20

21 **Declaration of Interest**

22 The authors report no declarations of interest.

23

1 **References**

- 2 Allen KJ, Griffiths MW. (2001). Effect of environmental and chemotactic stimuli
3 on the activity of the *Campylobacter jejuni* *flaA* σ^{28} promoter. FEMS
4 Microbiol Lett, 205, 43-48.
- 5 Abuoun M, Manning G, Cawthraw SA, Ridley A, Ahmed IH, Wassenaar TM,
6 Newell DG. (2005). Cytolethal distending toxin (CDT)-negative
7 *Campylobacter jejuni* strains and anti-CDT neutralizing antibodies are
8 induced during human infection but not during colonization in chickens.
9 Infect Immun, 73, 3053-3062.
- 10 Alemka A, Whelan S, Gough R, Clyne M, Gallagher ME, Carrington SD, Bourke
11 B. (2010). Purified chicken intestinal mucin attenuates *Campylobacter jejuni*
12 pathogenicity *in vitro*. J Med Microbiol 59, 898-903.
- 13 Ashgar SS, Oldfield NJ, Wooldridge KG, Jones MA, Irving GJ, Turner DP,
14 Ala'Aldeen DA (2007). CapA, an autotransporter protein of *Campylobacter*
15 *jejuni* mediates association with human epithelial cells and colonization of
16 the chicken gut. J Bacteriol, 189, 1856-1865.
- 17 Bar-Shira E, Friedman A. (2006). Development and adaptations of innate
18 immunity in the gastrointestinal tract of the newly hatched chick. Dev Comp
19 Immunol, 30, 930-941.
- 20 Barton C, Ng LK, Tyler SD, Clark CG. (2007). Temperate bacteriophages affect
21 pulsed-field gel electrophoresis patterns of *Campylobacter jejuni*. J Clin
22 Microbiol, 45, 386-391.

- 1 Beery JT, Hugdahl MB, Doyle MP. (1988). Colonization of gastrointestinal tracts
2 of chicks by *Campylobacter jejuni*. Appl Environ Microbiol, 54, 2365-2370.
- 3 Biswas D, Fernando U, Reiman C, Willson P, Potter A, Allan B. (2006). Effect of
4 cytolethal distending toxin of *Campylobacter jejuni* on adhesion and
5 internalization in cultured cells and in colonization of the chicken gut. Avian
6 Dis, 50, 586-593.
- 7 Biswas D, Fernando UM, Reiman CD, Willson PJ, Townsend HGG, Potter AA,
8 Allan BJ. (2007). Correlation between *in vitro* secretion of virulence-
9 associated proteins of *Campylobacter jejuni* and colonization of chickens.
10 Curr Microbiol, 54, 207-212.
- 11 Borrmann E, Berndt A, Hanel I, Kohler H. (2007). *Campylobacter*-induced
12 interleukin-8 responses in human intestinal epithelial cells and primary
13 intestinal chick cells. Vet Microbiol, 124, 115-124.
- 14 Boyd Y, Herbert EG, Marston KL, Jones MA, Barrow PA. (2005). Host genes
15 affect intestinal colonisation of newly hatched chickens by *Campylobacter*
16 *jejuni*. Immunogenetics, 57, 248-253.
- 17 Brisbin JT, Gong J, Sharif S. (2008). Interactions between commensal bacteria
18 and the gut-associated immune system of the chicken. Anim Health Res
19 Rev, 9, 101-110.
- 20 Brownlie R, Zhu J, Allan B, Mutwiri GK, Babiuk LA, Potter A, Griebel P. (2009).
21 Chicken TLR21 acts as a functional homologue to mammalian TLR9 in the
22 recognition of CpG oligodeoxynucleotides. Mol Immunol, 46, 3163-3170.

- 1 [Buckley AM, Wang JH, Hudson DL, Grant AJ, Jones MA, Maskell DJ, Stevens](#)
2 [MP. \(2010\). Evaluation of live-attenuated *Salmonella* vaccines expressing](#)
3 [*Campylobacter* antigens for control of *C. jejuni* in poultry. *Vaccine*, 28,](#)
4 [1094-1105.](#)
- 5 Byrne CM, Clyne M, Bourke B. (2007). *Campylobacter jejuni* adhere to and
6 invade chicken intestinal epithelial cells *in vitro*. *Microbiology*, 153, 561-
7 569.
- 8 Cawthraw S, Ayling R, Nuijten P, Wassenaar T, Newell DG. (1994). Isotype,
9 specificity, and kinetics of systemic and mucosal antibodies to
10 *Campylobacter jejuni* antigens, including flagellin, during experimental oral
11 infections of chickens. *Avian Dis*, 38, 341-349.
- 12 Cawthraw SA, Newell DG. (2010). Investigation of the presence and protective
13 effects of maternal antibodies against *Campylobacter jejuni* in chickens.
14 *Avian Dis*, 54, 86-93.
- 15 Clark CG, Ng LK. (2008). Sequence variability of *Campylobacter* temperate
16 bacteriophages. *BMC Microbiol*, 8, 49, doi:101186/1471-2180-8-49.
- 17 [Conlan AJK, Line JE, Hiatt K, Coward C, Van Diemen PM, Stevens MP, Jones](#)
18 [MA, Gog JR, Maskell DJ. \(2011\). Transmission and dose-response](#)
19 [experiments for social animals: a reappraisal of the colonization biology of](#)
20 [*Campylobacter jejuni* in chickens. *J R Soc Interface*,](#)
21 [doi:10.1098/rsif.2011.0125.](#)
- 22 Coward C, van Diemen PM, Conlan AJK, Gog JR, Stevens MP, Jones MA,
23 Maskell DJ. (2008). Competing isogenic *Campylobacter* strains exhibit

1 variable population structures *in vivo*. Appl Environ Microbiol, 74, 3857-
2 3867.

3 Cox NA, Hofacre CL, Bailey JS, Buhr RJ, Wilson JL, Hiatt KL, Richardson LJ,
4 Musgrove MT, Cosby DE, Tankson DJ, Vizzier YL, Cray PF, Vaughn LE,
5 Holt PS, Bourassaa DV. (2005). Presence of *Campylobacter jejuni* in
6 various organs one hour, one day, and one week following oral or
7 intracloacal inoculations of broiler chicks. Avian Dis, 49, 155-158.

8 Cox NA, Richardson LJ, Buhr RJ, Fedorka-Cray PJ. (2009). *Campylobacter*
9 species occurrence within internal organs and tissues of commercial caged
10 Leghorn laying hens. Poult Sci, 88, 2449-2456.

11 de Boer P, Wagenaar JA, Achterberg RP, van Putten JP, Schouls LM, Duim B.
12 (2002). Generation of *Campylobacter jejuni* genetic diversity *in vivo*. Mol
13 Microbiol, 44, 351-359.

14 de Zoete MR, Keestra AM, Roszczenko P, van Putten JPM. (2010). Activation of
15 human and chicken toll-like receptors by *Campylobacter* spp. Infect Immun,
16 78, 1229-1238.

17 EFSA. (2011). The European Union summary report on trends and sources of
18 zoonoses, zoonotic agents and food-borne outbreaks in 2009. EFSA J, 9,
19 2090.

20 EFSA. (2010), Panel on Biological Hazards (BIOHAZ). Scientific opinion on
21 quantification of the risk posed by broiler meat to human campylobacteriosis
22 in the EU EFSA J, 8, 1437 (89 pp), doi:102903/jefsa20101437.

- 1 Faragasan S. (2008). Evolution, development, mechanisms and function of IgA in
2 the gut. *Curr Opin Immunol*, 20, 170-177.
- 3 Flanagan RC, Neal-McKinney JM, Dhillon AS, Miller WG, Konkel ME. (2009).
4 Examination of *Campylobacter jejuni* putative adhesins leads to the
5 identification of a new protein, designated FlpA, required for chicken
6 colonization. *Infect Immun*, 77, 2399-2407.
- 7 Friedman A, Bar-shira E, Sklan D. (2003). Ontogeny of gut associated immune
8 competence in the chick. *Worlds Poult Sci J*, 59, 209-219.
- 9 Guerry P. (2007). *Campylobacter* flagella: not just for motility. *Trends Microbiol*,
10 15, 456-461.
- 11 Guerry P, Ewing CP, Hickey TE, Prendergast MM, Moran AP. (2000). Sialylation
12 of lipooligosaccharide core affects immunogenicity and serum resistance of
13 *Campylobacter jejuni*. *Infect Immun*, 68, 6656-6662.
- 14 Habib I, Louwen R, Uyttendaele M, Houf K, Vandenberg O, Nieuwenhuis EE,
15 Miller WG, van Belkum A, De Zutter L. (2009). Correlation between
16 genotypic diversity, lipooligosaccharide gene locus class variation and
17 Caco-2 cell invasion potential of *Campylobacter jejuni* isolates from chicken
18 meat and humans: contribution to virulotyping. *Appl Environ Microbiol*, 75,
19 4277-4288.
- 20 Hanel I, Borrmann E, Muller J, Muller W, Pauly B, Liebler-Tenorio EM, Schulze
21 F. (2009) Genomic and phenotypic changes of *Campylobacter jejuni* strains
22 after passage of the chicken gut. *Vet Microbiol*, 136, 121-129.

1 Hanninen ML, Hakkinen M, Rautelin H. (1999). Stability of related human and
2 chicken *Campylobacter jejuni* genotypes after passage through chick
3 intestine studied by pulsed-field gel electrophoresis. *Appl Environ*
4 *Microbiol*, 65, 2272-2275.

5 Hartley-Tassell LE, Shewell LK, Day CJ, Wilson JC, Sandhu R, Ketley JM,
6 Korolik V. (2010) Identification and characterization of the aspartate
7 chemosensory receptor of *Campylobacter jejuni*. *Mol Microbiol*, 75, 710-
8 730.

9 He H, Genovese KJ, Nisbet DJ, Kogut MH. (2006). Profile of Toll-like receptor
10 expressions and induction of nitric oxide synthesis by Toll-like receptor
11 agonists in chicken monocytes. *Mol Immunol*, 43, 783-789.

12 Hendrixson DR, DiRita VJ. (2004). Identification of *Campylobacter jejuni* genes
13 involved in commensal colonization of the chick gastrointestinal tract. *Mol*
14 *Microbiol*, 52, 471-484.

15 Hermans D, Van Deun K, Messens W, Martel A, Van Immerseel F, Haesebrouck
16 F, Rasschaert G, Heyndrickx M, Pasmans F. (2011a). *Campylobacter*
17 control in poultry by current intervention measures ineffective: urgent need
18 for intensified fundamental research. *Vet Microbiol*,
19 doi:10.1016/j.vetmic.2011.03.010.

20 [Hermans D, Van Deun K, Martel A, Van Immerseel F, Messens W, Heyndrickx](#)
21 [M, Haesebrouck F, Pasmans F. \(2011b\). Colonization factors of](#)
22 [Campylobacter jejuni in the chicken gut. *Vet Res*, 42, 82. doi:10.1186/1297-](#)
23 [9716-42-82.](#)

- 1 Ivanov II, Littman DR. (2011). Modulation of immune homeostasis by
2 commensal bacteria. *Curr Opin Microbiol*, 14, 106-114.
- 3 Jacobs-Reitsma WF, van de Giessen AW, Bolder NM, Mulder RW. (1995).
4 Epidemiology of *Campylobacter* spp at two Dutch broiler farms. *Epidemiol*
5 *Infect*, 114, 413-421.
- 6 Jennings JL, Sait LC, Perrett CA, Foster C, Williams LK, Humphrey TJ, Cogan
7 TA. (2011). *Campylobacter jejuni* is associated with but not sufficient to
8 cause vibronic hepatitis in chickens. *Vet Microbiol*, 149, 193-199.
- 9 Jeurissen SH, Janse EM, van Rooijen N, Claassen E. (1998). Inadequate anti-
10 polysaccharide antibody responses in the chicken. *Immunobiology*, 198,
11 385-395.
- 12 Kaiser P, Hughes S, Bumstead N. (1999). The chicken 9E3/CEF4 CXC
13 chemokine is the avian orthologue of IL-8 and maps to chicken
14 Chromosome 4 syntenic with genes flanking the mammalian chemokine
15 cluster. *Immunogenet*, 49, 673-684.
- 16 Kanungpean D, Kakuda T, Takai S. (2011). Participation of CheR and CheB in
17 chemosensory response of *Campylobacter jejuni*. *Microbiology*,
18 doi:101099/mic0047399-0.
- 19 [Karlyshev AV, Everest P, Linton D, Cawthraw S, Newell DG, Wren BW. \(2004\).](#)
20 [The *Campylobacter jejuni* general glycosylation system is important for](#)
21 [attachment to human epithelial cells and in the colonization of chicks.](#)
22 [Microbiology, 150, 1957-1964.](#)

- 1 Keeney KM, Finlay BB. (2011). Enteric pathogen exploitation of the microbiota-
2 generated nutrient environment of the gut. *Curr Opin Microbiol*, 14, 1-7.
- 3 Keestra AM, van Putten JPM. (2008). Unique properties of the chicken
4 TLR4/MD-2 complex: selective lipopolysaccharide activation of the
5 MyD88-dependent pathway. *J Immunol*, 181, 4354-4362.
- 6 Keestra AM, de Zoete MR, Bouwman LI, van Putten JPM. (2010). Chicken
7 TLR21 is an innate CpG DNA receptor distinct from mammalian TLR9. *J*
8 *Immunol*, 185, 460-467.
- 9 Konkel ME, Christensen JE, Dhillon AS, Lane AB, Hare-Sanford R, Schaberg
10 DM, Larson CL. (2007) *Campylobacter jejuni* strains compete for
11 colonization in broiler chicks. *Appl Environ Microbiol*, 73, 2297-2305.
- 12 Konkel ME, Klena JD, Rivera-Amill V, Monteville MR, Biswas D, Raphael B,
13 Mickelson J. (2004). Secretion of virulence proteins from *Campylobacter*
14 *jejuni* is dependent on a functional flagellar export apparatus. *J Bacteriol*,
15 186, 3296-3303.
- 16 Lam KM, DaMassa AJ, Morishita TY, Shivaprasad HL, Bickford AA. (1992).
17 Pathogenicity of *Campylobacter jejuni* for turkeys and chickens. *Avian Dis*,
18 36, 359-363.
- 19 Lamb-Rosteski JM, Kalischuk LD, Inglis GD, Buret AG. (2008). Epidermal
20 growth factor inhibits *Campylobacter jejuni*-induced claudin-4 disruption,
21 loss of epithelial barrier function, and *Escherichia coli* translocation. *Infect*
22 *Immun*, 76, 3390-3398.

- 1 Larson CL, Shah DH, Dhillon AS, Call DR, Ahn S, Haldorson GJ, Davitt C,
2 Konkel ME. (2008). *Campylobacter jejuni* invade chicken LMH cells
3 inefficiently and stimulate differential expression of the chicken CXCLi1
4 and CXCLi2 cytokines. *Microbiology*, 154, 3835-3847.
- 5 Li X, Swaggerty CL, Kogut MH, Chiang HI, Wang Y, Genovese KJ, He H, Stern
6 NJ, Pevzner IY, Zhou H. (2008) The paternal effect of *Campylobacter jejuni*
7 colonization in ceca in broilers. *Poult Sci*, 87, 1742-1747.
- 8 Li X, Swaggerty CL, Kogut MH, Chiang HI, Wang Y, Genovese KJ, He H, Zhou
9 H. (2010). Gene expression profiling of the local cecal response of genetic
10 chicken lines that differ in their susceptibility to *Campylobacter jejuni*
11 colonization. *PLoS One*, 5, e11827.
- 12 [Li X, Swaggerty CL, Kogut MH, Chiang HI, Wang Y, Genovese KJ, He H,
13 McCarthy FM, Burgess SC, Pevzner IY, Zhou H. \(2011\). Systemic response
14 to *Campylobacter jejuni* infection by profiling gene transcription in the
15 spleens of two genetic lines of chickens. *Immunogenetics*,
16 \[doi:10.1007/s00251-011-0557-1\]\(#\).](#)
- 17 Li YP, Ingmer H, Madsen M, Bang DD. (2008b). Cytokine responses in primary
18 chicken embryo intestinal cells infected with *Campylobacter jejuni* strains of
19 human and chicken origin and the expression of bacterial virulence-
20 associated genes. *BMC Microbiol*, 8, 107, doi:101186/1471-2180-8-107.
- 21 Lindblom G-B, Sjögren E, Kaijser B. (1986). Natural *Campylobacter* colonization
22 in chickens raised under different environmental conditions. *J Hyg Camb*,
23 96, 385-391.

1 Linde A, Ross CR, Davis EG, Dib L, Blecha F, Melgareo T. (2008). Innate
2 immunity and host defense peptides in veterinary medicine. *J Vet Intern*
3 *Med*, 22, 247-265.

4 Louwen R, Heikema A, van Belkum A, Ott A, Gilbert M, Ang W, Endtz HP,
5 Bergman MP, Nieuwenhuis EE. (2008). The sialylated lipooligosaccharide
6 outer core in *Campylobacter jejuni* is an important determinant for epithelial
7 cell invasion. *Infect Immun*, 76, 4431-4438.

8 MacPherson AJ, Uhr T. (2004). Induction of protective IgA by intestinal dendritic
9 cells carrying commensal bacteria. *Science*, 303, 1662-1665.

10 Manning G, Duim B, Wassenaar T, Wagenaar JA, Ridley A, Newell DG. (2001).
11 Evidence for a genetically stable strain of *Campylobacter jejuni*. *Appl*
12 *Environ Microbiol*, 67, 1185-1189.

13 Martins-Green M. (2001). The chicken chemotactic and angiogenic factor, cCAF,
14 a CXC chemokine. *Int J Biochem Cell Biol*, 33, 427-432.

15 Meade KG, Higgs R, Lloyd AT, Giles S, O'Farrelly C. (2009a). Differential
16 antimicrobial peptide gene expression patterns during early chicken
17 embryological development. *Dev Comp Immunol*, 33, 516-524.

18 Meade KG, Nirciandi F, Cahalane S, Reiman C, Allan B, O'Farrelly C. (2009b).
19 *Comparative in vivo* infection models yield insights on early host immune
20 response to *Campylobacter* in chickens. *Immunogenetics*, 61, 101-110.

21 Murphy C, Carrol C, Jordan KN. (2006). Environmental survival mechanisms of
22 the foodborne pathogen *Campylobacter jejuni*. *J Appl Microbiol*, 100, 623-
23 632.

- 1 Myszewski MA, Stern NJ. (1991). Phagocytosis and intracellular killing of
2 *Campylobacter jejuni* by elicited chicken peritoneal macrophages. Avian
3 Dis, 35, 750-755.
- 4 Newell DG. (2002). The ecology of *Campylobacter jejuni* in avian and human
5 hosts and in the environment. Int J Infect Dis, 6, 3S16-3S21.
- 6 Nielsen EM, Engberg J, Fussing V. (2001). Genotypic and serotypic stability of
7 *Campylobacter jejuni* strains during *in vitro* and *in vivo* passage. Int J Med
8 Microbiol, 291, 379-385.
- 9 [Oakland M, Jeon B, Sahin O, Shen Z, Zhang Q. \(2011\). Functional](#)
10 [characterization of a lipoprotein-encoding operon in *Campylobacter jejuni*.](#)
11 [Plos One, 6, e20084.doi:10.1371/journal.pone.0020084.](#)
- 12 Oza AN, Thwaites RT, Wareing DRA, Bolton FJ, Frost JA. (2002). Detection of
13 heat-stable antigens of *Campylobacter jejuni* and *C. coli* by direct
14 agglutination and passive hemagglutination. J Clin Microbiol, 40, 996-1000.
- 15 Pédrón T, Sansonetti P. (2004). Commensals, bacterial pathogens and intestinal
16 inflammation: an intriguing ménage à trois. Cell Host Microbe, 3, 344-347.
- 17 Peterson DA, McNulty NP, Guruge JL, Gordon JI. (2007). IgA response to
18 symbiotic bacteria as a mediator of gut homeostasis. Cell Host Microbe, 2,
19 328-339.
- 20 Phalipon A, Cardona A, Kraehenbuhl J-P, Edelman L, Sansonetti PJ, Corthésy B.
21 (2002). Secretory component: a new role in secretory IgA-mediated immune
22 exclusion *in vivo*. Immunity, 17, 107-115.

- 1 Ridley AM, Toszeghy MJ, Cawthraw SA, Wassenaar TM, Newell DG. (2008).
2 Genetic instability is associated with changes in the colonization potential of
3 *Campylobacter jejuni* in the avian intestine. J Appl Microbiol, 105, 95-104.
- 4 Ringoir DD, Korolik V. (2003). Colonisation phenotype and colonisation
5 potential differences in *Campylobacter jejuni* strains in chickens before and
6 after passage *in vivo*. Vet Microbiol, 92, 225-235.
- 7 Rosenquist H, Sommer HM, Nielsen, NL, Christensen BB. (2006). The effect of
8 slaughter operations on the contamination of chicken carcasses with
9 thermotolerant *Campylobacter*. Int J Food Microbiol, 108, 226-232.
- 10 Ruiz-Palacios GM, Escamilla E, Torres N. (1981). Experimental *Campylobacter*
11 diarrhea in chickens. Infect Immun, 34, 250-255.
- 12 Sahin O, Zhang Q, Meitzler JC, Harr BS, Morishita TY, Mohan R. (2001).
13 Prevalence, antigenic specificity, and bactericidal activity of poultry anti-
14 *Campylobacter* maternal antibodies. Appl Environ Microbiol, 67, 3951-
15 3957.
- 16 Sahin O, Luo N, Huang S, Zhang Q. (2003). Effect of *Campylobacter*-specific
17 maternal antibodies on *Campylobacter jejuni* colonization in young
18 chickens. Appl Environ Microbiol, 69, 5372-5379.
- 19 Sang FC, Shane SM, Yogasundram K, Hagstad HV, Kearney MT. (1989).
20 Enhancement of *Campylobacter jejuni* virulence by serial passage in chicks.
21 Avian Dis, 33, 425-430.

- 1 Sanyal SC, Islam KM, Neogy PK, Islam M, Speelman P, Huq MI. (1984).
2 *Campylobacter jejuni* diarrhea model in infant chickens. Infect Immun, 43,
3 931-936.
- 4 Scott AE, Timms AR, Connerton PL, El-Shibiny A, Connerton IF. (2007a).
5 Bacteriophage influence *Campylobacter jejuni* types populating broiler
6 chickens. Environ Microbiol, 9, 2341-2353.
- 7 Scott AE, Timms AR, Connerton PL, Loc Carrillo C, Adzfa Radzum K,
8 Connerton IF. (2007b). Genome dynamics of *Campylobacter jejuni* in
9 response to bacteriophage predation. PLoS Pathog, 3, e119,
10 doi:101371/journalppat0030119.
- 11 Scupham AJ. (2009). *Campylobacter* colonization of the turkey intestine in the
12 context of microbial community development. Appl Environ Microbiol, 75,
13 3564-3571.
- 14 Scupham AJ, Jones JA, Rettedal EA, Weber TE. (2010). Antibiotic manipulation
15 of intestinal microbiota to identify microbes associated with *Campylobacter*
16 *jejuni* exclusion in poultry. Appl Environ Microbiol, 76, 8026-8032.
- 17 Shaughnessy RG, Meade KG, Cahalane S, Allan B, Reiman C, Callanan JJ,
18 O'Farrelly C. (2009). Innate immune gene expression differentiates the early
19 avian intestinal response between *Salmonella* and *Campylobacter*. Vet
20 Immunol Immunopathol, 132, 191-198.
- 21 [Shaughnessy RG, Meade KG, McGivney BA, Allan B, O'Farrelly C. \(2011\).
22 Global gene expression analysis of chicken caecal response to
23 *Campylobacter jejuni*. Vet Immunol Immunopathol, 142, 64-71.](#)

- 1 Shoaf-Sweeney KD, Larson CL, Tang X, Konkel ME. (2008). Identification of
2 *Campylobacter jejuni* proteins recognized by maternal antibodies of
3 chickens. *Appl Environ Microbiol*, 74, 6867-6875.
- 4 Skanseng B, Trosvik P, Zimonja M, Johnsen G, Bjerrum L, Pedersen K, Wallin
5 N, Rudi K. (2007). Co-infection dynamics of a major food-borne zoonotic
6 pathogen in chicken. *PLoS Pathog*, 3, e157,
7 doi:101371/journalppat0030175.
- 8 Smith CK, Kaiser P, Rothwell L, Humphrey T, Barrow PA, Jones MA. (2005).
9 *Campylobacter jejuni*-induced cytokine responses in avian cells. *Infect*
10 *Immun*, 73, 2094-2100.
- 11 Smith CK, Abuoun M, Cawthraw SA, Humphrey TJ, Rothwell L, Kaiser P,
12 Barrow PA, Jones MA. (2008). *Campylobacter* colonization of the chicken
13 induces a proinflammatory response in mucosal tissues. *FEMS Immunol*
14 *Med Microbiol*, 54, 114-121.
- 15 Stahl M, Friis LM, Nothaft H, Liu X, Li J, Szymanski CM, Stintzi A. (2011). L-
16 fucose utilization provides *Campylobacter jejuni* with a competitive
17 advantage. *PNAS*, 108, 7194-7199.
- 18 Stern NJ, Meinersmann RJ, Cox NA, Bailey JS, Blankenship LC. (1990).
19 Influence of host lineage on cecal colonization by *Campylobacter jejuni* in
20 chickens. *Avian Dis*, 34, 602-606.
- 21 Stern NJ, Cox NA, Musgrove MT, Park CM. (2001). Incidence and levels of
22 *Campylobacter* in broilers after exposure to an inoculated seeder bird. *J*
23 *Appl Poult Res*, 10, 315-318.

- 1 Stern NJ. (2008). *Salmonella* species and *Campylobacter jejuni* cecal colonization
2 model in broilers. *Poult Sci*, 87, 2399-2403.
- 3 Tu QV, McGuckin MA, Mendz GL. (2008). *Campylobacter jejuni* response to
4 human mucin MUC2: modulation of colonization and pathogenicity
5 determinants. *J Med Microbiol*, 57, 795-802.
- 6 van de Giessen A, Mazurier SI, Jacobs-Reitsma W, Jansen W, Berkers P,
7 Ritmeester W, Wernars K. (1992) Study on the epidemiology and control of
8 *Campylobacter jejuni* in poultry broiler flocks. *Appl Environ Microbiol*, 58,
9 1913-1917.
- 10 Van Deun K, Haesebrouck F, Heyndrickx M, Favoreel H, Dewulf J, Ceelen L,
11 Dumez L, Messens W, Leleu S, Van Immerseel F, Ducatelle R, Pasmans F.
12 (2007) Virulence properties of *Campylobacter jejuni* isolates of poultry and
13 human origin. *J Medical Microbiol*, 56, 1284-1289.
- 14 Van Deun K, Pasmans F, Ducatelle R, Flahou B, Vissenberg K, Martel A, Van
15 den Broeck W, Van Immerseel F, Haesebrouck F. (2008). Colonization
16 strategy of *Campylobacter jejuni* results in persistent infection of the
17 chicken gut. *Vet Microbiol*, 130, 285-297.
- 18 van Dijk A, Veldhuizen EJA, Kalkhove SIC, Tjeerdsma-van Bokhoven JLM,
19 Romijn RA, Haagsman HP. (2007). The beta-defensin gallinacin-6 is
20 expressed in the chicken digestive tract and has antimicrobial activity
21 against food-borne pathogens. *Antimicrob Agents Chemother*, 51, 912-922.
- 22 van Gerwe T, Miflin JK, Templeton JM, Bouma A, Wagenaar JA, Jacobs-Reitsma
23 WF, Stegeman A, Klinkenberg D. (2009) Quantifying transmission of

1 *Campylobacter jejuni* in commercial broiler flocks. *Appl Environ Microbiol*,
2 75, 625-628.

3 Vegge CS, Brondsted L, Li YP, Bang DD, Ingmer H. (2009). Energy taxis drives
4 *Campylobacter jejuni* toward the most favorable conditions for growth.
5 *Appl Environ Microbiol*, 75, 5308-5314.

6 Widders PR, Thomas LM, Long KA, Tokhi MA, Panaccio M, Apos E. (1998).
7 The specificity of antibody in chickens immunised to reduce intestinal
8 colonisation with *Campylobacter jejuni*. *Vet Microbiol*, 64, 39-50.

9 Yamamoto M, Sato S, Hemmi H, Uematsu S, Hoshino K, Kaisho T, Takeuchi O,
10 Takeda K, Akira S. (2004). TRAM is specifically involved in the Toll-like
11 receptor 4-mediated MyD88-independent signaling pathway. *Nat Immunol*,
12 4, 1144-1150.

13 Zeng X, Xu F, Lin J. (2009). Molecular, antigenic, and functional characteristics
14 of ferric enterobactin receptor CfrA in *Campylobacter jejuni*. *Infect Immun*,
15 77, 5437-5448.

16 Zhu J, Meinersmann RJ, Hiett KL, Evans DL. (1999). Apoptotic effect of outer-
17 membrane proteins from *Campylobacter jejuni* on chicken lymphocytes.
18 *Curr Microbiol*, 38, 244-249.

19 Ziprin RL, Young CR, Byrd JA, Stanker LH, Hume ME, Gray SA, Kim BJ,
20 Konkel ME. (2001) Role of *Campylobacter jejuni* potential virulence genes
21 in cecal colonization. *Avian Dis*, 45, 549-557.

1 Ziprin RL, Young CR, Stanker LH, Hume ME, Konkel ME. (1999). The absence
2 of cecal colonization of chicks by a mutant of *Campylobacter jejuni* not
3 expressing bacterial fibronectin-binding protein. Avian Dis, 43, 586-589.
4