

**Running head: OBSERVING PAIN AND DETECTING TACTILE STIMULI**

Fibromyalgia patients and controls are equally accurate in detecting tactile stimuli while observing another in pain: an experimental study

Vandenbroucke, S.<sup>1</sup>, MSc, Crombez, G.<sup>1</sup>, PhD, Harrar, V.<sup>2</sup>, PhD, Brusselmans, G.<sup>3</sup>, MD, Devulder, J.<sup>3</sup>, MD, PhD, Spence, C.<sup>2</sup>, PhD, & Goubert, L.<sup>1</sup>, PhD.

1 Department of Experimental-Clinical and Health Psychology, Ghent University, Ghent, Belgium

2 Crossmodal Research Laboratory, Department of Experimental Psychology, Oxford University, Oxford, United Kingdom

3 Multidisciplinary Pain Clinic, Ghent University Hospital, Ghent, Belgium

**\* Correspondence:**

Sophie Vandenbroucke  
Ghent University  
Department of Experimental-Clinical and Health Psychology  
Henri Dunantlaan, 2,  
Ghent, 9000, Belgium  
Sophie.Vandenbroucke@ugent.be  
Tel: +32 (0)9 264 86 90  
Fax: +32 (0)9 264 64 71

Category: Research article

Number of text pages: 28

Number of tables: 3

Number of words: 8193

Number of figures: 2

Conflicts of Interest and Source of Funding: This research is supported by the Special Research Fund (BOF), Ghent University, grant number 01D3831. The authors have declared no conflicts of interest.

## OBSERVING PAIN AND DETECTING TACTILE STIMULI

2

## Abstract

Objective: This study investigated the effects of observing pain in others upon vicarious somatosensory experiences and the detection of somatosensory stimuli in both fibromyalgia patients (FM) and controls. The putative modulatory role of dispositional empathy, hypervigilance to pain, and central sensitization was examined.

Methods: FM patients (N=39) and controls (N=38) saw videos depicting pain-related (hands being pricked) and non-pain-related scenes, whilst occasionally experiencing vibrotactile stimuli themselves on the left, right, or both hands. Participants reported the location at which they felt a somatosensory stimulus. Tactile and visual scenes were presented in the same spatial location (congruent, e.g., left-left) or from opposite locations (incongruent, e.g., left-right). We calculated the proportion of correct responses, vicarious somatosensory experiences (i.e., trials on which an illusory somatosensory experience was reported while observing pain-related scenes), and neglect errors (i.e., only reporting the site congruent to the visual pain-related information when both hands had been stimulated).

Results: Observing another in pain resulted in an equal numbers of vicarious somatosensory experiences in both groups and facilitated the detection of tactile stimuli, especially during spatially congruent trials.

Counter to our expectations, this facilitation was not moderated by group. FM patients made fewer neglect errors. Hypervigilance for pain, dispositional empathy, and central sensitization did not exert a modulatory role.

Conclusion: Observing pain facilitates the detection of tactile stimuli in FM patients and controls. Overall, a low incidence of vicarious experiences was observed. Further research is needed to understand the role of attentional body focus in the elicitation of vicarious experiences.

FM: Fibromyalgia patients, CCE: congruency effect, TS: temporal summation

Keywords: vicarious experiences, visual enhancement, observation of pain, empathy, hypervigilance for pain

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

3

**Introduction**

Our senses do not operate independently of one another (Spence & Driver, 2004). For example, research has demonstrated that presenting visual information (e.g., a flash of light) may give rise to illusory experiences of touch (Lloyd, Mason, Brown, & Poliakoff, 2008; McKenzie, Poliakoff, Brown, & Lloyd, 2010). In particular, those individuals presenting with a large number of medically unexplained symptoms have been found to experience illusory tactile experiences (see Katzer, Oberfeld, Hiller, & Witthöft, 2011). Moreover, neuroimaging and neurophysiological studies have demonstrated that observing pain in others may activate brain areas similar to those activated when observers experience pain themselves (Jackson, Brunet, Meltzoff, & Decety, 2006; Osborn & Derbyshire, 2010). For example, those who experience vicarious pain (that is, an actual somatosensory experience in response to the observation of pain) show a hyperactivity of motor mirror neurons (enhanced motor-evoked potentials) to the observation of a needle penetrating the hand, relative to the needle having not yet penetrated the hand, as compared with controls (Fitzgibbon, Enticott, Bradshaw, Giummarra, Chou, Georgiou-Karistianis, & Fitzgerald, 2012a). These observations are intriguing as they indicate that tactile or nociceptive input may not be necessary to experience touch or pain.

Little research is yet available on the occurrence of vicarious somatosensory experiences and the mechanisms and conditions affecting this phenomenon (but see Fitzgibbon, Giummarra, Georgiou-Karistianis, Enticott, & Bradshaw, 2010; Fitzgibbon, et al., 2012b; Vandenbroucke et al., 2013). Fitzgibbon and colleagues (2010; 2012b) have put forward a neurobiological model to further our understanding of vicarious pain. They proposed several mechanisms to explain vicarious pain, such as hyperactivity of the somatosensory mirror systems, empathy or processes underlying empathy, central sensitization, hypervigilance to pain, and a history of chronic pain or trauma. Vision may not only induce vicarious somatosensory experiences, but may also influence the detection of tactile stimuli. For example, it has been demonstrated that simultaneously presenting a brief flash and a threshold-level tactile stimulus increases participants' ability to correctly perceive the tactile stimulus (i.e., increased number of 'hits'; Lloyd et al., 2008). From this perspective, the modulation of somatosensory experiences may represent a less extreme

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

4

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

variant of “illusory” experiences when observing another in pain. It has been argued that illusory experiences are akin to the kinds of misperceptions reported by patients with medically unexplained symptoms, and that similar processes are likely to be operating in each case (Lloyd et al., 2008).

In the present study, a variant of the crossmodal congruency task was used to investigate differences in vicarious somatosensory experiences between fibromyalgia patients (FM) and controls. FM patients were chosen as the clinical group because these patients suffer from medically unexplained symptoms, characterized by chronic widespread pain and central sensitization (see Staud et al., 2008, 2009), which have all been suggested as vulnerability factors in the production of vicarious and illusory sensations (see Fitzgibbon et al., 2010; 2012b; Katzer, Oberfeld, Hiller, & Witthöft, 2011). Both groups were presented two categories of videos in which pain-related situations (hands being pricked) or non-pain-related situations (e.g., a sponge being pricked) were shown. During this observation, the participants occasionally received vibrotactile stimuli themselves in the same spatial location (congruent trials) or in the opposite location (incongruent trials) as the visual stimuli. The participants were instructed to report the spatial location of the administered somatosensory stimuli as rapidly as possible. We examined whether the observation of pain-related scenes of a hand being pricked facilitated the detection of low-intensity vibrotactile stimuli compared to non-painful scenes. In contrary to our previous study (Vandenbroucke et al., 2013), instead of painful stimuli, we implemented non-painful vibrotactile stimuli near the perceptual threshold. This was done because Osborn and Derbyshire (2010) reported that most patients selected ‘tingling’ to describe their somatosensory vicarious experiences induced by observing pain. First, we hypothesized that the FM group would report more bodily illusions in response to the observation of pain (vicarious somatosensory experiences) than controls, as they have some of the suggested vulnerability factors to experience vicarious experiences, such as chronic pain, hypervigilance for pain, and central sensitization (Fitzgibbon et al., 2010, 2012b). We also explored whether there were any differences in neglect errors between FM patients and controls during the observation of pain-related videos (i.e. only reporting the site congruent to the visual information when both hands were stimulated). Second, we expected that the observation of pain-related visual scenes would facilitate the detection of vibrotactile stimuli as compared with non-pain-related scenes.

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

5

1  
2 Furthermore, we also expected to see a crossmodal congruency effect (CCE, that is, improved tactile acuity  
3  
4 in those conditions in which the visual and tactile stimuli were congruent). We hypothesized that this CCE  
5  
6 effect would be dependent on the type of visual information (pain-related and non-pain-related). As pain-  
7  
8 related visual stimuli may facilitate the detection of somatosensory stimuli, a higher CCE was expected  
9  
10 when pain-related visual stimuli were shown, as compared to non-pain-related visual stimuli. For  
11  
12 exploratory reasons, the effects and modulating role of dispositional empathy, hypervigilance to pain, and  
13  
14 central sensitization upon vicarious somatosensory experiences and general detectability were also  
15  
16 examined.  
17  
18  
19  
20  
21  
22

**Methods****Participants**

23  
24  
25  
26  
27  
28 Participants consisted of 39 patients with fibromyalgia (FM; 37 females; mean age=39.7 years,  
29  
30 SD=11.2, range 19-64 years) and a control group of 38 participants matched for age and sex (36 females;  
31  
32 mean age=38.3 years; SD=12.3; range 21-60 years). Fibromyalgia patients were recruited through the  
33  
34 Multidisciplinary Pain Clinic of Ghent University Hospital. Inclusion criteria included a diagnosis of  
35  
36 fibromyalgia (Wolfe et al., 2010), age between 18 and 65 years, and Dutch-speaking. Potential participants  
37  
38 were informed about the possibility to participate by means of a poster in the waiting room, information  
39  
40 given by their physician, and information letters. When they agreed to participate, they received a phone call  
41  
42 from the researcher providing details about the study. The fibromyalgia group reported pain complaints for,  
43  
44 on average, 10.01 years (SD=9.35 years). The mean score on the Widespread Pain Index (WPI) in the FM  
45  
46 group was 12.15 (SD: 2.72, range: 7-18); the mean score on the Severity Symptom scale (SS) scale was  
47  
48 9.64, (SD: 1.50, range: 6-12). Pain was reported on an average of 174 days (SD=21) over the last 6 months;  
49  
50  
51 46% reported a current poor state of health. All except one were Caucasian. Seventy-four percent were in  
52  
53 a relationship, 64% had children and 69% of them were not working because of the pain and received a  
54  
55 monthly allowance. Pain medication was used by 36.4% of the participants on the day of testing, especially  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

6

1  
2 in the FM group (69.2% of all FM patients). Twenty-six percent had a higher education (beyond the age of  
3  
4 18 years). On average, the FM group reported being unable to perform daily activities (work, household) on  
5  
6 101 days (SD=65) over the last 6 months. The control participants were recruited by means of  
7  
8 advertisements in the local newspapers. Inclusion criteria for the control participants were the absence of  
9  
10 chronic pain complaints or neurological or psychiatric conditions, Dutch-speaking, and aged between 18 and  
11  
12 65 years. Ninety-seven percent of the participants in the control group (N=38; 36 females; mean age=38.3  
13  
14 years, range 21-60 years) reported a good, very good, or excellent current state of health. Sixty-three percent  
15  
16 of the control participants had a relationship and 45% had children. The majority (82%) had had higher  
17  
18 education; 18% were unemployed. At the end of the experiment, the participants received 40 euro as  
19  
20 reimbursement for their expenses. The experiment lasted for approximately 1.5 hours and was part of a  
21  
22 larger protocol that had been approved by the Ethical Committee of the Ghent University Hospital.  
23  
24  
25  
26  
27

**Apparatus and stimuli**

28  
29  
30  
31 **Visual stimuli.** The visual stimuli consisted of two categories of videos (pain-related versus non-  
32  
33 pain related), each with a duration of 3000ms. The pain-related category included two scenes depicting a left  
34  
35 and right hand, with one of the two hands being pricked with a syringe or safety pin (2000ms after the onset  
36  
37 of the video). The non-pain related category also consisted of 2 scenes. In one scene, a left and right hand  
38  
39 was presented in which one of these hands was approached by a hand that was not holding an object (though  
40  
41 executing the same action as in the pain-related videos). In the second scene, one of the two hands was  
42  
43 replaced by a sponge that was pricked with a syringe. In this way, a human feature was always present in the  
44  
45 videos (e.g. a left or right hand). The penetration took place after 2000ms as in the first category. The  
46  
47 different scenes and the location of the sponge and movement were counterbalanced across videos. The  
48  
49 location of the penetration (left versus right hand) and type of category were counterbalanced across videos.  
50  
51 Videos were presented by INQUISIT Millisecond software (<http://www.millisecond.com>) on a Dell  
52  
53 computer with a 19-inch CRT-monitor.  
54  
55  
56  
57  
58  
59  
60

## OBSERVING PAIN AND DETECTING TACTILE STIMULI

7

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

**Somatosensory stimuli.** Vibrotactile stimuli (50 Hz, 50 ms) were delivered by means of two resonant-type tactors (C-2 tactor, Engineering Acoustics, Inc.) encased in a housing that was 3.05 cm in diameter and 0.79 cm high, with a skin contactor that was 0.76 cm in diameter. The somatosensory stimuli were delivered on the skin between the thumb and index finger on the back of the hand. All stimulus characteristics (amplitude, duration, and frequency) were controlled through a self-developed software program that was used to control the tactors. For each participant, the threshold intensity level was individually determined prior to the experiment (see Procedure-Preparation phase). Both hands were placed on the table in front of the screen and covered with a cardboard box so that they were not visible. Four different series of 20 stimuli/trials (two series for each hand) were randomly administered (80 stimuli/trials in total). First, a visual stimulus (an “X” in the middle of the screen, 1000ms duration) was presented combined with a somatosensory stimulus on the left or right hand. The participants were instructed to report whether they felt a somatosensory stimulus (“yes” or “no”). Responses were entered by the experimenter who pressed the corresponding response button on a keyboard. Each series started with a stimulus of 0.068 W. The intensity was decreased by 0.0002 W whenever the participants reported feeling the stimulus, and increased by 0.0002 W when no sensation was reported. After 80 trials, this resulted in a threshold intensity for each hand, which was based upon the mean intensity of the last stimuli of the two series for that particular hand. From these threshold intensities (threshold left hand:  $M=0.06W$ ,  $SD=0.006W$ , range: 0.004W-0.21W; threshold right hand:  $M=0.05W$ ,  $SD=0.008W$ , range: 0.006W-0.17W), 1/8 was subtracted (termed subthreshold) and added to the threshold (termed above threshold), which resulted in four different intensities (sub and above threshold, one for each hand; see Press, Taylor-Clarke, Kennett, & Haggard, 2004). Threshold intensities did not differ between groups (left hand:  $t(75)=-.25, p=.80$ ; right hand:  $t(75)=-.25, p=.80$ ).

51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Central sensitization: temporal summation.** Central sensitization was assessed using a temporal summation (TS) procedure (Staud, Craggs, Perlstein, Robinson, & Price, 2008). TS refers to an increased pain experience evoked by the repeated presentation of stimuli of the same intensity. Staud et al. (2009) has provided support for the presence of an alteration of central pain sensitivity in FM patients. The probe

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

8

1  
2 temperature was adjusted to each individual's heat pain sensitivity, which was determined during a  
3 preliminary phase (Staud et al., 2008) and was administered by means of a 'Contact Heat Evoked Potential  
4 Stimulator' (CHEPS) (Medoc Advanced Medical Systems, Ramat Yishai, Israel). During this preliminary  
5 phase of the study, a train of 6 stimuli at 0.33Hz were administered starting with peak pulse temperatures of  
6 47°C. After each pulse train, the participants reported the intensity of pain experienced between the first and  
7 last pulse by means of a 100-point Numeric Rating Scale (NRS; 0=no sensation; 100=intolerable pain). This  
8 intensity was subsequently raised until the participants achieved NRS ratings of  $45 \pm 10$  after 6 pulses. The  
9 participants were informed that the intensity could increase, decrease, or stay the same within each train of  
10 pulses. The test phase procedure consisted of a train of 6 heat pulses to the palm of the right hand in which  
11 the probe temperature was adjusted to each individual's heat pain sensitivity determined during the  
12 preliminary phase. Each train started with a 40s baseline followed by 6 pulses. The temperature of the  
13 thermal probe increased from baseline to peak temperature by 8°C/s, before returning to baseline at a rate of  
14 8°C/s. The duration of each heat pulse was always 3s (1.5s rise time; 1.5s return time; 0.33 Hz). The TS test  
15 phase procedure was repeated six times.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

**Self report measures**

35  
36  
37 The scale to rate the intensity of the different pulses during the acquisition of temporal summation  
38 ranged from 0 to 100 in increments of 5 (Vierck, Cannon, Fry, Maixner, &Whitsel, 1997) with verbal  
39 descriptors at intervals of 10: 10, warm; 20, a barely painful sensation; 30, very weak pain; 40, weak pain;  
40 50, moderate pain; 60, slightly strong pain; 70, strong pain; 80, very strong pain; 90; nearly intolerable pain;  
41 and 100, intolerable pain.  
42  
43  
44  
45  
46  
47

48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Vigilance to pain was assessed by the Dutch version of the Pain Vigilance and Awareness  
Questionnaire (PVAQ; McCracken, 1997, Roelofs, Peters, Muris, &Vlaeyen, 2002). This questionnaire  
consists of 16 items assessing awareness, consciousness and vigilance to pain on a six-point scale (0= never;  
5= always). Higher scores on the PVAQ are indicative of greater pain-related vigilance and awareness. The  
questionnaire can be used in both clinical (McCracken, 1997; Roelofs, Peters, McCracken, &Vlaeyen,



**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

9

2003) and non-clinical (McWilliams & Asmundson, 2001; Roelofs et al., 2002) samples. The Dutch version of the PVAQ is reliable and valid (Roelofs et al., 2002, 2003). Cronbach's alpha for the present study was 0.87.

Empathic disposition was assessed by means of the Dutch version of the Interpersonal Reactivity Index (IRI; Davis, 1983; De Corte et al., 2007). The questionnaire contains 28 items and consists of 4 subscales: Perspective Taking (i.e., cognitively taking the perspective of another, e.g., "I sometimes try to understand my friends better by imagining how things look from their perspective."), Fantasy (i.e., emotional identification with characters in books, movies etc., e.g., "When I watch a good movie, I can very easily put myself in the place of a leading character."), Empathic Concern (i.e., feeling emotional concern for others, e.g., "I am often quite touched by things that I see happen.") and Personal Distress (i.e., negative feelings in response to the distress of others, e.g., "When I see someone who badly needs help in an emergency, I go to pieces."). Each item is answered on a scale ranging from 1 ('does not describe me very well') to 5 ('describes me very well'). This questionnaire is reliable and valid (Davis et al., 1993; De Corte et al., 2007). Cronbach's alpha's in the current study were 0.84 (fantasy scale), 0.68 (empathic concern), 0.72 (personal distress), and 0.32 (perspective taking). The latter subscale was omitted from the analyses because of the low reliability of the scores.

Anxiety and depression was measured with the Dutch version of the Hospital Anxiety and Depression Scale (HADS; Spinhoven et al., 1997, Zigmond & Snaith, 1983) consisting of 14 items, of which 7 screen for symptoms of anxiety and 7 for symptoms of depression. Items are rated on a 4-point scale representing the degree of distress experienced during the previous week. Higher scores indicate higher feelings of anxiety and depression. In the present study Cronbach's alpha was .93.

Vicarious pain sensations in daily life were measured by means of four items adapted from Banissy et al. (2009). Participants were asked to indicate on an eleven point scale (0-10; totally disagree - totally agree) the extent to which they agreed with the questions: "Do you feel pain in your own body when you see someone accidentally bump into the corner of the table?", "Do you have the feeling that you are experiencing pain when you observe another person in pain?", "Do you feel bodily pain when you observe another person

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

10

1  
2 in pain?”, “Do you feel a physical sensation (e.g., tingling, stabbing) when you observe another person in  
3  
4 pain” (see Vandenbroucke et al., 2013). In the present study Cronbach’s alpha was .87.  
5  
6  
7

**Procedure**

8  
9  
10  
11 Upon arrival, the procedure started with signing the informed consent form. Subsequently, the  
12  
13 Fibromyalgia diagnostic criteria (Wolfe et al., 2010) were checked for each participant. All FM patients  
14  
15 fulfilled the Fibromyalgia diagnostic criteria (Wolfe et al., 2010). Thereafter, the participants were seated in  
16  
17 front of a table, about 60 cm away from the computer screen.  
18

**Behavioral paradigm.**

19  
20  
21  
22 *Preparation phase.* First, the detection threshold was determined for each hand separately. The  
23  
24 participants were informed that during the experiment they would feel subtle stimuli, varying in intensity  
25  
26 and length, on their left, right, or both hands. Participants were informed that different videos would be  
27  
28 presented which they needed to watch attentively. The hands of the participants were placed on the table and  
29  
30 covered by a cardboard box placed on the table in front of the screen. The participants were told that the  
31  
32 intensity of the somatosensory stimuli could vary across their hands and that there would also be trials  
33  
34 without any stimulus. In reality, only two fixed predetermined intensities with a fixed duration were applied  
35  
36 (threshold intensity  $\pm 1/8$ ) for each hand.  
37  
38

39  
40 *Experiment phase.* Each trial began with a fixation cross (1000 ms duration) presented in the middle of  
41  
42 the computer screen. Next, one of the scenes was presented. In 75% of the trials, a tactile stimulus was  
43  
44 delivered 2450ms after video onset to either the left hand, the right hand, or to both hands of the participant.  
45  
46 In line with Banissy and Ward (2007), the somatosensory stimulus was administered with a delay (in this  
47  
48 study 450ms after the visual image of the needle penetrating). This resulted in the following trial types:  
49  
50 congruent trials, incongruent trials, and trials in which no somatosensory stimuli were administered or in  
51  
52 which both of the participant’s hands received somatosensory stimuli. In congruent trials, the somatosensory  
53  
54 and visual stimuli were presented from the same spatial location (e.g., on the right). In the incongruent trials,  
55  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

11

1  
2 the somatosensory and visual stimuli were presented from opposite locations (e.g., one on the left and the  
3  
4 other on the right). The experiment started with 8 practice trials.

5  
6 The actual experimental phase consisted of three blocks of 64 trials, resulting in a total of 192 trials.  
7  
8 There were 48 congruent trials, 48 incongruent trials, 48 trials without sensory stimuli and 48 trials with  
9 somatosensory stimuli presented to both hands. The order of the trial types was randomized within each  
10 block and the intensity of the somatosensory stimuli (under and above threshold) were equally distributed  
11 within and across each block. An overview of all trial types is presented in Table 1. During each trial, the  
12 participants reported whether a physical experience was felt by reporting as rapidly as possible ‘YES’ and to  
13 discriminate the spatial location of the somatosensory stimuli by reporting “left”, “right” or “both” (see  
14 Figure 1). After the video had ended and 2000 ms elapsed, the word ‘next’ was presented on the screen.  
15 Then, the experimenter coded the response by pressing the corresponding response button (left, right, both  
16 or no response). In this manner, the time to respond was equal for every participant. The experiment took  
17 approximately 20 min.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 *Post-experiment phase.* After the experiment, participants filled out self-report scales  
32 measuring hypervigilance for pain (PVAQ) and empathic disposition (IRI). After a short break, the  
33 participants  
34 continued with the temporal summation measurement.  
35  
36  
37  
38  
39  
40  
41

**Statistical analysis**

42  
43  
44 The number of false alarms was calculated from the incongruent trials and from the trials without any  
45 somatosensory stimuli when erroneously a somatosensory stimulus was reported in the same spatial location  
46 as the visual stimulus. These false alarms were labeled ‘vicarious somatosensory experiences’ when the  
47 visual stimulus contained pain-related information. First, we tested whether the number of false alarms was  
48 dependent upon the type of video. As all participants observed both categories of videos and the number of  
49 false alarms during both categories of movies were not normally distributed, non-parametric analyses for  
50 related samples (Wilcoxon) were used. As we were particularly interested in those false alarms during pain-  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

12

1  
2 related videos, the number of vicarious somatosensory experiences was further selected as the dependent  
3  
4 variable.

5  
6 To test whether group predicted the number of vicarious somatosensory experiences, count  
7  
8 regression models were applied. The use of linear models was not appropriate due to the fact that the  
9  
10 frequency of responses had a skewed distribution that violated the normality assumption (Vives, Losilla, &  
11  
12 Rodrigo, 2006). Poisson regression is the basic model to analyze count data, but the variance of counts is  
13  
14 often larger than the mean (overdispersion). The Negative Binomial (NB) regression, a Poisson regression  
15  
16 with an overdispersion, may therefore fit the data better (e.g., Gardner, Mulvey, & Shaw, 1995). As count  
17  
18 data may additionally exhibit a lot of zero counts, zero-inflated extensions of both models, called Zero-  
19  
20 Inflated Poisson (ZIP) and Zero-Inflated NB (ZINB) models have been developed (see Karazsia & Van  
21  
22 Dulmen, 2010; Loeys, Moerkerke, De Smet, & Buysse, 2012). Deviance tests and the Vuong test were used  
23  
24 to select the best fitting count distribution for the dependent variable. After the best fitting count model was  
25  
26 chosen, a model with ‘group’ as predictor was added. In a further exploration of the data, central  
27  
28 sensitization, hypervigilance for pain, and dispositional empathy were added as a second predictor in  
29  
30 separate models to test whether they had a modulating role. Dummy coding was used for the categorical  
31  
32 variables and standardized z-scores for the continuous predictors. Regression coefficients were  
33  
34 exponentiated (eB) and called Rate Ratios (RRs). In percentages— $100 \times (eB - 1)$ —RRs reflect the  
35  
36 percentage decrease ( $RR < 1$ ) or increase ( $RR > 1$ ) in the expected frequency of vicarious somatosensory  
37  
38 experiences for each 1-unit increase in the independent variable.  
39  
40  
41  
42  
43

44  
45 Second, to investigate whether the observation of pain-related and non-pain-related scenes modulated  
46  
47 the detection of tactile stimuli, the proportion of correct responses (left versus right) for congruent and  
48  
49 incongruent trials for each category of visual information was calculated (pain-related versus non-pain-  
50  
51 related). A 2 (video category: pain-related versus non-pain-related) x 2 (congruency: congruent versus  
52  
53 incongruent) repeated measures ANOVA was performed, with congruency and type of video entered as a  
54  
55 within-participant variables and ‘group’ as a between-subject variable. In a further exploration of the data,  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

13

1  
2 central sensitization, hypervigilance for pain, and dispositional empathy were added as a covariate in  
3  
4 separate models to test whether they had a modulating role.

5  
6 The number of neglect errors was also calculated based upon those trials in which both hands were  
7  
8 stimulated, defined as reporting *only* the site congruent to the visual information and missing the fact that  
9  
10 there had been two tactile stimuli, one on each hand. Non-parametric analyses for related samples  
11  
12 (Wilcoxon) were used to test whether the number of neglect errors was dependent upon the type of video.  
13  
14 Count regression models were applied in which the dependent variable was the number of neglect errors  
15  
16 during pain-related visual information. After the best fitting count model was chosen, a model with ‘group’  
17  
18 as predictor was added. In a further exploration of the data, central sensitization, hypervigilance for pain,  
19  
20 and dispositional empathy were added as a second predictor in separate models to test whether they had a  
21  
22 modulating role. R (version 2.15.1) was used to fit the count models. Repeated measures were conducted  
23  
24 with an alpha < 0.05, using SPSS statistical software, version 21.0 for Windows.  
25  
26  
27

**Results****Descriptives**

28  
29  
30  
31  
32  
33 Mean scores, standard deviations, and correlations are presented in Tables 2 and 3. Because the  
34  
35 variables vicarious somatosensory experiences, vicarious pain during daily life, neglect errors, empathic  
36  
37 concern, and temporal summation (difference in reported intensity between first and last stimulus) did not  
38  
39 have a normal distribution (Kolmogorov-Smirnoff,  $p < .05$ ), Spearman correlations were computed for these  
40  
41 particular variables. A significant difference was found between FM patients and controls in fantasy scale  
42  
43 scores ( $t(75) = 3.49, p = .001$ ), PVAQ ( $t(75) = -4.27, p < .001$ ), and HADS ( $t(75) = -8.99, p < .001$ ), indicating that  
44  
45 FM patients were more hypervigilant for pain, obtained lower scores on the fantasy scale and were more  
46  
47 anxious and felt more depressed compared with control participants. Threshold intensities for the left hand  
48  
49 ( $t(75) = -.25, p = .80$ ) and right hand ( $t(75) = -.25, p = .80$ ) were similar for both groups. The control group  
50  
51 reported significantly more vicarious pain experiences during daily life than the FM group (Mann-Whitney,  
52  
53  $p = .03$ ). Regarding temporal summation, no differences in perceived intensity of the thermal stimuli were  
54  
55 found across both groups ( $t(75) = -1.29, p = .20$ ). The average reported intensity of the first stimulus ( $M = 33.86$ ;  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

14

1  
2  $SD=18.74$ ) and last stimulus ( $M=39.89$ ;  $SD=17.84$ ) over 6 trains was calculated. The average of the reported  
3  
4 intensity of the first stimulus was not normally distributed (Kolmogorov-Smirnov,  $p<.05$ ). Therefore, a  
5  
6  $\log_{10}$  transformation was performed for the reported intensity of first and last stimuli in the analysis. A  
7  
8 repeated measures ANOVA was performed including a within-participant variable stimulus (first versus  
9  
10 last) and between-participant variable group (FM versus control). The reported intensity of the last stimulus  
11  
12 was significant larger compared with the first ( $F(1,75)=28,94, p<.001$ ). No group x stimulus interaction was  
13  
14 observed ( $F(1,75)=2,4, p=.13$ ). In 2.5% of a total of 3648 trials, vicarious somatosensory experiences were  
15  
16 reported (90 vicarious somatosensory experiences from a total of 3648 trials). Of all vicarious  
17  
18 somatosensory experiences, 46.7% occurred in the FM group ( $n=42$ ) and 53.3% in the control group ( $n=48$ ). In  
19  
20 19.2% of the trials in which both hands were stimulated during the observation of pain-related stimuli,  
21  
22 neglect errors were made (350 from a total of 1824). Of all neglect errors, 41.1% occurred in the FM group  
23  
24 ( $n=144$ ) and 58.9% in the control group ( $n=206$ ). Data of 1 FM participant were excluded from the analyses  
25  
26 with regard to the crossmodal congruency task, as data on this task were missing.  
27  
28  
29  
30  
31  
32

**Vicarious somatosensory experiences**

33  
34  
35 Participants reported significantly more false alarms when scenes from the pain-related category  
36  
37 were shown, as compared to the non-pain-related category (Wilcoxon,  $p<.001$ ). This indicates that the type  
38  
39 of visual information (pain-related versus non-pain-related) is important as participants erroneously reported  
40  
41 more somatosensory stimuli in the same spatial location as the visual stimulus when it contained pain-  
42  
43 related information. To test the influence of group on the number of vicarious somatosensory experiences,  
44  
45 the NB model was found to be the best fitting count model ( $\chi^2[1, N=77]=54.38, p<.001$ ;  $V=-.79, p=.21$ ). In a  
46  
47 first step, group was added as a predictor. In contrary to our hypothesis, the results revealed that the number  
48  
49 of vicarious somatosensory experiences was not dependent upon group ( $p=.72$ ). In order to explore the role  
50  
51 of individual differences in PVAQ and the IRI, several additional models were run with PVAQ or IRI as a  
52  
53 second predictor and in interaction with group to explore its modulating role. No interactions were found  
54  
55 between group and EC ( $p=.86$ ), FS ( $p=.41$ ), PD ( $p=.93$ ), and temporal summation ( $p=.72$ ). A marginally  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

15

1  
2 significant interaction was found between group and PVAQ ( $p=.052$ ). For FM patients, the probability of  
3 making vicarious somatosensory errors decreased by 57% ( $RR=.43$ ) for every 1-unit increase in  
4 hypervigilance for pain. For the control group, the probability of making vicarious somatosensory errors  
5 increased by 7% ( $RR=1.07$ ) for every 1-unit increase in hypervigilance for pain. No main effect of  
6 hypervigilance for pain was found ( $p=.76$ ).  
7  
8  
9  
10  
11  
12  
13  
14

**Detection accuracy**

15  
16  
17 In line with our hypotheses, a 2 (video: pain-related versus non-pain-related) x 2 (congruency: congruent  
18 versus incongruent) repeated measures ANOVA with the between-participant variable 'group' (FM versus  
19 control) showed a main effect for video ( $F(1,74)=73.82, p<.001$ , Cohen's  $d=.46$ , [95% CI: .35, .57]). In  
20 general, pain-related videos resulted in better detection of tactile stimulation compared with non-pain related  
21 videos both in congruent trials ( $t(75)=8.44, p<.001$ , Cohen's  $d=.65$ , [95% CI: .48, .82]) as in incongruent  
22 trials ( $t(75)=4.10, p<.001$ , Cohen's  $d=.26$ , [95% CI: .14, .37]). Also, a main effect of congruency was found  
23 ( $F(1,74)=29.30, p<.001$ , Cohen's  $d=.27$ , [95% CI: .16, .38]). An interaction occurred between congruency  
24 and video: the CCE depended on the type of video presented ( $F(1,74)=17.08, p<.001$ , Cohen's  $d=.59$ , [95%  
25 CI: .26, .91]) (Figure 2). A paired sample t-test showed that the CCE was only significant for the pain-related  
26 videos ( $t(75)=-6.39, p<.001$ , Cohen's  $d=.45$ , [95% CI: .30, .61]), indicating that the increased detection  
27 accuracy in congruent trials compared with incongruent trials occurred only when pain-related videos were  
28 shown. The CCE was not significant for the non-pain related videos ( $t(75)=-1.4, p=.17$ ). No main effect  
29 occurred for group ( $F(1,74)=.42, p=.52$ ): Fibromyalgia patients were not more or less sensitive to the sensory  
30 stimuli. No interaction was found between group and video ( $F(1,74)=.01, p=.91$ ), between group and  
31 congruency ( $F(1,74)=.40, p=.53$ ), or between group, video and congruency ( $F(1,74)=.58, p=.45$ ).  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 Centered PVAQ and IRI subscales were entered separately as covariates. No main effects were found for  
52 PVAQ,  $F(1,73)=.18, p=.68$ , fantasy scale,  $F(1,73)=2.67, p=.11$ , personal distress,  $F(1,73)=.44, p=.51$ ,  
53 empathic concern,  $F(1,73)=.90, p=.35$ . Next, the centered difference between the first and the last intensity  
54  
55  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

16

1  
2 score (temporal summation) was added as a covariate in the above-described analyses. No main effect of  
3  
4 temporal summation upon the proportion of correct responses was found ( $F(1,73)=.54, p=.46$ ).  
5  
6  
7

**Neglect errors**

8  
9  
10  
11 Trials in which both hands were stimulated, with participants only reporting sensory experiences on the  
12 side congruent with the visual stimulus, provide additional information concerning somatosensory  
13 modulation. When both hands were stimulated, the participants tended to neglect the side that was  
14 incongruent with the visual stimulus more when scenes of the pain-related category were shown, as  
15 compared to the non-pain-related category (Wilcoxon,  $p<.001$ ); i.e., they reported significantly more often  
16 the side that was congruent with the visual stimulus when a pain-related situation was depicted compared  
17 with a non-pain-related visual situation. Next, the impact of group (FM versus control) was examined. The  
18 NB model was found to be the best fitting count model ( $\chi^2[1, N=77]=19.35, p<.001; V=.24, p=.40$ ). In a first  
19 step, group was added as a predictor. Results showed that the number of neglect errors during the  
20 observation of pain-related stimuli was dependent upon group ( $p=.02, RR=.70$ ). Noteworthy here is the fact  
21 that FM patients made 30% *less* neglect errors than the control group.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

35 In order to explore the role of individual differences in PVAQ, IRI, and central sensitization, several  
36 additional models were run with a second predictor and testing the interaction with group to explore its  
37 modulating role. No significant interactions were found with PVAQ ( $p=.64$ ), EC ( $p=.17$ ), FS ( $p=.43$ ), PD  
38 ( $p=.43$ ), or temporal summation ( $p=.24$ ).  
39  
40  
41  
42  
43  
44  
45

**Discussion**

46  
47  
48 The present study was designed to investigate (1) whether the observation of pain-related scenes  
49 elicits more vicarious somatosensory experiences in those patients suffering from FM compared with  
50 healthy controls; and (2) whether the observation of pain-related and non-pain-related scenes modulates the  
51 detection of tactile stimuli. Additionally, we explored the effects of potential moderating factors proposed by  
52 Fitzgibbon et al. (2010, 2012b), i.e., dispositional empathy, hypervigilance to pain, the presence of chronic  
53  
54  
55  
56  
57  
58  
59  
60



**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

17

1  
2 pain, and central sensitization. Participants were presented with a series of videos showing hands being  
3  
4 pricked and non-pain-related information such as a sponge being pricked whilst receiving occasionally near-  
5  
6 threshold vibrotactile stimuli themselves. In congruent trials, the somatosensory and visual stimuli were  
7  
8 applied to the same spatial location (e.g., on the right). In the incongruent trials, the somatosensory and  
9  
10 visual stimuli were presented from the opposite spatial location (e.g., left and right). Trials in which both of  
11  
12 the participant's hands were stimulated and trials without tactile stimulation were present. Participants were  
13  
14 required to report if and where they felt a somatosensory stimulus.  
15  
16

17  
18 In this study, only a small number of vicarious somatosensory experiences were observed (2.5%). In  
19  
20 the literature, percentages range from 1.6% for vicarious touch (Banissy et al., 2009), 16.2% for vicarious  
21  
22 pain in amputees (Fitzgibbon et al., 2010a), 6.6% (Vandenbroucke et al., 2013, study 1), 22.9%  
23  
24 (Vandenbroucke et al., 2013, study 2), and 30.0% for vicarious pain in a general population (Osborn &  
25  
26 Derbyshire, 2010). This variability is largely dependent upon the group investigated, and the criteria that are  
27  
28 used (questionnaires versus experimental paradigm). The percentage of reported vicarious pain in this study  
29  
30 is smaller than that reported in amputees (Fitzgibbon et al., 2010a), suggesting that prior trauma may be an  
31  
32 important modulator. Contrary to our hypotheses, no differences were found in vicarious somatosensory  
33  
34 experiences between the FM patients and the controls. In general, and across groups, the observation of pain  
35  
36 in another enhanced stimulus detection as compared to non-pain-related scenes in both the congruent and  
37  
38 incongruent trials. In line with our expectations, detection was better in congruent trials than in incongruent  
39  
40 trials only when pain-related information was shown. In general, neglect errors were more frequently made  
41  
42 (19.2%) compared with vicarious somatosensory experiences. FM patients made significantly fewer neglect  
43  
44 errors (30%) as compared with controls. Dispositional empathy, hypervigilance for pain, and central  
45  
46 sensitization had no modulating role upon the detection of vibrotactile stimuli, the experience of vicarious  
47  
48 experiences, or on neglect errors.  
49  
50  
51  
52

53  
54 Neglect errors were frequently observed in this study, which suggests that the observation of pain-  
55  
56 related information may modulate somatosensory experiences rather than induce illusory experiences. The  
57  
58 lower number of neglect errors in the FM group is intriguing and needs further exploration and elaboration.  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

18

1  
2 One possible explanation here is that an excessive attentional focus on the body may have come into play. It  
3  
4 is assumed that chronic pain patients are preoccupied with bodily cues signaling potential physical harm  
5  
6 (Crombez et al., 2013). In this way, the FM patients may have been less misled by the presence of visual  
7  
8 pain-related stimuli as their attention was focused on both hands, in contrast with controls who appear to  
9  
10 have been paying more attention to the site congruent to the visual pain-related information. This assumed  
11  
12 preoccupation with bodily cues may also explain the same number of vicarious somatosensory errors in both  
13  
14 groups. On the other hand, self-report of hypervigilance did not seem to modulate the number of neglect  
15  
16 errors. Another possibility may be that FM patients lack response inhibition as they detect vibrotactile  
17  
18 stimuli on both hands, whereas healthy controls tend to report only the vibrotactile stimulus congruent to the  
19  
20 visual stimulus and inhibit the detection of the incongruent vibrotactile stimulus. This is consistent with the  
21  
22 results of a study by Glass et al. (2011) reporting that FM patients showed lower activation in the inhibition  
23  
24 and attention networks and increased activation in other areas. Further research could explore whether this  
25  
26 inhibition theory played a role in the different number of neglect errors reported in the two groups tested  
27  
28 here.  
29  
30  
31  
32

33 Our findings corroborate previous research demonstrating that spatial coincidence plays a role in  
34  
35 multisensory integration (Spence, 2013). In the present study, the higher proportion of correct responses in  
36  
37 congruent as compared with incongruent trials, when pain-related information was shown, suggests that the  
38  
39 visual system may dominate somatosensation when visual and tactile processing provide conflicting  
40  
41 information (e.g., incongruent trials), or that vision may enhance sensitivity when providing similar  
42  
43 information (e.g., congruent trials). The finding that the congruency effect was only present when pain-  
44  
45 related scenes were shown attests to the relevance of the content of the visual information for tactile  
46  
47 sensitivity. That vision should dominate somatosensation may also explain the occurrence of neglect errors,  
48  
49 as attention may be more directed to the site congruent to the visual pain-related information. The content of  
50  
51 the visual information was relevant as the site congruent to the pain-related videos was more frequently  
52  
53 reported compared with non-pain related information, although both hands were stimulated.  
54  
55  
56  
57  
58  
59  
60

## OBSERVING PAIN AND DETECTING TACTILE STIMULI

19

1  
2 Our results are generally not supportive of Fitzgibbon et al.'s (2010, 2012b) model, in which  
3  
4 hypervigilance for pain, central sensitization, and the presence of chronic pain were suggested as precursors  
5  
6 of vicarious somatosensory experiences. In addition, controls reported even more vicarious pain experiences  
7  
8 during daily life compared with FM patients. A trend ( $p=.052$ ) suggested that, the more hypervigilant for  
9  
10 pain FM patients were, the less vicarious somatosensory experiences they reported during the experimental  
11  
12 paradigm in contrast to the control group in which more hypervigilance for pain was associated with more  
13  
14 vicarious somatosensory errors. This is in line with a study in which hypervigilance for pain was associated  
15  
16 with less vicarious somatosensory experiences in the pain responder group than in a non-pain responder  
17  
18 group (Vandenbroucke et al., 2013). Hypervigilance for pain may lead to a focus on the body involving a  
19  
20 higher sensitivity for somatosensory stimuli resulting in a better discrimination between false vicarious  
21  
22 experiences and actual bodily experiences. Further research is needed in order to understand the role of  
23  
24 hypervigilance in the elicitation of vicarious experiences in healthy controls and chronic pain patients. The  
25  
26 results are also not in line with those of Brown et al. (2010), who suggested that there might be an  
27  
28 interrelation between illusory tactile perceptions and the degree of pseudoneurological symptoms, nor with  
29  
30 Katzer et al. (2011) who suggested medically unexplained symptoms might be related to touch illusions,  
31  
32 because both groups in the present study reported a comparable number of vicarious somatosensory  
33  
34 experiences. Some previous studies have demonstrated that patients with FM have a hypersensitivity for  
35  
36 mechanical, cold and heat pain perception (Kosek et al., 1996; Smith et al., 2008) and mixed results exist for  
37  
38 non-painful sensations such as cold, warm and touch (Desmeules et al., 2003; Klauenberg et al., 2008). The  
39  
40 results of the present study show that threshold intensities for vibrotactile stimuli, although individually  
41  
42 determined, were not significantly different for both groups. In general, the results show that although FM  
43  
44 patients experience a lot of pain and medically unexplained symptoms, they are as good as controls at  
45  
46 detecting subtle vibrotactile stimuli on their hands despite seeing relevant pain-related scenes.  
47  
48  
49  
50  
51  
52

53 Some limitations of the present study deserve further consideration. First, vibrotactile stimuli were  
54  
55 administered instead of painful stimuli as in our previous study (Vandenbroucke et al., 2013). A study by  
56  
57 Osborn and Derbyshire (2010), found that most patients selected 'tingling' as a descriptor to describe the  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

20

1  
2 somatosensory vicarious experiences while observing pain. Therefore, we used near-threshold intensity  
3  
4 stimuli instead of painful stimuli in order to enhance the occurrence of vicarious somatosensory experiences,  
5  
6 which were consequently not labeled as vicarious pain. Further research could therefore include painful  
7  
8 stimuli to test whether the number of vicarious somatosensory experiences would remain the same. Second,  
9  
10 we included video clips showing hands being pricked. These videos depict less intense pain compared to the  
11  
12 images and movies used in the study by Osborn and Derbyshire (2010). Vicarious experiences may be  
13  
14 elicited more easily when very intense pain is observed. That said, participants in the present study reported  
15  
16 more false alarms during the observation of a subtle injury (the needle prick) as compared with control  
17  
18 videos, indicating that vicarious experiences can also be observed with low intensity pain-related stimuli.  
19  
20 Third, participants may have been more aroused when viewing the pain videos as compared to when  
21  
22 viewing the control videos. As pain captures attention and may induce threat, it may have been more  
23  
24 arousing than the control videos (an inherent feature of pain-related stimuli). Our aims were to investigate  
25  
26 pain videos and control videos, regardless of their arousal capacity. Fourth, in the non-painful videos, human  
27  
28 features were still present (e.g. hand(s)). It would be interesting to test whether the discrepancy in detection  
29  
30 accuracy while observing both videos would increase if all human features were to be removed during non-  
31  
32 painful videos, as tactile perception may be facilitated by simply viewing the body (Kennett, Taylor-Clarke,  
33  
34 & Haggard, 2001). Another limitation of the present study may be that both groups have different  
35  
36 educational levels (82% of the controls had a higher education compared with 26% in the FM group). It is  
37  
38 well known that socio-economic position is negatively associated with pain and general health (Lacey,  
39  
40 Belcher, & Croft, 2012). Further research could match groups regarding socio-economic demographics.  
41  
42  
43  
44  
45

46 In general, this study shows that FM patients and controls are equally accurate in detecting subtle  
47  
48 somatosensory stimuli while observing another in pain. The results further indicate that chronic pain may  
49  
50 not act as a vulnerability factor for the presence of vicarious experiences as suggested by Fitzgibbon et al.  
51  
52 (2010, 2012b). The lower number of neglect errors in FM patients suggest that they stay focused upon  
53  
54 bodily processes even when observing another's pain, and more so than control participants. More research  
55  
56  
57  
58  
59  
60

**OBSERVING PAIN AND DETECTING TACTILE STIMULI**

is needed to explain this discrepancy between controls and FM patients (e.g. accounting for attentional or disinhibition mechanisms).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

OBSERVING PAIN AND DETECTING TACTILE STIMULI

**Acknowledgments**

This research was supported by the Special Research Fund (BOF), Ghent University, grant number 01D3831.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## OBSERVING PAIN AND DETECTING TACTILE STIMULI

23

## References

- 1  
2  
3  
4  
5  
6  
7 Banissy, M.J., Cohen Kadosh, R., Maus, G.W., Walsh, V., Ward, J., 2009. Prevalence, characteristics and  
8 a neurocognitive model of mirror-touch synaesthesia. *Experimental Brain Research*, 198, 261-272.  
9  
10  
11 Banissy, M.J., & Ward, J. (2007). Mirror-touch synesthesia is linked with empathy. *Nature*  
12 *Neuroscience*, 10, 815-816.  
13  
14  
15 Brown, R.J., Brunt, N., Poliakoff, E., & Lloyd, D.M. (2010). Illusory touch and tactile perception in  
16 somatoform dissociators. *Journal of Psychosomatic Research*, 69, 241-248.  
17  
18  
19 Crombez, G., Van Ryckeghem, D.M.L., Eccleston, C., & Van Damme, S. (2013). Attentional bias to pain-  
20 related information: a meta-analysis. *Pain*, 154, 497-510.  
21  
22  
23  
24 Davis, M.H. (1983). Measuring individual differences in empathy: Evidence for a multidimensional  
25 approach. *Journal of Personality and Social Psychology*, 44, 113-126.  
26  
27  
28 De Corte, K., Buysse, A., Verhofstadt, L.L., Roeyers, H., Ponnet, K., & Davis, M. (2007). Measuring  
29 empathic tendencies: reliability and validity of the Dutch version of the interpersonal reactivity index.  
30 *Psychologica Belgica*, 47, 235-260.  
31  
32  
33  
34 Desmeules, J.A., Cedraschi, C., Rapiti, E., Baumgartner, E., Finckh, A., Cohen, P., Dayer, P., & Vischer,  
35 T.L. (2003). Neurophysiologic evidence for a central sensitization in patients with fibromyalgia.  
36 *Arthritis and Rheumatism*, 48, 1420-1429.  
37  
38  
39  
40 Fitzgibbon, B.M., Enticott, P.G., Bradshaw, J.L., Giummarra, M.J., Chou, M., Georgiou-Karistianis, N., &  
41 Fitzgerald, P.B. (2012a). Enhanced corticospinal response to observed pain in pain synaesthetes.  
42 *Cognitive Affective Behavioral Neuroscience*, 12, 406-418.  
43  
44  
45  
46 Fitzgibbon, B.M., Enticott, P.G., Rich, A.N., Giummarra, M.J., Georgiou-Karistianis, N., & Bradshaw,  
47 J.L. (2012b). Mirror-sensory synaesthesia: Exploring 'shared' sensory experiences as synaesthesia.  
48 *Neuroscience and Biobehavioral Reviews*, 36, 645-657.  
49  
50  
51  
52 Fitzgibbon, B.M., Giummarra, M.J., Georgiou-Karistianis, N., Enticott, P.G., & Bradshaw, J.L. (2010).  
53 Shared pain: from empathy to synaesthesia. *Neuroscience & Biobehavioral Reviews*, 34, 500-512.  
54  
55  
56 Gardner, W., Mulvey, E., & Shaw, E. (1995). Regression analyses of counts and rates: Poisson,  
57 overdispersed Poisson, and negative binomial models. *Psychological Bulletin*, 118, 392-404.  
58  
59  
60

## OBSERVING PAIN AND DETECTING TACTILE STIMULI

24

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Glass, J.M., Williams, D.A., Fernandez-Sanchez, M.L., Kairys, A., Barjola, P., Heitzeg, M.M., Clauw, D.J., & Schmidt-Wilcke, T. (2011). Executive functioning in chronic pain patients and healthy controls: different cortical activation during response inhibition in fibromyalgia. *Journal of Pain, 12*, 1219-1229.
- Jackson, P.L, Brunet, E., Meltzoff, A.N., &Decety, J. (2006). Empathy examined through the neural mechanisms involved in imaging how I feel versus how you feel pain. *Neuropsychologia, 44*, 752-761.
- Karazsia, B.T., & van Dulmen, M.H.M. (2010). Modeling infrequent outcomes: Illustrations using prospective predictors of pediatric injuries. In H. Schuster & W. Metzger (Eds.), *Biometrics: Methods, applications and analyses* (pp. 1-27). New York, NY: Nova Science.
- Katzer, A., Oberfeld, D., Hiller, W., &Witthöft, M. (2011). Tactile perceptual processes and their relationship to medically unexplained symptoms and health anxiety. *Journal of PsychosomaticResearch, 71*, 335–341.
- Kennett, S., Taylor-clarke, M. & Haggard, P. (2001). Noninformative vision improves the spatial resolution of touch in humans. *Current Biology, 11*, 118-1191.
- Klauenberg, S., Maier, C., Assion, H.J., Hoffman, A., Krumova, E.K., Magerl, W., Scherens, A., Treede, R.D., &Juckel, G. (2008). Depression and changed pain perception; hints for a central disinhibition mechanism. *Pain, 140*, 332-343.
- Kosek, E., Ekholm, J., & Hansson, P. (1996). Sensory dysfunction in fibromyalgia patients with implications for pathogenic mechanisms. *Pain, 68*, 375-383.
- Lacey, R.J., Belcher, J., & Croft, P.R. (2012). Does life course socio-economic position influence chronic disabling pain in older adults? A general population study. *European Journal of Public Health, 23*, 534-540.
- Lloyd, D.M., Mason, L., Brown, R.J., &Poliakoff, E. (2008). Development of a paradigm for measuring somatic disturbance in clinical populations with medically unexplained symptoms. *Journal ofPsychosomatic Research, 64*, 21-24.
- Loeys, T., Moerkerke, B., De Smet, O., Buysse, A. (2012). The analysis of zero-inflated count data: beyond zero-inflated Poisson regression. *Britisch Journal of Mathematical and Statistical Psychology, 65*, 163-180
- McCracken, L.M. (1997). Attention to pain in persons with chronic pain: a behavioral approach. *BehaviorTherapy, 28*, 271-284.



## OBSERVING PAIN AND DETECTING TACTILE STIMULI

25

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- McKenzie, K.J., Poliakoff, E., Brown, R.J., & Lloyd, M. (2010). Now you feel it, now you don't: How robust is the phenomenon of illusory tactile experience? *Perception*, *39*, 839-850.
- McWilliams, L.A., & Asmundson, G.J.G. (2001). Assessing individual differences in attention to pain: psychometric properties of the pain vigilance and awareness questionnaire modified for a non-clinical pain sample. *Personality and Individual Differences*, *31*, 239-246.
- Osborn, J., & Derbyshire, S.W.G. (2010). Pain sensation evoked by observing injury in others. *Pain*, *148*, 268-274.
- Press, C., Taylor-Clarke, M., Kennett, S., & Haggard, P. (2004). Visual enhancement of touch in spatial body representation. *Experimental Brain Research*, *154*, 238-245.
- Roelofs, J., Peters M.L., McCracken, L., & Vlaeyen, J.W.S. (2003). The pain vigilance and awareness questionnaire (PVAQ): further psychometric evaluation in fibromyalgia and other chronic pain syndromes. *Pain*, *101*, 299-306.
- Roelofs, J., Peters, M.L., Muris, P., & Vlaeyen, J.W.S. (2002). Dutch version of the pain vigilance and awareness questionnaire: validity and reliability in a pain-free population. *Behavioral Research and Therapy*, *40*, 1081-1090.
- Smith, W.S., Tooley, E.M., Montague, E.Q., Robinson, A.E., Cospers, C.J., & Mullins, P.G. (2008). Habituation and sensitization to heat and cold pain in women with fibromyalgia and healthy controls. *Pain*, *140*, 420-428.
- Spence, C. (2013). Just how important is spatial coincidence to multisensory integration? Evaluating the spatial rule. *Annals of the New York Academy of Sciences*, *1296*, 31-49.
- Spence, C., & Driver, J. (Eds.) (2004). Crossmodal space and crossmodal attention. Oxford: Oxford University Press.
- Spinhoven, Ph., Ormel, J., Sloekers, P.P.A., Kempen, G.J.M., Speckens, A.E.M & Van Hemert, A.M. (1997). A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. *Psychological Medicine*, *27*, 363-370
- Staud, R., Bovee, C.E., Robinson, M.E., & Price, D.D. (2009). Cutaneous C-fiber pain abnormalities of fibromyalgia patients are specifically related to temporal summation. *Pain*, *139*, 315-323.

## OBSERVING PAIN AND DETECTING TACTILE STIMULI

26

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Staud, R., Craggs, J.B., Perlstein, W.M., Robinson, M.E., & Price, D.D. (2008). Brain activity associated with slow temporal summation of C-fiber evoked pain in fibromyalgia patients and healthy controls. *European Journal of Pain*, 12, 1078-1089.
- Vandenbroucke, S., Crombez, G., Van Ryckeghem, D.M.L, Brass, M, Van Damme, S., &Goubert, L. (2013). Vicarious pain while observing another in pain: an experimental approach. *Frontiers in HumanNeuroscience*, 7, 265.
- Vierck, C.J., Cannon, R.L., Fry, G., Maixner, W., &Whitsel, B.L. (1997). Characteristics of temporal summation of second pain sensations elicited by brief contact of glabrous skin by a preheated thermode. *Journal of Neurophysiology*, 78, 992-1002.
- Vives, J., Losilla, J.M., & Rodrigo, M.F. (2006). Count data in psychological applied research. *Psychological Reports*, 98, 821-835.
- Wolfe, F., Clauw, D.J., Fitzcharles, M., Goldenberg, D.L., Katz, R.S., Mease, P., Russell, A.S., Russell, I.J., Winfield, J.B., &Yunus, M.B. (2010). The American college of rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Research*, 62, 600-610.
- Zigmond, A.S., &Snaith, R.P. (1983). The hospital anxiety and depression scale. *Acta PsychiatricaScandinavica*, 67, 361-370.

OBSERVING PAIN AND DETECTING TACTILE STIMULI

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Tables

Table 1. Detection accuracy for both groups and all video types.

Table 2. Pearson/Spearman correlations of all measures.

Table 3. Mean scores and standard deviations of all measures.

OBSERVING PAIN AND DETECTING TACTILE STIMULI

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figures

Figure 1. Example of a possible trial.

Figure 2. The relationship between congruency and video.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

	INCONGRUENT				CONGRUENT				NO TACTILE STIMULATION				BOT
Reported site	Correct site	Opposite site (=visual site)	Both hands	No hands	Correct site	Opposite site to visual and tactile	Both hands	No hands	Site congruent tovisual	Opposite site to visual	Both hands	Correct No hands	Visua site
Visual pain/ Control group	41.45%	1.43% <i>Vicarious error</i>	1.32% <i>Vicarious error</i>	55.81%	53.62%	.33%	.55%	45.50%	2.52% <i>Vicarious error</i>	.66%	.22%	96.60%	22.5% <i>Neg error</i>
Visual pain/ FM group	40.02%	1.21% <i>Vicarious error</i>	2.08% <i>Vicarious error</i>	56.69%	49.12%	.44%	2.30%	48.14%	1.32% <i>Vicarious error</i>	.99%	.44%	97.26%	15.7% <i>Neg error</i>
Visual control/ Control group	36.62%	.33%	.66%	62.39%	38.60%	.11%	.44%	60.86%	.88%	.55%	0%	98.58%	12.2% <i>Neg error</i>
Visual control/ FM group	33.33%	.22%	.11%	66.34%	35.42%	.33%	1.21%	63.05%	.66%	.66%	.11%	98.58%	11.1% <i>Neg error</i>

# Attention, Perception, & Psychophysics

Table 2. Pearson/Spearman correlations of all measures.

---

	2.	3.	4.	5.	6.	7.	8.	9.
1. Vicarioussomatosensoryerrors	.32**	-.03	-.08	.08	.10	-.00	-.02	.09
2. Neglect errors (pain-related videos)	-	-.03	-.05	.16	-.01	-.16	-.29*	-.01
3. Hypervigilance (PVAQ)		-	-.02	-.30**	-.04	.06	.40**	-.01
4. Empathic concern			-	.22	.16	-.06	.07	-.01
5. Fantasy scale				-	.14	-.43**	-.33**	.17
6. Personal distress					-	.00	.21	.12
7. Temporal summation, (intensity last-first stimuli)						-	.04	-.11
8. Hospital and Anxiety Scale (HADS)							-	-.09
9. Vicarious pain experiences during daily life								-

---


Note. Pain Vigilance and Awareness Questionnaire (PVAQ). \* p<0.05; \*\*p<0.01

Table 3. Mean scores and standard deviations of all measures.

	M (SD)	M(SD)	M(SD)
	FM	comparison	total
	group	group	group
1. Vicarioussomatosensoryerrors	1.11 (1.67)	1.26 (2.30)	1.18 (1.99)
2. Neglect errors (pain-related videos)	3.79 (2.73)	5.42 (2.85)	4.61 (2.89)
3. Hypervigilance (PVAQ)	41.98 (10.07)	30.12 (14.04)	36.12 (13.50)
4. Empathic concern	20.43 (4.27)	19.95 (4.54)	20.19 (4.38)
5. Fantasy scale	12.46 (5.96)	17.35 (6.32)	14.87 (6.58)
6. Personal distress	11.89 (5.86)	11.32 4.34	11.61 (5.14)
7. Temporalsummation, (intensity last-first stimuli)	7.57 (11.18)	6.46 (6.40)	6.04 (9.22)
8. HospitalAnxietyand DepressionScale (HADS)	18.70 (7.13)	6.01 (5.03)	12.44 (8.86)
9. Vicariouspainexperiences duringdaily life	3.21 (4.86)	6.14 (8.33)	4.63 (6.89)

Note. Pain Vigilance and Awareness Questionnaire (PVAQ).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



215x111mm (96 x 96 DPI)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

