

Impact of pile-driving of offshore monopile foundations on young sea bass

Debusschere Elisabeth^{1,5}, Bert De Coensel², Dominique Adriaens³, Aline Bajek⁴, Dick Botteldooren², Kris Hostens¹, Jan Vanaverbeke⁵, Sofie Vandendriessche¹, Karl Van Ginderdeuren¹, Magda Vincx⁵ and Steven Degraer^{6,5}

¹ Institute for Agricultural and Fisheries Research, Animal Sciences, bio-environmental research group, Ankerstraat 1, B-8400 Oostende, Belgium
E-mail: Elisabeth.Debusschere@ilvo.vlaanderen.be

² Ghent University, Department of Information Technology, Research Group Acoustics, Sint-Pietersnieuwstraat 41, B-9000 Ghent, Belgium

³ Ghent University, Biology Department, Evolutionary Morphology of Vertebrates, K.L. Ledeganckstraat 35, B-9000 Ghent, Belgium

⁴ Marine de Gravelines, Voie des enrochements, 59820 Gravelines, France

⁵ Marine Biology Section, Biology Department, Ghent University, Krijgslaan 281 S8, B-9000 Ghent, Belgium

⁶ Operational Directorate Natural Environment, Marine Ecology and Management Section, Royal Belgian Institute of Natural Sciences, Gulledele 100, B-1200 Brussel, Belgium and Marine Biology Section, Biology Department, Ghent University, Krijgslaan 281 S8, 9000 Ghent, Belgium

The construction of offshore wind farms generates impulsive sounds which are potentially harmful for marine fish. To fill the gaps in the assessment of the lethal and stress sound level thresholds for fish, we carried out a 'worst-case scenario' field experiment on board of a piling vessel. European sea bass of 68 and 115 days old were exposed to pile-driving sound at 45m from the monopile driven in the seabed. Fish were exposed to strikes with a sound exposure level between 181 and 188dB re $1\mu\text{Pa}^2\cdot\text{s}$, and an acoustic particle velocity level of 95dB re $1(\text{nm}\cdot\text{s}^{-1})^2\cdot\text{s}$. The number of strikes ranged from 1739 to 3067, resulting in a cumulative sound exposure level ranging from 215 to 222dB re $1\mu\text{Pa}^2\cdot\text{s}$, and a cumulative acoustic particle velocity level of 130dB re $1(\text{nm}\cdot\text{s}^{-1})^2\cdot\text{s}$. No difference in immediate or delayed mortality up to 14 days after exposure between control and exposed fish groups was found. Fish exposed to pile-driving had a depressed respiration compared to fish in the control groups. This so-called hypo-activity displayed by the exposed fish is an indicator for stress. The lack of mortality in our field experiments confirms the results of lab experiments carried out by other researchers. Although, the wide diversity in physiology in fish species complicates generalizations, the results indicate that pile-driving of windmill monopile foundations creates sound pressure levels that are below the lethal sound threshold for juvenile fish, but above the stress sound threshold, at least for juvenile sea bass.