

## Slippage of DNA polymerase I during synthesis of ds-cDNA

R.Devos, J.Tavernier and W.Fiers

BIOGENT, J. Plateaustraat 22, 9000 Gent and Laboratorium voor Moleculaire Biologie, Rijksuniversiteit Gent, Ledeganckstraat 35, 9000 Gent, Belgium  
Submitted December 23, 1987

Synthesis of ds-cDNA copies of mRNA usually involves oligo-dT primed first strand synthesis followed by the RNase H-DNA polymerase I mediated second strand synthesis (1, 2). The mechanism whereby under these conditions the RNA strand is replaced with DNA probably involves nicking - followed by displacement of the RNA strand. Few intermediate length second strand products are generated starting from homogeneous cDNA.RNA hybrid and no ligase is required to synthesize full length ds-cDNA. This means that within a short time, only a small RNA segment complementary to the 5' region of the cDNA is left over by the RNase H attack to function as a primer for the DNA polymerase I. Using unlabelled globin cDNA.RNA hybrid for synthesis of ds-cDNA some product is also generated using the 5' cDNA-hairpin loop as a primer. Such molecules migrate in an alkaline agarose gel with a size twice that of the original cDNA (Figure). In addition to the full length second strand globin DNA, we also observed products between 100<sub>3</sub> and 300 nucleotides in length in the reactions containing [ $\alpha$ -<sup>32</sup>P]dATP (or [ $\alpha$ -<sup>32</sup>P] dTTP, not shown) as a labelled triphosphate (lane 2, Figure). These products are absent in the reactions containing [ $\alpha$ -<sup>32</sup>P] dCTP (or [ $\alpha$ -<sup>32</sup>P] dGTP, not shown) as the labelled triphosphate (lane 3 and 4, Figure). This means that the DNA polymerase can initiate a slippage poly-dA.poly-dT synthesis starting on the oligo-dT, or on the poly-dT generated by slippage of the reverse transcriptase during the first strand synthesis (3, 4).




Figure : Analysis of second strand globin cDNA products synthesized according to Gubler and Hoffman (2) in the presence of different [ $\alpha$ -<sup>32</sup>P] dNTPs. Samples were analyzed on a 2% alkaline agarose gel. Markers were 5' <sup>32</sup>P-labeled HaeIII fragments of  $\phi$ X174 DNA. Lane 1 : markers; Lane 2 : 2nd strand synthesized in the presence of [ $\alpha$ -<sup>32</sup>P] dATP; Lane 3 : [ $\alpha$ -<sup>32</sup>P] dCTP; Lane 4 : [ $\alpha$ -<sup>32</sup>P] dCTP, DNA ligase and  $\beta$ -NAD.

## REFERENCES :

- (1) Okayama, H. and Berg, P. (1982) Mol. Cell. Biol. 2 : 161-170.
- (2) Gubler, U. and Hoffman, J. (1983) Gene 25 : 263-269.
- (3) Falvey, A.K. et al. (1976) Nucleic Acid Res. 3 : 79-88.
- (4) Devos R. et al. (1977) Eur. J. Biochem. 79 : 419-432.